

VIOREL VÂJĂITU

The analyticity of q -concave sets of locally finite Hausdorff $(2n - 2q)$ measure

Annales de l'institut Fourier, tome 50, n° 4 (2000), p. 1191-1203

http://www.numdam.org/item?id=AIF_2000__50_4_1191_0

© Annales de l'institut Fourier, 2000, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

THE ANALYTICITY OF q -CONCAVE SETS OF LOCALLY FINITE HAUSDORFF $(2n-2q)$ -MEASURE

by Viorel VÂJĂITU

1. Introduction.

Let A be a closed subset of a complex space X . The question of finding reasonable assumptions on A which guarantee its analyticity has been studied over the years by various authors.

Hartogs [14] considered a continuous function $f : D \rightarrow \mathbb{C}$, where $D \subset \mathbb{C}^n$ is open, and showed that the graph G_f of f in $D \times \mathbb{C}$ is pseudoconcave (*i.e.*, the complement of G_f in $D \times \mathbb{C}$ is locally Stein) if and only if f is holomorphic, that is G_f is analytic.

Grauert revealed in his thesis [13] a new interesting aspect of the above question bringing into play thin complements of complete Kähler domains. This topic was afterwards thoroughly studied by Diederich and Fornæss ([6], [7]) and Ohsawa [19].

On the other hand, Hirschowitz [15] settled the case when X is non-singular and A is pseudoconcave of locally finite Hausdorff $(2n-2)$ -measure, where n is the complex dimension of X .

In this article, using q -convexity with corners we introduce the notion of q -concavity. (See §2 for definition. Note that for $q = 1$ we recover the usual pseudoconcavity as used in [15] and [18].) For instance, if X is a complex manifold of pure dimension n and $A \subset X$ is an analytic subset

Keywords: q -convexity – q -concavity – Hausdorff measure – Analytic set.
Math. classification: 32F10 – 32C25.

such that every irreducible of it has dimension $\geq n-q$, then A is q -concave [20]. Two more examples are given at the end of Section 2.

Our main result in this note, which establishes a converse of the above result due to M. Peternell and generalizes Hirschowitz's theorem already quoted above, is the following:

THEOREM 1. — *Let X be a complex space of pure dimension n and q a positive integer less than n . If $A \subset X$ is a q -concave subset such that its Hausdorff $(2n-2q)$ -measure is locally finite, then A is analytic of pure dimension $n-q$.*

As an application (see also Example 2 in Section 2) we have:

COROLLARY 1. — *Let T be a closed positive current of bidimension (q, q) on a complex manifold M . If the Hausdorff $2q$ -measure of $\text{Supp}(T)$ is locally finite, then $\text{Supp}(T)$ is an analytic subset of M of pure dimension q .*

On the other hand, using [16], Theorem 1 yields the following removability theorem. (For $q = 1$ we recover the main result in [2].)

THEOREM 2. — *Let M be a complex manifold of pure dimension n , q a positive integer less than n , $E \subset M$ a closed subset of locally finite Hausdorff $(2n-2q)$ -measure, and f a meromorphic mapping from $M \setminus E$ into a complex space Y . If E does not contain any $(n-q)$ -dimensional analytic subset of M and Y possesses the meromorphic extension property in bidimension $(q, n-q)$ (e.g., if Y is q -complete), then f is continued to a meromorphic mapping from M into Y .*

The organization of this paper is as follows. After a preliminary section, we give in §3 the proofs of Theorems 1 and 2. The last section, §4, establishes connections with the q -pseudoconcavity notion introduced by M. Peternell [20].

2. Preliminaries.

(●) Let T be a metric space and S a subset of T . For $p > 0$ and $\varepsilon > 0$ let $h_\varepsilon^p(S)$ denote the infimum of all (infinite) sums of the form $\sum \delta(S_n)^p$ where $S = \cup S_n$ is an arbitrary decomposition of S with $\delta(S_n) < \varepsilon$ for all n ($\delta = \text{diameter}$). For $p > 0$ the Hausdorff p -measure h^p is defined by

$h^p(S) = \sup_{\varepsilon > 0} h_\varepsilon^p(S) \leq +\infty$. We define $h^0(S)$ to be equal to the cardinality of S . The usual notion of k -dimensional volume in a Riemannian manifold agrees with h^k up to a constant factor depending only on n (for positive integers k). Thus, if A is a pure k -dimensional analytic set in a domain in \mathbb{C}^n , then $h^{2k}(A)$ is equal to a universal constant (depending on k) times the Riemannian volume of the set of regular points of A . For a detailed discussion on Hausdorff measure, see [11].

(•) The definition of q -convexity is the same as in [1], namely; a function $\varphi \in C^\infty(D, \mathbb{R})$, where $D \subset \mathbb{C}^n$ is an open subset, said to be q -convex if its *Levi form*

$$\mathcal{L}_\varphi(z)(\xi) := \sum_{i,j=1}^n \frac{\partial^2 \varphi}{\partial z_i \partial \bar{z}_j}(z) \xi_i \bar{\xi}_j, \quad \xi \in \mathbb{C}^n,$$

has at least $n-q+1$ positive (> 0) eigenvalues for every $z \in D$. This definition can be carried over to complex spaces by local restriction.

Let X be a complex space. X is said to be q -complete if there exists a q -convex function $\varphi \in C^\infty(X, \mathbb{R})$ which is *exhaustive*, i.e., the sublevel sets $\{x \in X; \varphi(x) < c\}$, $c \in \mathbb{R}$, are relatively compact in X . We choose the normalization such that 1-complete spaces correspond to Stein spaces.

Following [8] and [20] a function $\varphi \in C^0(X, \mathbb{R})$ is said to be q -convex with corners on X if every point of X admits an open neighborhood U on which there are finitely many q -convex functions f_1, \dots, f_k such that $\varphi|_U = \max(f_1, \dots, f_k)$. Denote by $F_q(X)$ the set of all functions q -convex with corners on X .

We say that X is q -complete with corners if there exists an exhaustion function $\varphi \in F_q(X)$.

DEFINITION 1. — Let X be a complex space. A subset A of X is said to be q -concave (in X) if A is closed and every point of A has an open neighborhood Ω such that $\Omega \setminus A$ is q -complete with corners.

From [24] (see also [25]) we deduce immediately:

COROLLARY 2. — Let $\pi : X \rightarrow Y$ be a finite surjective holomorphic map of complex spaces and $A \subset Y$ a closed subset. Then A is q -concave in Y if and only if $\pi^{-1}(A)$ is q -concave in X .

Subsequently we give some facts on q -completeness with corners which allow us to reduce the proof of Theorem 1 to the case when X is a domain in \mathbb{C}^n .

PROPOSITION 1. — *Let Y be an analytic set in a complex space X . If Y is q -complete with corners, then Y has a neighborhood system of open sets which are q -complete with corners.*

Proof. — By ([3], Lemma 3) if $\varphi \in F_q(Y)$ and $\eta \in C^0(Y, \mathbb{R})$, $\eta > 0$, then there exists an open neighborhood V of Y in X and $\psi \in F_q(V)$ such that $|\psi - \varphi| < \eta$ on Y . The method of Colţoiu ([4], Theorem 2) or Demailly ([5], the proof of Theorem 1, p. 287) can easily be adapted to our case. \square

PROPOSITION 2. — *Let X be a complex space and φ, ψ be continuous exhaustion functions on X such that there is an open neighborhood Ω of the set $\{\varphi = \psi\}$ in X with $\varphi \in F_p(\Omega \cup \{\varphi < \psi\})$ and $\psi \in F_q(\Omega \cup \{\psi < \varphi\})$. Then X is $(p + q)$ -complete with corners.*

Proof. — Let $\Lambda := \{\lambda \in C^\infty(\mathbb{R}, \mathbb{R}); \lambda' > 0, \lambda'' \geq 0\}$. For $\lambda \in \Lambda$ define $\Phi_\lambda : X \rightarrow \mathbb{R}$ by

$$\Phi_\lambda := 1 / (\exp(-\lambda(\varphi)) + \exp(-\lambda(\psi))).$$

It is straightforward to see that Φ_λ is exhaustive for X and it is $(p + q)$ -convex with corners on Ω . Now we let $\varepsilon > 0$ be continuous on X such that $\{|\varphi - \psi| \leq \varepsilon\} \subset \Omega$; define $W_- = \{\varphi - \psi \leq -\varepsilon\}$ and $W_+ = \{\varphi - \psi \geq \varepsilon\}$. Clearly W_-, W_+ are closed subsets of X and $W_- \cup W_+ \cup \Omega = X$. The proof is concluded if we show the next

CLAIM. — *There is $\lambda \in \Lambda$ such that Φ_λ is p -convex with corners on W_- and q -convex with corners on W_+ .*

But this follows by adjusting the arguments in [22]. We omit the details. \square

PROPOSITION 3. — *Let U, V be open subsets of a complex space X such that U is p -complete with corners and V is q -complete with corners. Then $U \cup V$ is $(p + q)$ -complete with corners.*

Proof. — Consider exhaustion functions $f \in F_p(U)$ and $g \in F_q(V)$ for U and V respectively. Let $a \in C^\infty(U, \mathbb{R})$ with $0 \leq a \leq 1$, $a(x) = 1$ if $x \in U \setminus V$ or $x \in U \cap V$ and $f(x) \leq g(x) + 1$; $a(x) = 0$ if $x \in U \cap V$ and $f(x) > g(x) + 2$. Set $D := U \cup V$. Define φ on D by setting

$$\varphi = \begin{cases} f & \text{on } U \setminus V, \\ af + (1 - a)(1 + g) & \text{on } U \cap V, \\ 1 + g & \text{on } V \setminus U. \end{cases}$$

Then φ is continuous and exhaustive for D .

Let $b \in C^\infty(V, \mathbb{R})$ with $0 \leq b \leq 1$, $b(x) = 1$ if $g(x) \leq \varphi(x) + 1$ and $b(x) = 0$ if $g(x) > \varphi(x) + 2$. Define ψ on D by setting

$$\psi = \begin{cases} bg + (1 - b)(1 + \varphi) & \text{on } V, \\ 1 + \varphi & \text{on } U \setminus V. \end{cases}$$

Then ψ is continuous and exhaustive for D .

Finally, it is easy to see that $S := \{\psi < 1 + \varphi\} \subset V$ and $\psi = g$ on S ; hence $\psi \in F_q(S)$. Similarly, $T := \{\varphi < 1 + \psi\} \subset U$ and $\varphi = f$ on T ; so $\varphi \in F_p(T)$. The conclusion then follows from Proposition 2. \square

COROLLARY 3. — *Let A and B be p -concave and q -concave sets in the complex spaces X and Y respectively. Then $A \times B$ is $(p + q)$ -concave in $X \times Y$.*

Proof. — Since the assertion is local, we may assume that X and Y are Stein spaces, $X \setminus A$ is p -complete with corners, and $Y \setminus B$ is q -complete with corners. Then $X \times Y \setminus A \times B = X \times (Y \setminus B) \cup (X \setminus A) \times Y$ is $(p + q)$ -complete with corners by Proposition 3. \square

For a complex space X we introduce [20] the set $G_q(X)$ as follows: For $x_o \in X$ let $G_q(x_o)$ be the set of all functions $g : X \rightarrow \mathbb{R}$ such that there are: an open neighborhood U of x_o (which may depend on g) and $f \in F_q(U)$ with $f(x_o) = g(x_o)$ and $f \leq g|_U$. Then put

$$G_q(X) := C^0(X, \mathbb{R}) \cap \bigcap_{x \in X} G_q(x).$$

Clearly $F_q(X) \subseteq G_q(X) \subset C^0(X, \mathbb{R})$.

Note that given an open set $D \Subset X$, an $\varepsilon > 0$, and a function $g \in G_q(X)$, there is a function $h \in F_q(D)$ such that $|h - g| < \varepsilon$ on D . See [20], Lemma 1. But we cannot use this fact and the classical perturbation procedure (see for instance [8]) to get a globally defined h since we do not know that given $v \in G_q(X)$ and $\theta \in C^\infty(X, \mathbb{R})$ there is $\varepsilon_o > 0$ such that $v + \lambda\theta \in G_q(X)$ for every $\lambda \in \mathbb{R}$, $|\lambda| < \varepsilon_o$. However we can avoid this difficulty since we show:

LEMMA 1. — *The set $F_q(X)$ is dense in $G_q(X)$ in the sense that given an arbitrary $g \in G_q(X)$ and $\eta \in C^0(X, \mathbb{R})$, $\eta > 0$, there is $f \in F_q(X)$ such that $|f - g| < \eta$.*

Proof. — We do this in three steps.

Step 1). Fix $x \in X$ and $\varepsilon > 0$. By definition there is an open neighborhood Ω of x and $\varphi \in F_q(\Omega)$ with $\varphi(x) = g(x)$ and $\varphi \leq g$ on Ω . Let W, U be open neighborhoods of x , $W \subseteq U \subseteq \Omega$, such that $\varphi \geq g - \varepsilon$ on U ; then let $\theta \in C^\infty_0(U, \mathbb{R})$, $\theta = -1$ on ∂W and $\theta(x) = 1$. If $c > 0$ is small enough, then $\psi := \varphi + c\theta \in F_q(U)$, $\psi < g$ on ∂W , $\psi > g$ on a neighborhood V of x in W , and $|\psi - g| < 2\varepsilon$ on U .

Step 2). The above step shows that for all compact subsets K, L of X , L a neighborhood of K and $\varepsilon > 0$, there are: a finite set of indices I (which depends on K and L), open sets $V_i \subseteq W_i \subseteq U_i \subseteq L$ such that $\{V_i\}_{i \in I}$ cover K , functions $f_i \in F_q(U_i)$ with $|f_i - g| < 2\varepsilon$ on W_i , $f_i > g$ on V_i and $f_i < g$ on ∂W_i .

Step 3). Let $\{K_\nu\}_{\nu \in \mathbb{N}}$ be an exhaustion sequence for X by compact sets, $K_0 = \emptyset$ (by convention set $K_{-1} = \emptyset$), and $K_\nu \subset \text{int}(K_{\nu+1})$ for all ν . For each ν apply Step 2 to $K = K_\nu \setminus \text{int}(K_{\nu-1})$, $L = K_{\nu+1} \setminus \text{int}(K_{\nu-2})$, and $\varepsilon = (\min_L \eta)/2$. We therefore obtain open sets $V_{i\nu} \subseteq W_{i\nu} \subseteq U_{i\nu}$ such that the family $\{W_{i\nu}\}$ is locally finite, $\{V_{i\nu}\}_{i\nu}$ is a covering of X , and functions $f_{i\nu} \in F_q(U_{i\nu})$ as in Step 2 from above. Then define $f : X \rightarrow \mathbb{R}$ by $f(x) = \max\{f_{i\nu}(x); x \in W_{i\nu}\}$, where the maximum is taken over all indices i, ν such that $W_{i\nu} \ni x$. It is straightforward to see that f is continuous, $f \in F_q(X)$, and $g < f < g + \eta$. □

Remark. — It can be shown that for $q > \dim(X)$ the set $F_q(X)$ is dense in the above sense even in $C^0(X, \mathbb{R})$.

From ([20], Lemma 4) we quote:

LEMMA 2. — *Let U be a complex space, V a complex manifold of pure dimension r , and $f \in F_{q+r}(U \times V)$ such that $\sup f < \infty$. Consider $g : U \rightarrow \mathbb{R}$ defined by*

$$g(x) = \sup\{f(x, y); y \in V\}, \quad x \in U.$$

Assume that for some $x_0 \in U$ there is $y_0 \in V$ with $g(x_0) = f(x_0, y_0)$. Then $g \in G_q(x_0)$.

The key proposition for the proof of Theorem 1 is:

PROPOSITION 4. — *Let X and Y be complex manifolds such that Y is of pure dimension r and p -complete with corners. Let A be a $(q + r)$ -concave subset in $X \times Y$ such that the natural projection $\pi : A \rightarrow X$ is*

proper. Then $\pi(A)$ is $(q + p - 1)$ -concave in X . In particular, if Y is Stein (i.e. $p = 1$), then $\pi(A)$ is q -concave.

Proof. — Set $m := q + p - 1$. We may assume without any loss in generality that X is Stein. The statement of the proposition follows from the next claim.

CLAIM. — For every relatively compact Stein open subset U of X , the set $U \setminus \pi(A)$ is m -complete with corners.

In order to show this, consider a relatively compact open subset V of Y which is p -complete with corners and such that $\pi^{-1}(\overline{U} \times \pi(A)) \subset \overline{U} \times V$. Then $K := \overline{U} \times \partial V$ is compact and disjoint from A . Now, since $U \times Y \setminus A$ is $(m + r)$ -complete with corners by [20], there exists an exhaustion function $\psi \in F_{m+r}(U \times Y \setminus A)$.

Let $\lambda := \max_K \psi$ and define $\sigma : U \setminus \pi(A) \rightarrow \mathbb{R}$ by setting

$$\sigma(x) = \max\{\psi(x, y), y \in \overline{V}\}, x \in U \setminus \pi(A).$$

Clearly σ is continuous. Consider θ be a 1-convex exhaustion function on U and then define $\varphi : U \setminus \pi(A) \rightarrow \mathbb{R}$ by setting

$$\varphi = \theta + \max(\lambda, \sigma).$$

Then φ is continuous and exhaustive. To conclude the proof, in view of Lemma 1, it suffices to show that $\varphi \in G_m(x)$ for ever $x \in U \setminus \pi(A)$. Indeed, two cases may occur:

a) If $\sigma(x) > \lambda$, then $\sigma \in G_m(x)$ by Lemma 2. Since $\varphi = \sigma + \theta$ on a neighborhood of x , we get $\varphi \in G_m(x)$.

b) If $\sigma(x) \leq \lambda$, then $\theta(x) + \lambda = \varphi(x)$ and since $\lambda + \theta \leq \varphi$ on $U \setminus \pi(A)$, $\varphi \in G_1(x)$, *a fortiori*, $\varphi \in G_m(x)$.

The proof is complete. \square

(\bullet) Denotes by $\Delta^k(t)$ the open polydisc in \mathbb{C}^k of polyradius (t, \dots, t) centered at the origin. Let n and q be positive integers such that $q < n$. We define the $(q, n - q)$ Hartogs figure in $\mathbb{C}^n = \mathbb{C}^q \times \mathbb{C}^{n - q}$ to be the open set $H_q \subset \mathbb{C}^n$ given by

$$H_q := \left((\Delta^q(1) \setminus \overline{\Delta^q(t)}) \times \Delta^{n - q}(1) \right) \cup \left(\Delta^q(1) \times \Delta^{n - q}(s) \right)$$

where $0 < t, s < 1$. Put $\widehat{H}_q := \Delta^n(1)$, i.e. the envelope of holomorphy of H_q .

Following [16] we say that a complex space Y possesses the meromorphic extension property (in bidimension $(q, n-q)$) if every meromorphic map $f : H_q \rightarrow Y$ extends to a meromorphic map $\hat{f} : \hat{H}_q \rightarrow Y$.

By [16] every q -complete complex space possesses a meromorphic extension property in bidimension $(q, n-q)$ for every integer $n > q$.

DEFINITION 2. — M be a complex manifold of pure dimension n . We say that a closed subset $A \subset M$ is pseudoconcave of order q if for every injective holomorphic map $f : \hat{H}_q \rightarrow M$ such that $f(H_q) \cap A$ is empty, the set $f(\hat{H}_q) \cap A$ is also empty.

In this set-up, a variant of Proposition 4 for $Y = \mathbb{C}^r$ is straightforward. See ([10], Lemma 3.6).

Also by ([24], Corollary 5) one has: *A closed subset A of a pure dimensional complex manifold is pseudoconcave of order q if and only if A is q -concave.*

Pseudoconcavity of order q is easier to handle; though it does not suit to complex spaces. One has the next examples:

1) Let M be a Stein manifold of pure dimension n and $K \subset M$ a compact set. Then $\hat{K} \setminus K$ is $(n-1)$ -concave in $X \setminus K$. (See [23].)

2) The support of a closed positive current of bidegree (q, q) on a pure dimensional complex manifold is q -concave. (This follows by [12], Corollary 2.6 and the above remark.)

3. Proof of Theorems 1 and 2.

Proof of Theorem 1.

We remark that it suffices to show that A is analytic and for this we distinguish three steps.

Step 1). — Here we reduce the proof to the case when $X \subset \mathbb{C}^n$ is open. For this we need:

LEMMA 3. — *Let Z be a complex space, $X \subset Z$ an analytic subset, and $A \subset X$ a closed subset (not necessarily analytic). If A is q -concave in X and X is r -concave in Z , then A is $(q+r)$ -concave in Z .*

Proof. — Let $x_o \in A$ and U be a Stein open neighborhood of x_o in Z such that $U \setminus X$ is r -complete with corners and $(U \setminus A) \cap X$ is q -complete with corners. Since $(U \setminus A) \cap X$ is analytic in $U \setminus A$, there is by Proposition 1 an open subset Ω of $U \setminus A$ which is q -complete with corners and contains $(U \setminus A) \cap X$. Therefore $U \setminus A = (U \setminus X) \cup \Omega$ is $(q+r)$ -complete with corners by Proposition 3. \square

To complete Step 1, we let $x \in A$, then take a coordinate patch $\iota : U \rightarrow D \subset \mathbb{C}^N$ around $x \in X$ with D Stein; hence U is isomorphic to the closed analytic subset $\iota(U)$ of D , hence $\iota(A \cap U)$ is q -concave in $\iota(U)$. Put $p := q + N - n$. Note that $N - p = n - q$. Therefore $\iota(A \cap U)$ is p -concave in D by Lemma 3 since $\iota(U)$ is $(N - n)$ -concave in D . On the other hand, $\iota(A \cap U)$ as a closed subset of D has its Hausdorff $(2N - 2p)$ -measure locally finite.

Step 2). — We give here some general facts for further reduction of the proof of Theorem 1.

Let $E \subset \mathbb{C}^n$ be a locally closed set with $h^{2n-2q+1}(E) = 0$ and suppose $0 \in E$. Then there is a complex $(n-q)$ -plane Γ through 0 such that $h^1(E \cap \Gamma) = 0$ ([21], Lemma 2). Hence for a suitable unitary transformation σ of \mathbb{C}^n we have $h^1(\sigma(E) \cap (\mathbb{C}^{n-q} \times \{0\})) = 0$. By ([21], Corollary 2), $\sigma(E) \cap (\partial B(r) \times \{0\})$ is empty for (h^1) -almost all $r > 0$. (Here $B(r)$ denotes the open unit ball in \mathbb{C}^{n-q} of radius r .) Since $\sigma(E)$ is also locally closed in \mathbb{C}^n and $0 \in \sigma(E)$, there is $r > 0$ arbitrary small and a polydisc P in \mathbb{C}^q centered at the origin such that $\sigma(E) \cap (\overline{B(r)} \times \overline{P})$ is closed in $\overline{B(r)} \times \overline{P}$ and $\sigma(E) \cap (\partial B(r) \times \overline{P})$ is empty. In particular, the canonically induced projection map π from $\sigma(E) \cap (B(r) \times P)$ into $B(r)$ is proper.

If furthermore $h^{2n-2q}(E) < \infty$, then $\pi^{-1}(z)$ is finite for (h^{2n-2q}) -almost all $z \in B(r)$ ([21], Corollary 4).

Recall that a set $\Gamma \subset \mathbb{C}^n$ is said to be *locally pluripolar* if for every $a \in \Gamma$ there is a connected neighborhood $U \ni a$ and a plurisubharmonic function φ on U , $\varphi \neq -\infty$, such that $\Gamma \cap U \subset \{\varphi = -\infty\}$. In fact, if Γ is locally pluripolar then by [17] one can take $U = \mathbb{C}^n$, so Γ is pluripolar. Note that for $n = 1$ pluripolarity of a set in \mathbb{C} means that it is of *zero-capacity* as used in [18]. Also it is easy to check that for $U \subset \mathbb{C}^n$ open and $S \subset \mathbb{C}^n$ of zero Lebesgue measure, the set $U \setminus S$ is not pluripolar.

Step 3). — Here we conclude the proof.

By Steps 1, 2, and Proposition 4 it remains to show the next lemma.

LEMMA 4. — *Let $U \subset \mathbb{C}^{n-q}$ be an open set, Δ the open unit disc in \mathbb{C} , and $A \subset U \times \Delta^q$ a closed subset such that the canonical projection $\pi : A \rightarrow U$ is proper. If A is q -concave and $\pi^{-1}(z)$ is finite for z in a non pluripolar subset of U , then A is analytic of pure dimension $n-q$.*

Proof. — For $q = 1$ this is precisely the lemma due to Hartogs-Oka-Nishino [18]. For $q > 1$ we proceed as follows. Notice that it suffices to show the analyticity of A . In order to do this we let $p_j : \Delta^q \rightarrow \Delta$, $j = 1, \dots, q$, denote the projection onto the j^{th} component of Δ^q , then let $\sigma_j : A \rightarrow U \times \Delta$ naturally induced by p_j . Then σ_j is proper and Proposition 4 implies that $\sigma_j(A)$ is 1-concave in $U \times \Delta$ for all indices $j = 1, \dots, q$. Furthermore if we consider $\pi_j : \sigma_j(A) \rightarrow U$ canonically induced, we arrive at the case $q = 1$. So the sets $\sigma_j(A)$ are analytic for all j .

Now, if $\iota : U \times \Delta^q \rightarrow (U \times \Delta) \times \dots \times (U \times \Delta)$ (the product is taken q -times) is given by $\iota(z, t_1, \dots, t_q) = ((z, t_1), \dots, (z, t_q))$, then $A = \iota^{-1}(\sigma_1(A) \times \dots \times \sigma_q(A))$, whence the lemma. Thus the proof of Theorem 1. \square

Proof of Theorem 2.

Denote by $A^0 :=$ the set of points $x \in A$ such that f extends meromorphically onto a neighborhood of x . Then $A' := A \setminus A^0$ is closed and as the complement to A is locally connected in M these local meromorphic continuations of f in points of A^0 glue together to a unique meromorphic map from $M \setminus A'$ into Y .

Now, we assert that A' is pseudoconcave of order q . For this we let $\Phi : \widehat{H}_q \rightarrow M$ be an injective holomorphic map with $\Phi(H_q) \cap A' = \emptyset$. Then $f \circ \Phi$ is meromorphic from H_q into Y , hence it extends to \widehat{H}_q ; therefore f extends over $\Phi(\widehat{H}_q)$, and by definition $\Phi(\widehat{H}_q) \subset A^0$; whence the desired assertion.

Finally, by Theorem 1, if A' is not the empty set, then A' is analytic of pure dimension $n-q$. But this contradicts the hypothesis, whence the proof. \square

4. A final remark.

Motivated by M. Peternell's work ([20], §7) we give:

DEFINITION 3. — *Let X be a complex space of pure dimension n . A closed subset A of X is said to be q -pseudoconcave if there is an analytic subset $B \subset X$ such that*

$$1) \overline{A \setminus B} = A.$$

2) *For each point $x \in A \setminus B$ there is a locally closed analytic subset Y of X which passes through x , $Y \subset A$, and Y is a complex manifold of dimension $n - q$.*

As an example, if A is analytic and $\dim_x A \geq n - q$, $\forall x \in A$, then A is q -pseudoconcave.

Let now r be a non-negative integer and suppose X is purely dimensional. We say that X has property (E_r) , if there is $\varphi \in F_{n+r}(X \times X \setminus \Delta_X)$, where Δ_X is the diagonal set of $X \times X$, such that $\varphi(x_\nu, x) \rightarrow +\infty$ if $x_\nu \rightarrow x$, $x_\nu \neq x$, $\forall x \in X$. Condition (E_r) holds locally on X if every point of X admits an open neighborhood U which satisfies (E_r) .

The next proposition is an easy consequence of ([20], Lemma 9).

PROPOSITION 5. — *Let X be a pure dimensional complex space such that (E_r) holds locally. Then every q -pseudoconcave subset of X is $(q + r)$ -concave.*

The importance of the condition (E_r) resides in the fact that, for example, if a Stein space X fulfils (E_0) , then every locally Stein open subset of X is Stein. It is easy to check for a Stein manifold that (E_0) holds. However, this fails, in general, if we allow singularities. For example, we let X be the Segre cone in \mathbb{C}^4 , $X = \{xy = zw\}$. Clearly the hypersurface $A = \{x = z = 0\}$ is 1-pseudoconcave. Now, if (E_0) would hold locally on X , then A will be 1-concave; and as X has isolated singularities $X \setminus A$ will be Stein. But this is absurd since $X \setminus A$ is biholomorphic to $(\mathbb{C}^2 \setminus \{0\}) \times \mathbb{C}$.

COROLLARY 4. — *If X is a complex manifold, then every q -pseudoconcave subset of X is also q -concave.*

Example 3. — For every positive integer q there is an open subset X of \mathbb{C}^{q+1} and a q -concave subset $A \subset X$ which is **not** q -pseudoconcave.

To do this we consider a compact subset K of \mathbb{C}^2 such that $\widehat{K} \setminus K$ contains no analytic disc. See [26] for the existence of K . Put $X := (\mathbb{C}^2 \setminus K) \times \mathbb{C}^{q-1}$ and $A := (\widehat{K} \setminus K) \times \{0\}$. Then A is **not** q -pseudoconcave in X ; however, by Example 1 in §2 and Corollary 3 it is easily seen that $\widehat{K} \setminus K$ is q -concave in X . \square

The corresponding version of Theorem 1 reads:

THEOREM 3. — *Let A be a closed subset of a pure n -dimensional complex space X such that A is q -pseudoconcave and its Hausdorff $(2n-2q)$ -measure is locally finite. Then A is analytic of pure dimension $n-q$.*

Proof. — If $\iota : U \rightarrow D$ is a local path of X , where D is an open subset of \mathbb{C}^N , then $\iota(A \cap U)$ is $(N-n+q)$ -pseudoconcave in D . Now we conclude by the above corollary and Theorem 1. \square

Acknowledgements. — A part of this work has been supported by an ANSTI grant No. 5232/1999.

BIBLIOGRAPHY

- [1] A. ANDREOTTI, H. GRAUERT, Théorèmes de finitude pour la cohomologie des espaces complexes, *Bull. Soc. Math. France*, 90 (1962), 193–259.
- [2] E.M. CHIRKA, On the removable singularities for meromorphic mappings, *Ann. Polon. Math.*, 70 (1998), 43–47.
- [3] M. COLTOIU, n -concavity of n -dimensional complex spaces, *Math. Z.*, 210 (1992), 203–206.
- [4] M. COLTOIU, Complete locally pluripolar sets, *J. reine angew. Math.*, 412 (1990), 108–112.
- [5] J.-P. DEMAILLY, Cohomology of q -convex spaces in top degrees, *Math. Z.*, 204 (1990), 283–295.
- [6] K. DIEDERICH, J.-E. FORNÆSS, Thin complements of complete Kähler domains, *Math. Ann.*, 259 (1982), 331–341.
- [7] K. DIEDERICH, J.-E. FORNÆSS, On the nature of thin complements of complete Kähler domains, *Math. Ann.*, 268 (1984), 475–495.
- [8] K. DIEDERICH, J.-E. FORNÆSS, Smoothing q -convex functions and vanishing theorems, *Invent. Math.*, 82 (1985), 291–305.
- [9] K. DIEDERICH, J.-E. FORNÆSS, Smoothing q -convex functions in the singular case, *Math. Ann.*, 273 (1986), 665–671.
- [10] G. DLOUSSKY, Analyticité séparée et prolongement analytique, *Math. Ann.*, 286 (1990), 153–170.

- [11] H. FEDERER, Geometric measure theory, Berlin-Heidelberg-New York, Springer, 1969.
- [12] J.-E. FORNÆSS, N. SIBONY, Oka's inequality for currents and applications, *Math. Ann.*, 301 (1995), 399–419.
- [13] H. GRAUERT, Charakterisierung der Holomorphiegebiete durch die vollständige Kählersche Metrik, *Math. Ann.*, 131 (1965), 38–75.
- [14] F. HARTOGS, Über die aus der singulären Stellen einer analytischen Funktion mehrerer Veränderlichen bestehende Gebilde, *Acta Math.*, 32 (1909), 57–79.
- [15] A. HIRSCHOWITZ, Entre les hypersurfaces et les ensembles pseudoconcaves, *Ann. Scuola Norm. Sup. Pisa*, 27 (1973), 873–887.
- [16] S. IVASHKOVICH, A. SILVA, The Hartogs type extension theorem for meromorphic mappings into q -complete complex spaces, *Boll. U.M.I.*, (8) 2-B (1999), 251–261.
- [17] B. JOSEFSON, On the equivalence between locally polar and globally polar in \mathbb{C}^n , *Arkiv för Mat.*, 16 (1978), 109–115.
- [18] T. NISHINO, Sur les ensembles pseudoconcaves, *J. Math. Kyoto Univ.*, 1 (1961/62), 225–245.
- [19] T. OHSAWA, Analyticity of complements of complete Kähler domains, *Proc. Japan Acad.*, 56, Ser. A, (1980), 484–487.
- [20] M. PETERNELL, Continuous q -convex exhaustion functions, *Invent. Math.*, 85 (1986), 249–262.
- [21] B. SHIFFMAN, On the removal of singularities for analytic sets, *The Mich. Math. J.*, 15-16 (1968/69), 111–120.
- [22] V. VĂJĂITU, q -completeness and q -concavity of the union of open subspaces, *Math. Z.*, 221 (1996), 217–229.
- [23] V. VĂJĂITU, On \mathcal{P} -complete morphisms of complex spaces, *Geometric Complex Analysis*, Proc. the third International Research Institute, Math. Soc. Japan, Hayama 1995; Eds. J. Noguchi, H. Fujimoto, J. Kajiwara, and T. Ohsawa, pag. 653–665.
- [24] V. VĂJĂITU, Invariance of q -completeness with corners under finite holomorphic surjective maps, *Bull. Belg. Math. Soc.*, 5 (1998), 713–718.
- [25] V. VĂJĂITU, A Levi problem for continuous strongly q -plurisubharmonic functions, *C. R. Acad. Sci. Paris*, 328 (1999), 573–578.
- [26] J. WERMER, Polynomially convex hull and analyticity, *Arkiv för Matem.*, 20 (1982), 129–135.

Manuscrit reçu le 16 juin 1999,
accepté le 17 janvier 2000.

Viorel VĂJĂITU,
Institute of Mathematics of the Romanian Academy
P.O. Box 1–764
RO–70700 Bucharest (Romania).
vvajaitu@stoilow.imar.ro