

ANNALES DE L'INSTITUT FOURIER

LEO SARIO

Completeness and existence of bounded biharmonic functions on a riemannian manifold

Annales de l'institut Fourier, tome 24, n° 1 (1974), p. 311-317

<http://www.numdam.org/item?id=AIF_1974__24_1_311_0>

© Annales de l'institut Fourier, 1974, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

COMPLETENESS AND EXISTENCE OF BOUNDED BIHARMONIC FUNCTIONS ON A RIEMANNIAN MANIFOLD

by Leo SARIO (¹)

A.S. Galbraith has communicated to us the following intriguing problem : Does the completeness of a manifold imply, or is it implied by, the emptiness of the class H^2B of bounded nonharmonic biharmonic functions ? Among all manifolds considered thus far in biharmonic classification theory (cf. Bibliography), those that are complete fail to carry H^2B -functions, and one might suspect that this is always the case. We shall show, however, that there do exist complete manifolds of any dimension that carry H^2B -functions. Moreover, there exist both complete and incomplete manifolds not permitting these functions, and, trivially, incomplete manifolds possessing them.

We attach a Bibliography of recent work in the field.

1. Let C be the totality of complete Riemannian manifolds M , characterized by an infinite distance of any point of M to the ideal boundary. Denote by $\Theta_{H^2B}^N$ and $\widetilde{\Theta}_{H^2B}^N$ the classes of N -manifolds, $N \geq 2$, for which $H^2B = \emptyset$ or $H^2B \neq \emptyset$, respectively.

THEOREM 1. — $C \cap \widetilde{\Theta}_{H^2B}^N \neq \emptyset$ for every N .

Proof. — Take the N -cylinder

$$|x| < \infty, \quad |y| \leq 1, \quad i = 1, 2, \dots, N - 1,$$

with each face $y_i = 1$ identified with $y_i = -1$, so as to obtain a covering space of the N -torus in the same manner as a conventional cylinder is a covering surface of the torus. Let T be this N -cylinder with the Riemannian metric

(¹) The work was sponsored by the U.S. Army Research Office-Durham, Grant DA-ARO-D-31-124-71-G181, University of California, Los Angeles.

$$ds^2 = \mu^{-2}(x) dx^2 + \mu^{4/(N-1)}(x) \sum_{i=1}^{N-1} dy_i^2$$

where

$$\mu(x) = (2 + x^2)^{\frac{1}{2}} \log(2 + x^2).$$

To see that $T \in C$, it suffices to show, in view of the symmetry, that $\int_0^\infty \mu^{-1}(x) dx = \infty$. The verification is immediate :

$$\begin{aligned} \int_0^\infty (2 + x^2)^{-\frac{1}{2}} \log^{-1}(2 + x^2) dx &> \frac{1}{2} \int_0^\infty (2 + x)^{-1} \log^{-1}(2 + x) dx \\ &= \frac{1}{2} \int_0^\infty \log \log(2 + x) = \infty. \end{aligned}$$

We introduce the function

$$u(x) = \int_0^x \mu^{-3}(t) \int_0^t \mu(s) \int_0^s \mu^{-3}(r) dr ds dt.$$

The Laplace-Beltrami operator $\Delta = d\delta + \delta d$ gives

$$\Delta u = -g^{-\frac{1}{2}} (g^{\frac{1}{2}} g^{xx} u')' = -\mu^{-1} (\mu \mu^2 u')' = -\int_0^x \mu^{-3}(r) dr$$

and

$$\Delta^2 u = -\mu^{-1} (\mu \mu^2 (-\mu^{-3}))' = 0.$$

Thus u is nonharmonic biharmonic.

To see that u is bounded it suffices to show that it is so for $x > 0$. For all $s > 0$,

$$\int_0^s \mu^{-3}(r) dr = \int_0^s (2 + r^2)^{-3/2} \log^{-3}(2 + r^2) dr = o(1),$$

and for all $t > 0$,

$$\begin{aligned} \int_0^t \mu(s) \int_0^s \mu^{-3}(r) dr ds &< c \int_0^t (2 + s^2)^{\frac{1}{2}} \log(2 + s^2) ds \\ &< 2c \int_0^t (2 + s) \log(2 + s) ds \\ &= c \left[(2 + t)^2 \log(2 + t) \right. \\ &\quad \left. - \frac{1}{2} (2 + t)^2 + \text{const.} \right]. \end{aligned}$$

Here and later c is a constant, not always the same. We let [] stand for the expression in brackets and obtain

$$u(x) < c \int_0^x (2 + t^2)^{-3/2} \log^{-3}(2 + t^2) [] dt.$$

The dominating term in the integrand is majorized by

$$\frac{1}{2} t^{-3} \log^{-3} t \times (2 + t)^2 \log(2 + t).$$

The integral from 1 to $x > 1$ is bounded, and consequently so is u for all x .

This completes the proof of Theorem 1.

2. The following simple example, valid for $N \geq 3$, is perhaps also of interest. Let

$$T : |x| < \infty, \quad |y| \leq \pi, \quad |z_i| \leq 1, \quad i = 1, \dots, N - 2,$$

with the metric

$$ds^2 = dx^2 + e^{-x} dy^2 + e^{(2e^x-x)/(N-2)} \sum_{i=1}^{N-2} dz_i^2,$$

the opposite faces again identified by pairs. Clearly $T \in C$.

The function

$$u = \cos y$$

belongs to $H^2 B$. In fact,

$$\begin{aligned} \Delta u &= -e^{-x+x} (e^{e^x-x} e^x) (-\cos y) \\ &= e^x \cos y, \end{aligned}$$

and

$$\Delta^2 u = -e^{-x+x} [(e^{e^x-x} e^x)' \cos y + e^{e^x-x} e^x e^x (-\cos y)] = 0.$$

Thus $T \in C \cap \widetilde{\Theta}_{H^2 B}$.

3. The reason that we are only interested in nonharmonic biharmonic functions is, of course, that completeness is known not

to exclude bounded harmonic functions (Nakai-Sario [6]). For $N \geq 3$, we insert here a simple proof of this fact.

Take the N -cylinder

$$T: |x| < \infty, \quad |y| \leq 1, \quad |z_i| \leq 1, \quad i = 1, \dots, N - 2,$$

with the metric

$$ds^2 = dx^2 + e^{2x^2} dy^2 + \sum_{i=1}^{N-2} dz_i^2.$$

Trivially $T \in C$. The function

$$h(x) = \int_0^x e^{-t^2} dt$$

is harmonic,

$$\Delta h = -e^{-x^2} (e^{x^2} e^{-x^2})' = 0.$$

It also is bounded and, in fact, even Dirichlet finite :

$$D(h) = c \int_{-\infty}^{\infty} e^{-2x^2} e^{x^2} dx < \infty.$$

4. We return to nonharmonic biharmonic functions.

THEOREM 2. — $C \cap \Theta_{H^2 B}^N \neq \emptyset$ for every N .

Proof. — The Euclidean N -space $E^N \in C$. Every biharmonic function u has an expansion in spherical harmonics S_{nm}

$$u = \sum_{n=0}^{\infty} \sum_{m=1}^{m_n} (a_{nm} r^{n+2} + b_{nm} r^n) S_{nm}.$$

If $u \in H^2 B$, then

$$\int_{|x|=r} u S_{nm} d\omega = c(a_{nm} r^{n+2} + b_{nm} r^n)$$

is bounded in r , hence $a_{nm} = b_{nm} = 0$ for all n , except for b_{01} . Therefore u is constant.

5. In view of $u = r^2 \in H^2 B$ on the Euclidean N -ball, we have trivially $\tilde{C} \cap \tilde{\Theta}_{H^2 B}^N \neq \emptyset$ for every N , with \tilde{C} the totality of incomplete Riemannian manifolds. It remains to show :

THEOREM 3. — $C \cap \Theta_{H^2B}^N \neq \emptyset$ for every N .

Proof. — Let E_α^N be the N -space $0 < r < \infty$ with the metric

$$ds = r^\alpha |dx|,$$

α a constant. It is known (Sario-Wang [19, 21]) that if $N \geq 4$, $E_\alpha^N \in \Theta_{H^2B}$ for every α ; $E_\alpha^2 \in \Theta_{H^2B}$ if and only if $\alpha \neq -1 \mp n/2$, $n = 1, 2, \dots$; $E_\alpha^3 \in \Theta_{H^2B}$ if and only if $\alpha \neq -1 \mp \left[\frac{1}{2} n(n+1)\right]^{\frac{1}{2}}$.

On the other hand, $E_\alpha^N \in \widetilde{C}$ for every α , hence the theorem.

The author is sincerely indebted to Professor Cecilia Wang for a painstaking checking of the manuscript.

BIBLIOGRAPHY

- [1] D. Hada, L. Sario, C. Wang, Dirichlet finite biharmonic functions on the Poincaré N -ball, *J. Reine Angew. Math.* (to appear).
- [2] D. Hada, L. Sario, C. Wang, N -manifolds carrying bounded but no Dirichlet finite harmonic functions, *Nagoya Math. J.* (to appear).
- [3] Y.K. Kwon, L. Sario, B. Walsh, Behavior of biharmonic functions on Wiener's and Royden's compactifications, *Ann. Inst. Fourier* (Grenoble) 21 (1971), 217-226.
- [4] N. Mirsky, L. Sario, C. Wang, Bounded polyharmonic functions and the dimension of the manifold, *J. Math. Kyoto Univ.*, 13 (1973), 529-535.
- [5] M. Nakai, Dirichlet finite biharmonic functions on the plane with distorted metrics, (to appear).
- [6] M. Nakai, L. Sario, Completeness and function-theoretic degeneracy of Riemannian spaces, *Proc. Nat. Acad. Sci.*, 57 (1967), 29-31.
- [7] M. Nakai, L. Sario, Biharmonic classification of Riemannian manifolds, *Bull. Amer. Math. Soc.*, 77 (1971), 432-436.
- [8] M. Nakai, L. Sario, Quasiharmonic classification of Riemannian manifolds, *Proc. Amer. Math. Soc.*, 31 (1972), 165-169.

- [9] M. Nakai, L. Sario, Dirichlet finite biharmonic functions with Dirichlet finite Laplacians, *Math. Z.*, 122 (1971), 203-216.
- [10] M. Nakai, L. Sario, A property of biharmonic functions with Dirichlet finite Laplacians, *Math. Scand.*, 29 (1971), 307-316.
- [11] M. Nakai, L. Sario, Existence of Dirichlet finite biharmonic functions, *Ann. Acad. Sci. Fenn. A.I.*, 532 (1973), 1-34.
- [12] M. Nakai, L. Sario, Existence of bounded biharmonic functions, *J. Reine Angew. Math.*, 259 (1973), 147-156.
- [13] M. Nakai, L. Sario, Existence of bounded Dirichlet finite biharmonic functions, *Ann. Acad. Sci. Fenn. A.I.*, 505 (1972), 1-12.
- [14] M. Nakai, L. Sario, Biharmonic functions on Riemannian manifolds, *Continuum Mechanics and Related Problems of Analysis*, Nauka, Moscow, 1972, 329-335.
- [15] L. Sario, Biharmonic and quasiharmonic functions on Riemannian manifolds, *Duplicated lecture notes* 1968-70, University of California, Los Angeles.
- [16] L. Sario, M. Nakai, Classification Theory of Riemann Surfaces, Springer-Verlag, 1970, 446 pp.
- [17] L. Sario, C. Wang, The class of (p, q) -biharmonic functions, *Pacific J. Math.*, 41 (1972), 799-808.
- [18] L. Sario, C. Wang, Counterexamples in the biharmonic classification of Riemannian 2-manifolds, *Pacific J. Math.* (to appear).
- [19] L. Sario, C. Wang, Generators of the space of bounded biharmonic functions, *Math. Z.*, 127 (1972), 273-280.
- [20] L. Sario, C. Wang, Quasiharmonic functions on the Poincaré N-ball, *Rend. Mat.* (to appear).
- [21] L. Sario, C. Wang, Riemannian manifolds of dimension $N \geq 4$ without bounded biharmonic functions, *J. London Math. Soc.* (to appear).
- [22] L. Sario, C. Wang, Existence of Dirichlet finite biharmonic functions on the Poincaré 3-ball, *Pacific J. Math.*, 48 (1973), 267-274.
- [23] L. Sario, C. Wang, Negative quasiharmonic functions, *Tôhoku Math. J.*, 26 (1974), 85-93.
- [24] L. Sario, C. Wang, Radial quasiharmonic functions, *Pacific J. Math.*, 46 (1973), 515-522.

COMPLETENESS AND EXISTENCE OF BOUNDED BIHARMONIC FUNCTIONS

- [25] L. Sario, C. Wang, Parabolicity and existence of bounded biharmonic functions, *Comm. Math. Helv.* 47, (1972), 341-347.
- [26] L. Sario, C. Wang, Positive harmonic functions and biharmonic degeneracy, *Bull. Amer. Math. Soc.*, 79 (1973), 182-187.
- [27] L. Sario, C. Wang, Parabolicity and existence of Dirichlet finite biharmonic functions, *J. London Math. Soc.* (to appear).
- [28] L. Sario, C. Wang, Harmonic and biharmonic degeneracy, *Kodai Math. Sem. Rep.*, 25 (1973), 392-396.
- [29] L. Sario, C. Wang, M. Range, Biharmonic projection and decomposition, *Ann. Acad. Sci. Fenn. A.I.*, 494 (1971), 1-14.
- [30] C. Wang, L. Sario, Polyharmonic classification of Riemannian manifolds, *Kyoto Math. J.*, 12 (1972), 129-140.

Manuscrit reçu le 22 mai 1973
accepté par M. Brelot.

Leo SARIO
Department of Mathematics
University of California
Los Angeles, Calif. (USA)