



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Mathématique

Jeffrey D. Carlson

The K-theory of the conjugation action

Volume 359, issue 7 (2021), p. 795-796

Published online: 2 September 2021

<https://doi.org/10.5802/crmath.235>



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Les Comptes Rendus. Mathématique sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org
e-ISSN : 1778-3569



Geometry and Topology / *Géométrie et Topologie*

The K-theory of the conjugation action

Jeffrey D. Carlson^a

^a Department of Mathematics, Imperial College London, 180 Queen's Gate, London SW7 2AZ, UK

E-mail: j.carlson@imperial.ac.uk

Abstract. In 1999, Brylinski and Zhang computed the complex equivariant K-theory of the conjugation self-action of a compact, connected Lie group with torsion-free fundamental group. In this note we show it is possible to do so in under a page.

Manuscript received 7th April 2021, revised and accepted 9th June 2021.

Brylinski and Zhang [2] showed that if G is a compact, connected Lie group with torsion-free fundamental group, then the equivariant K-theory of its conjugation action G^{Ad} is isomorphic to the ring $\Omega_{RG/\mathbb{Z}}^*$ of Grothendieck differentials on the complex representation ring RG of G . Their proof uses results on holomorphic differentials on complex manifolds, a reduction to the case G is a torus, and some algebraic geometry. We show a more concrete and arguably more natural expression for the ring $K_G^*(G^{\text{Ad}})$ can be obtained rapidly using only Hodgkin's Künneth spectral sequence [5], in the same manner they already use it, and elementary algebraic considerations. We then show this purely algebraic isomorphism admits a satisfying geometric interpretation, and remark finally that this geometric version gives back Brylinski and Zhang's description in terms of Grothendieck differentials at no added cost.

Theorem (Brylinski–Zhang [2, Thm. 3.2]). *Let G be a compact, connected Lie group with torsion-free fundamental group. Then $K_G^*(G^{\text{Ad}})$ is isomorphic to $RG \otimes K^*G$ as an RG -algebra. Under this identification, the forgetful map $f: K_G^*(G^{\text{Ad}}) \rightarrow K^*G$ becomes reduction with respect to the augmentation ideal IG of RG .*

Proof. Write G^{bi} for G under the $(G \times G)$ -action $(h, k) \cdot g = h g k^{-1}$. The orbit space of $G^{\text{bi}} \times G^{\text{bi}}$ under the restricted, free diagonal action of $1 \times G$ is $(G \times 1)$ -equivariantly diffeomorphic to G^{Ad} via $(g', g) \mapsto g' g^{-1}$, so when $X = Y = G^{\text{bi}}$, Hodgkin's $(\mathbb{Z} \times \mathbb{Z}/2)$ -graded Künneth spectral sequence $\text{Tor}_{R(G \times G)}(K_{G \times G}^* X, K_{G \times G}^* Y) \implies K_{G \times G}^*(X \times Y)$ reduces to $\text{Tor}_{RG \otimes RG}(RG, RG) \implies K_G^*(G^{\text{Ad}})$. Here the two structure maps $RG \otimes RG \rightarrow RG$ are both the multiplication of RG . Recall [5, Prop. 11.1] that under our hypotheses, RG is the tensor product of a polynomial ring on generators $y_i \in IG$ and a Laurent polynomial ring on generators $t_j \in 1 + IG$. Let P be the free abelian group on generators q_i and w_j and let $\gamma: P \rightarrow IG$ be the linear map taking q_i to y_i and w_j to $t_j - 1$. Then an $(RG)^{\otimes 2}$ -module resolution of RG is given by $RG \otimes \Lambda P \otimes RG$, with differential the derivation vanishing on $RG \otimes \mathbb{Z} \otimes RG$ and sending $1 \otimes z \otimes 1$, for $z \in P$, to $1 \otimes 1 \otimes \gamma(z) - \gamma(z) \otimes 1 \otimes 1$. To compute the Tor, apply $-\otimes_{RG \otimes RG} RG$ to this resolution to obtain the CDGA $RG \otimes \Lambda P$ with differential, RG in bidegree $(0, 0)$, and P in bidegree $(-1, 0)$.

The spectral sequence collapses because the differentials d_r for $r \geq 2$ send all generators into the right half-plane. Since $\pi_1 G$ is torsion-free and $X = G$ is locally contractible of finite covering dimension, the spectral sequence strongly converges to the intended target. Hence $RG \otimes \Lambda P$ is the graded algebra associated to a filtration $(F_p)_{p \leq 0}$ of $K_G^*(G^{\text{Ad}})$ with $F_0 \cong RG$ and $F_{-1}/F_0 \cong RG \otimes P$. Since RG and ΛP are both free abelian, there is no additive extension problem, so $K_G^*(G^{\text{Ad}})$ is also free abelian as a group. Let \tilde{z}_k be elements in F_{-1} lifting $1 \otimes q_i$ and $1 \otimes w_j$ under the isomorphism $F_{-1}/F_0 \cong RG \otimes P$. Then the \tilde{z}_k anticommute with each other because they lie in $K_G^1(G^{\text{Ad}})$ and square to 0 since $K_G^*(G^{\text{Ad}})$ contains no 2-torsion, and by induction, they generate $K_G^*(G^{\text{Ad}})$ as an RG -algebra, so $K_G^*(G^{\text{Ad}}) = RG \otimes \Lambda[\tilde{z}_k]$.

To see the forgetful map f is as claimed, note that forgetting the $(G \times 1)$ -action on G^{bi} induces a map to the spectral sequence $\text{Tor}_{RG}(\mathbb{Z}, \mathbb{Z}) \Rightarrow K^*G$, which again collapses by lacunary considerations. Computing $\text{Tor}_{RG}(\mathbb{Z}, \mathbb{Z}) \cong \Lambda P$ with the resolution $\Lambda P \otimes RG$ of \mathbb{Z} shows the map $E_2(f): RG \otimes \Lambda P \rightarrow \Lambda P$ is reduction modulo IG and $\Lambda P \cong K^*G$. \square

Remark. We can be completely explicit about the exterior generators. As observed by Hodgkin [4, Thm. A], the injection $U(n) \hookrightarrow U := \varinjlim U(n)$ induces an additive map $\beta: RG \rightarrow K^1G$ descending to a group isomorphism between the module $IG/(IG)^2$ of indecomposables of RG and the module PK^*G of primitives of the exterior Hopf algebra $K^*G \cong \Lambda PK^*G$. In particular, a set of generators is given by $\beta(\lambda_i) = \beta(\lambda_i - \dim \lambda_i)$ for λ_i lifts in G of the fundamental representations of the commutator subgroup G' and $\beta(t_j) = \beta(t_j - 1)$ for $t_j: G \rightarrow G/G' \rightarrow U(1)$ circular coordinate functions of the torus $G^{\text{ab}} = G/G' \cong (S^1)^{\text{rk}G - \text{rk}G'}$. Let $Q = \{\lambda_i, t_j\}_{i,j}$.

The map β in fact factors as $f \circ \beta^{\text{Ad}}$ for a map $\beta^{\text{Ad}}: RG \rightarrow K_G^*(G^{\text{Ad}})$, already giving surjectivity of f since PK^*G generates $K^*G = \Lambda PK^*G$ as a ring. Atiyah [1, Lem. 2, pf.] described β^{Ad} and hence β geometrically: given a representation $\rho: G \rightarrow U(n)$, we can build a representative E of $\beta(\rho)$ via the clutching construction, taking with two trivial bundles $CG \times \mathbb{C}^n$ over the cone CG on G and gluing them along $G \times \mathbb{C}^n$ via the relation $(g, v) \sim (g, \rho(g)v)$ to obtain a bundle over the suspension $CG \cup_G CG$. The action $h \cdot (g, v) = (hgh^{-1}, \rho(h)v)$ of G on $G^{\text{Ad}} \times \mathbb{C}^n$ preserves this relation and so induces a G -action on E making it a G -equivariant bundle over the suspension of G^{Ad} .

The RG -module structure on $K_G^*(X)$ is always given by $\sigma \cdot [E] = [\alpha(\sigma) \otimes E]$, where if $\sigma: G \rightarrow \text{Aut} V$ is a representation, then $\alpha(\sigma)$ is the trivial bundle $X \times V$ equipped with the diagonal G -action. Thus $\sigma \otimes \prod_k \beta(\rho_k) \mapsto \alpha(\sigma) \cdot \prod_k \beta^{\text{Ad}}(\rho_k)$ for $\rho_k \in Q$ gives an explicit isomorphism $RG \otimes K^*G \xrightarrow{\sim} K_G^*(G^{\text{Ad}})$.

Remark. The first paragraph of our proof is a variant of Brylinski–Zhang’s §4 [2]. Once the ring structure is determined as in our proof’s second paragraph, replacing §§5–6, one knows their map $\phi: \Omega_{RG/\mathbb{Z}}^* \rightarrow K_G^*(G^{\text{Ad}})$ from the ring of Grothendieck differentials is an isomorphism as soon as one knows it is a well-defined RG -algebra map [2, Prop 3.1], for the class $\beta^{\text{Ad}}(\rho) \in K_G^1(G^{\text{Ad}})$ from the previous remark is in fact the same as¹ Brylinski–Zhang’s $\phi(d\rho)$, so that $f \circ \phi$ takes a basis $\{d\rho: \rho \in Q\}$ of the free RG -module $\Omega_{RG/\mathbb{Z}}^1$ to a \mathbb{Z} -basis $\{\beta(\rho): \rho \in Q\}$ of PK^*G .

References

- [1] M. F. Atiyah, “On the K -theory of compact Lie groups”, *Topology* **4** (1965), no. 1, p. 95-99.
- [2] J.-L. Brylinski, B. Zhang, “Equivariant K -theory of compact connected Lie groups”, *K-Theory* **20** (2000), no. 1, p. 23-36.
- [3] C.-K. Fok, “The Real K -theory of compact Lie groups”, *SIGMA, Symmetry Integrability Geom. Methods Appl.* **10** (2014), article no. 022 (26 pages).
- [4] L. Hodgkin, “On the K -theory of Lie groups”, *Topology* **6** (1967), no. 1, p. 1-36.
- [5] ———, “The equivariant Künneth theorem in K -theory”, in *Topics in K-theory*, Lecture Notes in Mathematics, vol. 496, Springer, 1975, p. 1-101.

¹ the corrected version of—see Fok [3, Rmk. 2.8.1]