



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Mathématique

Bertrand Cloez and Pierre Gabriel

On an irreducibility type condition for the ergodicity of nonconservative semigroups

Volume 358, issue 6 (2020), p. 733-742

Published online: 2 October 2020

<https://doi.org/10.5802/crmath.92>



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



Les Comptes Rendus. Mathématique sont membres du
Centre Mersenne pour l'édition scientifique ouverte
www.centre-mersenne.org
e-ISSN : 1778-3569



Partial Differential Equations, Functional Analysis / *Équations aux dérivées partielles, Analyse fonctionnelle*

On an irreducibility type condition for the ergodicity of nonconservative semigroups

Sur une condition de type irréductibilité pour l'ergodicité des semi-groupes non conservatifs

Bertrand Cloez^a and Pierre Gabriel^b

^a MISTEA, INRAE, Montpellier SupAgro, Univ. Montpellier, 2 place Pierre Viala, 34060 Montpellier, France

^b Laboratoire de Mathématiques de Versailles, UVSQ, CNRS, Université Paris-Saclay, 45 Avenue des États-Unis, 78035 Versailles cedex, France
E-mails: bertrand.cloez@inrae.fr, pierre.gabriel@uvsq.fr

Abstract. We propose a simple criterion, inspired from the irreducible aperiodic Markov chains, to derive the exponential convergence of general positive semigroups. When not checkable on the whole state space, it can be combined to the use of Lyapunov functions. It differs from the usual generalization of irreducibility and is based on the accessibility of the trajectories of the underlying dynamics. It allows to obtain new existence results of principal eigenelements, and their exponential attractiveness, for a nonlocal selection-mutation population dynamics model defined in a space-time varying environment.

Résumé. Nous proposons une condition simple, inspirée des notions d'irréductibilité et d'apériodicité pour les chaînes de Markov, qui permet d'assurer la convergence exponentielle de semi-groupes positifs généraux. Lorsque celle-ci ne s'applique pas sur tout l'espace, elle peut être localisée via l'utilisation de fonctions de Lyapunov. Elle diffère des généralisations habituelles de l'irréductibilité et est basée sur la notion d'accessibilité des trajectoires sous-jacentes. Finalement, cette condition nous permet d'obtenir de nouveaux résultats d'existence d'éléments propres, et les bornes de convergence exponentielle associées, pour un modèle de sélection-mutation en environnement changeant.

2020 Mathematics Subject Classification. 47A35, 35B40, 47D06, 60J80, 92D15, 92D25.

Funding. B.C. has received the support of the Chair “Modélisation Mathématique et Biodiversité” of VEOILIA-Ecole Polytechnique-MnHn-FX and the ANR project MESA (ANR-18-CE40-006), funded by the French Ministry of Research.

Manuscript received 19th September 2019, revised 2nd July 2020, accepted 27th June 2020.

1. The main result

We are interested in the long time behavior of positive semigroups in *weighted signed measures* spaces. Before dealing with these semigroups, let us start by defining more precisely what we mean by weighted signed measures.

Let \mathcal{X} be a measurable space. We work in weighted signed measures spaces on \mathcal{X} . More precisely for a weight function $V : \mathcal{X} \rightarrow (0, \infty)$ we denote by $\mathcal{M}_+(V)$ the set of positive measures on \mathcal{X} which integrate V and we define the space of weighted signed measures as

$$\mathcal{M}(V) = \mathcal{M}_+(V) - \mathcal{M}_+(V)$$

the smallest vector space with positive cone $\mathcal{M}_+(V)$, see [3] for a rigorous construction as a quotient space. Basically, an element μ of $\mathcal{M}(V)$ is the difference $\mu = \mu_+ - \mu_-$ of two positive measures $\mu_+, \mu_- \in \mathcal{M}_+(V)$ which are mutually singular (Hahn–Jordan decomposition). It acts on the Banach space

$$\mathcal{B}(V) = \left\{ f : \mathcal{X} \rightarrow \mathbb{R} \text{ measurable, } \|f\|_{\mathcal{B}(V)} := \sup_{x \in \mathcal{X}} |f(x)|/V(x) < +\infty \right\}$$

through

$$\mu(f) = \int_{\mathcal{X}} f \, d\mu_+ - \int_{\mathcal{X}} f \, d\mu_-.$$

The space $\mathcal{M}(V)$ is a Banach space for the weighted total variation norm

$$\|\mu\|_{\mathcal{M}(V)} = \sup_{\|f\|_{\mathcal{B}(V)} \leq 1} \mu(f).$$

Notice that, with these definitions, the standard total variation norm is $\|\cdot\|_{\text{TV}} = \|\cdot\|_{\mathcal{M}(\mathbb{1})}$, where $\mathbb{1}$ stands for the mapping $x \mapsto 1$.

We consider positive semigroups $(M_t)_{t \geq 0}$ acting on $\mathcal{M}(V)$ on the left ($\mu \mapsto \mu M_t$) and on $\mathcal{B}(V)$ on the right ($f \mapsto M_t f$), which enjoy the classical duality relation $\mu(M_t f) = (\mu M_t)(f)$. We present sufficient irreducibility type conditions which, possibly combined to Lyapunov type conditions, ensure the so-called *asynchronous exponential behavior* of the semigroup:

$$\mu M_t \sim e^{\lambda t} \mu(h) \gamma \quad \text{as } t \rightarrow \infty,$$

where $\lambda \in \mathbb{R}$, h is a positive function, and γ is a positive measure.

We start with the global case, i.e. conditions verified on the whole state space \mathcal{X} .

Hypothesis 1. *The weight function is given by $V = \mathbb{1}$. There exist $\tau > 0$ and $C \geq 1$ such that $C^{-1} \leq M_s \mathbb{1} \leq C$ for all $s \in [0, \tau]$, and there exist $c > 0$ and a family of probability measures $(\sigma_{x,y})_{x,y \in \mathcal{X}}$ over $[0, \tau]$ such that*

$$\forall x, y \in \mathcal{X}, \quad \delta_x M_\tau(\cdot) \geq c \int_0^\tau \delta_y M_{\tau-s}(\cdot) \sigma_{x,y}(ds), \quad (\text{H1})$$

$$\sup_{x, x' \in \mathcal{X}} \inf_{y \in \mathcal{X}} \|\sigma_{x,y} - \sigma_{x',y}\|_{\text{TV}} < 2. \quad (\text{H2})$$

Let us do some comments on these two assumptions.

Condition (H1) ensures the existence of a time τ such that, with positive probability and uniformly with respect to any initial positions x and y in \mathcal{X} , the trajectories issued from x intersect at time τ the trajectories issued from y at some random times $s \in [0, \tau]$. Even though they are not comparable in general, this *path crossing* condition has connections with the notion of irreducibility of Markov processes, which means basically that for any x and y there is a deterministic time at which the trajectories issued from x reach y with positive probability [16].

Consider for instance the semigroup $(M_t)_{t \geq 0}$ of a time continuous finite Markov chain $(X_t)_{t \geq 0}$:

$$M_t f(x) = \mathbb{E}[f(X_t) \mid X_0 = x].$$

Irreducibility induces the existence of a random time $T(x, y)$ such that $X_{T(x,y)} = y$ and the strong Markov property gives

$$M_t f(x) \geq \mathbb{E}[f(X_t) \mathbb{1}_{T(x,y) \leq t} \mid X_0 = x] = \mathbb{E}[M_{t-T(x,y)} f(y) \mathbb{1}_{T(x,y) \leq t} \mid X_0 = x].$$

Assumption (H1) then holds with $\sigma_{x,y}$ being the law of $T(x, y)$ conditioned on being positive.

For Markov processes on infinite state spaces however, irreducibility does not imply (H1) in general, due to the requirement in (H1) for the probability of intersection and the crossing maximal time τ to be uniform in x and y . Actually there are few examples of Markov processes on infinite state spaces which are irreducible since in most cases the probability of reaching a point y at a deterministic time starting from an arbitrary position x is zero. Similarly, the concept of irreducibility for positive semigroups, see [15, Definition C-III.3.1 p. 306] for instance, which proves very useful when working in spaces of functions, is rarely applicable in spaces of measures, exactly for the same reasons as for Markov processes. Condition (H1) can then be seen as a useful alternative to irreducibility since it allows to cover many more interesting examples.

Consider for instance a Markov process on a compact topological state space \mathcal{X} . If the random hitting time $T(x, y)$ has a positive density $\varepsilon(x, y, \cdot)$ with respect to the Lebesgue measure which is continuous on its three components, then the associated semigroup satisfies (H1) and (H2). A simple and typical example satisfying (H1) but not (H2) is given by the periodic (and irreducible) semigroup $(M_t)_{t \geq 0}$ defined for every $f \in \mathcal{B}(\mathbb{1})$, $t \geq 0$ and $x \in \mathcal{X} = [0, 1]$ by

$$M_t f(x) = f(x + t - \lfloor x + t \rfloor),$$

where $\lfloor \cdot \rfloor$ stands for the integer part. For this semigroup, (H1) is satisfied only with $(\sigma_{x,y})_{0 \leq x, y \leq 1}$ a family of Dirac masses, so that (H2) does not hold. A more sophisticated example (still irreducible) can be found in [5], see also the comments in Section 2.

Condition (H2) is thus an aperiodicity assumption, reminiscent from the aperiodicity of Markov chains. It can also be seen as a coupling condition since it means that, uniformly in x and x' in \mathcal{X} , the trajectories issued from x and x' intersect at time τ with positive probability (the intersection point being on a trajectory issued from some $y \in \mathcal{X}$ in the time interval $[0, \tau]$).

In Hypothesis 1 the conditions are satisfied on the whole state space, which can be too much to ask in many applications, typically when the state space is not compact. In that case Assumptions (H1) and (H2) can be localized through the use of Lyapunov functions, in the spirit of [3, 9]. It leads to the following set of assumptions where, for two functions $f, g : \Omega \rightarrow \mathbb{R}$, the notation $f \lesssim g$ means that $f \leq Cg$ on Ω for some $C > 0$.

Hypothesis 2. *There exists a function $\psi : \mathcal{X} \rightarrow (0, \infty)$, a subset $K \subset \mathcal{X}$, a time $\tau > 0$, and constants $\beta > \alpha > 0$ and $\theta > 0$ such that*

$$\psi \leq V \text{ on } \mathcal{X} \text{ and } V \lesssim \psi \text{ on } K; \quad MV \lesssim V \text{ and } M\psi \gtrsim \psi \text{ on } [0, \tau] \times \mathcal{X}, \tag{A0}$$

$$M_\tau V \leq \alpha V + \theta \mathbb{1}_K \psi, \tag{A1}$$

$$M_\tau \psi \geq \beta \psi, \tag{A2}$$

and there exist $c > 0$ and a family of probability measures $(\sigma_{x,y})_{x,y \in K}$ over $[0, \tau]$ such that

$$\forall f \in \mathcal{B}(V), f \geq 0, \forall x, y \in K, \quad \frac{M_\tau f(x)}{\psi(x)} \geq c \int_0^\tau \frac{M_{\tau-s} f(y)}{\psi(y)} \sigma_{x,y}(ds), \tag{H1'}$$

$$\sup_{x, x' \in K} \inf_{y \in K} \int_0^\tau \frac{M_{\tau-s}(\psi \mathbb{1}_K)(y)}{\psi(y)} (\sigma_{x,y} \wedge \sigma_{x',y})(ds) > 0. \tag{H2'}$$

We are now ready to state our main result.

Theorem 3. *Assume that Hypothesis 1 or Hypothesis 2 is verified. Then there exists a unique triplet $(\gamma, h, \lambda) \in \mathcal{M}_+(V) \times \mathcal{B}_+(V) \times \mathbb{R}$ of eigenelements of $(M_t)_{t \geq 0}$ with $\gamma(h) = \|h\|_{\mathcal{B}(V)} = 1$, i.e. satisfying for all $t \geq 0$*

$$\gamma M_t = e^{\lambda t} \gamma \quad \text{and} \quad M_t h = e^{\lambda t} h.$$

Moreover, there exist $C, \omega > 0$ such that for all $t \geq 0$ and $\mu \in \mathcal{M}(V)$,

$$\|e^{-\lambda t} \mu M_t - \mu(h)\gamma\|_{\mathcal{M}(V)} \leq C \|\mu\|_{\mathcal{M}(V)} e^{-\omega t}. \quad (1)$$

In [3], the authors give a set Assumption A of conditions which they prove to be equivalent to the convergence (1). These conditions are the same as Hypothesis 2 where (H1') and (H2') are replaced by a Doeblin type coupling condition (A3) and an assumption (A4) which is basically a control on the ratio $M_t \psi(x)/M_t \psi(y)$ uniform in $x, y \in K$ and $t \geq 0$. The main advantage of Hypothesis 2 compared to Assumption A in [3] is that (H1') is much easier to check than (A4) on examples. It will be illustrated on an example coming from evolutionary biology in Section 2.

The idea for proving Theorem 3 is to show that (H1') implies a slightly stronger control than (A4) and, combined with (H2'), a slightly weaker version of (A3). Then we verify that the proof in [3] can be adapted to these small variations of (A3)–(A4).

Note that, contrary to Assumption A in [3], Hypothesis 2 is sufficient but not necessary for ensuring the convergence (1), see for instance the comments on the example in Section 2. It is similar to the situation of finite Markov chains for which irreducibility and aperiodicity are a powerful sufficient but not necessary condition for proving convergence to an equilibrium. Perron–Frobenius theory states that it becomes equivalent to the convergence only by adding the assumption that the Markov chain is undecomposable, in the sense that it has only one recurrent class.

Proof of Theorem 3. We will prove that Hypothesis 2 implies the existence of a constant $d > 0$ such that for all x, y in K and all $t \geq 0$

$$\frac{M_t \psi(x)}{\psi(x)} \geq d \frac{M_t \psi(y)}{\psi(y)}, \quad (2)$$

and the existence of a constant $\tilde{c} > 0$ and a family $(\nu_{x,x'})_{x,x' \in K}$ of probability measures such that for all x, x' in K and all $y \in \{x, x'\}$

$$\frac{\delta_y M_t(\psi \cdot)}{M_t \psi(y)} \geq \tilde{c} \nu_{x,x'}, \quad (3)$$

and that Hypothesis 1 implies these same inequalities with K replaced by \mathcal{X} and $\psi = \mathbb{1}$. Such estimates, which first appeared in [8], are a powerful tool to derive the exponential ergodicity of nonconservative semigroups, as attested by the further developments in [2, 3, 9].

When (2) and (3) are verified on \mathcal{X} with $\psi = \mathbb{1}$, and $t \mapsto \|M_t \mathbb{1}\|_\infty$ is locally bounded, we can use the proof of [2, Theorem 3.5] to get (1) with $V = \mathbb{1}$. Indeed (2) readily implies [2, Assumption (H2)] and (3) is similar to [2, Assumption (H1)] with the difference that the coupling measure ν is independent of x, x' . But it is not an issue since the proof in [2] also works with a family $(\nu_{x,x'})$. This was already noticed in [8], where Assumption (A1') is the analogue to (3).

When (2) and (3) are verified in $K \subset \mathcal{X}$ and ψ is as in Hypothesis 2, we can use the proof in [3] to obtain (1). Similarly as above, (2) implies [3, Assumption (A4)] and (3) is the same as [3, Assumption (A4)] except that in (3) the coupling measure depends on x, x' . The proof in [3] can be adapted to this slight variation by: replacing the quantity $\nu(M_{k\tau} \psi / \psi)$ by $\inf_K(M_{k\tau} \psi / \psi)$ in the definition of the Lyapunov functions V_k and using accordingly (2) instead of [3, Assumption (A4)]; letting ν depend on x, x' in the proof of [3, Lemma 3.2] and using (3) instead of [3, Assumption (A3)]

It thus only remains to check that Hypothesis 2 implies (2) and (3), and that Hypothesis 1 implies the same inequalities with $K = \mathcal{X}$ and $\psi = \mathbb{1}$. Since the proof for Hypothesis 1 is simpler, we only give it for Hypothesis 2 (replace below K by \mathcal{X} and ψ by $\mathbb{1}$ for Hypothesis 1).

By virtue of Assumption (A0), the condition (2) is clearly verified for $t \in [0, \tau]$. When $t > \tau$, using $f = M_{t-\tau}\psi$ in (H1') ensures that for all $x, y \in K$

$$\frac{M_t\psi(x)}{\psi(x)} \geq c \int_0^\tau \frac{M_{t-s}\psi(y)}{\psi(y)} \sigma_{x,y}(ds). \tag{4}$$

By Assumption (A0), there exists a constant $C > 0$ such that for all $y \in \mathcal{X}$, $t \geq 0$ and $s \in [0, \tau]$

$$\frac{M_t\psi(y)}{M_{t-s}\psi(y)} \leq C \frac{M_{k\tau}V(y)}{M_{k\tau}\psi(y)}, \tag{5}$$

where $k = \lfloor \frac{t-s}{\tau} \rfloor$. Using Assumptions (A1) and (A2) we can prove as in [3, Lemma 4.1] that for all $y \in \mathcal{X}$

$$\frac{M_{k\tau}V(y)}{M_{k\tau}\psi(y)} \leq \left(\frac{\alpha}{\beta}\right)^k \frac{V(y)}{\psi(y)} + \frac{\theta}{\beta - \alpha}. \tag{6}$$

Since $\beta > \alpha$ and $V \lesssim \psi$ on K by (A0), we deduce from (4), (5) and (6) that Condition (2) is verified for a constant $d > 0$ independent of $x, y \in K$ and $t \geq 0$.

We turn now to (3). Due to (A0), there exists $C > 0$ such that $M_\tau\psi \leq C\psi$ on K . Hypothesis (H1') then yields that for every $x, y \in K$

$$\frac{\delta_x M_\tau(\psi \cdot)}{M_\tau\psi(x)} \geq \frac{c}{C} \int_0^\tau \frac{\delta_y M_{\tau-s}(\psi \cdot)}{\psi(y)} \sigma_{x,y}(ds). \tag{7}$$

By Hypothesis (H2') there exists $\epsilon > 0$ such that for all $x, x' \in K$ we can find $y \in K$ such that

$$m_{x,x',y} := \int_0^\tau \frac{M_{\tau-s}(\psi \mathbb{1}_K)(y)}{\psi(y)} (\sigma_{x,y} \wedge \sigma_{x',y})(ds) \geq \epsilon.$$

Setting

$$v_{x,x'} = \frac{1}{m_{x,x',y}} \int_0^\tau \frac{M_{\tau-s}(\psi \mathbb{1}_K \cdot)(y)}{\psi(y)} (\sigma_{x,y} \wedge \sigma_{x',y})(ds),$$

we deduce from (7) that (3) is satisfied with $\tilde{c} = c\epsilon/C$. □

2. An application in population dynamics

We apply our method to the following nonlocal equation with drift:

$$\partial_t u(t, x) + \partial_x u(t, x) = \int_{\mathbb{R}} u(t, y) Q(y, dx) dy + a(x)u(t, x), \quad x \in \mathbb{R}. \tag{8}$$

The solutions describe the density of traits in a population with nonlocal mutation in a space-time varying environment, encoding for instance the influence of a climate change through a spatial shift of the coefficient, see [4, 12] for motivations. In the recent work [12], Coville and Hamel address the Perron spectral problem associated to Equation (8) in a slightly more general case, namely with a possibly non constant transport speed $q(x)$ and on a domain Ω which can be an open subinterval of \mathbb{R} . In the present paper we choose to limit ourselves to the case $q \equiv 1$ and $\Omega = \mathbb{R}$ in order to remain concise. However, the method can be applied to other situations: to the price of additional technicalities for $q \neq 1$ since the flow of the transport part may not be explicit, or on the contrary with less calculations when $\Omega = (r_1, r_2)$ is a bounded interval ($-\infty < r_1 < r_2 < +\infty$) since no Lyapunov condition is needed in that case (i.e. Hypothesis 1 is verified).

We assume that $a : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function such that

$$\bar{a} = \sup_{x \in \mathbb{R}} a(x) < +\infty \quad \text{and} \quad \lim_{x \rightarrow \pm\infty} a(x) = -\infty \tag{9}$$

and $x \mapsto Q(x, \cdot)$ is a weak* continuous function $\mathbb{R} \rightarrow \mathcal{M}_+(\mathbb{1})$ which satisfies

$$\exists \epsilon, \kappa_0 > 0, \forall x \in \mathbb{R}, \quad Q(x, dy) \geq \kappa_0 \mathbb{1}_{(x-\epsilon, x+\epsilon)}(y) dy \quad (10)$$

$$\bar{Q} = \sup_{x \in \mathbb{R}} Q(x, \mathbb{R}) < +\infty. \quad (11)$$

The assumptions on Q are typically verified by a convolution kernel $Q(x, y) = J(x - y)$ with J a finite positive measure such that $J(dz) \geq \kappa_0 \mathbb{1}_{(-\epsilon, \epsilon)}(z) dz$. They are less restrictive than in [12], since we do not require the measures $Q(x, \cdot)$ to have a bounded and compactly supported Lebesgue density. Our potential function a satisfies a confining assumption $a(\pm\infty) = -\infty$ (no longer required if working in a bounded subinterval of \mathbb{R}), while it is supposed to be bounded in [12]. The reason is that in [12], only the existence of (λ, h) is addressed, whereas we prove the existence of (λ, h, γ) and the asynchronous exponential behavior of the solutions to the Cauchy problem. The well-posedness of Equation (8) completed with an initial condition $u(0, \cdot) = u_0$ in $\mathcal{M}(\mathbb{1})$ or in the subspace $L^1(\mathbb{R}, dx)$ is a standard result. Our method allows us to prove the following result stated in the L^1 framework, which is more usual in the partial differential equations community, but it is also valid for measure solutions by replacing the L^1 norm by the total variation norm.

Theorem 4. *Under Assumptions (9), (10) and (11), there exist constants $C, \omega > 0$ and a unique eigentriplet $(\lambda, \gamma, h) \in \mathbb{R} \times L^1 \times C_b^1$ with $\gamma \geq 0, h > 0$ and $\int h \gamma = \|h\|_\infty = 1$, such that for any initial condition $u_0 \in L^1(\mathbb{R}, dx)$ the corresponding solution $u(t, x)$ of Equation (8) verifies*

$$\left\| u(t, \cdot) e^{-\lambda t} - \left(\int_{\mathbb{R}} h(x) u_0(x) dx \right) \gamma \right\|_{L^1} \leq C \|u_0\|_{L^1} e^{-\omega t}.$$

The method of proof consists in applying our general result in Theorem 3 to the semigroup $(M_t)_{t \geq 0}$ associated to Equation (8), which is defined through the Duhamel formula

$$M_t f(x) = f(x+t) e^{\int_x^{x+t} a(s) ds} + \int_0^t e^{\int_x^{x+s} a(s') ds'} \int_{\mathbb{R}} M_{t-s} f(y) Q(x+s, dy) ds. \quad (12)$$

It is a standard result, see for instance [2, 3] for more details on a similar model, that the Duhamel formula (12) indeed defines a positive semigroup $(M_t)_{t \geq 0}$ in $\mathcal{B}(\mathbb{1})$, but also in $C_b(\mathbb{R})$ and in $C_b^1(\mathbb{R})$. When $f \in C_b^1(\mathbb{R})$ it additionally holds that $(t, x) \mapsto M_t f(x)$ is continuously differentiable and satisfies

$$\partial_t M_t f = \mathcal{L} M_t f = M_t \mathcal{L} f,$$

where

$$\mathcal{L} f(x) = f'(x) + a(x) f(x) + \int_{\mathbb{R}} f(y) Q(x, dy).$$

This right action of the semigroup provides the unique solutions to the dual equation

$$\partial_t \varphi(t, x) = \partial_x \varphi(t, x) + a(x) \varphi(t, x) + \int_{\mathbb{R}} \varphi(t, y) Q(x, dy)$$

and thus the left action, defined by duality through $(\mu M_t)(f) = \mu(M_t f)$, yields the unique measure solutions to the direct equation (8). The well-posedness of the direct equation in $L^1(\mathbb{R}, dx)$ ensures that if μ has a Lebesgue density, then so does μM_t for all $t \geq 0$.

For applying Theorem 3, we check that $(M_t)_{t \geq 0}$ verifies Hypothesis 2. Equation (8) is a nice example to illustrate the conditions (H1') and (H2'), as we will explain now. The drift term, which represents the spatial shift of the space-time varying environment, is crucial for verifying (H1'). If we delete it, the situation is more involved: the result of Theorem 4 is still true when typically $1/(\bar{a} - a)$ is not locally integrable [1, 6, 10, 14], otherwise it may not hold due to the presence of a Dirac mass in γ , see [7, 11]. Assumption (10) is crucial for verifying (H2'). If we consider for instance the singular kernel $Q(x, \cdot) = \delta_{x-1} + \delta_{x+1}$, then the convergence in Theorem 4 does not hold and a periodic asymptotic behavior takes place, similarly as in [5, 13].

Before proving Theorem 4, we start with a useful strong positivity result about M_t .

Lemma 5. *Let $x_1, x_2 \in \mathbb{R}$ with $x_1 < x_2$. Then for any $y_1, y_2 \in \mathbb{R}$ with $y_1 < y_2$ and any $\tau > 0$, there exists $\eta > 0$ such that*

$$M_\tau \mathbb{1}_{[x_1, x_2]} \geq \eta \mathbb{1}_{[y_1, y_2]}.$$

Proof of Lemma 5. Iterating once the Duhamel formula (12) we get for all $t > 0$

$$\begin{aligned} M_t \mathbb{1}_{[x_1, x_2]}(x) &\geq e^{t \inf_{(x, x+t)} a} \int_0^t \int_{\mathbb{R}} \mathbb{1}_{[x_1, x_2]}(y+t-s) Q(x+s, dy) ds \\ &\geq \kappa_0 t e^{t \inf_{(x, x+t)} a} \int_{x+t-\epsilon}^{x+t+\epsilon} \mathbb{1}_{[x_1, x_2]}(z) dz \end{aligned}$$

from which we deduce that $M_t \mathbb{1}_{[x_1, x_2]}(x) \geq \text{const} > 0$ for all $x \in [x_1 - t - \epsilon/2, x_2 - t + \epsilon/2]$. Letting n be a positive integer and considering $t = \tau/n$ we get that $M_t \mathbb{1}_{[x_1, x_2]}(x) \geq c_0 > 0$ for all $x \in [x_1 - \tau/n - \epsilon/2, x_2 - \tau/n + \epsilon/2]$, and by iteration $M_\tau \mathbb{1}_{[x_1, x_2]} \geq c_0^n \mathbb{1}_{[x_1 - \tau - n\epsilon/2, x_2 - \tau + n\epsilon/2]}$. The conclusion follows by choosing n large enough. \square

Proof of Theorem 4. We proceed in two steps. First we check that on any bounded set K , Hypotheses (H1') and (H2') are satisfied for any $\tau > 0$ (and any function ψ bounded from above and below by a positive constant over K). Then for Assumptions (A0)–(A2), we use $V = \mathbb{1}$ and build a suitable function ψ such that the sublevel sets of V/ψ are bounded.

Step 1. Let K be a bounded set and $\tau \in (0, \epsilon/2)$. We will build by induction two families $(\sigma_{x,y}^{t,n})$ and $(c_{x,y}^{t,n})$ indexed by $n \in \mathbb{N}$, $t \in [0, \tau]$, $x \in \mathbb{R}$ and $y \in [x + t - n\epsilon/2, x + t + n\epsilon/2]$ such that: $\sigma_{x,y}^{t,n}$ is a probability measure on $[0, t]$ which has a positive Lebesgue density $\mathfrak{s}_{x,y}^{t,n}$ when $n \geq 1$, $c_{x,y}^{t,n}$ is a positive constant, $(x, y, t, s) \mapsto \mathfrak{s}_{x,y}^{t,n}(s)$ and $(x, y, t) \mapsto c_{x,y}^{t,n}$ are continuous functions, and for all $f \geq 0$

$$M_t f(x) \geq c_{x,y}^{t,n} \int_0^t M_{t-s} f(y) \sigma_{x,y}^{t,n}(ds). \tag{13}$$

The integer n essentially represents a number of jumps required for reaching y by starting from x . Once these families are built, we choose n large enough so that $K \subset [x + \tau - n\epsilon/2, x + \tau + n\epsilon/2]$ for every $x \in K$. Positivity and continuity of $(x, y, t, s) \mapsto \mathfrak{s}_{x,y}^{t,n}(s)$ and $(x, y, t) \mapsto c_{x,y}^{t,n}$ then guarantee, together with Lemma 5, that Hypotheses (H1') and (H2') are satisfied with

$$\sigma_{x,y} = \sigma_{x,y}^{\tau,n} \quad \text{and} \quad c = \inf_{x,y \in K} c_{x,y}^{\tau,n}$$

for any function ψ bounded from above and below by a positive constant on K . Clearly if (H1') and (H2') are satisfied for any $\tau \in (0, \epsilon/2)$, they are also satisfied for any $\tau > 0$. Let us now give the details of the induction.

For $n = 0$, $y = x + t \geq x$ and the Duhamel formula (12) ensures that for any $f \geq 0$

$$M_t f(x) \geq f(y) e^{\int_x^y a(s) ds}. \tag{14}$$

This transcribes that y can be reached from x by only following the drift, without doing any jump, and it gives (13) with $\sigma_{x,y}^{t,0} = \delta_t$ and $c_{x,y}^{t,0} = e^{\int_x^y a(s) ds}$.

For $n = 1$, we start by noting that replacing f by $M_{t'-(y-x)} f$ in (14) gives

$$M_{t'} f(x) \geq e^{\int_x^y a(s) ds} M_{t'-(y-x)} f(y)$$

for any $t' \geq y - x \geq 0$. Using this inequality with $t' = t - s$ and $x = z$ together with Assumption (10) in the Duhamel formula (12) we get for all $y \in [x + t - \epsilon/2, x + t + \epsilon/2]$, $t \in [0, \tau]$,

$$\begin{aligned} M_t f(x) &\geq \int_0^t e^{\int_x^{x+s} a(s') ds'} \int_{\mathbb{R}} M_{t-s} f(z) Q(x + s, dz) ds \\ &\geq \kappa_0 \int_0^t e^{\int_x^{x+s} a} \int_{x+s-\epsilon}^{x+s+\epsilon} M_{t-s} f(z) dz ds \\ &\geq \kappa_0 \int_0^t e^{\int_x^{x+s} a} \int_{y+s-t}^y e^{\int_z^y a} M_{t-s-(y-z)} f(y) dz ds \\ &\geq \kappa_0 \int_0^t M_{t-s'} f(y) \left(\int_0^{s'} e^{\int_x^{x+s'} a + \int_{y+s'-s}^y a} ds' \right) ds'. \end{aligned}$$

This gives (13) when defining

$$c_{x,y}^{t,1} = \kappa_0 \int_0^t \int_0^s e^{\int_x^{x+s'} a + \int_{y+s'-s}^y a} ds' ds > 0$$

and $\sigma_{x,y}^{t,1}(ds) = \mathfrak{s}_{x,y}^{t,1}(s) ds$ with

$$\mathfrak{s}_{x,y}^{t,1}(s) = \frac{\kappa_0}{c_{x,y}^{t,1}} \int_0^s e^{\int_x^{x+s'} a + \int_{y+s'-s}^y a} ds' > 0.$$

For $n \rightarrow n + 1$, the induction hypothesis ensures that for any $0 \leq s \leq t \leq \tau$ and $z \in [y - t + s - n\epsilon/2, y - t + s + n\epsilon/2]$

$$M_{t-s} f(z) \geq c_{z,y}^{t-s,n} \int_s^t M_{t-s'} f(y) \mathfrak{s}_{z,y}^{t-s,n}(s' - s) ds'.$$

Injecting this inequality in the Duhamel formula (12) yields that for all $y \in [x + t - (n + 1)\epsilon/2, x + t + (n + 1)\epsilon/2]$, $t \in [0, \tau]$,

$$\begin{aligned} M_t f(x) &\geq \kappa_0 \int_0^t e^{\int_x^{x+s} a} \int_{x+s-\epsilon}^{x+s+\epsilon} M_{t-s} f(z) dz ds \\ &\geq \kappa_0 \int_0^t e^{\int_x^{x+s} a} \int_{\max(x+s-\epsilon, y-t+s-n\epsilon/2)}^{\min(x+s+\epsilon, y-t+s+n\epsilon/2)} c_{z,y}^{t-s,n} \int_s^t M_{t-s'} f(y) \mathfrak{s}_{z,y}^{t-s,n}(s' - s) ds' dz ds \\ &\geq \kappa_0 \int_0^t M_{t-s'} f(y) \left(\int_0^{s'} e^{\int_x^{x+s'} a} \int_{\max(x+s'-\epsilon, y-t+s'-n\epsilon/2)}^{\min(x+s'+\epsilon, y-t+s'+n\epsilon/2)} c_{z,y}^{t-s',n} \mathfrak{s}_{z,y}^{t-s',n}(s' - s) dz ds' \right) ds', \end{aligned}$$

which gives (13) for $n + 1$ with

$$c_{x,y}^{t,n+1} = \kappa_0 \int_0^t \int_0^s e^{\int_x^{x+s'} a} \int_{\max(x+s'-\epsilon, y-t+s'-n\epsilon/2)}^{\min(x+s'+\epsilon, y-t+s'+n\epsilon/2)} c_{z,y}^{t-s',n} \mathfrak{s}_{z,y}^{t-s',n}(s - s') dz ds' ds > 0$$

and $\sigma_{x,y}^{t,n+1}(ds) = \mathfrak{s}_{x,y}^{t,n+1}(s) ds$ with

$$\mathfrak{s}_{x,y}^{t,n+1}(s) = \frac{\kappa_0}{c_{x,y}^{t,n+1}} \int_0^s e^{\int_x^{x+s'} a} \int_{\max(x+s'-\epsilon, y-t+s'-n\epsilon/2)}^{\min(x+s'+\epsilon, y-t+s'+n\epsilon/2)} c_{z,y}^{t-s',n} \mathfrak{s}_{z,y}^{t-s',n}(s - s') dz ds' > 0.$$

Step 2. Let us fix $\tau > 0$. We use drift conditions, similarly as in [3, Section 2.2]. More precisely, we will prove that the functions

$$V = \mathbb{1} \quad \text{and} \quad \psi_0(x) = ((1 - (x/x_0)^2)_+)^2$$

satisfy

$$\mathcal{L}V \leq \alpha_0 V + \theta_0 \psi_0 \quad \text{and} \quad \mathcal{L}\psi_0 \geq \beta_0 \psi_0$$

for some $x_0 \geq \epsilon$ and some constants $\alpha_0 = \beta_0 - 1$ and $\theta_0 > 0$. From these drift conditions, we get that the function $\phi = V - \theta_0 \psi_0$ verifies

$$\mathcal{L}\phi \leq \alpha_0 V + \theta_0 \psi_0 - \theta_0 \beta_0 \psi_0 = \alpha_0 \phi$$

and, since $\partial_t M_t \phi = M_t \mathcal{L} \phi$, we deduce from Grönwall's lemma that $M_\tau \phi \leq e^{\alpha_0 \tau} \phi$, which gives

$$M_\tau V \leq e^{\alpha_0 \tau} (V - \theta_0 \psi_0) + \theta_0 M_\tau \psi_0 \leq e^{\alpha_0 \tau} V + \theta_0 M_\tau \psi_0.$$

We then prove the existence of $\zeta > 0$ such that $M_\tau \psi_0 \leq \zeta V$ and we set $\psi = \zeta^{-1} M_\tau \psi_0$, so that $\psi > 0$ due to Lemma 5, $\psi \leq V$, and

$$M_\tau V \leq e^{\alpha_0 \tau} V + \theta_0 \zeta \psi. \tag{15}$$

Besides, $\mathcal{L} \psi_0 \geq \beta_0 \psi_0$ yields $M_{2\tau} \psi_0 \geq e^{\beta_0 \tau} M_\tau \psi_0$, which also reads

$$M_\tau \psi \geq e^{\beta_0 \tau} \psi.$$

Choosing

$$R > \frac{\theta_0 \zeta}{e^{\alpha_0 \tau} (e^\tau - 1)} \quad \text{and} \quad K = \{V \leq R\psi\}$$

we obtain from (15) that

$$M_\tau V \leq \left(e^{\alpha_0 \tau} + \frac{\theta_0 \zeta}{R} \right) V + \theta_0 \zeta \mathbb{1}_K \psi$$

and Assumptions (A1) and (A2) are verified with $\theta = \theta_0 \zeta$ and

$$\alpha = e^{\alpha_0 \tau} + \frac{\theta_0 \zeta}{R} < e^{(\alpha_0 + 1)\tau} = e^{\beta_0 \tau} = \beta.$$

Finally we check that $\{V \leq R\psi\}$ is bounded, which ends the verification of Assumptions (A0)–(A2).

Now we show that $\mathcal{L} \psi_0 \geq \beta_0 \psi_0$ for some β_0 . We have

$$\begin{aligned} \mathcal{L} \psi_0(x) &= -4 \frac{x}{x_0^2} (1 - (x/x_0)^2)_+ + a(x) \psi_0(x) + \int_{-x_0}^{x_0} (1 - (y/x_0)^2)^2 Q(x, dy) \\ &\geq -\frac{4}{x_0} (1 - (x/x_0)^2)_+ + \left[\inf_{(-x_0, x_0)} a \right] \psi_0(x) + \kappa_0 x_0 \int_{-1}^1 (1 - y^2)^2 \mathbb{1}_{(x-\epsilon, x+\epsilon)}(x_0 y) dy \\ &\geq -\frac{4}{x_0} (1 - (x/x_0)^2)_+ + \left[\inf_{(-x_0, x_0)} a \right] \psi_0(x) + \kappa_0 \mathbb{1}_{(-x_0, x_0)}(x) x_0 \int_{1-\epsilon/x_0}^1 (1 - y)^2 (1 + y)^2 dy \\ &\geq -\frac{4}{x_0} (1 - (x/x_0)^2)_+ + \left[\inf_{(-x_0, x_0)} a \right] \psi_0(x) + \frac{8\kappa_0 \epsilon^3}{15 x_0^2} \mathbb{1}_{(-x_0, x_0)}(x), \end{aligned}$$

since for $r \in [0, 1]$ we have $\int_{1-r}^1 (1 - y)^2 (1 + y)^2 dy = \int_0^r z^2 (2 - z)^2 dz = r^3 (\frac{4}{3} - r + \frac{r^2}{5}) \geq \frac{8r^3}{15}$. If $2\kappa_0 \epsilon^3 \geq 15x_0$ we get

$$\mathcal{L} \psi_0 \geq \left[\inf_{(-x_0, x_0)} a \right] \psi_0.$$

Otherwise we split, for $|x| \geq \sqrt{1 - (2\kappa_0 \epsilon^3)/(15x_0)} x_0$,

$$\mathcal{L} \psi_0(x) \geq \left[\inf_{(-x_0, x_0)} a \right] \psi_0(x),$$

and for $|x| \leq \sqrt{1 - (2\kappa_0 \epsilon^3)/(15x_0)} x_0$, we have $\sqrt{\psi_0(x)} \geq \frac{2\kappa_0 \epsilon^3}{15x_0}$ which yields $\sqrt{\psi_0(x)} \leq \frac{15x_0}{2\kappa_0 \epsilon^3} \psi_0(x)$ and thus

$$\mathcal{L} \psi_0(x) \geq \left[-\frac{30}{\kappa_0 \epsilon^3} + \inf_{(-x_0, x_0)} a + \frac{8\kappa_0 \epsilon^3}{15 x_0^2} \right] \psi_0(x).$$

At the end, in any case and for any $x_0 \geq \epsilon$,

$$\mathcal{L} \psi_0 \geq \left[-\frac{30}{\kappa_0 \epsilon^3} + \inf_{(-x_0, x_0)} a \right] \psi_0 := \beta_0 \psi_0.$$

Next, for $V = \mathbb{1}$, we have

$$\mathcal{L} V(x) = a(x) + Q(x, \mathbb{R}) \leq a(x) + \bar{Q}.$$

Choosing $r_0 > 0$ such that $a(x) < -\bar{Q} + \beta_0 - 1 := -\bar{Q} + \alpha_0$ when $|x| > r_0$ we get

$$\mathcal{L} V(x) \leq \alpha_0 V(x) + (\bar{a} + \bar{Q}) \mathbb{1}_{(-r_0, r_0)}(x).$$

So if we choose $x_0 \geq \sqrt{2}r_0$ in the definition of ψ_0 we get $\mathbb{1}_{(-r_0, r_0)} \leq 4\psi_0$ and thus

$$\mathcal{L}V \leq \alpha_0 V + 4(\bar{a} + \bar{Q})\psi_0 := \alpha_0 V + \theta_0 \psi_0.$$

Besides, since $\mathcal{L}V \leq \bar{a} + \bar{Q}$, we have by Grönwall's lemma that

$$M_\tau \psi_0 \leq M_\tau V \leq e^{-(\bar{a} + \bar{Q})\tau} V := \zeta V.$$

Finally $K = \{V \leq R\psi\}$ is bounded because ψ tends to zero at $\pm\infty$: from the Duhamel formula (12) we have

$$M_\tau \psi_0(x) \leq \psi_0(x+t) e^{\int_x^{x+t} a(s) ds} + \bar{Q} \int_0^\tau e^{\int_x^{x+s} a(s') ds'} e^{(t-s)(\bar{a} + \bar{Q})} ds$$

which yields the result since $\lim_{x \rightarrow \pm\infty} a(x) = -\infty$.

Conclusion. We have checked Hypothesis 2 so the result is a direct application of Theorem 3, the regularity of γ following from the convergence in total variation norm and the invariance of L^1 under the semigroup, and the regularity of h being obtained by usual arguments from Equation (12). \square

Acknowledgments

The authors are very grateful to Vincent Bansaye and Aline Marguet for useful discussions about the results of the present note.

References

- [1] M. Alfaro, P. Gabriel, O. Kavian, "Confining integro-differential equations from evolutionary biology: ground states and long time dynamics", in preparation.
- [2] V. Bansaye, B. Cloez, P. Gabriel, "Ergodic behavior of non-conservative semigroups via generalized Doeblin's conditions", *Acta Appl. Math.* **166** (2020), no. 1, p. 29-72.
- [3] V. Bansaye, B. Cloez, P. Gabriel, A. Marguet, "A non-conservative Harris ergodic theorem", <https://arxiv.org/abs/1903.03946>, 2019.
- [4] H. Berestycki, O. Diekmann, C. J. Nagelkerke, P. A. Zegeling, "Can a species keep pace with a shifting climate?", *Bull. Math. Biol.* **71** (2009), no. 2, p. 399-429.
- [5] É. Bernard, M. Doumic, P. Gabriel, "Cyclic asymptotic behaviour of a population reproducing by fission into two equal parts", *Kinet. Relat. Models* **12** (2019), no. 3, p. 551-571.
- [6] R. Bürger, "Perturbations of positive semigroups and applications to population genetics", *Math. Z.* **197** (1988), no. 2, p. 259-272.
- [7] R. Bürger, I. M. Bomze, "Stationary distributions under mutation-selection balance: structure and properties", *Adv. Appl. Probab.* **28** (1996), no. 1, p. 227-251.
- [8] N. Champagnat, D. Villemonais, "Exponential convergence to quasi-stationary distribution and Q -process", *Probab. Theory Relat. Fields* **164** (2016), no. 1-2, p. 243-283.
- [9] ———, "General criteria for the study of quasi-stationarity", <https://arxiv.org/abs/1712.08092>, 2017.
- [10] J. Coville, "On a simple criterion for the existence of a principal eigenfunction of some nonlocal operators", *J. Differ. Equations* **249** (2010), no. 11, p. 2921-2953.
- [11] ———, "Singular measure as principal eigenfunction of some nonlocal operators", *Appl. Math. Lett.* **26** (2013), no. 8, p. 831-835.
- [12] J. Coville, F. Hamel, "On generalized principal eigenvalues of nonlocal operators with a drift", *Nonlinear Anal., Theory Methods Appl.* **193** (2019), article ID 111569 (20 pages).
- [13] P. Gabriel, H. Martin, "Periodic asymptotic dynamics of the measure solutions to an equal mitosis equation", <https://arxiv.org/abs/1909.08276>, 2019.
- [14] F. Li, J. Coville, X. Wang, "On eigenvalue problems arising from nonlocal diffusion models", *Discrete Contin. Dyn. Syst.* **37** (2017), no. 2, p. 879-903.
- [15] R. J. Nagel (ed.), *One-parameter semigroups of positive operators*, Lecture Notes in Mathematics, vol. 1184, Springer, 1986.
- [16] J. R. Norris, *Markov chains*, Cambridge Series in Statistical and Probabilistic Mathematics, vol. 2, Cambridge University Press, 1998, reprint of 1997 original.