

# Far from equilibrium steady states of 1D-Schrödinger–Poisson systems with quantum wells I

V. Bonnaillie-Noël<sup>a,\*</sup>, F. Nier<sup>b</sup>, Y. Patel<sup>b</sup>

<sup>a</sup> IRMAR, UMR–CNRS 6625, ENS Cachan Bretagne, Université Rennes 1, UEB, av. Robert Schuman, 35170 Bruz, France

<sup>b</sup> IRMAR, UMR–CNRS 6625, Université Rennes 1, Campus de Beaulieu, 35042 Rennes, France

Received 17 January 2007; received in revised form 3 May 2007; accepted 31 May 2007

Available online 17 October 2007

---

## Abstract

We describe the asymptotics of the steady states of the out-of-equilibrium Schrödinger–Poisson system, in the regime of quantum wells in a semiclassical island. After establishing uniform estimates on the nonlinearity, we show that the nonlinear steady states lie asymptotically in a finite-dimensional subspace of functions and that the involved spectral quantities are reduced to a finite number of so-called asymptotic resonant energies. The asymptotic finite dimensional nonlinear system is written in a general setting with only a partial information on its coefficients. After this first part, a complete derivation of the asymptotic nonlinear system will be done for some specific cases in a forthcoming article [V. Bonnaillie–Noël, F. Nier, M. Patel, Far from equilibrium steady states of 1D-Schrödinger–Poisson systems with quantum wells II, Prépublications IRMAR, 2007].

© 2007 Elsevier Masson SAS. All rights reserved.

MSC: 34L25; 34L30; 34L40; 65L10; 65Z05; 81Q20; 82D37

Keywords: Schrödinger–Poisson system; Asymptotic analysis; Multiscale problems

---

## 1. Introduction

### 1.1. Motivation

This analysis is motivated by the study of quantum electronic transport in semiconductor heterostructures, like resonant tunnelling diodes. It is modelled on the basis of a mean field Hartree type description of the electrostatic interaction of particles, known as the Schrödinger–Poisson system. The modelling of resonant tunnelling diodes includes the following characteristic features:

- (1) Steady electronic currents are observed. This can be achieved only within the modelling of out-of-equilibrium quantum systems.

---

\* Corresponding author.

E-mail addresses: [Virginie.Noel-Bonnaillie@univ-rennes1.fr](mailto:Virginie.Noel-Bonnaillie@univ-rennes1.fr) (V. Bonnaillie-Noël), [Francis.Nier@univ-rennes1.fr](mailto:Francis.Nier@univ-rennes1.fr) (F. Nier).

- (2) The  $I$ – $V$  curves of such devices present negative differential resistance. We are in a far from equilibrium regime, for which the linear response theory is questionable.
- (3) A very rich nonlinear phenomenology can be observed in such devices, with hysteresis phenomena (see [21,29]) and even steadily oscillating currents (see [22]).
- (4) The general wisdom about these systems says that the nonlinear effects are governed by a small number of resonant states.

This article is a part of a larger program, namely the understanding of the nonlinear dynamics of these out-of-equilibrium quantum systems. One issue is to prove rigorously that a simple Schrödinger–Poisson system in a far from equilibrium regime, that is when the steady states show a strong anisotropy in the momentum variable at the quantum scale, can lead to multiple solutions to the nonlinear stationary problem with nontrivial bifurcation diagrams. A first check was provided by Jona-Lasinio, Presilla and Sjöstrand in [21,29]. A second issue which goes definitely further than those previous works is the explanation of the production of complex bifurcation diagrams in terms of the geometry of the potential, which requires an accurate analysis of tunnel effects.

The present work was achieved on the basis of former works by the second author and of the PhD thesis of the third author. This analysis led the three authors to the introduction of some reduced model which happens to be very efficient in the numerical simulation of realistic devices (see [5]). Only the first part of the mathematical analysis is provided here and complements will be presented in a forthcoming article [6].

The points (1) and (2) above are now well understood. A presentation can be done within a Landauer–Büttiker approach (see [7,23,8,4]) which involves the scattering states. This modelling allows a strong anisotropy of the occupation number with respect to the momentum and it definitely differs from the approach where the density matrix looks like a function of the Hamiltonian (see [2,3]). This latter modelling (and probably the entropy maximising approach of [10] as well) better suits the situation of little variations from the thermodynamical equilibrium, ends with corrected drift-diffusion models and cannot produce multiple solutions due to monotonicity properties. It should be noted that all these modellings consider the reservoirs as fixed objects which only provide some kind of inhomogeneous boundary conditions, in comparison with the theoretical analysis of nonequilibrium steady states widely studied within the framework of the von Neumann algebraic approach of statistical physics and which concerns the evolution of the full system, small system plus reservoirs (see for example [20]).

For our model, a complete general functional framework which catches the proper nonlinear steady states and provides a well defined nonlinear dynamics was provided in [26], after using a phase-space approach with some specific tools of the time dependent approach in scattering theory.

Besides the building of a proper functional framework, those models became even more interesting after the articles of Jona-Lasinio, Presilla and Sjöstrand [21,29] where convincing heuristic arguments and calculations on those simple nonlinear systems were provided as an explanation for observed hysteresis phenomena, in agreement with point (3). Later the question arose whether a complete explanation from an asymptotic analysis on the Schrödinger–Poisson system or whether new nonlinear phenomena could be predicted in some more complex geometric setting like a multiple wells problem. For instance, no real explanation is provided in [21,29] for the presence or the absence of hysteresis phenomena according to the geometry of the barrier potentials. Our reduced model (see [27,5] and forthcoming article [6]) provides such an explanation, with additional results.

Finally point (4) provides the relevant asymptotics. Resonant states are effective when the imaginary part of resonances are small. Such a behaviour can be achieved when the potential barrier are high or large and it is well formulated within a semiclassical asymptotics (small parameter  $\hbar \rightarrow 0$ , imaginary part of resonances of order  $\mathcal{O}(e^{-c/\hbar})$ ). Nevertheless a full semiclassical asymptotics with  $\mathcal{O}(1)$  large wells would lead to a large number of resonant states within a fixed energy interval. Point (4) can be fulfilled by considering quantum wells in a semiclassical island. The introduction of the small parameter  $\hbar > 0$  as a rescaled Fermi-length as well as a full justification of this asymptotic regime within the presentation of realistic devices has been done in [5].

From a mathematical point of view, this problem presents two specific difficulties.

- A nonusual multiple wells problem has to be considered: it is not exactly a semiclassical problem and it is nonlinear.
- The introduction of resonances requires the implementation of a complex deformation and the study of non self-adjoint operators.

Fortunately, the one-dimensional framework provides some simplifications or accurate estimates which allow a complete analysis. First a uniform control on the nonlinear potential with the help of some monotony principles can be obtained in  $W^{1,\infty}$ . Hence the nonlinear potential can be replaced by an  $h$ -dependent potential, with uniform bounds in  $W^{1,\infty}$ . Some standard arguments of the semiclassical analysis for resonances (see [15]), for multiple wells (see [16, 17]), or for the Breit–Wigner formula (see [13]) have to be adapted. Again the weak regularity is partly compensated by the fact that we work on a 1D problem. Throughout this text proofs which are exactly the same as in the usual semiclassical setting are not reproduced. Precise references are given for these technical parts. Nevertheless some details have to be checked in order to ensure that these techniques can be adapted with the quantum wells and the limited regularity of the nonlinear semiclassical potential. The 1D Schrödinger–Poisson system studied here admits natural *a priori* regularity estimates, uniform with respect to the small parameter  $h \rightarrow 0$ . This leads asymptotically to a perfect splitting of the quantum and classical scales.

### 1.2. Quantum framework

In the whole study, the framework is the following:  $h > 0$  denotes the semiclassical parameter obtained in realistic cases as a rescaled Fermi length (see [5]) and  $I := [a, b]$  is a given compact interval of the real line. Let  $P_B^h$  be the Schrödinger operator on the real line:

$$P_B^h := -h^2 \frac{d^2}{dx^2} + \mathcal{B}, \quad \mathcal{B} \equiv \mathcal{B}_I + \mathcal{B}_\infty, \tag{1.1}$$

where

$$\mathcal{B}_I(x) := -B \frac{x-a}{b-a} \mathbf{1}_{[a,b]}(x), \quad \mathcal{B}_\infty(x) := -B \cdot \mathbf{1}_{[b,+\infty)}(x), \tag{1.2}$$

and  $B$  is a nonnegative constant. The potential  $\mathcal{B}$  simply describes the applied bias. The reference Hamiltonian is the self-adjoint realisation in the Hilbert space  $L^2(\mathbb{R})$  of  $P_B^h$ :

$$D(H_B^h) = H^2(\mathbb{R}), \quad \forall u \in D(H_B^h), \quad H_B^h u := P_B^h u. \tag{1.3}$$

Since several self-adjoint (or non self-adjoint) closures of the same differential operator associated with various boundary value problems will be considered, the notation  $P$  refers to the differential operators acting on  $C_0^\infty$ , while  $H$  will be used for its realisation as an unbounded operator on  $L^2$ .

We restrict our analysis in this work to operators in the form

$$P^h[V] := P_B^h + V, \quad V \in L^\infty(I), \tag{1.4}$$

and denote by  $H^h[V]$  the self-adjoint realisation in  $L^2(\mathbb{R})$  of  $P^h[V]$ :

$$D(H^h[V]) = H^2(\mathbb{R}), \quad \forall u \in D(H^h[V]), \quad H^h[V]u := P^h[V]u, \tag{1.5}$$

after identifying  $V \in L^\infty(I)$  with  $V(x)\mathbf{1}_I(x) \in L^\infty(\mathbb{R})$ .

Of particular interest is the case where the potential  $V = V^h$  depends on the small parameter  $h$  and describes quantum wells in an island with cliffs (i.e. the classically forbidden region ends with discontinuities of the potential). It splits into

$$V^h := V_0 + V_{NL}^h, \quad V_0 := \tilde{V}_0 - W^h, \quad \tilde{V}_0, V_{NL}^h \in W^{1,\infty}(I). \tag{1.6}$$

The function  $\tilde{V}_0$ , which models the island potential, can be any nonnegative Lipschitz function independent of  $h$ . Practically it is simply a constant potential on  $I$ ,  $\tilde{V}_0(x) = V_0 \mathbf{1}_I(x)$  with  $V_0 \in \mathbb{R}_+$ . The function  $W^h$ , which described the quantum wells, is defined by

$$W^h(x) := \sum_{i=1}^N w_i \left( \frac{x - c_i}{h} \right). \tag{1.7}$$

In this definition of  $W^h$ , the positions  $(c_i)_{i=1}^N$  are  $N$  given points in  $(a, b)$  and  $w_i$  are nonnegative  $L^\infty$ -functions supported in the interval  $[-\kappa, \kappa]$ , with  $\kappa > 0$  fixed. We denote by  $U^h$  the support of the function  $W^h$  and  $U := \bigcup_{i=1}^N \{c_i\}$  the region where the quantum wells concentrate, and set  $c_0 := a$ ,  $c_{N+1} := b$  (see Fig. 1).




Fig. 1. Total potential  $\mathcal{B} + V^h - W^h$ .

**Assumption 1.** Suppose that

$$\Lambda_0 := \inf_{x \in I} \tilde{V}_0(x) + \mathcal{B}(x) > 0, \tag{1.8}$$

and fix the parameters  $\Lambda_*$  and  $\Lambda^*$  so that  $0 < \Lambda_* < \Lambda^* < \Lambda_0$ .

We will focus on the energy range  $\lambda \in [\Lambda_*, \Lambda^*]$ .

Finally the function  $V_{NL}^h$  describes the mean field nonlinear potential which takes into account the repulsive electrostatic interaction. It will be given as a solution to the Poisson equation on  $I = [a, b]$  and will satisfy

$$\forall h > 0, \quad V_{NL}^h \in W^{1,\infty}(I), \quad V_{NL}^h \geq 0. \tag{1.9}$$

Such Hamiltonians are used in the modelling of quantum electronic transport in mesoscopic structures like resonant tunnelling diodes (RTD) or super-lattices. The nonlinear steady states can be studied within a Landauer–Büttiker approach: see [7,23,8,4] or [26] for possible functional frameworks concerned with the extension to the nonlinear analysis including the nonlinear dynamics. This approach involves the scattering wave functions and requires the analysis of the continuous spectrum of  $H^h[V]$ . Since for any potential  $V \in L^\infty(I)$ ,  $H^h[V]$  is a compactly supported  $L^\infty$ -perturbation of the reference Hamiltonian  $H_B^h$  or the Hamiltonian with step potential  $-h^2\Delta + \mathcal{B}_\infty$ , the limiting absorption principle holds. By standard arguments (see [32,28]) one even gets the absence of embedded eigenvalues

$$\forall h > 0, \quad \sigma_{\text{ess}}(H^h[V]) = \sigma_{\text{ac}}(H^h[V]) = [-B; \infty), \tag{1.10}$$

and the scattering states of  $H^h[V]$  are indeed well defined for any  $V \in L^\infty(I)$ .

**Remark 1.** Under the nonnecessary additional assumption

$$\forall i \in \{1, \dots, N\}, \quad \tilde{V}_0(c_i) + \inf \sigma(-\Delta - w_i) > 0, \tag{1.11}$$

one can even check like in Theorem 3.4 or Theorem 3.6 that there is no eigenvalue at all for  $h > 0$  small enough (and  $V_{NL}^h \geq 0$ );

$$\sigma(H^h[V]) = \sigma_{\text{ac}}(H^h[V]) = [-B, +\infty).$$

We focus on the energies  $\lambda \in [\Lambda_*, \Lambda^*]$ .

We consider the incoming scattering states  $\psi_-^h(k, \cdot)$  of the Hamiltonian  $H^h[V]$  parameterised by the wave vector  $k$  (we omit to write the dependence with respect to the potential for scattering states). They provide a diagonalisation of  $H^h[V]$  over the continuous spectrum (see formula (1.19)). Precisely, introduce first the dispersion relation associated with the reference Hamiltonian  $H_B^h$ .

**Definition 1.1.** Set for  $k \in \mathbb{R}^*$

$$\lambda_k := \begin{cases} k^2 & \text{if } k > 0, \\ k^2 - B & \text{if } k < 0. \end{cases} \tag{1.12}$$

This dispersion relation (1.12) gives, for the wave vector  $k$ , the energy  $\lambda_k$  of the incoming plane wave represented by  $\psi_-^h(k, \cdot)$ . Again, we are mostly interested in the  $k$ 's such that  $\lambda_k \in [\Lambda_*, \Lambda^*]$ .

By definition, the incoming generalised eigenfunction  $\psi_-^h(k, \cdot)$  defined for  $k \in \mathbb{R}$  solves the differential equation:

$$P^h \psi_-^h(k, \cdot) = \lambda_k \psi_-^h(k, \cdot), \tag{1.13}$$

with the normalisation (of incoming plane waves)

$$\text{for } k > 0, \quad \psi_-(k, x) = \begin{cases} e^{i\frac{kx}{h}} + r_k e^{-i\frac{kx}{h}} & \text{for } x < a, \\ t_k e^{i\frac{(\lambda_k+B)^{1/2}x}{h}} & \text{for } x > b, \end{cases} \tag{1.14}$$

$$\text{for } k < 0, \quad \psi_-(k, x) = \begin{cases} t_k e^{-i\frac{(\lambda_k)^{1/2}x}{h}} & \text{for } x < a, \\ e^{i\frac{kx}{h}} + r_k e^{-i\frac{kx}{h}} & \text{for } x > b. \end{cases} \tag{1.15}$$

The square root  $z^{1/2}$  is chosen with the ramification along the half-line  $i\mathbb{R}_-$  in order to ensure that  $e^{-i(\lambda_k)^{1/2}x}$  decays exponentially as  $x \rightarrow -\infty$  when  $\lambda_k \in (-B, 0)$ .

These coefficients determine the scattering matrix  $(r_k, t_k)$  for positive energies  $\lambda_k > 0$ . They are linked for  $\lambda_k > 0$  by the relation

$$|r_k|^2 + \sqrt{\frac{\lambda_k}{\lambda_k + B}} |t_k|^2 = 1, \quad \lambda_k > 0. \tag{1.16}$$

Since the wave vector  $k$  is a log-derivative, this normalisation of the wave functions can be written in terms of boundary conditions at  $x = a$  and  $x = b$ , in this specific one-dimensional case fitting with realistic problems:

$$\begin{aligned} [h\partial_x + i\lambda_k^{1/2}]_{|x=a} u &= 2ik e^{i\frac{ka}{h}}, \\ [h\partial_x - i(\lambda_k + B)^{1/2}]_{|x=b} u &= 0, \quad \text{for } k > 0, \end{aligned} \tag{1.17}$$

and

$$\begin{aligned} [h\partial_x + i\lambda_k^{1/2}]_{|x=a} u &= 0, \\ [h\partial_x - i(\lambda_k + B)^{1/2}]_{|x=b} u &= 2ik e^{i\frac{kb}{h}}, \quad \text{for } k < 0. \end{aligned} \tag{1.18}$$

Thus the problem over the real line is reduced to a boundary problem on  $I$  with boundary conditions depending on the spectral parameter (1.17)–(1.18). These boundary conditions are exact transparent boundary conditions. This setting makes rather easy the complex deformation argument used in the analysis of resonances (see [1,15] or [19] for a more general introduction). Here considering a complex  $\lambda_k$  around any positive value is easily implemented because the coefficients on the boundary conditions at  $x = a$  and  $x = b$  depend holomorphically on  $\lambda_k$  (or  $k$ ).

We end this section with three elementary properties:

1. With this normalisation, it appears that for any nonnegative continuous function  $\theta$  on  $[\Lambda_*, \Lambda^*]$ , the operator  $\mathbf{1}_I \theta(H^h[V]) \mathbf{1}_I$  is an integral operator. Moreover the kernel is given by

$$\mathbf{1}_I \theta(H^h[V]) \mathbf{1}_I [x, y] = \int_{\mathbb{R}} \theta(\lambda_k) \psi_-^h(k, x) \overline{\psi_-^h(k, y)} \frac{dk}{2\pi h}, \quad (x, y) \in I \times I. \tag{1.19}$$

2. Note that because of the regularity of  $\psi_-^h$ , it follows by Mercer's theorem (see [30, Theorem 3.5]) that this operator is trace-class, with a trace equal to the diagonal integral.
3. Note also that because the solutions to the ODE (1.13) in the interval  $I$  is a 2-dimensional linear subspace, say  $\mathcal{S}_{\lambda_k} \subset H^2(a, b)$ , conditions (1.17)–(1.18) form an affine system in  $\mathcal{S}_{\lambda_k}$ . Resonances around positive energies correspond to the exceptional complex values of  $\lambda_k = z$  for which the continuous linear functionals defining this system are proportional.

### 1.3. Schrödinger–Poisson system

Here we are interested in the study of the stationary case. We first fix the profile of the incoming beam of electrons over the structure between  $a$  and  $b$ .

**Notation 1.** Fix a continuous nonnegative function  $k \mapsto g(k)$  such that  $g(k) = 0$  if  $\lambda_k \notin (\Lambda_*, \Lambda^*)$ , see (1.12).

A beam of electrons corresponds to a superposition of scattering states with density  $g$ . The electronic density is described by the measure  $dn_g[V]$  defined by

$$dn_g[V](x) := \int_{\mathbb{R}} g(k) |\psi_{-}^h(k, x)|^2 \frac{dk}{2\pi h}. \tag{1.20}$$

It is convenient to introduce the function  $g(K_{-}^h)$  of the asymptotic momentum operator defined (see [11,26] for a more general presentation) according to:

$$g(K_{-}^h)[x, y] = \int_{\mathbb{R}} g(k) \psi_{-}^h(k, x) \overline{\psi_{-}^h(k, y)} \frac{dk}{2\pi h}.$$

Its localised version  $\mathbf{1}_I g(K_{-}^h) \mathbf{1}_I$  has the integral kernel

$$\mathbf{1}_I g(K_{-}^h) \mathbf{1}_I[x, y] = \int_{\mathbb{R}} g(k) \mathbf{1}_I(x) \psi_{-}^h(k, x) \overline{\psi_{-}^h(k, y)} \mathbf{1}_I(y) \frac{dk}{2\pi h}. \tag{1.21}$$

The operator  $g(K_{-}^h)$  is a density matrix and the density fulfills the weak formulation

$$\forall \varphi \in C^0(I), \quad \int_I \varphi(x) dn_g[V](x) = \text{Tr}[\mathbf{1}_I g(K_{-}^h) \mathbf{1}_I \varphi]. \tag{1.22}$$

Note that in the particular case where  $g(k)$  is a function of the energy, i.e.  $g(k) \equiv \theta(\lambda_k)$ ,  $g(K_{-}^h)$  is a function of the Hamiltonian

$$g(K_{-}^h) = \theta(H^h). \tag{1.23}$$

Functions of the Hamiltonian can be viewed as equilibrium states (and even thermodynamical equilibrium states when  $\theta$  is decreasing). For such states, the current through the device is null. Hence out-of-equilibrium steady states with a nonvanishing current have to be described with a function  $g(k)$  which is not a function of the energy. In order to make this situation clear, we assume the next possibly extendible assumption (see [5] for an easy generalisation towards more realistic problems).

**Assumption 2.** Fix a nonnegative function  $\theta \in C_c^0((\Lambda_*, \Lambda^*))$  and assume that

$$g(k) = 1_{k>0} \cdot \theta(\lambda_k). \quad \text{In particular, } 0 \leq g(k) \leq \theta(\lambda_k). \tag{1.24}$$

The Schrödinger–Poisson system is an Hartree model which includes the self-consistent electrostatic potential within the device ( $a \leq x \leq b$ ). Hence the nonlinear potential  $V_{NL}^h$  is a solution to

$$\begin{cases} H^h[V^h] = H_B^h + \tilde{V}_0 - W^h + V_{NL}^h, \\ -\Delta V_{NL}^h = dn_g[V^h], \quad V_{NL}^h(a) = V_{NL}^h(b) = 0. \end{cases} \tag{1.25}$$

Note that the assumption  $g \geq 0$  yields  $dn_g[V^h] \geq 0$  and  $V_{NL}^h \geq 0$ .

It is known, (see [4,26]), that the system (1.25) admits solutions, for fixed  $h > 0$ . Furthermore with the absence of negative eigenvalues provided by the condition (1.11), it is easily checked that the solutions to (1.25) are the only steady states of the nonlinear dynamics studied in [26].

Yet, uniform estimates with respect to  $h$  are not given in [26]. Here we are interested in the structure of the set of asymptotic solutions as  $h \rightarrow 0$ . A first step consists in getting *a priori* estimates on the semi-linear problem. This is

performed in Section 2. Since for a given  $h > 0$  the density  $dn_g[V^h]$  is a bounded positive measure, we introduce the following spaces:

**Definition 1.2.** Call  $(\mathcal{M}_b(I), \|\cdot\|_b)$  the Banach space of bounded complex measures on  $[a, b]$  and set

$$BV_0^2(I) := \{V \in \mathcal{C}^0(I) \mid V'' \in \mathcal{M}_b(I), V(a) = 0 = V(b)\}, \tag{1.26}$$

with the norm  $\|V\| := \|V\|_\infty + \|V''\|_b$ .

With this norm,  $BV_0^2(I)$  is a Banach space continuously embedded in  $W^{1,\infty}(I)$  and compactly embedded in the Hölder spaces  $\mathcal{C}^{0,\alpha}(I)$  for  $\alpha \in (0, 1)$ .

1.4. Results

**Theorem 1.3.** Consider problem (1.25). Then for  $h > 0$  sufficiently small:

- (i) The family of potentials  $(V_{NL}^h)_{h>0}$  is uniformly bounded in  $L^\infty(I)$ .
- (ii) The family of measures  $(dn_g[V^h])_{h>0}$  is bounded in  $\mathcal{M}_b(I)$  and the family  $(V_{NL}^h)_{h>0}$  is bounded in  $BV_0^2(I)$ .
- (iii) Consequently, the family of potentials  $(V_{NL}^h)_{h>0}$  is bounded in  $W^{1,\infty}(I)$  and relatively compact in the Hölder space  $\mathcal{C}^{0,\alpha}(I)$  for any  $\alpha \in (0, 1)$ .

After this, we try to identify the weak\* possible limits  $dn_g^0$  of the measure  $dn_g[V^h]$ . Owing to the boundedness stated in Theorem 1.3(ii), we shall make the next simplifying assumption which makes sense after possibly extracting a subsequence  $(h_n)_{n \in \mathbb{N}}$ .

**Assumption 3.** The convergence

$$dn_g[V^h] \xrightarrow{h \rightarrow 0} dn_g^0$$

holds for the weak topology of  $\mathcal{M}_b(I) = \mathcal{C}^0(I)'$ .

The following notations for the total potential

$$\mathcal{V}^h := V^h + \mathcal{B} = \tilde{V}_0 + V_{NL}^h - W^h + \mathcal{B}, \tag{1.27}$$

and for the total potential with filled wells

$$\tilde{\mathcal{V}}^h := \mathcal{V}^h + W^h = \tilde{V}_0 + V_{NL}^h + \mathcal{B}, \tag{1.28}$$

are convenient. The solution to

$$-\Delta V = dn_g^0, \quad V(a) = V(b) = 0 \tag{1.29}$$

is denoted  $V_{NL}^0$  and we set

$$\tilde{\mathcal{V}}^0 := \tilde{V}_0 + V_{NL}^0 + \mathcal{B}. \tag{1.30}$$

Theorem 1.3 has the next consequence.

**Corollary 1.4.** Make Assumption 3. Then the filled potential  $\tilde{\mathcal{V}}^h$  is uniformly bounded in  $W^{1,\infty}(I)$  and converges in  $\mathcal{C}^{0,\alpha}(I)$  to  $\tilde{\mathcal{V}}^0$  as  $h \rightarrow 0$  for any  $\alpha < 1$ . Moreover if the second derivative  $\partial_x^2 \tilde{V}_0$  is a bounded measure, the weak convergence

$$\partial_x^2 \tilde{\mathcal{V}}^h \xrightarrow{h \rightarrow 0} \partial_x^2 \tilde{\mathcal{V}}^0 = \partial_x^2 \tilde{V}_0 - dn_g^0$$

also holds in  $\mathcal{M}_b(I)$ .

**Remark 2.** Note that the solution of the asymptotic Poisson equation does not depend on the possible mass of  $dn_g^0$  concentrated on  $x = a$  or  $x = b$ . Indeed the asymptotic potential  $V_{NL}^0$  forgets any boundary layer and the boundary value problem (1.29) is equivalently written with the restricted measure  $dn_g^0|_{(a,b)}$ .

The idea leading to an accurate description of the asymptotic density  $dn_g^0$  is the following: suppose in a first step that the wells are filled, that is  $W^h = 0$  and  $\mathcal{V}^h = \tilde{\mathcal{V}}^h$ . In the classical picture, the incoming particles of energy  $\lambda_k \leq \Lambda^*$  are reflected by the cliffs, so one expects that  $dn_g^0 \equiv 0$  in  $(a, b)$ . The introduction of the wells  $W^h$  generates trapped quantum states transformed into resonant states after the interaction with the continuous spectrum. The tunnel effect allows these states to be occupied in a stationary setting. Besides, the quantum wells with an  $\mathcal{O}(h)$ -diameter produce two interesting effects. Firstly the density will asymptotically concentrate like delta-functions in positions around the  $c_i$ 's. Secondly the resonant energies attached to one well are separated by  $\mathcal{O}(1)$  gaps (see Remark 3 below). With a finite number of wells, this asymptotics implements the general wisdom that the nonlinear system is essentially governed by *finite* number of resonant states of the system (point (4) of our introduction). The relevancy of this asymptotics, with quantum wells in a semiclassical island, has been carefully checked in [5] with numerical data fitting with realistic situations.

To state our results we need the notion of asymptotic resonant energy.

**Notation 2.** Denote, for  $i = 1, \dots, N$ , by  $\sigma_i$  the set of the eigenvalues of the Hamiltonian  $-\Delta - w_i$  on the real line

$$\sigma_i := \{e_k^i\}_{k \in K_i} \subset (-\infty, 0), \quad K_i \subset \mathbb{N}, \quad i = 1, \dots, N. \tag{1.31}$$

**Definition 1.5.** We will say that  $\lambda \in \mathbb{R}$  is an asymptotic resonant energy for the potential  $\mathcal{V}^h$  if and only if

$$\lambda \in \mathcal{E}_0 := \bigcup_{i=1}^N \mathcal{E}_i, \quad \mathcal{E}_i := \sigma_i + \tilde{\mathcal{V}}^0(c_i). \tag{1.32}$$

Moreover, we define the multiplicity  $m_\lambda$  of the asymptotic resonant energy  $\lambda$  as

$$m_\lambda := \#J_\lambda, \quad \text{where } J_\lambda := \{i \in \{1, \dots, N\} \text{ s.t. } \lambda \in \mathcal{E}_i\}. \tag{1.33}$$

Finally, for  $i = 1, \dots, N$ , we will say that the well  $c_i$  is resonant at the energy  $\lambda$  (or  $\lambda$ -resonant) if and only if  $i \in J_\lambda$ .

**Remark 3.** The set  $\sigma_i + \tilde{\mathcal{V}}^0(c_i)$  is nothing but the set of the eigenvalues of the Hamiltonian  $\hat{H}_i^1 := -\Delta - w_i + \tilde{\mathcal{V}}^0(c_i)$  on  $\mathbb{R}$ , which is unitarily equivalent to the Hamiltonian  $\hat{H}_i^h := -h^2 \Delta - w_i(\cdot/h) + \tilde{\mathcal{V}}^0(c_i)$ .

**Theorem 1.6.** Make the Assumptions 1 and 3 and fix a nonnegative function  $\theta \in C_c^0((\Lambda_*, \Lambda^*))$  and assume the convergence of  $\tilde{\mathcal{V}}^h$  stated in Corollary 1.4. Let  $dn_g[V^h]$  be the density defined according to (1.20) and Assumption 2 or by taking  $g(k) = \theta(\lambda_k)$ . Then the weak limit  $dn_g^0$  satisfies

$$dn_g^0|_{(a,b)} = \sum_{\lambda \in \mathcal{E}_0} \sum_{i \in J_\lambda} t_i^\lambda \theta(\lambda) \delta_{x=c_i}, \tag{1.34}$$

with the following specifications:

- (i) In the case of a function of the Hamiltonian, that is  $g(k) = \theta(\lambda_k)$ , all the  $t_i^\lambda$ 's are equal to 1 for every  $\lambda \in \mathcal{E}_0$  and  $i \in J_\lambda$ .
- (ii) If  $g(k) = 1_{k>0} \cdot \theta(\lambda_k)$ , then for every  $\lambda \in \mathcal{E}_0$  and  $i \in J_\lambda$ ,  $t_i^\lambda$  lies in the interval  $[0, 1]$ .

Finally, the asymptotic nonlinear potential  $V_{NL}^0$  which solves (1.29) is an affine function on each interval  $[c_i, c_{i+1}]$ ,  $i = 0, \dots, N$ .

Note that the sum is a finite sum, since the set  $\mathcal{E}_0 \cap \text{supp } \theta$  is finite. Observe immediately that point (ii) follows from (i) because if one denotes

$$\theta_\lambda(k) := \theta(\lambda_k) \tag{1.35}$$



one has  $0 \leq dn_g[V^h] \leq dn_{\theta_\lambda}[V^h]$ , and (ii) is obtained by Theorem 1.3 and Poisson’s equation (1.25). Moreover, the nonlinearity asymptotically lies in a finite dimensional subspace  $\mathcal{A}$  of  $\mathcal{C}^0(I)$ :

$$\mathcal{A} := \{V \in \mathcal{C}^0(I) \text{ s.t. } V|_{\partial I} = 0 \text{ and } V|_{[c_i, c_{i+1}]} \text{ is affine, } i = 0, \dots, N\}. \tag{1.36}$$

In this finite dimensional space, the asymptotic nonlinear system can be written either with the coordinate system  $(V(c_i))_{i=1, \dots, N} \in \mathbb{R}^N$  or with the more convenient one  $(-V'(c_i + 0) + V'(c_i - 0))_{i=1, \dots, N}$  proportional to the collection of total charges in the wells.

Theorem 1.6(i) gives a mean to compute the potential  $V_{NL}^0$  in the particular case where  $g$  is a function of the Hamiltonian. In the anisotropic case (ii) the determination of the  $t_i^\lambda$ ’s relies on a discussion on the Agmon distance between the wells. A forthcoming paper [6] will be dedicated to the analysis of these coefficients.

In order to prove the results, we adopt the following strategy: as the problem is semi-linear, we get *a priori* estimates for the nonlinear potential (Section 2), and reduce the analysis to the linear analysis of the Hamiltonian  $H^h[V^h]$  with uniform estimates on the potential  $(V^h)_{h>0}$ . Useful results on the Dirichlet problem in the interval  $I$  with accurate estimates of the resolvent kernel are reviewed in Section 3. The analysis of resonances starts in Section 4 and Section 5 and ends in Section 6 with a version of the Breit–Wigner formula for the local density of states.

## 2. A priori estimates

We first prove some estimates for self-adjoint realisations of  $P^h$  on  $\Omega = \mathbb{R}$  or  $\Omega$  an open sub-interval of  $I$ .

Consider the differential operator  $P^h$  defined by (1.4), for any  $B \geq 0$  with (1.6)–(1.9), and let  $\tilde{P}^h$  be defined by

$$\tilde{P}^h[V^h] := P^h[V^h] + W^h \equiv -h^2 \frac{d^2}{dx^2} + \tilde{V}^h.$$

**Remark 4.** The  $\tilde{\cdot}$  symbol recurrently refers to the situation where the wells are filled. According to our general convention the letter  $P$  refers to the differential operator while  $H$  refers to some closed realisation as an unbounded operator.

**Proposition 2.1.** Fix a nonnegative smooth function  $\hat{\theta} \in \mathcal{C}_0^\infty(\mathbb{R})$ , and call  $H_\Omega^h$  (resp.  $\tilde{H}_\Omega^h$ ) the self-adjoint realisation on  $L^2(\Omega)$  of  $P^h$  (resp.  $\tilde{P}^h$ ) with domain  $H_0^1(\Omega) \cap H^2(\Omega)$ . Then, for any given compact subset  $K \subset \mathbb{R}$ , and  $h > 0$ , the operators  $\mathbf{1}_K \hat{\theta}(H_\Omega^h) \mathbf{1}_K$  and  $\mathbf{1}_K \hat{\theta}(\tilde{H}_\Omega^h) \mathbf{1}_K$  are trace-class. Moreover the estimate

$$\text{Tr}[\mathbf{1}_K \hat{\theta}(H_\Omega^h) \mathbf{1}_K] - \text{Tr}[\mathbf{1}_K \hat{\theta}(\tilde{H}_\Omega^h) \mathbf{1}_K] \leq C_K (1 + \|\tilde{V}^h\|_{L^\infty})$$

holds with a constant  $C_K$  independent of  $h \in (0, h_0)$ .

**Proof.** In dimension 1 and for any fixed  $h > 0$ , these operators are trace class (see [30]). For the comparison, we use the Dynkin–Helffer–Sjöstrand formula (see [9,18,24]):

$$\hat{\theta}(H_\Omega^h) = \frac{1}{2i\pi} \int_{\mathbb{C}} \frac{\partial \tilde{\theta}}{\partial \bar{z}}(z) (z - H_\Omega^h)^{-1} dz \wedge d\bar{z}, \tag{2.1}$$

where  $\tilde{\theta}$  is a compactly supported almost-analytic extension of  $\hat{\theta}$ . Apply the second resolvent formula for  $z \notin \mathbb{R}$  and write with  $\tilde{P}^h - P^h = W^h$ :

$$\mathbf{1}_K (z - H_\Omega^h)^{-1} \mathbf{1}_K - \mathbf{1}_K (z - \tilde{H}_\Omega^h)^{-1} \mathbf{1}_K = -\mathbf{1}_K (z - H_\Omega^h)^{-1} W^h (z - \tilde{H}_\Omega^h)^{-1} \mathbf{1}_K. \tag{2.2}$$

Introduce a smooth cut-off function  $\chi$ , equal to 1 on a fixed neighbourhood of  $U^h$  if  $\Omega \neq \mathbb{R}$ , and take  $\chi \equiv 1$  if  $\Omega = \mathbb{R}$ . Write the r.h.s. of (2.2)

$$[\mathbf{1}_K (z - H_\Omega^h)^{-1} \chi][W^h (i + h^2 \Delta)^{-1}][(i + h^2 \Delta) \chi (z - \tilde{H}_\Omega^h)^{-1} \mathbf{1}_K], \tag{2.3}$$

where  $-\Delta$  denotes the free Laplacian on  $\mathbb{R}$ . By the spectral theorem, the first factor of (2.3) is a bounded operator with norm  $\mathcal{O}(|\text{Im}(z)|^{-1})$  uniformly w.r.t.  $h > 0$ . Since the operator  $[W^h (i + h^2 \Delta)^{-1}]$  is unitarily equivalent to  $W^{h=1} (i + \Delta)^{-1}$ , it is trace class uniformly with respect to  $h, z$ .

For the last factor, the decomposition

$$(i + h^2 \Delta)\chi(z - \tilde{H}_\Omega^h)^{-1} = (i + h^2 \Delta)\chi(i + h^2 \Delta_\Omega)^{-1} [1 + (i - z + \tilde{V}^h)(z - \tilde{H}_\Omega^h)^{-1}],$$

leads to

$$\|(i + h^2 \Delta)\chi(z - \tilde{H}_\Omega^h)^{-1}\| \leq C_K \frac{\langle z \rangle}{|\text{Im}(z)|} (1 + \|\tilde{V}^h\|_{L^\infty}). \quad \square$$

Proposition 2.1 says that the quantum wells can be forgotten for a uniform global estimate of the density of states. Thanks to a monotony principle shown in [25], one can prove the following result:

**Proposition 2.2.** *Consider the Schrödinger–Poisson system (1.20)–(1.25). Then the family of potentials  $(V_{NL}^h)_{h>0}$  is uniformly bounded in  $L^\infty$ .*

**Proof.** For a given function  $F$ , we will denote by  $F_\lambda$  the function  $k \mapsto F(\lambda_k)$  (see (1.12) for the definition of  $\lambda_k$ ). By assumption on the shape of the incoming beam of electrons, one has:

$$0 \leq g(k) \leq \theta_\lambda(k), \tag{2.4}$$

so we will first study the density of particles corresponding to the equilibrium state described by  $\theta_\lambda$ , that is the measure  $dn_{\theta_\lambda}[V^h]$ . The proof consists in controlling the total mass of the measures by similar quantities relative to other Hamiltonians. In dimension 1, the regularity provided by the Poisson equation with bounded measure as a right-hand side allows the integration by parts

$$\frac{1}{2} \int_a^b \left( \frac{dV_{NL}^h}{dx} \right)^2 dx = \int_a^b V_{NL}^h dn_g[V^h](x) \leq \int_a^b V_{NL}^h dn_{\theta_\lambda}[V^h](x). \tag{2.5}$$

Choose a nonnegative smooth compactly supported function  $\hat{\theta} \in C_0^\infty(\mathbb{R})$  decreasing over  $(-B, \Lambda^*)$  and with support included in  $(-\infty, \Lambda^*)$  such that

$$0 \leq \theta \leq \hat{\theta}. \tag{2.6}$$

We get by positivity of  $V_{NL}^h$  and the expression of the measure in (1.20)

$$\frac{1}{2} \int_a^b \left( \frac{dV_{NL}^h}{dx} \right)^2 dx \leq \int_a^b V_{NL}^h dn_{\theta_\lambda}[V^h](x) \leq \int_a^b V_{NL}^h dn_{\hat{\theta}_\lambda}[V^h](x). \tag{2.7}$$

Set

$$V_2^h := V^h - V_{NL}^h \equiv \tilde{V}_0 - W^h, \tag{2.8}$$

and consider the Hamiltonian  $H_2^h := H_B^h + V_2^h$ . Apply the monotony principle (see Appendix B) with  $H_1 = H_2^h = H_B^h + V_2^h$  and  $H_2 = H_B^h + V^h$ : the last term of (2.7) is bounded by

$$\begin{aligned} \int_a^b V_{NL}^h dn_{\hat{\theta}_\lambda}[V^h](x) &\leq \int_a^b V_{NL}^h dn_{\hat{\theta}_\lambda}[V_2^h](x) \\ &\leq \|V_{NL}^h\|_{L^\infty(I)} \int_a^b dn_{\hat{\theta}_\lambda}[V_2^h](x). \end{aligned} \tag{2.9}$$

Applying Proposition 2.1 gives, coming back to (2.8)

$$\int_a^b dn_{\hat{\theta}_\lambda}[\tilde{V}_0 - W^h](x) \leq C + \int_a^b dn_{\hat{\theta}_\lambda}[\tilde{V}_0](x), \tag{2.10}$$

the constant  $C$  being independent of  $h$  since the potential  $\tilde{V}_0$  does not depend on  $h$ . Finally, we need an upper bound for the density of particles in the island  $I$  in the case of the potential  $\tilde{V}_0 + \mathcal{B}$ . For this, we reduce the problem to the case of the constant potential on  $I$  and equal to  $\Lambda^*$ . Apply again the monotony principle with  $H_1 = H_B^h - \mathcal{B} + \Lambda^*$  and  $H_2 = H_B^h + \tilde{V}_0$ . Since  $H_2 - H_1 = \tilde{V}_0 + \mathcal{B} - \Lambda^* =: \delta V$  is larger than  $\Lambda_0 - \Lambda^* > 0$  according to (1.8), one has uniformly on  $I$

$$\delta V(x) > \inf_I(\tilde{V}_0 + \mathcal{B}) - \Lambda^* \geq \Lambda_0 - \Lambda^* =: \alpha > 0, \quad \text{and} \quad \delta V(x) \leq \|\tilde{V}_0\|_{L^\infty}. \tag{2.11}$$

By writing  $dn_{\hat{\theta}_\lambda}^*$  for the measure  $dn_{\hat{\theta}_\lambda}[\Lambda^* - \mathcal{B}_I]$ , the inequality

$$\alpha \int_a^b dn_{\hat{\theta}_\lambda}[\tilde{V}_0] \leq \int_a^b \delta V \cdot dn_{\hat{\theta}_\lambda}[\tilde{V}_0] \leq \int_a^b \delta V \cdot dn_{\hat{\theta}_\lambda}^* \leq \|\tilde{V}_0\|_{L^\infty} \int_a^b dn_{\hat{\theta}_\lambda}^*,$$

implies

$$0 \leq \int_a^b dn_{\hat{\theta}_\lambda}[\tilde{V}_0] \leq \frac{\|\tilde{V}_0\|_{L^\infty}}{\alpha} \int_a^b dn_{\hat{\theta}_\lambda}^*. \tag{2.12}$$

Since  $\int_a^b dn_{\hat{\theta}}^*$  is a constant not depending on  $h$  (see Appendix D for explicit formulas), we get, combining (2.7), (2.10) and (2.12)

$$\frac{1}{2} \|V_{NL}^h\|_{H_0^1}^2 \leq \left( C + \frac{\|\tilde{V}_0\|_{L^\infty}}{\alpha} \int_a^b dn_{\hat{\theta}}^* \right) \|V_{NL}^h\|_{L^\infty}. \tag{2.13}$$

We conclude with the standard embedding of  $H_0^1$  in  $L^\infty$ .  $\square$

Theorem 1.3 gathers the results of Proposition 2.2 with the next result.

**Proposition 2.3.** *The family of measures  $(dn_g[V^h])_h$  is uniformly bounded in  $\mathcal{M}_b(I)$ . It follows that the family of potentials  $(V_{NL}^h)$  is bounded in  $BV_0^2(I)$ . In particular it is a relatively compact family in every Hölder space  $C^{0,\alpha}(I)$ ,  $\alpha \in (0, 1)$ .*

**Proof.** By definition of  $dn_{\theta_\lambda}$  and simple comparison, one gets

$$\int_I dn_g[V^h] \leq \int_I dn_{\theta_\lambda}[V^h] = \text{Tr}[\mathbf{1}_I \theta(H^h) \mathbf{1}_I] \leq \text{Tr}[\mathbf{1}_I \hat{\theta}(H^h) \mathbf{1}_I].$$

Apply again Proposition 2.1, since the family of potentials is uniformly bounded in  $L^\infty$ . Again the uniform boundedness of the right-hand side with respect to  $h > 0$  comes from (2.9), (2.10), (2.12) and Appendix D.  $\square$

### 3. Results on the Dirichlet problem

From now on, we systematically make Assumption 3 and reduce the analysis to a linear analysis of  $H^h[V^h]$ .

For the contribution of the resonances in the evaluation of spectral quantities, the idea consists in considering the non self-adjoint boundary value problem with complex coefficients in the boundary conditions (1.17)–(1.18) as a perturbation of the homogeneous Dirichlet problem.

#### 3.1. Some notations

In order to measure the error, we shall use several standard tools:

1) The  $h$ -dependent  $H^s$ -norms:

$$\|u\|_{s,h}^2 := \sum_{k \leq s} \|h^k \partial_x^k u\|_{L^2(I)}^2, \quad (u \in H^s(I)) \tag{3.1}$$

will be used mainly with  $s = 0, 1, 2$ .

2) The Agmon distance is defined for any potential  $V \in L^\infty(I)$  according to

**Definition 3.1.** For an energy  $\lambda \in \mathbb{R}$  and a potential  $V$ , we define the Agmon distance by:

$$\forall x, y \in I, \quad d(x, y; V, \lambda) = \left| \int_x^y \sqrt{(V(t) - \lambda)_+} dt \right|. \tag{3.2}$$

For our estimates, we should take  $V = \mathcal{V}^h$ . Yet, it is equivalent to work with the distance relative to the potential  $\tilde{\mathcal{V}}^h$  since the support of  $W^h$  is included in a finite union of intervals with diameter  $2\kappa h$ .

Moreover owing to the lower bound

$$\forall \lambda \in [\Lambda_*, \Lambda^*], \quad \forall x \in I, \quad \inf_{h>0, x \in I} \tilde{\mathcal{V}}^h(x) - \lambda \geq \Lambda_0 - \Lambda^* =: \delta > 0, \tag{3.3}$$

all the Agmon distances (depending on  $\tilde{\mathcal{V}}^h$ ) are uniformly equivalent to the usual Euclidean distance.

3) Finally in the analysis of the tunnel effect, it is usual to introduce the estimates within the next setting.

**Definition 3.2.** For an  $h$ -dependent vector  $f(h)$  in a normed space  $E$  with norm  $\|\cdot\|_E$  and a positive real valued function  $g(h)$ , we write

$$f(h) = \tilde{\mathcal{O}}(g(h)) \quad (\text{as } h \rightarrow 0) \tag{3.4}$$

if there exists  $\eta_0 > 0$  such that

$$\forall \eta \in (0, \eta_0), \quad \exists C_\eta > 0, \quad \forall h \in (0, h_0), \quad \|f(h)\|_E \leq C_\eta e^{\frac{\eta}{h}} g(h).$$

### 3.2. Decay estimate

Like in Proposition 2.1,  $\Omega$  denotes an open interval in  $I$  and  $H_\Omega^h$  the self-adjoint Dirichlet realisation of  $P^h[V^h]$  with domain  $H_0^1(\Omega) \cap H^2(\Omega)$ .

We shall use the following result about the decay of the eigenfunctions of  $H_\Omega^h$ .

**Proposition 3.3.** Suppose that  $U_\Omega := \{c_1, \dots, c_N\} \cap \Omega$  is not empty. For every  $h > 0$  sufficiently small, let  $\lambda^h \in (\Lambda_*, \Lambda^*)$  be an eigenvalue of  $H_\Omega^h$  and  $\phi^h$  an  $L^2$ -normalised corresponding eigenfunction:

$$(H_\Omega^h - \lambda^h)\phi^h = 0.$$

Then, the estimate

$$\forall x \in \Omega, \quad \left| \frac{d^j}{dx^j} \phi^h(x) \right| \leq Ch^{-2j-1} e^{-\frac{\tilde{d}_h(x, U_\Omega)}{h}}, \quad j \in \{0, 1\},$$

holds with  $C > 0$  uniform w.r.t.  $h \in (0, h_0)$  if  $\tilde{d}_h$  stands for the Agmon distance for the potential  $\tilde{\mathcal{V}}^h$  at the energy  $\lambda^h$ .

**Remark 5.** Note that contrary to the general use, we do not introduce at this level the  $\tilde{\mathcal{O}}$  but an accurate estimate made possible in this simple one-dimensional case. This accurate estimate will be combined in the proof of Theorem 3.4 with the uniform Lipschitz estimate on  $\tilde{\mathcal{V}}^h$  (see especially (3.11)–(3.13)). This provides a complete splitting between the semiclassical and quantum scale in spite of a limited regularity assumption.

**Proof.** Set  $\Omega = [\alpha, \beta]$ .

1) Let us begin with the estimate of  $\phi^h(x)$ .

Apply the Agmon identity of Appendix A with  $P = P^h$ ,  $z = \lambda^h$ ,  $u_1 = u_2 = \phi^h$  and  $\varphi(x) = \tilde{d}_h(x, U_\Omega)$  where  $\phi^h$  is an eigenfunction of  $H_\Omega^h$  with eigenvalue  $\lambda^h$ . Since  $\mathcal{V}^h - \lambda^h - \varphi'^2 = -W^h$ , the inequalities  $\varphi = \mathcal{O}(h)$  in  $U^h$  and  $\|\phi^h\|_{L^2} = 1$  imply

$$e^{\pm \frac{\varphi}{h}} = \mathcal{O}(1) \quad \text{in } U^h \quad \text{and} \quad \int (\mathcal{V}^h - \lambda^h - \varphi'^2) |v^h|^2 = \mathcal{O}(1).$$

From the Agmon identity, we deduce an estimate for  $v^h = e^{\varphi/h} \phi^h$ :

$$\left\| h \frac{dv^h}{dx} \right\|_{L^2} = \mathcal{O}(1).$$

Since  $v^h(\alpha) = v^h(\beta) = 0$ , it follows

$$\|v^h\|_{L^2} + \left\| \frac{dv^h}{dx} \right\|_{L^2} = \mathcal{O}\left(\frac{1}{h}\right).$$

This implies

$$\|v^h\|_{L^\infty} = \mathcal{O}\left(\frac{1}{h}\right),$$

and therefore

$$\forall x \in \Omega, \quad |\phi^h(x)| \leq \frac{C}{h} e^{-\tilde{d}_h(x, U_\Omega)}.$$

2) For the estimate of  $d\phi^h/dx$ , we use the equation

$$\begin{cases} -h^2 \frac{d^2 \phi^h}{dx^2} + \mathcal{V}^h \phi^h = \lambda^h \phi^h, \\ \phi^h(\alpha) = \phi^h(\beta) = 0. \end{cases}$$

As  $\phi^h \in C^1([\alpha, \beta])$ , there exists  $c \in (\alpha, \beta)$  such that  $\frac{d\phi^h}{dx}(c) = 0$ . The function  $g$  defined by  $g = e^{\varphi/h} d\phi^h/dx$  satisfies

$$\begin{cases} h^2 g' = h\varphi' e^{\frac{\varphi}{h}} \frac{d\phi^h}{dx} + h^2 e^{\frac{\varphi}{h}} \frac{d^2 \phi^h}{dx^2}, \\ h^2 g(c) = 0. \end{cases}$$

Using the equation satisfied by  $\phi^h$ , we deduce

$$\begin{aligned} h^2 g' &= h\varphi' (e^{\frac{\varphi}{h}} \phi^h)' - |\varphi'|^2 e^{\frac{\varphi}{h}} \phi^h + (\mathcal{V}^h - \lambda^h) e^{\frac{\varphi}{h}} \phi^h \\ &= h\varphi' \frac{dv^h}{dx} - |\varphi'|^2 v^h + (\mathcal{V}^h - \lambda^h) v^h. \end{aligned}$$

Then  $\|h^2 g'\|_{L^2} = \mathcal{O}(1/h)$ . Cauchy–Schwarz inequality gives the  $L^\infty$ -estimate for  $g : |g(x)| \leq C/h^3$  for any  $x \in [\alpha, \beta]$  and also of  $d\phi^h/dx$ :

$$\forall x \in \Omega = [\alpha, \beta], \quad \left| \frac{d\phi^h}{dx}(x) \right| \leq \frac{C}{h^3} e^{-\tilde{d}_h(x, U_\Omega)}. \quad \square$$

**Remark.** When the potential is regular, a better estimate like

$$\forall x \in \Omega, \quad |\phi^h(x)| \leq Ch^{-\frac{1}{2}} e^{-\tilde{d}_h(x, U_\Omega)/h},$$

holds and even a complete WKB expansion is possible. Here the low regularity and the concentration of the quantum wells prevent from such an accurate result.

### 3.3. Spectrum for one single well

From the spectral viewpoint, we are interested in localising the eigenvalues of  $H_\Omega^h$  in the limit  $h \rightarrow 0$ . The first result concerns the problem with one well.

**Theorem 3.4.** *Let  $\Omega$  be a sub-interval of  $(a, b)$  containing exactly one well  $c_i$ ,  $i \in \{1, \dots, N\}$ . Then:*

- (i) *Every eigenvalue of  $H_\Omega^h$  in  $(\Lambda_*, \Lambda^*)$  converges, and the limit belongs to the set  $\mathcal{E}_i$  (see (1.32)).*
- (ii) *For every  $\lambda_0 \in (\Lambda_*, \Lambda^*) \cap \mathcal{E}_i$  and any fixed small enough  $\varepsilon > 0$ , the Dirichlet Hamiltonian  $H_\Omega^h$  has exactly one eigenvalue in  $[\lambda_0 - \varepsilon, \lambda_0 + \varepsilon]$  for  $h \in (0, h_\varepsilon)$ .*

**Proof.** Call  $\{\lambda_1^h, \dots, \lambda_r^h\}$  the eigenvalues of  $H_\Omega^h$  in the interval  $[\Lambda_*, \Lambda^*]$ , and  $\phi_1^h, \dots, \phi_r^h$  an orthonormal system of corresponding eigenfunctions. Because of Proposition 2.1, since the rank of the spectral projections are given by traces of functions of  $H_\Omega^h$  one has:

$$r = \mathcal{O}(1), \quad h \rightarrow 0$$


(take for  $\theta$  a smooth version of the function  $\mathbf{1}_{[\varepsilon, \Lambda_0]}$ ,  $\varepsilon > 0$  small). The idea is to use the ellipticity of the problem, and the scaling of the wells in order to replace the potential  $\tilde{\mathcal{V}}^h$  near a well by a constant one. Let  $\hat{H}^h$  be the Hamiltonian with domain  $H^2(\mathbb{R})$  given by:

$$\forall u \in D(\hat{H}^h), \quad \hat{H}^h u := \hat{P}^h u, \quad \hat{P}^h := -h^2 \frac{d^2}{dx^2} + \tilde{\mathcal{V}}^h(c_i) \cdot \mathbf{1} - w_i \left( \frac{x - c_i}{h} \right). \tag{3.5}$$

This Hamiltonian is unitarily equivalent to  $-\Delta + \tilde{\mathcal{V}}^h(c_i) - w_i(\cdot - c_i)$ , whose eigenvalues are the set

$$\mathcal{E}_i^h := \mathcal{E}_i + \alpha_i^h, \quad \alpha_i^h = \tilde{\mathcal{V}}^h(c_i) - \tilde{\mathcal{V}}^0(c_i) \rightarrow 0, \quad h \rightarrow 0. \tag{3.6}$$

Since  $\|\tilde{\mathcal{V}}^h - \tilde{\mathcal{V}}^0\|_{C^0} \rightarrow 0$  when  $h \rightarrow 0$ , for any  $\lambda_0 \in [\Lambda_*, \Lambda^*] \cap \mathcal{E}_i$  there exists  $\varepsilon_0 > 0$  such that  $\hat{H}^h$  has exactly one eigenvalue in  $(\lambda_0 - \varepsilon_0, \lambda_0 + \varepsilon_0)$ . To analyse the spectrum of  $H^h$  in the whole set  $[\Lambda_*, \Lambda^*]$ , we choose, for each  $\lambda_0$ , two numbers  $\varepsilon_0^+ > 0$ ,  $\varepsilon_0^- > 0$  such that the intervals  $(\lambda_0 - \varepsilon_0^-, \lambda_0 + \varepsilon_0^+)$  are disjoint and their union covers a compact neighbourhood of  $[\Lambda_*, \Lambda^*]$  and such that  $\hat{H}^h$  has no eigenvalue in each annulus  $\{\varepsilon_0 < |\lambda - \lambda_0| < 2 \min\{\varepsilon_0^+, \varepsilon_0^-\}\}$ .



Fix  $\eta > 0$  and let  $\chi$  be a smooth cut-off function supported in  $\Omega$  such that  $\chi = 1$  if  $d(x, \partial\Omega) \geq 2\eta$  and  $\chi = 0$  if  $d(x, \partial\Omega) \leq \eta$ . Owing to the exponential decay of the  $\phi_j^h$ 's stated in Proposition 3.3, the estimate

$$\langle \chi \phi_j^h, \chi \phi_k^h \rangle_{L^2(\Omega)} = \delta_{jk} + \mathcal{O}(e^{-\frac{c_0}{h}}), \quad j, k \in \{1, \dots, r\}, \tag{3.7}$$

holds for some  $c_0 > 0$  independent on  $h > 0$  and  $\eta > 0$ .

For any  $j \in \{1, \dots, r\}$ , the function  $\chi \phi_j^h$  belongs to the domain of  $\hat{H}^h$  with the identity

$$\hat{P}^h \chi \phi_j^h = \lambda_j^h \chi \phi_j^h + [P^h, \chi] \phi_j^h + (\tilde{\mathcal{V}}^h(c_i) - \tilde{\mathcal{V}}^h(x)) \chi \phi_j^h. \tag{3.8}$$

Owing to the exponential decay of  $\phi_j^h$ , the commutator term satisfies:

$$\|[P^h, \chi] \phi_j^h\|_{L^2(\Omega)} = \mathcal{O}(h^{-1} e^{-\frac{\tilde{d}_h(c_i, \partial\Omega) - 2\eta}{h}}), \tag{3.9}$$

where  $\tilde{d}_h$  is the Agmon distance for  $\tilde{\mathcal{V}}^h$  at the energy  $\lambda_i^h$ . Because the potential  $\tilde{\mathcal{V}}^h$  is greater than  $\Lambda_0$  and  $\lambda_i^h \leq \Lambda^* < \Lambda_0$ , the r.h.s. in (3.9) is of order  $\mathcal{O}(e^{-c'/h})$  with  $c'$  independent of the potential and the energy.

For the last term of the r.h.s. of (3.8), just write for  $\varepsilon > 0$

$$[\tilde{\mathcal{V}}^h(c_i) - \tilde{\mathcal{V}}^h(x)]\chi\phi_j^h = \mathbf{1}_{|x-c_i|\leq\varepsilon} \cdot [\tilde{\mathcal{V}}^h(c_i) - \tilde{\mathcal{V}}^h(x)]\chi\phi_j^h + \mathbf{1}_{|x-c_i|>\varepsilon} \cdot [\tilde{\mathcal{V}}^h(c_i) - \tilde{\mathcal{V}}^h(x)]\chi\phi_j^h. \tag{3.10}$$

Since the family of potentials  $(\tilde{\mathcal{V}}^h)_{h>0}$  is  $W^{1,\infty}(I)$ -bounded, the first term is treated by writing

$$\|\mathbf{1}_{|x-c_i|\leq\varepsilon} \cdot [\tilde{\mathcal{V}}^h(c_i) - \tilde{\mathcal{V}}^h(x)]\chi\phi_j^h\|_{L^2(\Omega)} \leq \varepsilon \sup\|\tilde{\mathcal{V}}^h\|_{W^{1,\infty}}\|\chi\phi_j^h\|_{L^2(\Omega)} = \mathcal{O}(\varepsilon), \tag{3.11}$$

and again by the accurate decay estimates of Proposition 3.3, the second term is estimated by

$$\|\mathbf{1}_{|x-c_i|>\varepsilon} \cdot (\tilde{\mathcal{V}}^h(c_i) - \tilde{\mathcal{V}}^h(x))\chi\phi_j^h\|_{L^2(\Omega)} = \mathcal{O}(e^{-\frac{c'_0\varepsilon}{h}}). \tag{3.12}$$

We choose

$$\varepsilon := h^\alpha, \quad \alpha \in (0, 1), \tag{3.13}$$

and we obtain by combining (3.12), (3.11), (3.9), (3.8)

$$\forall j = 1, \dots, r, \quad \hat{P}^h \chi\phi_j^h = \lambda_j^h \chi\phi_j^h + \mathcal{O}(h^\alpha) \quad \text{in } L^2(\Omega). \tag{3.14}$$

Fix  $\delta > 0$  such that  $\hat{H}^h$  has no eigenvalue in  $\{\varepsilon_0^+ < \lambda - \lambda_0 < \varepsilon_0^+ + \delta\} \cup \{-\varepsilon_0^- - \delta < \lambda - \lambda_0 < -\varepsilon_0^-\}$  and apply Proposition C.1 (see Appendix C) to  $A = \hat{H}^h$ ,  $[\lambda_-, \lambda^+] = [\lambda_0 - \varepsilon_0^-, \lambda_0 + \varepsilon_0^+]$ ,  $N = r$ ,  $a = \delta > 0$ ,  $\mu_j = \lambda_j^h$ ,  $\psi_j = \chi\phi_j^h$ , from which we conclude

$$\tilde{d}(\text{span}\{\chi\phi_1^h, \dots, \chi\phi_r^h\}, \mathbf{1}_{[\lambda_0 - \varepsilon_0^-, \lambda_0 + \varepsilon_0^+]}) (\hat{H}^h) \leq \left(\frac{r}{1 + o(1)}\right)^{1/2} \frac{\varepsilon}{a} = \mathcal{O}(h^\alpha). \tag{3.15}$$

This last estimate forces  $H^h$  to have at most one eigenvalue in  $[\lambda_0 - \varepsilon_0^-, \lambda_0 + \varepsilon_0^+]$ ,  $r \leq 1$ , when  $h > 0$  is small enough.

We finish by proving (i) and (ii). Notice that the asymptotic value  $\lambda_0$  is exactly the eigenvalue of the Hamiltonian  $-h^2 d^2/dx^2 + \tilde{\mathcal{V}}^0(c_i) - w_i((\cdot - c_i)/h)$ , unitarily equivalent to the  $h$ -independent operator  $-d^2/dx^2 + \tilde{\mathcal{V}}^0(c_i) - w_i$ , and let  $\hat{\phi}_0^h$  denote a corresponding normalised eigenvector. Such a wave function  $\hat{\phi}_0^h$  is an eigenvector of  $\hat{H}^h$  for the eigenvalue  $\lambda_0 + \alpha_i^h$  (see (3.6)). Estimates similar to (3.9), (3.11), (3.12) lead to

$$P^h \hat{\phi}_0^h = (\lambda_0 + \alpha_i^h)\chi\hat{\phi}_0^h + \mathcal{O}(h^\alpha) \quad \text{in } L^2(\Omega). \tag{3.16}$$

Apply again Proposition C.1 in a small interval centred around  $\lambda_0 + \alpha_i^h$  in the following way: since  $\hat{H}^h$  has at most one eigenvalue in  $[\lambda_0 - \varepsilon_0^-, \lambda_0 + \varepsilon_0^+]$ , it is easy to choose a convenient parameter  $a$  in Proposition C.1 (Appendix C) by a simple argument of counting: set  $L_j := [jh^{\alpha/2}, (j+1)h^{\alpha/2}[$ , and  $K_j := -L_j \cup L_j$ . If  $\{\lambda_0 + \alpha_i^h\} + K_1$  contains the eigenvalue, one defines  $I_h = [\lambda_0 + \alpha_i^h - 2h^{\alpha/2}, \lambda_0 + \alpha_i^h + 2h^{\alpha/2}]$ , else  $I_h = [\lambda_0 + \alpha_i^h - h^{\alpha/2}, \lambda_0 + \alpha_i^h + h^{\alpha/2}]$ . This furnishes an interval  $I_h$  of diameter  $\mathcal{O}(h^{\alpha/2})$  around  $\lambda_0 + \alpha_i^h$  and a real  $a = a(h) > 0$  of order  $h^{\alpha/2}$  leading again with Proposition C.1 to

$$\tilde{d}(\text{span}(\chi\hat{\phi}_0^h), \mathbf{1}_{I_h}(H_\Omega^h)) = \mathcal{O}(h^{\alpha/2}). \tag{3.17}$$

This yields  $r = 1$  and the convergence of the eigenvalue to  $\lambda_0$ .  $\square$

**Remark 6.** It follows that the well  $c_i$  is  $\lambda$ -resonant if and only if there exists a domain  $\Omega$  containing  $c_i$  such that for any open set  $\omega \subset \Omega$  the Dirichlet operator  $H_\omega^h$  has an eigenvalue converging to  $\lambda$  as  $h$  goes to 0.

### 3.4. Spectrum in the multiple wells case

A way of studying the spectral properties of the multiple wells Dirichlet problem consists in decoupling it into  $N$  one-well problems. Following [14] or [17], a good choice of open sets is the following: fix  $\lambda \in [\Lambda_*, \Lambda^*]$ , and if  $\tilde{d}_h$  (resp.  $\tilde{d}_0$ ) denotes the Agmon distance at the energy  $\lambda$  for the potential  $\tilde{\mathcal{V}}^h$  (resp.  $\tilde{\mathcal{V}}^0$ ), we define

$$S_1 := \min_{j \neq k} \tilde{d}_h(c_j, c_k) \quad (= S_1(h)), \tag{3.18}$$

and for a fixed small enough  $\eta > 0$ ,

$$\Omega_i := I \setminus \bigcup_{k \neq i} \{x \in I, \tilde{d}_0(x, c_k) \leq \eta\}, \quad i = 1, \dots, N. \tag{3.19}$$

The  $h$ -dependence of  $S_1$  recalled between the parentheses of (3.18) is omitted in the sequel.

Note that these open sets are not disjoint and  $\Omega_i$  contains only the well  $c_i$ . The use of the distance  $\tilde{d}_0$  makes sure that they do not depend on  $h$  although the  $h$ -dependence would be well controlled.

We first eliminate the nonresonant wells before giving a result similar to Theorem 3.4.

**Proposition 3.5.** *Let  $\lambda$  be an asymptotic resonant energy and suppose that the well  $c_i$  is not  $\lambda$ -resonant. Then there exists a positive constant  $c$  such that for any eigenvalue  $\lambda^h \in (\lambda - c, \lambda + c)$ , one has*

$$\forall x \in (c_i - c, c_i + c), \quad |\phi^h(x)| \leq e^{-\frac{c}{h}}, \quad h \rightarrow 0$$

where  $\phi^h$  is an  $L^2$ -normalised eigenfunction of  $H_I^h$  for the eigenvalue  $\lambda^h$ .

In plain words, eigenfunctions for eigenvalues converging to  $\lambda$  are exponentially small in the non  $\lambda$ -resonant wells.

**Proof.** Since  $\lambda$  is not a resonant energy for the well  $c_i$ , we can choose the open set  $\omega$  containing the only well  $c_i$  and the compact energy interval  $\Lambda \ni \lambda$  such that for  $h > 0$  sufficiently small, the Dirichlet operator  $H_\omega^h$  has no spectrum in  $\Lambda$  (see Remark 6). For a smooth cut-off function  $\theta$  supported in  $\omega$  and equal to 1 on a  $\delta$ -neighbourhood of  $c_i$  ( $\delta > 0$  small), one has

$$P^h \theta \phi^h = \lambda^h \theta \phi^h + [P^h, \theta] \phi^h. \tag{3.20}$$

The residual term satisfies by Proposition 3.3 the decay estimate

$$\|[P^h, \theta] \phi^h\|_{L^2(I)} \leq C_\delta e^{-\frac{c_\delta}{h}}, \quad c_\delta > 0, \quad h \rightarrow 0.$$

Note that the vector  $\theta \phi^h$  is not zero.

Apply again Proposition C.1 in a compact interval strictly contained in  $\Lambda$  and  $a > 0$  not depending on  $h > 0$ . If we denote by  $F$  the spectral subspace for  $H_\omega^h$  associated to this compact interval, it follows

$$\tilde{d}(\text{span}\{\theta \phi^h\}, F) \leq \frac{1}{\|\theta \phi^h\|} \frac{C_\delta e^{-\frac{c_\delta}{h}}}{a}. \tag{3.21}$$

Since  $F$  is null by choice of  $\Lambda$ , it follows by properties of the distance  $\tilde{d}$  that the l.h.s. of (3.21) is greater than 1. This provides an  $L^2$ -estimate of  $\theta \phi^h$ . The  $H^2$  regularity of a solution to (3.20) provides the pointwise estimate in  $(c_i - \delta, c_i + \delta)$ . Finally choose the constant  $c > 0$  small enough.  $\square$

The analogous to Theorem 3.4 writes

**Theorem 3.6.** *Recall that  $H_\omega^h$  denotes the Dirichlet realisation of  $P^h$  to the open set  $\omega$ . Then, for  $h > 0$  sufficiently small:*

- (i) *After ordering, every eigenvalue of  $H_I^h$  in  $(\Lambda_*, \Lambda^*)$  converges as  $h \rightarrow 0$  and the limit belongs to the set  $\mathcal{E}_0$  (see (1.32)).*
- (ii) *For every  $\lambda \in (\Lambda_*, \Lambda^*) \cap \mathcal{E}_0$  and any small enough  $\varepsilon > 0$ , the operators  $H_I^h$  has exactly  $m_\lambda$  eigenvalue(s) in  $[\lambda - \varepsilon, \lambda + \varepsilon]$  as soon as  $h < h_\varepsilon$ .  
Call them  $\lambda_i^h$  ( $i \in J_\lambda$ ).*
- (iii) *Fix such a  $\lambda$ . Let  $(\Omega_i)_{i \in J_\lambda}$  be the subdomains of  $I$  defined in (3.19). Call  $(\psi_i^h)_{i \in J_\lambda}$  normalised eigenvectors associated to the unique eigenvalue of  $H_{\Omega_i}^h$  converging to  $\lambda$ . There exists a unitary matrix  $(p_{i,j}^h)_{1 \leq i, j \leq m_\lambda}$  such that in  $L^2(I)$*

$$\forall i \in J_\lambda, \quad \phi_i^h - \sum_{j \in J_\lambda} p_{i,j}^h \psi_j^h = \tilde{O}(e^{-\frac{S_1}{h}}),$$

with  $S_1$  defined according to (3.18).



**Proof.** It suffices to follow the proof in [14, pp. 34–35], while Proposition 3.5 guarantees that the nonresonant wells are negligible in the decay estimates (see also [28, p. 148] for details). □

### 3.5. Resolvent estimates

Let us briefly recall the decay results of the kernel of the resolvents. Fix  $\eta > 0$  ( $\eta$  small) and for a point  $p \in (a, b)$ , let  $\chi_p$  denote a smooth cut-off function supported in the set  $\{|x - p| \leq \eta\}$ .

Like in [17, p. 143] (see also [12] or [28, p. 135] for this specific case), the combination of the Agmon estimate (see Appendix A) with the spectral theorem provides in the one well-case ( $N = 1$ ) the following estimates

$$\forall z \notin \sigma(H_I^h), \quad \|\chi_x(H_I^h - z)^{-1}\chi_y\| \leq C_\eta \frac{e^{-\frac{\tilde{d}_h(x,y)+C\eta}{h}}}{\min(r_h, 1)}, \tag{3.22}$$

where  $r_h = \text{dist}(z, \sigma(H_I^h))$ , and  $\tilde{d}_h$  is the Agmon distance for the potential  $\tilde{V}^h$  at the energy  $\lambda := \text{Re}(z)$ .

A straightforward adaptation of the analysis of the multiple wells Dirichlet problem carried out in [16], [17, p. 147] or [28, p. 151] provides the same estimate for  $N > 1$ .

**Proposition 3.7.** *For  $h$  in  $(0, h_0)$ ,  $h_0$  small enough, consider  $z_h \in \mathbb{C} \setminus \sigma(H_I^h)$  such that there exists  $\lambda_0 \in [\Lambda_*, \Lambda^*]$  with  $z_h \rightarrow \lambda_0$  as  $h \rightarrow 0$  and set  $\lambda_h = \text{Re}(z_h)$  and  $r_h = \text{dist}(z_h, \sigma(H_I^h))$ . If  $r_h \geq e^{-S_1/2h}$  with  $S_1 := \min_{k \neq l} \tilde{d}_h(c_k, c_l)$ , then the kernel of the resolvent  $(H_I^h - z_h)^{-1}$  satisfies*

$$|(H_I^h - z_h)^{-1}[x, y]| = \frac{\tilde{O}(e^{-\frac{\tilde{d}_h(x,y)}{h}})}{\min(r_h, 1)},$$

with uniform constants with respect to  $x, y \in I$  and where  $\tilde{d}_h$  is the Agmon distance for the potential  $\tilde{V}^h$  at the energy  $\lambda_h := \text{Re}(z_h)$ .

**Proof.** Let  $\theta$  be a  $C^\infty$  even function supported in a neighbourhood  $[-3\eta, 3\eta]$  and equal to 1 on  $[-\eta, \eta]$  where  $\eta$  and  $\Omega_i$  are linked by relation (3.19). We define

$$\theta_i(x) := \theta(x - c_i), \quad \chi_i(x) = 1 - \sum_{j \neq i} \theta_j(x), \quad \forall i = 1, \dots, N. \tag{3.23}$$

Let  $\tilde{\chi}_i$  be  $C^\infty$  functions with support in  $\Omega_i$  defined in (3.19) such that

$$\sum_{i=1}^N \tilde{\chi}_i = 1.$$

Introduce

$$T_i(z) := (H_{\Omega_i}^h - z)^{-1} \quad \text{and} \quad R_0 := \sum_{i=1}^N \chi_i T_i(z) \tilde{\chi}_i.$$

This leads to

$$\begin{aligned} (H_I^h - z)R_0 &= \sum_{i=1}^N \chi_i \tilde{\chi}_i + \sum_{i=1}^N [P^h, \chi_i] T_i(z) \tilde{\chi}_i \\ &= 1 + \sum_{i=1}^N [P^h, \chi_i] T_i(z) \tilde{\chi}_i \\ &= 1 - \sum_{i=1}^N \sum_{k \neq i} [P^h, \theta_k] T_i(z) \tilde{\chi}_i, \end{aligned}$$

owing to  $\chi_i \tilde{\chi}_i = \tilde{\chi}_i$  and (3.23). We have to study the convergence of the series  $\sum_{n \geq 0} R_0 \varepsilon^n$  with

$$\varepsilon = \sum_{i=1}^N \sum_{k \neq i} [P^h, \theta_k] T_i(z) \tilde{\chi}_i.$$

We notice that  $\tilde{\chi}_i [P^h, \theta_k]$  is equal to 0 as soon as  $k \neq i$  and if  $k = i$ , this term is  $[P^h, \theta_k]$ . Hence

$$R_0 \varepsilon^n = \sum_{i_0=1}^N \sum_{i_1 \neq i_0}^N \dots \sum_{i_{n-1} \neq i_n}^N \chi_{i_0} T_{i_0} [P^h, \theta_{i_1}] T_{i_1} [P^h, \theta_{i_2}] T_{i_2} \dots [P^h, \theta_{i_n}] T_{i_n} \tilde{\chi}_{i_n}.$$

Since the function  $\theta_k$  is localised in a neighbourhood of the well  $c_k$ , we can write for  $s = 0, \dots, N - 1$

$$[P^h, \theta_{i_s}] T_{i_s}(z) [P^h, \theta_{i_{s+1}}] = [P^h, \theta_{i_s}] \chi_{i_s} T_{i_s}(z) \chi_{i_{s+1}} [P^h, \theta_{i_{s+1}}].$$

This last relation allows to use results on the one-well problem (3.22):

$$\| \chi_{i_s} T_{i_s}(z) \chi_{i_{s+1}} \| \leq C_\eta \frac{e^{-\frac{\tilde{d}_h(x,y) - C_\eta}{h}}}{\min(r_h, 1)}.$$

This leads to the following estimate

$$\| \chi_{x_0} R_0 \varepsilon^n \chi_{y_0} \| \leq C_\eta^{n+1} \frac{e^{-\frac{\varphi_n(x_0, y_0) - n C_\eta}{h}}}{\min(r_h, 1)^{n+1}},$$

where  $\varphi_n(x_0, y_0) = \min_{i_0, \dots, i_n} d(y_0, c_{i_n}) + d(c_{i_n}, c_{i_{n-1}}) + \dots + d(c_{i_1}, c_{i_0}) + d(c_{i_0}, x_0)$ . In fact, the function  $\varphi_n$  is the length of the shortest way from  $y$  to  $x$  going through  $n$  different wells. The quantity  $\varphi_n$  is bounded from below according to

$$\varphi_n(x_0, y_0) \geq d(x_0, y_0) + n S_1.$$

Hence the serie is convergent under the assumption  $r_h \geq e^{-S_1/2h}$  and we can write

$$\chi_{x_0} (H_I^h - z)^{-1} \chi_{y_0} = \sum_{n \geq 0} \chi_{x_0} R_0 \varepsilon^n \chi_{y_0}.$$

Appendix E provides the pointwise estimates.  $\square$

**Corollary 3.8.** *If  $r_h \geq C^{-1} h^C$  for some  $C > 0$ , then*

$$|(z - H_I^h)^{-1}[x, y]| = \tilde{O}(e^{-\tilde{d}_h(x,y)}).$$

Another consequence is the improved pointwise estimate for the eigenfunctions of the Dirichlet problem ([17, p.138] or [28, p. 153]):

**Proposition 3.9.** *For every  $h > 0$  sufficiently small, let  $\lambda^h$  belong to  $(\Lambda_*, \Lambda^*)$  and let  $\phi^h$  be an  $L^2$ -normalised corresponding eigenfunction of  $H_\Omega^h$ . Suppose that  $\lambda^h \rightarrow \lambda_0 \in \mathcal{E}_0 \cap (\Lambda_*, \Lambda^*)$ . Then the estimates*

$$\forall x \in \Omega, \quad \left| \frac{d^j}{dx^j} \phi^h(x) \right| = \tilde{O}(e^{-\frac{\tilde{d}_0(x, U_0)}{h}}), \quad j \in \{0, 1\},$$

hold when  $\tilde{d}_0$  stands for the Agmon distance for the potential  $\tilde{V}^0$  at the energy  $\lambda_0$  and  $U_0 = \bigcup_{i \in J_{\lambda_0}} \{c_i\}$  for the set of  $\lambda_0$ -resonant wells.

**Remark 7.** Here the  $\tilde{O}$ -writing of the estimates allows to replace the  $h$ -dependent quantities,  $\tilde{V}^h$ ,  $\tilde{d}_h$  and  $\lambda_h$  by their asymptotic values  $\tilde{V}^0$ ,  $\tilde{d}_0$  and  $\lambda_0$ .

### 4. Complex deformation

#### 4.1. A reduced Stone's formula

The results of Theorem 1.6 are derived from a good information about the asymptotic local density of states associated with functions of the Hamiltonian. According to Stone's formula and the limiting absorption principle, a possible method is the computing of a quite precise expression of the resolvent, since for  $\lambda \in [\Lambda_*, \Lambda^*] \subset \sigma_{ac}(H^h)$  ( $H^h = H^h[V^h]$ ):

$$\frac{1}{2i\pi} \mathbf{1}_I [(H^h - (\lambda + i0))^{-1} - (H^h - (\lambda - i0))^{-1}] \mathbf{1}_I = \mathbf{1}_I \frac{\partial E}{\partial \lambda}(\lambda) \mathbf{1}_I, \tag{4.1}$$

and of its meromorphic extension through the spectral half-line  $(0, \infty) \subset [-B, \infty)$ , in order to take into account the contribution of resonant states.

We will focus on this meromorphic extension from the upper-half plane while the corresponding results for the extension from the lower-half plane are easily carried over after changing  $i$  into  $-i$ .

**Resolvent.** Fix  $z \in \mathbb{C}$ ,  $\text{Im}(z) > 0$  and consider the problem with unknown  $u \in H^2(\mathbb{R})$ :

$$(P^h - z)u = f, \quad f \in L^2(I), \quad z \in \mathbb{C}, \quad \text{Im}(z) > 0, \quad \text{Re}(z) \in (\Lambda_*, \Lambda^*). \tag{4.2}$$

Again because the potential is constant on both sides of the interval  $I$ , the problem with unknown  $u \in H^2(\mathbb{R})$ :

$$(P^h - z)u = f, \quad f \in L^2(I),$$

can be explicitly solved outside  $I$ , and the condition  $u \in L^2$  eliminates exponentially growing modes. It is easy to check that this condition is exactly given by (1.17)–(1.18) when  $\text{Im}(z) > 0$ . Precisely, we can write the next statement.

**Proposition 4.1.** *Let  $z$  belong to  $\mathbb{C}$  with  $\text{Im}(z) > 0$  and  $\text{Re}(z) \in (\Lambda_*, \Lambda^*)$ . Consider the linear functionals  $T_a(z)$ ,  $T_b(z)$  on  $H^2(I)$  given by:*

$$T_a(z)u := [h\partial_x + iz^{1/2}]_{|x=a}u, \quad T_b(z)u := [h\partial_x - i(z + B)^{1/2}]_{|x=b}u,$$

and the closed unbounded operator  $H_z^h$  defined by

$$D(H_z^h) := \{u \in H^2(I) \text{ s.t. } T_a(z)u = T_b(z)u = 0\}, \quad \forall u \in D(H_z^h), \quad H_z^h u := P^h u.$$

Then the restriction on  $I$  of the solution to Eq. (4.2) is  $(H_z^h - z)^{-1}f$ . In other words:

$$\mathbf{1}_I (H^h - z)^{-1} \mathbf{1}_I = (H_z^h - z)^{-1}, \quad \text{Im}(z) > 0, \quad \text{Re}(z) \in (\Lambda_*, \Lambda^*).$$

**Remark 8.**

1. We will check that for such  $z$ 's, operator  $H_z^h - z$  is invertible (see Proposition 4.2 and Proposition 5.2 below).
2. Note that since the solutions on  $I$  of the homogeneous equation associated with (4.2) make a linear 2-dimensional subspace of  $H^2(I)$ , the injectivity of operator  $(H_z^h - z)$  is equivalent to the independence of the functionals  $T_a(z)$ ,  $T_b(z)$ .
3. By replacing  $i$  by  $-i$  in the definitions of the functionals  $T_a(z)$  and  $T_b(z)$ , one obtains the corresponding boundary conditions for  $\text{Im}(z) < 0$ .

#### 4.2. Resonances

In our one-dimensional situation, it is quite simple to detect the resonances as poles of the scattering matrix. According to the end of Subsection 4.1, one states

**Proposition 4.2.** *Let  $z$  be a complex number such that  $\text{Re}(z) > 0$ . Then  $z$  is a resonance of the operator  $P$  if and only if  $H_z^h - z$  is not injective.*

Indeed, the non-injectivity of  $H_z^h - z$  is equivalent to the fact that the linear functionals are proportional, so the normalisation given in (1.14)–(1.15) is not performable.

**Remark 9.** The anti-resonances are defined similarly after considering the meromorphic extension from the lower half-plane  $\{\text{Im}(z) < 0\}$  while changing  $i$  into  $-i$  in the transparent boundary conditions (see Remark 8).

### 4.3. Analysis of the resolvent

Recall that since we are interested in getting the spectral density inside the island  $I$ , Proposition 4.1 allows to work with  $H_z^h - z$  in place of  $H^h - z$ . Moreover, because Theorem 3.6 ensures that the set  $\mathcal{E}^0$  of asymptotic resonant energies is discrete, we will make the following reduction:

**Assumption 4.** Suppose that the set  $[\Lambda_*, \Lambda^*]$  contains exactly one asymptotic resonant energy  $\lambda_0 \in (\Lambda_*, \Lambda^*)$ . Recall that  $m_{\lambda_0}$  denotes its multiplicity according to (1.33) and that  $(\lambda_j^h)_{1 \leq j \leq m_{\lambda_0}}$  are the ordered eigenvalues of  $H_I^h$  lying in  $[\Lambda_*, \Lambda^*]$  (and converging to  $\lambda_0$ ).

Introduce

$$\Omega_h := \{z \in \mathbb{C} \text{ s.t. } \text{Re}(z) \in K_h, \text{Im}(z) \in [-4h, 4h]\}, \tag{4.3}$$

$$\text{with } K_h := [\lambda_0 - \alpha^h, \lambda_0 + \alpha^h], \text{ and } \alpha^h := 4 \max\{h, |\lambda_0 - \lambda_j^h|, j = 1, \dots, m_{\lambda_0}\}. \tag{4.4}$$

The parameter  $z$  is assumed to satisfy

$$z \in \Omega_h.$$

Proposition 3.9 indicates that from the spectral viewpoint, around a resonant energy the nonresonant wells do not matter. We adapt to this remark the filled well Hamiltonians

$$\tilde{H}_I^h = H_I^h + W^h \quad \text{and} \quad \tilde{H}_z^h = H_z^h + W^h. \tag{4.5}$$

For given  $\lambda \in (\Lambda_*, \Lambda^*)$ , set

$$W_\lambda^h := \sum_{i \in J_\lambda} w_i \left( \frac{-c_i}{h} \right), \quad U_\lambda^h := \text{supp } W_\lambda^h. \tag{4.6}$$

Define

$$\tilde{H}_I^h(\lambda) := H_I^h + W_\lambda^h \quad \text{and} \quad \tilde{H}_z^h(\lambda) := H_z^h + W_\lambda^h, \tag{4.7}$$

the operators associated to respectively the Dirichlet and transparent problems with the  $\lambda$ -resonant wells filled. The parameter  $\lambda$  remains fixed as  $h \rightarrow 0$  and those definitions lead to

$$\tilde{H}_\bullet^h(\lambda) = H_\bullet^h$$

when  $\lambda \neq \lambda_0$  and

$$\tilde{H}_\bullet^h(\lambda_0) = H_\bullet^h + W_{\lambda_0}^h.$$

In particular,  $\tilde{H}_I^h(\lambda_0)$  has no eigenvalue in  $[\Lambda_*, \Lambda^*]$ .

An accurate analysis of the resolvent  $(H_z^h - z)^{-1}$  starts with essentially two steps:

1. Eliminate the nonresonant wells: we show that  $\tilde{H}_z^h(\lambda_0) - z$  is invertible for all  $z \in \Omega_h$ .
2. Check that for  $z$  far from  $\lambda_0$ ,  $H_z^h - z = \tilde{H}_z^h(\lambda) - z$ ,  $\lambda \neq \lambda_0$ , is invertible.

Hence the notation  $\tilde{H}_z^h(\lambda)$  is convenient for a compact formulation of different results.

**Proposition 4.3.** *Make Assumption 4 and fix any  $\lambda \in [\Lambda_*, \Lambda^*]$ .*

(i) For any  $z \in \Omega_h$  if  $\lambda = \lambda_0$  (resp.  $z \in [\Lambda_*, \Lambda^*] \times [-4h, 4h]$  and  $\text{dist}(z, \lambda_0) > \alpha^h/2$  or  $|\text{Im}(z)| \geq 2h$  if  $\lambda \neq \lambda_0$ ), the operator  $\tilde{H}_z^h(\lambda) - z$  is invertible. The kernel of the resolvent is estimated by

$$|(\tilde{H}_z^h(\lambda) - z)^{-1}[x, y]| = \tilde{O}(e^{-\frac{\tilde{d}(x,y)}{h}}),$$

where  $\tilde{d}$  stands for the Agmon distance for the potential  $\tilde{V}^h$  at the energy  $\text{Re}(z)$ . Moreover the constants can be chosen uniform with respect to  $x, y \in I$  and  $z$ .

(ii) For any function  $\varphi \in C_c^0((a, b))$ ,  $(\tilde{H}_z^h(\lambda) - z)^{-1}\varphi$  belongs to the space  $\mathcal{L}^1$  of trace-class operators for  $z \in \Omega_h$  if  $\lambda = \lambda_0$  (resp.  $z \in [\Lambda_*, \Lambda^*] \times [-4h, 4h]$  and  $\text{dist}(z, \lambda_0) > \alpha^h/2$  or  $|\text{Im}(z)| \geq 2h$  if  $\lambda \neq \lambda_0$ ), with the estimate

$$\|(\tilde{H}_z^h(\lambda) - z)^{-1}\varphi\|_{\mathcal{L}^1} \leq C_\varphi h^{-2}.$$

**Remark 10.** In particular, applying (i) with  $\lambda = \lambda_0$ , gives, since  $H_z^h(\lambda) = H_z^h$  and using Proposition 4.2 that  $P^h$  has no resonance in the set

$$\left\{ z \in \Omega_h, |\text{Im}(z)| > 2h \text{ or } \text{dist}(z, \lambda_0) \geq \frac{\alpha_h}{2} \right\}.$$

**Proof.** The first statement will be proved in three steps (a), (b) and (c) where the last two ones are very similar.

(i)-(a) We start with the strongly elliptic problem: suppose that  $\lambda = \lambda_0$ ,  $z \in \Omega_h$  and  $J_{\lambda_0} = \{1, \dots, N\}$ , that is  $\tilde{H}_z^h(\lambda_0) = \tilde{H}_z^h$  (every well is filled). We use the Agmon identity of Appendix A where  $\varphi$  is a  $C^1(I)$ -function satisfying the eiconal condition:

$$\inf_{h>0, x \in I} \tilde{V}^h(x) - \text{Re}(z) - \varphi'^2(x) \geq m > 0,$$

and we take the real part of both sides. Since  $z \in \Omega_h$  is possibly complex, there are boundary terms in the Agmon estimates (see Appendix A) but their coefficients are  $\mathcal{O}(h^3)$ . For  $z \in \Omega_h$  and with the condition  $\Lambda_0 - \Lambda^* > 0$  according Assumption 1, the coercivity of the variational formulation with the transparent conditions (see Proposition 4.1) is easily checked when  $h > 0$  is small enough: Taking  $\varphi \equiv 0$  provides the existence of the resolvent and uniform bounds.

Taking  $\varphi$  with the above eiconal condition provides the weighted estimate

$$\forall f \in L^2(I), \quad \|e^{\frac{\varphi}{h}}(\tilde{H}_z^h - z)^{-1}f\|_{1,h} \leq C \|e^{\frac{\varphi}{h}}f\|_{L^2}.$$

The case  $\varphi \equiv (1 - \eta)\tilde{d}(\cdot, y)$  for fixed  $y \in (a, b)$  (which satisfies the eiconal condition) implies (i) in this specific case. The pointwise estimate of the Schwartz kernel of the resolvent is obtained after Appendix E.

(i)-(b) In the weaker case,  $\lambda = \lambda_0$ ,  $z \in \Omega_h$ ,  $J_{\lambda_0} \neq \{1, \dots, N\}$ , the problem is neither self-adjoint nor strongly elliptic. Only the wells in  $U_{\lambda_0}^h = \text{supp } W_{\lambda_0}^h$  according to (4.6) are filled and the other nonresonant wells are left. We use an approximation argument with the latter estimate. Set

$$S_0^z := \tilde{d}(U^h \setminus U_{\lambda_0}^h, \partial I) \tag{4.8}$$

where  $\tilde{d}$  is the Agmon distance for the potential  $\tilde{V}^h$  and the energy  $\text{Re}(z)$ . Introduce, for  $\eta > 0$  small, the cut-off functions  $\chi, \tilde{\psi}$  such that  $0 \leq \chi, \tilde{\psi} \leq 1$ ,  $\chi \equiv 1$  in the set  $\{x \in I, \tilde{d}(x, U^h \setminus U_{\lambda_0}^h) \leq S_0^z - \eta\}$ ,  $\tilde{\psi} \equiv 1$  in the set  $\{\tilde{d}(x, \tilde{U}^h \setminus U_{\lambda_0}^h) \leq (S_0^z - \eta)/2\}$ ,  $\chi \equiv 0$  in  $\{\tilde{d}(x, U^h \setminus U_{\lambda_0}^h) \geq S_0^z - \eta/2\}$  and  $\tilde{\psi} \equiv 0$  in the set  $\{\tilde{d}(x, \tilde{U}^h \setminus U_{\lambda_0}^h) \geq (S_0^z + \eta)/2\}$ .

Choose

$$R(\lambda_0) := (\tilde{H}_z^h - z)^{-1}(1 - \tilde{\psi}) + \chi(\tilde{H}_I^h(\lambda_0) - z)^{-1}\tilde{\psi} \tag{4.9}$$

as an approximate right inverse for  $\tilde{H}_z^h(\lambda_0) - z$ : Actually  $\tilde{H}_z^h(\lambda_0)$  is replaced by the corresponding Dirichlet Hamiltonian around the remaining non  $\lambda_0$ -resonant wells. Note that  $R(\lambda_0)$  is well defined since for  $z \in \Omega_h$ ,  $z$  is uniformly far away from the spectrum of  $\tilde{H}_I^h(\lambda_0)$ .

A straightforward computation using  $\tilde{H}_z^h(\lambda_0)\chi = \tilde{H}_I^h(\lambda_0)\chi$  and  $\chi\tilde{\psi} = \tilde{\psi}$  gives

$$(\tilde{H}_z^h(\lambda_0) - z)R(\lambda_0) = 1 - \varepsilon, \quad \varepsilon := \varepsilon_0 + \varepsilon_1, \tag{4.10}$$

where

$$\varepsilon_0 := \tilde{W}_\lambda^h(\tilde{H}_z^h - z)^{-1}(1 - \tilde{\psi}), \quad \varepsilon_1 := -[P^h, \chi](\tilde{H}_I^h(\lambda_0) - z)^{-1}\tilde{\psi}. \tag{4.11}$$

With the estimate about  $(\tilde{H}_z^h - z)^{-1}$  and the control of the resolvent  $(\tilde{H}_I^h(\lambda_0) - z)^{-1}$  of the Dirichlet Hamiltonian provided by Proposition 3.7 with the uniform lower bound  $\text{dist}(z, \sigma(H_I^h(\lambda_0))) \geq c > 0$ , one deduces the inequality

$$\|\varepsilon_0\| + \|\varepsilon_1\| \leq C_\eta e^{\frac{-S_0^z + c\eta}{2h}}, \tag{4.12}$$

in the operator norm.

The relation

$$(\tilde{H}_z^h(\lambda_0) - z)R(\lambda_0) = 1 - \varepsilon, \quad \|\varepsilon\| \leq C_\eta e^{\frac{-S_0^z + c\eta}{2h}} \tag{4.13}$$

ensures the injectivity of  $(\tilde{H}_z^h(\lambda_0) - z)$  and provides a right inverse after using the Neumann series for  $(1 - \varepsilon)^{-1}$ .

Similarly, setting

$$L(\lambda_0) := (1 - \tilde{\psi})(\tilde{H}_z^h - z)^{-1} + \tilde{\psi}(\tilde{H}_I^h(\lambda_0) - z)^{-1}, \tag{4.14}$$

leads to

$$L(\lambda_0)(\tilde{H}_z^h(\lambda_0) - z) = 1 + \varepsilon', \quad \|\varepsilon'\| \leq C_\eta e^{\frac{-S_0^z + c\eta}{2h}}, \tag{4.15}$$

and provides a left inverse for  $\tilde{H}_z^h(\lambda_0) - z$ .

The estimate of the kernel of the resolvent is obtained after considering the first terms in the expansion series defining the inverse

$$\chi_x \cdot R(\lambda_0) \sum_{k=0}^{\infty} \varepsilon^k \cdot \chi_y.$$

The estimate for  $k = 0$  is clear according to the estimates of the kernels (part (a) and Proposition 3.7) appearing in the definition of  $R(\lambda_0)$ . For  $k \geq 1$ , note first, since  $\tilde{\psi}[P^h, \chi] = 0$  and  $(1 - \tilde{\psi})\tilde{W}_\lambda^h = 0$  that by computing the terms corresponding to  $k = 1, k = 2$  and then by induction, the general term splits for any  $k \geq 1$  into two terms, namely

$$\chi_x R(\lambda) \varepsilon^k \chi_y = \chi_x (\tilde{H}_z^h - z)^{-1} \left( \prod_{j=1}^k \varepsilon_{[j]} \right) \chi_y + \chi_x (\tilde{H}_I^h(\lambda_0) - z)^{-1} \left( \prod_{j=1}^k \varepsilon_{[j+1]} \right) \chi_y, \tag{4.16}$$

where  $[\ell]$  stands for the class of  $\ell \pmod{2}$ . Each term involves  $k + 1$  resolvents, which induces a prefactor  $(C_\eta e^{\frac{c\eta}{2h}})^{k+1}$  in the estimate

$$\forall k \geq 1, \quad \|\chi_x \cdot R(\lambda) \varepsilon^k \cdot \chi_y\| \leq (C_\eta e^{\frac{c\eta}{2h}})^{k+1} e^{-\frac{\varphi_k(x,y)}{h}},$$

with

$$\varphi_k(x, y) = \min\{L_k(x, y), L_k(y, x)\}, \quad L_k(x, y) = \tilde{d}(x, \partial I) + (k - 1)\frac{S_0^z}{2} + \tilde{d}(y, \tilde{U}_\lambda^h).$$

We conclude, since  $\varphi_k(x, y) \geq \tilde{d}(x, y) + (k - 2)S_0^z$ , that the series is convergent (the convergence is uniform w.r.t.  $z \in \Omega_h$ ). Again the pointwise estimate is provided by Appendix E.

(i)-(c) To finish the proof of (i), it remains the case  $\lambda \neq \lambda_0, \text{dist}(z, \lambda_0) \geq \alpha_h/2$  or  $|\text{Im}(z)| \geq 2h$ . The strategy is essentially the same as in (i)-(b): we replace  $H_z^h = \tilde{H}_z^h(\lambda)$  by  $\tilde{H}_z^h$  far away from the wells and by  $\tilde{H}_I^h(\lambda) = H_I^h$  around non  $\lambda$ -resonant wells, which are all the wells. Consider this time

$$S_0^z := d(U^h, \partial I), \quad \text{with } U^h = \text{supp } W^h,$$

and  $\chi, \psi$  such that  $0 \leq \chi, \psi \leq 1, \chi \equiv 1$  in the set  $\{x \in I, \tilde{d}(x, U^h) \leq S_0^z - \eta\}, \psi \equiv 1$  in the set  $\{\tilde{d}(x, U^h) \leq (S_0^z - \eta)/2\}$  and  $\psi \equiv 0$  in the set  $\{\tilde{d}(x, U^h) \geq (S_0^z + \eta)/2\}$ . Choose as an approximate right inverse (well defined for  $z \in \Omega_h$  such that  $|\text{Im}(z)| > h$  or  $\text{dist}(z, \Lambda_0) \geq \alpha_h/2$ )

$$R = (\tilde{H}_z^h - z)^{-1}(1 - \psi) + \chi(H_I^h - z)^{-1}\psi,$$

and as an approximate left inverse

$$L = (1 - \psi)(\tilde{H}_z^h - z)^{-1} + \psi(H_I^h - z)^{-1}.$$

One obtains again a norm-convergent series thanks to resolvent estimates and the pointwise estimates of the kernel are derived from Appendix E.

(ii) We start again like for (i) by the case where  $\lambda = \lambda_0$ ,  $J_{\lambda_0} = \{1, \dots, N\}$ . For  $H_0^h$  being the Dirichlet  $h$ -Laplacian on  $I$ , write, since  $(H_0^h + i)\varphi = (\tilde{H}_z^h + i - \tilde{\nu}^h)\varphi$ :

$$\varphi(\tilde{H}_z^h - z)^{-1} = (H_0^h + i)^{-1}\varphi[1 + (z + i - \tilde{\nu}^h)](\tilde{H}_z^h - z)^{-1} + (H_0^h + i)^{-1}[P^h, \varphi](\tilde{H}_z^h - z)^{-1}. \tag{4.17}$$

One sees that the first term of the r.h.s. of (4.17) is trace-class with the announced estimates because  $(H_0^h + i)^{-1}$  is trace-class whereas the second factor is uniformly bounded. For the last term, use again that  $(H_0^h + i)^{-1}$  is trace-class and the fact that we obtained estimates for  $(\tilde{H}_z^h - z)^{-1}$  in the  $H^{1,h}$ -norm. The result follows by taking the adjoint. In the case  $\lambda = \lambda_0$ ,  $z \in \Omega_h$  and  $m_{\lambda_0} < N$ , use the series  $R(\lambda_0) \sum_{k=0}^{\infty} \varepsilon^k$  to see that

$$(\tilde{H}_z^h(\lambda_0) - z)^{-1} = [(\tilde{H}_z^h - z)^{-1}(1 - \tilde{\psi}) + \chi(\tilde{H}_I^h(\lambda_0) - z)^{-1}\tilde{\psi}][1 + \mathcal{O}(e^{-\frac{\varepsilon}{h}})], \tag{4.18}$$

and notice that the first factor is trace-class. Finally, one has something similar for  $\lambda \neq \lambda_0$  and suitable  $z$

$$(H_z^h - z)^{-1} = [(\tilde{H}_z^h - z)^{-1}(1 - \tilde{\psi}) + \chi(H_I^h - z)^{-1}\tilde{\psi}][1 + \mathcal{O}(e^{-\frac{\varepsilon}{h}})]. \quad \square \tag{4.19}$$

### 5. Localising resonances

The formalism of Grushin’s Problem provides a convenient way to treat simultaneously the question of the invertibility of the operator  $(H_z^h - z)$  raised in the latter section, and (through a perturbative formulation) to localise the resonances of  $P^h$ . We refer the reader to the appendix of [15] or to [31] for a general presentation of this technique. Fix the reference energy to the value  $\lambda_0 \in (\Lambda_*, \Lambda^*)$  and work in the set  $\Omega_h$  defined in (4.3). Denote by  $\lambda_1^h, \dots, \lambda_n^h$  the eigenvalues of  $H_I^h$  converging to  $\lambda_0$  (they lie in  $K_h$ ), and  $\phi_1^h, \dots, \phi_{m_{\lambda_0}}^h$  a corresponding orthonormal system of eigenvectors. Start by writing the Grushin’s problem for the Dirichlet realisation  $H_I^h$ :

$$\begin{cases} (H_I^h - z)u + R_0^- u^- = v, \\ R_0^+ u = v^+, \end{cases} \tag{5.1}$$

with

$$(u, u^-) \in D(H_I^h) \times \mathbb{C}^{m_{\lambda_0}}, \quad (v, v^+) \in L^2(I) \times \mathbb{C}^{m_{\lambda_0}},$$

$$R_0^- : \mathbb{C}^{m_{\lambda_0}} \longrightarrow L^2(I), \quad u^- := \begin{pmatrix} u_1^- \\ \vdots \\ u_{m_{\lambda_0}}^- \end{pmatrix} \mapsto R_0^- u^- := \sum_{j=1}^{m_{\lambda_0}} u_j^- \phi_j^h, \tag{5.2}$$

and

$$R_0^+ : L^2(I) \longrightarrow \mathbb{C}^{m_{\lambda_0}}, \quad u \mapsto R_0^+ u := \begin{pmatrix} \langle u, \phi_1^h \rangle_{L^2} \\ \vdots \\ \langle u, \phi_{m_{\lambda_0}}^h \rangle_{L^2} \end{pmatrix}. \tag{5.3}$$

Set  $F'' := \text{span}\{\phi_j^h\}_{j=1}^n$ ,  $F' := (F'')^\perp$ . This problem is invertible and the solution is given, with obvious notations by

$$\begin{cases} u' = (H_I^{h'} - z)^{-1}v', \\ u'' = \sum_{j=1}^{m_{\lambda_0}} \langle u, \phi_j^h \rangle \phi_j^h = \sum_{j=1}^{m_{\lambda_0}} v_j^+ \phi_j^h, \\ u_j^- = \langle v, \phi_j^h \rangle + (z - \lambda_j^h)v_j^+, \quad j = 1, \dots, m_{\lambda_0}, \end{cases} \tag{5.4}$$

where  $H_I^{h'}$  denotes the restriction of  $H_I^h$  to  $F'$ . In terms of operators

$$\begin{cases} u = E_0(z)v + E_0^+ v^+, \\ u^- = E_0^- v + E_0^{-+}(z)v^+, \end{cases} \tag{5.5}$$

with

$$E_0(z)v = (H_I^h - z)^{-1} \Pi_I^h v, \quad E_0^+ v^+ = \sum_{j=1}^{m_{\lambda_0}} v_j^+ \phi_j^h,$$

$$E_0^- v = \begin{pmatrix} \langle v, \phi_1^h \rangle_{L^2} \\ \vdots \\ \langle v, \phi_{m_{\lambda_0}}^h \rangle_{L^2} \end{pmatrix}, \quad E_0^{-+}(z)v^+ = \text{diag}(z - \lambda_j^h)v^+,$$

and  $\Pi_I^h$  is the orthogonal projector onto  $F'$ :

$$\Pi_I^h v := \left( 1 - \sum_{j=1}^{m_{\lambda_0}} |\phi_j^h\rangle \langle \phi_j^h| \right) v. \tag{5.6}$$

Finally, write

$$\mathcal{H}_I^h(z) := \begin{pmatrix} H_I^h - z & R_0^- \\ R_0^+ & 0 \end{pmatrix}, \quad \mathcal{E}_I^h(z) := (\mathcal{H}_I^h(z))^{-1} = \begin{pmatrix} E_0(z) & E_0^+ \\ E_0^- & E_0^{-+}(z) \end{pmatrix}. \tag{5.7}$$

We perturb the problem in order to obtain the resonant problem. Like in the proof of Proposition 4.3, set

$$S_0 := \tilde{d}_0(U_{\lambda_0}^h, \partial I), \tag{5.8}$$

where  $\tilde{d}_0$  is the Agmon distance for the potential  $\mathcal{V}^0$  at the energy  $\lambda_0$ . For  $\eta > 0$  small, fix two smooth cut-off functions  $\chi, \psi$  such that  $0 \leq \chi, \psi \leq 1$ ,  $\chi \equiv 1$  in the set  $\{x \in I, d(x, U_{\lambda_0}^h) \leq S_0 - \eta\}$ ,  $\psi \equiv 1$  in the set  $\{d(x, U_{\lambda_0}^h) \leq (S_0 - \eta)/2\}$  and  $\psi \equiv 0$  in the set  $\{d(x, U_{\lambda_0}^h) \geq (S_0 + \eta)/2\}$ . Define

$$\mathcal{H}(z; h) := \begin{pmatrix} H_z^h - z & \chi R_0^- \\ R_0^+ & 0 \end{pmatrix}, \quad z \in \Omega_h. \tag{5.9}$$

Far from the resonant wells,  $H_z^h$  looks like  $\tilde{H}_z^h(\lambda_0)$  and around the wells the Dirichlet problem (with all the wells) is a good approximation of  $H_z^h$ . This leads to set

$$\mathcal{F}(z; h) := \begin{pmatrix} \chi E_0 \psi + (\tilde{H}_z^h(\lambda_0) - z)^{-1} (1 - \psi) & \chi E_0^+ \\ E_0^- \psi & E_0^{-+} \end{pmatrix}. \tag{5.10}$$

One shows that

$$\mathcal{H}(z; h) \mathcal{F}(z; h) = 1 + \mathcal{K}(z; h)$$

and  $\mathcal{K}$  satisfies the estimate

$$\mathcal{K}(z; h) = \begin{pmatrix} \tilde{\mathcal{O}}(e^{-\frac{S_0}{2h}}) & \tilde{\mathcal{O}}(e^{-\frac{S_0}{h}}) \\ \tilde{\mathcal{O}}(e^{-\frac{S_0}{2h}}) & \tilde{\mathcal{O}}(e^{-\frac{2S_0}{h}}) \end{pmatrix}. \tag{5.11}$$

More precise computations with the second order expansion of the Neumann series and using the resolvent estimates of Proposition 4.3 can be done. When all the wells are resonant,  $m_{\lambda_0} = N$ , details are given by the direct transcription of [15, pp. 117–128]. The more general case was treated in [28, pp. 178–189].

**Proposition 5.1.** *With the notations (4.3) and (5.8) and for  $z \in \Omega_h$ , the operator is invertible, and the inverse is given by the norm convergent series*

$$\mathcal{H}(z; h)^{-1} = \mathcal{F}(z; h) \sum_{j=0}^{\infty} (-1)^j \mathcal{K}^j(z; h) = \begin{pmatrix} E(z; h) & E^+(z; h) \\ E^-(z; h) & E^{-+}(z; h) \end{pmatrix},$$

with

$$E^{-+}(z) = E_0^{-+} + \tilde{\mathcal{O}}(e^{-\frac{2S_0}{h}}).$$

Moreover, it is uniformly norm-bounded holomorphic function of  $z \in \Omega_h$ .



Within the Grushin problem approach, the invertibility of  $H_z^h - z$  is reduced to the question of invertibility of the finite-dimensional block  $E^{-+}(z)$  (see the Schur complement formula (6.7)). In particular, considering  $\det(E^{-+}(z))$  leads to the next standard approximation result of resonances by Dirichlet eigenvalues.

**Proposition 5.2.** *Take the notation (4.3) and (5.8). The operator  $P^h$  has exactly  $m_{\lambda_0}$  resonances (counted with multiplicity)  $z_1^h, \dots, z_{m_{\lambda_0}}^h$  in  $\Omega_h$ . They satisfy*

$$\forall j \in \{1, \dots, m_{\lambda_0}\}, \quad |z_j^h - \lambda_j^h| = \tilde{\mathcal{O}}(e^{-\frac{2S_0}{h}})$$

and have negative imaginary parts.

### 6. Local density of states

We end the proof of Theorem 1.6 by considering the asymptotic behaviour of the density associated with a function of the energy.

**Proposition 6.1.** *Let  $\theta \in C_c^0((\Lambda_*, \Lambda^*))$  and keep the notations (4.4) under Assumptions 1, 3 and 4. The particle density  $dn_{\theta_\lambda}[V^h]$  defined for  $g(k) = \theta(\lambda_k)$  satisfies the following weak\* asymptotics in  $\mathcal{M}_b((a, b))$ : For all  $\varphi \in C_c^0((a, b))$ ,*

$$\lim_{h \rightarrow 0} \int_a^b \varphi(x) dn_{\theta_\lambda} = \lim_{h \rightarrow 0} \text{Tr}[\theta(H^h)\varphi] = \lim_{h \rightarrow 0} \text{Tr}[(\theta \cdot 1_{K_h})(H^h)\varphi] = \sum_{i \in J_{\lambda_0}} \theta(\lambda_0)\varphi(c_i). \tag{6.1}$$

This result which is a Breit–Wigner type formula for the density of states like in [13] will be proved in two steps: 1) eliminating the nonresonant energies; 2) specifying the contribution of resonant states.

#### 6.1. Eliminating the nonresonant energies

We first check that the density goes to 0 in  $(a, b)$  as  $h$  goes to 0 when all the wells are filled, that is for  $\tilde{H}^h$ , and reduce the more general nonresonant energy problem to this case after using an approximate resolvent provided by (4.18)–(4.19). We start with a simple accurate estimate.

**Proposition 6.2.** *Let  $\tilde{\psi}_-^h(k, \cdot)$  be the incoming scattering states of  $\tilde{H}^h$ , such that  $\lambda_k \in [\Lambda_*, \Lambda^*]$ . The function  $\tilde{\psi}_-^h(k, \cdot)$  is uniformly bounded with respect to  $x \in [a, b]$  and  $k$ . Moreover one has the uniform pointwise estimate*

$$\begin{aligned} \tilde{\psi}_-^h(k, x) &= \mathcal{O}(h^{-1/2}e^{-\frac{\tilde{d}_h(a,x)}{h}}), \quad k > 0, \\ \text{and } \tilde{\psi}_-^h(k, x) &= \mathcal{O}(h^{-1/2}e^{-\frac{\tilde{d}_h(b,x)}{h}}), \quad k < 0, \end{aligned}$$

where  $\tilde{d}_h$  stands for the Agmon distance for the potential  $\tilde{V}^h$  at the energy  $\lambda_k$ .

**Proof.** We focus on the case  $k > 0$  (if  $k < 0$ , just swap  $a$  and  $b$ ). Start by noticing that for given  $k$ , the function  $A_k^h : x \mapsto |\tilde{\psi}_-^h(k, x)|^2$  satisfies

$$h^2 \frac{d^2}{dx^2} A_k^h = 2|h\partial_x \tilde{\psi}_-^h(k, \cdot)|^2 + 2(\tilde{V}^h - \lambda_k)|\tilde{\psi}_-^h(k, \cdot)|^2 \geq 0. \tag{6.2}$$

It follows that the function  $h\partial_x A_k^h$  is increasing on  $I$ . But the scattering condition (1.17) says that this function vanishes at  $x = b$ . So the function  $A_k^h$  is convex and decreasing on  $I$ . Now, it suffices to show that the family  $(A_k^h(a))_k$  is uniformly bounded. But it equals

$$A_k^h(a) = |\tilde{\psi}_-^h(k, a)|^2 = |e^{i\frac{ka}{h}} + r_k e^{-i\frac{ka}{h}}|^2, \tag{6.3}$$

which is bounded according to (1.16).

Use the Agmon estimate of Appendix A with  $V = \tilde{\mathcal{V}}^h$ ,  $z = \lambda_k$ ,  $u = v = \tilde{\psi}_-^h(k, \cdot)$  and  $\varphi = \tilde{d}_h(a, x)$ . Since  $\tilde{P}^h u = zu$ , and  $V - \varphi^2 - z = 0$ , this leads after taking the real part to

$$\|h\partial_x(e^{\frac{\varphi}{h}}\tilde{\psi}_-^h(k, \cdot))\|_{L^2(I)}^2 \leq h^2 e^{\frac{2\varphi(a)}{h}} |\operatorname{Re}(h\partial_x\tilde{\psi}_-^h(k, a)\overline{\tilde{\psi}_-^h(k, a)})| + h^2 e^{\frac{2\varphi(b)}{h}} |\operatorname{Re}(h\partial_x\tilde{\psi}_-^h(k, b)\overline{\tilde{\psi}_-^h(k, b)})| \tag{6.4}$$

$$\leq 2|k|A_k^h(a)^{1/2} = \mathcal{O}(1). \tag{6.5}$$

Writing

$$e^{\frac{\varphi(x)}{h}}\tilde{\psi}_-^h(k, x) = \tilde{\psi}_-^h(k, a) + h^{-1} \int_a^x h\partial_x(e^{\frac{\varphi(t)}{h}}\tilde{\psi}_-^h(k, t)) dt,$$

and Schwarz’s inequality yield the result.  $\square$

**Corollary 6.3.** Assume  $\theta \in C_c^0((\Lambda_*, \Lambda^*))$  and  $\varphi \in C_c^0((a, b))$ . The operator  $\theta(\tilde{H}^h)\varphi$  is trace-class with a trace estimated by

$$\operatorname{Tr}[\theta(\tilde{H}^h)\varphi] = \tilde{\mathcal{O}}(e^{-\frac{c \operatorname{dist}(\operatorname{supp}\varphi, \partial I)}{h}}),$$

where  $\operatorname{dist}(x, y) = |x - y|$  and  $c$  is a positive constant. The family of measures  $(dn_{\theta_\lambda}[\tilde{V}^h])_{h>0}$  weakly converges to 0 in  $\mathcal{M}_b((a, b))$ .

**Proof.** The function  $\varphi$  can be assumed nonnegative. We write

$$\begin{aligned} \int_a^b \varphi(x) dn_{\theta_\lambda}[\tilde{V}^h](x) &= \operatorname{Tr}[\varphi^{1/2}\theta(\tilde{H}^h)\varphi^{1/2}] \\ &= \int_{\mathbb{R}} \int_a^b \theta(\lambda_k) |\tilde{\psi}_-^h(k, x)|^2 \varphi(x) \frac{dk}{2\pi h}, \end{aligned}$$

after using the expression of the kernel of  $\theta(\tilde{H}^h)$ . Proposition 6.2 combined with the fact that the Agmon distance  $\tilde{d}_h$  associated with  $\tilde{\mathcal{V}}$  and an energy  $\lambda \in (\Lambda_*, \Lambda^*)$  is uniformly equivalent to the Euclidean distance, yields the result after integration.  $\square$

Thanks to this result one easily gets rid of nonresonant energies.

**Proposition 6.4.** Consider the energy interval  $K_h$  defined in (4.4) and set  $\tilde{\theta}^h(\lambda) := (1 - 1_{K_h}(\lambda)) \cdot \theta(\lambda)$ . Then in restriction to  $(a, b)$ , the measure  $dn_{\tilde{\theta}^h}^h$  weakly converges to 0 as  $h$  goes to 0:

$$\forall \varphi \in C_c^0((a, b)), \lim_{h \rightarrow 0} \operatorname{Tr}(\tilde{\theta}^h(H^h)\varphi) = 0.$$

**Proof.** We again assume again  $\varphi \geq 0$  and apply Stone’s formula in order to compute the trace of  $\varphi^{1/2}\mathbf{1}_I\tilde{\theta}^h(H^h)\mathbf{1}_I\varphi^{1/2}$ . By referring to Proposition 4.1 and by using successively (4.18)–(4.19) one obtains

$$\operatorname{Tr}(\tilde{\theta}^h(H^h)\varphi) = \operatorname{Tr}(\tilde{\theta}^h(\tilde{H}^h)(1 - \tilde{\psi})\varphi) + \operatorname{Tr}(\chi\tilde{\theta}^h(H_I^h)\tilde{\varphi}) + \mathcal{O}(h^{-2}e^{-\frac{c}{h}}), \quad h \rightarrow 0. \tag{6.6}$$

The first term can be estimated by

$$0 \leq \operatorname{Tr}(\tilde{\theta}^h(\tilde{H}^h)(1 - \tilde{\psi})\varphi) \leq \operatorname{Tr}(\theta(\tilde{H}^h)(1 - \tilde{\psi})\varphi),$$

with a right-hand side converging to 0 by Corollary 6.3. Meanwhile the second term cancels since  $H_I^h$  has no spectrum on the support of  $\tilde{\theta}^h$ . This finishes the proof.  $\square$

### 6.2. Contribution of resonant states

Let us first go back to the Grushin problem introduced in Section 5. According to Proposition 5.1, and estimates (5.11) we have

$$\mathcal{H}(z; h)^{-1} := \begin{pmatrix} E(z) & E^+(z) \\ E^-(z) & E^{-+}(z) \end{pmatrix} = \mathcal{F}(z; h) \begin{pmatrix} 1 + \varepsilon(z) & \varepsilon^+(z) \\ \varepsilon^-(z) & 1 + \varepsilon^{-+}(z) \end{pmatrix},$$

with  $\varepsilon^\bullet(z) = \tilde{\mathcal{O}}(e^{-S_0/2h})$  uniformly in  $z \in \Omega_h$ . This implies

$$[H_z^h - z]^{-1} = E(z) - E^+(z)(E^{-+}(z))^{-1}E^-(z). \tag{6.7}$$

Coming back to the definition (5.10) of  $\mathcal{F}^h(z)$ , this can be improved into

$$E(z) = (\tilde{H}_z^h(\lambda_0) - z)^{-1}(1 - \psi)(1 + \varepsilon) + \chi E_0(z)\psi(1 + \varepsilon) + \chi E_0^+\varepsilon^-, \tag{6.8}$$

$$E^+(z) = \chi E_0^+ + (\tilde{H}_z^h(\lambda_0) - z)^{-1}(1 - \psi)\varepsilon^+ + \chi E_0^+\varepsilon^{-+} + \chi E_0(z)\psi\varepsilon^+, \tag{6.9}$$

$$E^-(z) = E_0^-\psi + E_0(z)\psi\varepsilon + E_0^{-+}(z)\varepsilon^-, \tag{6.10}$$

$$E^{-+}(z) = E_0^{-+}(z) + \tilde{\mathcal{O}}(e^{-\frac{2S_0}{h}}). \tag{6.11}$$

We are ready to apply Stone’s formula with a complex deformation of the integration contour. Before this, we write under an adapted form the polar part coming from (6.11).

**Lemma 6.5.** *Set  $\tilde{\Omega}_h := [\lambda_0 - \alpha^h/2, \lambda_0 + \alpha^h/2] \times [-2ih, 2ih]$ . For  $z$  in  $\Omega_h \setminus \tilde{\Omega}_h$ , there exist a constant  $c > 0$  and a matrix-valued meromorphic function  $G$  such that*

$$E^{-+}(z)^{-1} = E_0^{-+}(z)^{-1} + G(z), \quad \|G(z)\| = \mathcal{O}(e^{-\frac{c}{h}}), \quad h \rightarrow 0.$$

**Proof.** Fix any matrix-norm on  $\mathbb{C}^{m_{\lambda_0}}$  and use again (6.11) to see that

$$E^{-+}(z) = (1 + F(z)E_0^{-+}(z)^{-1})E_0^{-+}(z), \quad z \neq \lambda_j^h, \\ \|F(z)\| = \mathcal{O}(e^{-\frac{2S}{h}}), \quad 0 < S < S_0 \text{ for } z \in \Omega_h \setminus \tilde{\Omega}_h. \tag{6.12}$$

Because of the expression of  $E_0^{-+}(z)$ ,

$$\|F(z)E_0^{-+}(z)^{-1}\| = \mathcal{O}(e^{-\frac{2S}{h}}) \left( \min_{j=1, \dots, m_{\lambda_0}} |z - \lambda_j^h| \right)^{-1}.$$

For  $z \neq \lambda_j^h, j = 1, \dots, m_{\lambda_0}$ ,

$$E^{-+}(z)^{-1} = E_0^{-+}(z)^{-1} [1 + F(z)E_0^{-+}(z)^{-1}]^{-1}, \tag{6.13}$$

and the condition  $z \in \Omega_h \setminus \tilde{\Omega}_h$  implies  $\min_{j=1, \dots, m_{\lambda_0}} |z - \lambda_j^h| \geq h$ . Therefore, the Neumann expansion of  $[1 + F(z)E_0^{-+}(z)^{-1}]^{-1}$  converges, which yields the result.  $\square$

We can end the proof of Theorem 1.6 with the

**Proof of Proposition 6.1.** Owing to Proposition 6.4 it is enough to consider the trace

$$\mathbf{1}_I(\mathbf{1}_{K_h} \cdot \theta)(H^h)\mathbf{1}_I\varphi.$$

According to Stone’s formula and Proposition 4.1 one gets for nonnegative functions  $\theta \in \mathcal{C}_c^0((\Lambda_*, \Lambda^*))$ , and  $\varphi \in \mathcal{C}_c^0(I)$ ,

$$\mathbf{1}_I(\mathbf{1}_{K_h} \cdot \theta)(H^h)\mathbf{1}_I\varphi = \frac{-1}{2i\pi} \int_{K_h+i0} \theta(\lambda)(\lambda - H_\lambda^h)^{-1}\varphi d\lambda + \frac{1}{2i\pi} \int_{K_h-i0} \theta(\lambda)(\lambda - H_\lambda^h)^{-1}\varphi d\lambda, \tag{6.14}$$




Fig. 2. Application of Stone's formula. Resonances lie on the second sheet and close to  $\lambda_0$  (semi-circle gray).

where  $(H_z^h - z)^{-1}$  denotes the (meromorphic continuation from the lower-half complex plane) of the truncated resolvent  $\mathbf{1}_I(H^h - z)^{-1}\mathbf{1}_I$ , corresponding to the anti-resonant boundary conditions (see Remark 9).

For fixed  $\varepsilon > 0$ , consider the contour  $C_\varepsilon$  made by the segments  $(\Lambda_* + i\varepsilon, \Lambda^* + i\varepsilon) \cap \Omega_h$  and  $(\Lambda_* - i\varepsilon, \Lambda^* - i\varepsilon) \cap \Omega_h$  scoured in opposite way, the first one by real parts increasing (see Fig. 2). This contour is homotopic to the union of the circle  $\gamma_h$  and the contour  $C'_\varepsilon$  (depicted in Fig. 2) which lies on the square root Riemann surface ramified along  $\mathbb{R}_+$ . Part of the deformation takes place on the second sheet where resonances appear as poles. Meanwhile in the lower half-plane (first sheet) the resolvent is given by the anti-resonant boundary conditions (see Remark 9). The operator corresponding to these dual transparent boundary conditions is denoted by  $H_z^h$  and its resolvent,  $[H_z^h - z]^{-1}$ , has the same properties as  $[H_z^h - z]^{-1}$ , up to the sign of imaginary parts. Since for any given function  $\varphi \in \mathcal{C}_c^0((a, b))$ , the functional  $\theta \mapsto \text{Tr}[\theta(H^h)\varphi]$  defines a nonnegative measure while the right-hand side  $\sum_{i \in J_{\lambda_0}} \theta(\lambda_0)\varphi(c_i)$  of (6.1) is also a positive functional of  $\theta$ , the function  $\theta$  can be replaced by a polynomial approximation on the interval  $[\Lambda_*, \Lambda^*]$ . Use polynomial approximations from below (resp. from above) in order to get a lower bound (resp. upper bound) of the limit in (6.1). Hence we can assume that  $\theta$  is a polynomial function on  $[\Lambda_*, \Lambda^*]$ , which allows the complex deformation of the contour integral.

We first integrate the polar part. Consider first the integral over  $\gamma_h$ , which involves only  $(H_z^h - z)^{-1}$ . Use expression (6.7) first. Let us note immediately that  $E(z)$  is a holomorphic function in a neighbourhood of  $\gamma_h$ , its integral is null. Moreover (6.9) and (6.10) say

$$E^+(z) = \chi E_0^+ + \tilde{\mathcal{O}}(e^{-\frac{S_0}{2h}}), \quad E^-(z) = E_0^- \psi + \tilde{\mathcal{O}}(e^{-\frac{S_0}{2h}}). \tag{6.15}$$

These estimates hold in the norm of trace-class operators since these operators are of finite-rank.

On the contour  $\gamma_h$ , one has

$$E^{-+}(z)^{-1} = E_0^{-+}(z)^{-1} + G(z), \quad G(z) = \mathcal{O}(1), \tag{6.16}$$

so coming back to (6.7)

$$(z - H_z^h)^{-1} = -E(z) + \chi E_0^+ E_0^{-+}(z)^{-1} E_0^- \psi + \mathcal{O}(e^{-\frac{S_0}{2h}}), \quad h \rightarrow 0. \tag{6.17}$$

We integrate over  $\gamma_h$  and we obtain with  $\theta(\lambda_j^h) = \theta(\lambda_0) + o(1)$ :

$$\int_{\gamma_h} \theta(z)(H_z^h - z)^{-1} \varphi \frac{dz}{2i\pi} = 0 + \theta(\lambda_0)\chi E_0^+ E_0^- \psi \varphi + o(1)\|\varphi\|_\infty. \tag{6.18}$$

Note that  $E_0^+ E_0^-$  is nothing but the orthogonal projector on the Dirichlet states  $\sum_{j=1}^{m_{\lambda_0}} |\phi_j^h\rangle\langle\phi_j^h|$ . By taking the trace and using its cyclicity, one has with the approximation of the Dirichlet states by superpositions of the eigenfunctions of the one-well problem in Theorem 3.6

$$\begin{aligned} \operatorname{Tr} \left[ \chi \sum_{j=1}^{m_{\lambda_0}} |\phi_j^h\rangle \langle \phi_j^h| \psi \varphi \right] &= \operatorname{Tr} \left[ \sum_{j=1}^{m_{\lambda_0}} |\phi_j^h\rangle \langle \phi_j^h| \psi \varphi \right] \\ &= \sum_{j=1}^{m_{\lambda_0}} \langle \phi_j^h, \phi_j^h \psi \varphi \rangle_{L^2} \\ &= \sum_{j \in J(\lambda_0)} \varphi(c_j) + o(1) \|\varphi\|_\infty. \end{aligned}$$

Let us come to the contour  $C'_\varepsilon$  of which the projection on  $\mathbb{C}$  lies in  $\Omega_h \setminus \tilde{\Omega}_h$ . Note that the polar part coming from  $(H_z^h - z)^{-1}$  is to be treated with the integral of the polar part coming from  $(H_z^h - z)^{-1}$ . Since (with obvious notations)

$$E'^{-+}(z)^{-1} - E^{-+}(z)^{-1} = E'^{-+}(z)^{-1} (E^{-+}(z) - E'^{-+}(z)) E^{-+}(z)^{-1},$$

Lemma 6.5 implies that the difference is exponentially small because the resonances and anti-resonances are at distance greater than  $h$  from  $C'_\varepsilon$ .

It remains the holomorphic part over  $C'_\varepsilon$ . Because the polar part is treated, one can compute this integral after the inverse homotopy leading back to  $C_\varepsilon$ . But coming back to the expansion series (6.8) of  $E(z)$  (resp.  $E'(z)$ ) with main term given by  $\tilde{H}_z^h$  (resp.  $\tilde{H}'_z^h$ ), the application of Stone’s formula gives that the contribution of these terms is zero by Proposition 6.4.  $\square$

### Appendix A. Agmon identity

Here we just give the basic energy identity.

**Lemma A.1.** *Let  $\Omega := (\alpha, \beta)$  be an open interval,  $V \in L^\infty(\omega)$ ,  $z \in \mathbb{C}$  and  $\varphi$  a Lipschitz real function on  $\Omega$ . Denote by  $P$  the Schrödinger operator  $P := -h^2 d^2/dx^2 + V$ . Then for any  $u_1, u_2$  in  $H^2(\Omega)$ , and setting  $v_j := e^{\varphi/h} u_j$  one has:*

$$\begin{aligned} \int_\alpha^\beta e^{\frac{2\varphi}{h}} (P - z) u_1 \bar{u}_2 \, dx &= \int_\alpha^\beta h v_1' \overline{h v_2'} \, dx + \int_\alpha^\beta (V - z - \varphi'^2) v_1 \bar{v}_2 \, dx + \int_\alpha^\beta h \varphi' (v_1' \bar{v}_2 - v_1 \bar{v}_2') \, dx \\ &\quad + h^2 (e^{\frac{2\varphi(\alpha)}{h}} u_1' \bar{u}_2(\alpha) - e^{\frac{2\varphi(\beta)}{h}} u_1' \bar{u}_2(\beta)). \end{aligned}$$

This identity is obtained after conjugation of  $hd/dx$  by  $e^{\varphi/h}$  and integration by parts.

### Appendix B. Monotony principle

A small variation of [25] provides the next result.

**Proposition B.1.** *For  $i = 1, 2$ , let  $V_i$  be two nonnegative functions in  $L^\infty(I)$  and  $H_i := H_B^h + V_i$ . Consider a function  $F \in \mathcal{S}(\mathbb{R})$  which is decreasing on  $[-B, +\infty)$ . Write  $F_\lambda(k) = F(\lambda_k)$  and define  $dn_{F_\lambda}$  according to (1.35) and (1.20). Then the inequality*

$$\int_I (V_2 - V_1) dn_F[V_2] \leq \int_I (V_2 - V_1) dn_F[V_1]$$

holds.

This inequality is a convexity inequality which is a key ingredient in the analysis of thermodynamical equilibria of Schrödinger–Poisson systems (see [24,25]). Here the assumption  $V_i \geq 0$  ensures  $\sigma(H_i) \subset [-B, +\infty)$ . The convexity inequality with a continuous spectrum has been proved in [25], with the assumption that the potential is 0 at infinity. Here the different values 0 and  $-B$  for  $x < a$  and  $x > b$  do not bring any additional difficulties in this simple one-dimensional problem.

### Appendix C. Spectral approximation

We refer the reader to [14,16] for the details. Recall that if  $E$  and  $F$  are two given closed subspaces of a Hilbert space  $\mathcal{H}$ , with orthogonal projections  $\Pi_E$  and  $\Pi_F$ , the nonsymmetric distance from  $E$  to  $F$ , denoted by  $\vec{d}(E, F) \in [0, 1]$  is the norm of operator  $(1 - \Pi_F)\Pi_E$ , and if  $\vec{d}(E, F) < 1$ ,  $\Pi_F$  induces on  $E$  a continuous injection on its range with bounded inverse. Moreover, if at the same time  $\vec{d}(F, E) < 1$ , the latter distances are equal. In particular  $E$  and  $F$  have same dimension.

**Proposition C.1.** *Let  $A$  be an unbounded self-adjoint operator on  $\mathcal{H}$  and  $\Lambda := [\lambda_-, \lambda_+] \subset \mathbb{R}$ . Suppose that there exist  $\varepsilon > 0$ ,  $N$  linearly independent vectors  $\psi_1, \dots, \psi_N$  in the domain of  $A$ ,  $\mu_1, \dots, \mu_N$ ,  $N$  real numbers in  $\Lambda$  such that  $A\psi_j = \mu_j\psi_j + r_j$ , with  $\|r_j\| \leq \varepsilon$ . If  $A$  has no spectrum in  $\{x, 0 < \text{dist}(x, \Lambda) \leq 2a\}$  for some  $a > 0$ , then the subspaces  $E := \text{Span}(\psi_1, \dots, \psi_N)$  and  $F$  equal to the spectral subspace  $\mathbf{1}_\Lambda(A)\mathcal{H}$  verify*

$$\vec{d}(E, F) \leq \left(\frac{N}{\rho_*}\right)^{1/2} \frac{\varepsilon}{a},$$

where  $\rho_*$  is the smallest eigenvalue of the Gram matrix with entries  $\langle \psi_i, \psi_j \rangle$ .

In particular if  $A$  is known to have only discrete spectrum and if the directed distance  $\vec{d}(E, F)$  can be proved in this way to be smaller than 1, then  $A$  has at least  $N$  eigenvalues lying in  $\Lambda$ .

### Appendix D. Scattering states for the barrier

**Proposition D.1.** *Let  $V_0$  denote the constant potential  $V_0(x) \equiv \Lambda^*$  on  $I$ . Set  $H_0^h := -h^2\Delta + V_0 - B \cdot \mathbf{1}_{(b, \infty)}$  and let  $\{\psi_-^h(k, \cdot)\}_k$  denote its scattering states. Set  $S_k := \sqrt{(\Lambda^* - \lambda_k)}$ ,  $\lambda_k < \Lambda^*$ . Then one has as  $h \rightarrow 0$ , and uniformly for  $x \in I$ , for  $k > 0$*

$$\begin{aligned} |\psi_-^h(k, x)|^2 &= \frac{4k^2}{\Lambda^*} e^{-\frac{S_k(x-a)}{h}} \left(1 + \mathcal{O}\left(e^{-\frac{2S_k(b-x)}{h}}\right)\right), \\ |\psi_-^h(-k, x)|^2 &= \frac{4k^2}{\Lambda^* + B} e^{\frac{S_k(x-b)}{h}} \left(1 + \mathcal{O}\left(e^{-\frac{2S_k(x-a)}{h}}\right)\right). \end{aligned}$$

It suffices to solve explicitly on  $I$  the system characterising  $\psi_-^h(k, \cdot)$  on the explicit basis of solutions to the ODE (since the potential is constant on  $I$ ). Use the scattering conditions (1.17)–(1.18). These conditions are still valid when  $\lambda < 0$  because of the choice of the square root indeed. Finally the computation reduces to the solving of 2 by 2 affine systems. We just give the final result.

### Appendix E. Pointwise estimate for the resolvent

The next result shows that no Lipschitz regularity is necessary in dimension 1 in order to transform weighted  $L^2$ -estimates into pointwise estimates of the Green functions. Once the weighted  $L^2$ -estimates are obtained from the Agmon identity of Appendix A, it suffices to use the equation after the regularisation of the Agmon distance which is possible because the  $\tilde{\mathcal{O}}$  estimates can absorb little exponential errors.

**Proposition E.1.** *Let  $H = -h^2\Delta + \mathcal{V}$  be a closed operator with  $\mathcal{V} \in L^\infty(I)$ ,  $I = [a, b]$ ,  $D(H) \subset H^2(I)$ , with dual  $H'$  and  $D(H') \subset H^2(I)$ . Fix  $z \in \mathbb{C}$  such that  $z \notin \sigma(H)$  for all  $h \in (0, h_0)$ . We assume that there is a distance  $d \in C^0(I \times I)$ , such that the resolvent estimate*

$$\|\chi_x(z - H)^{-1}\chi_y\|_{\mathcal{L}(L^2)} \leq C_\eta A(h) e^{-\frac{d(x,y)+\eta}{h}}$$

holds for all  $(x, y, h) \in I \times I \times (0, h_0)$  as soon as  $\eta \in (0, \eta_0)$ , with  $\eta_0 > 0$  small enough and  $\chi_p$  generically denotes a cut-off function supported in  $|x - p| = \mathcal{O}(\eta)$ . Then the pointwise estimate

$$|(z - H)^{-1}[x, y]| = \tilde{\mathcal{O}}\left(A(h) e^{-\frac{d(x,y)}{h}}\right)$$

holds with uniform constants with respect to  $(x, y, h) \in I \times I \times (0, h_0)$ .

**Proof.** Let  $y_0 \in I$  be fixed. Consider a smooth function  $\varphi \in C^\infty(I)$  which is an approximation of  $d(x, y_0)$ , such that  $\|\varphi - d(\cdot, y_0)\|_{L^\infty} \leq \eta$  and  $f \in L^2(I)$ .

Let  $u$  be the solution to  $(H - z)u = \chi_{y_0} f$  so that

$$e^{\frac{\varphi}{h}}(-h^2 \Delta + \mathcal{V} - z)e^{-\frac{\varphi}{h}}(e^{\frac{\varphi}{h}} u) = e^{\frac{\varphi}{h}} \chi_{y_0} f.$$

By defining  $v = e^{\varphi/h} u$ , the assumption leads to the estimate

$$\|v\|_{L^2} \leq C_\eta A(h) e^{\frac{c\eta}{h}} \|\chi_{y_0} f\|. \tag{E.1}$$

Using the relation

$$e^{\frac{\varphi}{h}}(-h^2 \partial_x^2 + V - z)e^{-\frac{\varphi}{h}} = -h^2 \partial_x^2 + 2h\varphi' \partial_x + h\varphi'' + V - (\varphi')^2 - z,$$

we can write

$$[C - h^2 \partial_x^2 + 2h\varphi' \partial_x]v = e^{\frac{\varphi}{h}} \chi_{y_0} f + Cv + h\varphi'' v - (V - (\varphi')^2 - z)v, \tag{E.2}$$

where  $C$  is a strictly positive constant large enough. The regularity of  $\varphi$  implies

$$\|v\|_{H^{2,h}} \leq C_\eta e^{\frac{c\eta}{h}} \|\chi_{y_0} \tilde{f}\|_{L^2}.$$

In dimension one,  $H^{2,h}$  is continuously embedded in  $C^0([a, b])$ . Thus the application  $f \mapsto e^{\varphi/h}(H - z)^{-1}e^{-\varphi/h} \chi_{y_0} f$  is continuous from  $L^2([a, b])$  onto  $C^0([a, b])$  with the above uniform estimate.

By duality,  $\chi_{y_0} e^{-\varphi/h}(H' - z)^{-1}e^{-\varphi/h}$  is continuous from  $(\mathcal{M}_b(I), \|\cdot\|_b)$  onto  $L^2$ .

By changing  $y_0$  into  $x_0$  and  $H$  into  $H'$ , this says that  $v_1 = \chi_{x_0} e^{d(x_0, y_0 - c\eta)/h}(H - z)^{-1} \delta_{y_0}$  has an  $L^2(I)$ -norm bounded by  $C_\eta A(h)$ . A bootstrap with (E.2) leads to the uniform estimate of  $|v_1(x)|$ , which yields the pointwise resolvent estimate.  $\square$

### Acknowledgements

Part of this work has been carried out while the second author was visiting the Weierstrass Institute for Applied Analysis and Stochastics in Berlin. He takes the opportunity to thank them for their hospitality. This work has benefited from many discussions with many people. Among them, we thank N. Ben Abdallah, H.C. Kaiser, T. Koprucki, F. Méhats, H. Neidhardt, G. Perelman, C. Presilla, J. Rehberg and J. Sjöstrand.

### References

[1] E. Balslev, J.M. Combes, Spectral properties of many-body Schrödinger operators with dilatation-analytic interactions, *Commun Math. Phys.* 22 (1971) 280–294.  
 [2] M. Baro, H.-Chr. Kaiser, H. Neidhardt, J. Rehberg, A quantum transmitting Schrödinger–Poisson system, *Rev. Math. Phys.* 16 (3) (2004) 281–330.  
 [3] M. Baro, H.-Chr. Kaiser, H. Neidhardt, J. Rehberg, Dissipative Schrödinger–Poisson systems, *J. Math. Phys.* 45 (1) (2004) 21–43.  
 [4] N. Ben Abdallah, P. Degond, P.A. Markowich, On a one-dimensional Schrödinger–Poisson scattering model, *Z. Angew. Math. Phys.* 48 (1) (1997) 135–155.  
 [5] V. Bonnaillie-Noël, F. Nier, M. Patel, Computing the steady states for an asymptotic model of quantum transport in resonant heterostructures, *J. Comp. Phys.* 219 (2006) 644–670.  
 [6] V. Bonnaillie-Noël, F. Nier, M. Patel, Far from equilibrium steady states of 1D-Schrödinger–Poisson systems with quantum wells II, *Prépublications IRMAR*, 2007.  
 [7] M. Büttiker, Y. Imry, R. Landauer, S. Pinhas, Generalized many-channel conductance formula with application to small rings, *Phys. Rev. B* 31 (1985) 6207–6215.  
 [8] F. Chevoir, B. Vinter, Scattering assisted tunneling in double barriers diode: scattering rates and valley current, *Phys. Rev. B* 47 (1993) 7260–7274.  
 [9] E.B. Davies, *Spectral Theory and Differential Operators*, Cambridge Studies in Advanced Mathematics, vol. 42, Cambridge University Press, Cambridge, 1995.  
 [10] P. Degond, F. Mehats, C. Ringhofer, Quantum hydrodynamic models derived from the entropy principle, *Contemp. Math.* 371 (2005) 107–131.  
 [11] J. Dereziński, C. Gérard, *Asymptotic Completeness of Classical and Quantum N-Particles Systems*, Texts and Monographs in Physics, Springer-Verlag, 1997.  
 [12] M. Dimassi, J. Sjöstrand, *Spectral Asymptotics in the Semi-Classical Limit*, London Mathematical Society Lecture Note Series, vol. 268, Cambridge University Press, 1999.

- [13] C. Gérard, A. Martinez, Semiclassical asymptotics for the spectral function of long-range Schrödinger operators, *J. Funct. Anal.* 84 (1) (1989) 226–254.
- [14] B. Helffer, *Semi-Classical Analysis for the Schrödinger Operator and Applications*, Lecture Notes in Mathematics, vol. 1336, Springer-Verlag, 1988.
- [15] B. Helffer, J. Sjöstrand, Résonances en limite semi-classique, *Mém. Soc. Math. France* 24–25 (1986).
- [16] B. Helffer, J. Sjöstrand, Multiple wells in the semi-classical limit I, *Comm. Partial Differential Equations* 9 (4) (1984) 337–408.
- [17] B. Helffer, J. Sjöstrand, Puits Multiples en limite semi-classique II. Interaction moléculaire. Symétries. Perturbation, *Ann. Inst. H. Poincaré Phys. Théor.* 42 (2) (1985) 127–212.
- [18] B. Helffer, J. Sjöstrand, Analyse semiclassique pour l'équation de Harper, *Mém. Soc. Math. France* 34 (1988).
- [19] P.D. Hislop, I.M. Sigal, *Introduction to Spectral Theory with Applications to Schrödinger Operators*, Applied Mathematical Sciences, vol. 113, Springer-Verlag, New York, 1996.
- [20] V. Jakšić, C.-A. Pillet, Non-equilibrium steady states of finite quantum systems coupled to thermal reservoirs, *Commun. Math. Phys.* 226 (1) (2002) 131–162.
- [21] G. Jona-Lasinio, C. Presilla, J. Sjöstrand, On Schrödinger equations with concentrated nonlinearities, *Ann. Phys.* 240 (1) (1995) 1–21.
- [22] J. Kastrup, R. Klann, H. Grahn, K. Ploog, L. Bonilla, J. Galàn, M. Kindelan, M. Moscoso, R. Merlin, Self-oscillations of domains in doped GaAs–Al–As superlattices, *Phys. Rev. B* 52 19 (1995) 13761–13764.
- [23] R. Landauer, Spatial variation of currents and fields due to localized scatterers in metallic conduction, *IBM J. Res. Develop.* 1 (1957) 223–231.
- [24] F. Nier, A variational formulation of Schrödinger–Poisson systems in dimension  $d \leq 3$ , *Comm. Partial Differential Equations* 18 (7–8) (1993) 1125–1147.
- [25] F. Nier, Schrödinger–Poisson systems in dimension  $d \leq 3$ : the whole-space case, *Proc. Roy. Soc. Edinburgh Sect. A* 123 (6) (1993) 1179–1201.
- [26] F. Nier, The dynamics of some quantum open systems with short-range nonlinearities, *Nonlinearity* 11 (4) (1998) 1127–1172.
- [27] F. Nier, M. Patel, Nonlinear asymptotics for quantum out-of-equilibrium 1D systems: reduced models and algorithms, in: Blanchard, Dell'Antonio (Eds.), *Multiscale Methods in Quantum Mechanics: Theory and Experiment*, Birkhäuser, 2004, pp. 99–111.
- [28] M. Patel, Développement de modèles macroscopiques pour des systèmes quantiques non-linéaires hors-équilibre, Ph.D. Thesis, Université de Rennes 1, 2005.
- [29] C. Presilla, J. Sjöstrand, Transport properties in resonant tunneling heterostructures, *J. Math. Phys.* 37 (10) (1996) 4816–4844.
- [30] B. Simon, *Trace Ideals and Their Applications*, London Mathematical Society Lecture Note Series, vol. 35, Cambridge University Press, 1979.
- [31] J. Sjöstrand, M. Zworski, Elementary linear algebra for advanced spectral problems, <http://math.berkeley.edu/~zworsky/>, December 2003.
- [32] D. Yafaev, *Mathematical Scattering Theory, General Theory*, Translation of Mathematical Monographs, vol. 105, Amer. Math. Soc., 1992.