



Partial Differential Equations

A singular asymptotic behavior of a transport equation

Philippe Michel

DMI, Institut Camille-Jordan, École centrale de Lyon, 36, avenue Guy-de-Collongue, 69134 Ecully cedex, France

Received 24 August 2007; accepted after revision 11 December 2007

Available online 14 January 2008

Presented by Jean-Michel Bony

Abstract

We consider a simple conservative transport equation where the speed is strictly decreasing. The monotonicity property of the speed rate leads to a singular asymptotic behavior and the concentration of the mass of the solution at a point. Thus, a model which contains a transport structure with monotone decay of the speed rate can be reduced by using the result of convergence to a Dirac mass. It is useful in the case where we have to simulate numerous nonlinear PDEs containing such a structure. Indeed, the concentration of the mass makes the variable in which the mass concentrate useless and thus we lose a dimension. The gain in time calculus is important when the number of equations is large. *To cite this article: P. Michel, C. R. Acad. Sci. Paris, Ser. I 346 (2008).*

© 2007 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Résumé

États asymptotiques singuliers pour certaines équations de transports. On s'intéresse à des équations de transport dans lesquelles la vitesse de transport est strictement décroissante. Cette propriété de monotonie entraîne la concentration de la masse de la solution en un point et, par conséquent, un comportement asymptotique singulier. D'autre part, il est possible de réduire un modèle contenant une structure de transport avec la condition de monotonie (dans une direction) de la vitesse de transport. En utilisant la convergence de la solution en une masse de Dirac suivant la direction du transport, on rend cette direction inutile et on peut éliminer une variable d'espace. Cela réduit le coût numérique lorsqu'il faut simuler un certains nombres d'EDP contenant une telle structure. Le gain est d'autant plus important que le nombre d'équations est grand. *Pour citer cet article : P. Michel, C. R. Acad. Sci. Paris, Ser. I 346 (2008).*

© 2007 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Version française abrégée

Cette Note est consacré à l'étude du comportement asymptotique de solutions d'équations de transport dans lesquelles la vitesse de transport est décroissante. Plus précisément, pour n solution positive de

$$\frac{\partial}{\partial t}n + \nabla(Vn) = 0, \tag{1}$$

avec V régulière et strictement monotone,

E-mail address: philippe.michel@ec-lyon.fr.

$$\langle V(t, x) - V(t, y), x - y \rangle < -\epsilon \|x - y\|^2, \quad \forall t \geq 0, \forall x \neq y \in \mathbb{R}^m, \quad (2)$$

avec $\epsilon > 0$. Toutes les solutions de (1) ont le même type de comportement asymptotique décrit par les solutions “Dirac” le long des trajectoires

$$\frac{\partial}{\partial t} z = V(t, z). \quad (3)$$

Proposition 0.1. *Supposons que (2) soit vérifié et soit n solution positive de (1) avec $\int n(0, x) dx = 1$ alors la distance de Wasserstein entre $n dx$ et $\delta_{x=z(t)}$ tends vers 0 exponentiellement*

$$d_{W_2}(n(t, x) dx, \delta_{x=z(t)}) \leq C e^{-\epsilon t}. \quad (4)$$

Cette proposition peut être étendue à des équations de renouvellement contenant une telle structure de transport comme par exemple

$$\begin{cases} \frac{\partial}{\partial t} n + \frac{\partial}{\partial a} n + \frac{\partial}{\partial x} (Vn) = 0, \\ n(t, 0, x) = 2n(t, 1, x). \end{cases} \quad (5)$$

Dans ce cas, on a la suivante :

Proposition 0.2. *Si (2) est satisfaite et n solution positive de (5) avec $\int n(0, a, x) 2^a dx = 1$ alors on a*

$$\begin{aligned} \int n(t, a, x) 2^{a-t} da dx &= 1, \quad \forall t \geq 0, \\ \frac{d}{dt} \int n(t, a, x) 2^{a-t} |x - z(t, a)|^2 da dx &= 2 \int n 2^{a-t} \langle V(t, x) - V(t, z), x - z \rangle da dx < 0, \end{aligned} \quad (6)$$

avec z solution d’une équation de transport avec condition aux bords périodique. Plus précisément, la quantité $\int n(t, a, x) 2^{a-t} |x - z(t, a)|^2 da dx$ converge exponentiellement vers 0.

Cela implique qu’il y a concentration de la masse autour de z et par conséquent, en temps long, on peut s’affranchir de la variable x . On peut considérer que $n \sim \delta_{x=z(t)} m(t, a)$ avec m solution de l’équation de renouvellement

$$\begin{cases} \frac{\partial}{\partial t} m + \frac{\partial}{\partial a} m = 0, \\ m(t, 0) = 2m(t, 1). \end{cases} \quad (7)$$

La perte d’une variable d’espace rend le système plus facile à implémenter et le gain en temps de calcul est appréciable lorsqu’il faut simuler un grand nombre d’EDP non linéaires couplées (voir [7]).

1. Introduction

In this Note, we are interested in the asymptotic behavior of some transport equations when the transport speed is monotonic. More precisely, we consider n positive solution to the conservative transport equation (see [1,4–6])

$$\frac{\partial}{\partial t} n + \nabla(Vn) = 0, \quad (8)$$

where $V \in C^1([0, \infty) \times \mathbb{R}^m, \mathbb{R}^m)$ satisfies the monotonicity condition

$$\langle V(t, x) - V(t, y), x - y \rangle < -\epsilon \|x - y\|^2, \quad \forall t \geq 0, \forall x \neq y \in \mathbb{R}^m, \quad (9)$$

where ϵ is strictly positive. We show that the limit of $n(t, \cdot)$ when $t \rightarrow \infty$ is a Dirac measure supported by the trajectories of the characteristic equation (independently of its initial data)

$$\frac{\partial}{\partial t} z = V(t, z). \quad (10)$$

To evaluate how $n(t, \cdot)$ converges to this singular limit, we introduce the Wasserstein distance (see [15]) between measures

$$d_{W_p}(\mu, \nu) := \left(\inf_{\eta \in \Gamma(\mu, \nu)} \left\{ \iint_{(-\infty, \infty)^2} |x - y|^p \, d\eta(x, y) \right\} \right)^{1/p}, \tag{11}$$

where $\Gamma(\mu, \nu)$ denotes the collection of all measures on $(-\infty, \infty)^2$ with marginals μ and ν . We notice that

$$d_{W_p}(\mu, \nu) \leq \left(\iint_{(-\infty, \infty)^2} |x - y|^p \, d\mu(x) \, d\nu(y) \right)^{1/p}. \tag{12}$$

Then we extend the convergence result to a singular limit, to a renewal equation of McKendrick–Von Foerster type. In this context, we use this result to reduce the dimension in a renewal model [7].

2. Transport equation: main result

We present here the main result of convergence of n to a Dirac mass:

Proposition 2.1. *Assume (9) and n positive solution to (8) with $\int n(0, x) \, dx = 1$, then the Wasserstein distance between $n \, dx$ and $\delta_{x=z(t)}$ vanishes*

$$\lim_{t \rightarrow \infty} d_{W_2}(n(t, x) \, dx, \delta_{x=z(t)}) = 0. \tag{13}$$

Proof. Under the assumption (9), we find directly that

$$\int n(t, x) \, dx = \int n(0, x) \, dx = 1, \tag{14}$$

$$\frac{d}{dt} \int n(t, x) \|x - z(t)\|^2 \, dx = 2 \int n(t, x) \langle V(t, x) - V(t, z), x - z \rangle \, dx < 0. \tag{15}$$

Since V is strictly decreasing, there exists $\epsilon > 0$ such that,

$$\langle V(t, x) - V(t, z), x - z \rangle \leq -\epsilon \|x - z(t)\|^2.$$

Therefore, we have

$$\frac{d}{dt} \int n(t, x) \|x - z(t)\|^2 \, dx \leq -2\epsilon \int n(t, x) \|x - z(t)\|^2 \, dx, \tag{16}$$

and using Gronwall inequality, we have exponential convergence to 0,

$$\lim_{t \rightarrow \infty} \int n(t, x) \|x - z(t)\|^2 \, dx = 0.$$

Using (12) for $p = 2$ and (16) we prove that the Wasserstein distance between $n \, dx$ and $\delta_{x=z(t)}$ converges to 0 when $t \rightarrow \infty$. \square

We notice that, in dimension one ($m = 1$), we easily see the convergence of n to a Dirac mass. Indeed, in this case, we have directly that

$$\frac{d}{dt} \int n(t, x) (x - z(t))_+^2 \, dx = 2 \int n 1_{x \geq z(t)} (V(t, x) - V(t, z(t)), x - z(t)) \, dx < 0, \tag{17}$$

and thus the support of $n(t, \cdot)$ must decrease. More precisely, under the assumption on the support of $n(0, \cdot)$

$$\text{Supp } n(0, \cdot) \subset [x_0, x_1], \tag{18}$$

we have the following lemma:

Lemma 2.2. *Assume (18) and let w and z be a solution to (10) with the initial condition $w(0) = x_0$ and $z(0) = x_1$. Then, under the monotonicity condition (9), we find that the support of $n(t, \cdot)$ belongs to the interval $[w(t), z(t)]$ which length converges to 0,*

$$\text{Supp } n(t, \cdot) \subset [w(t), z(t)], \tag{19}$$

and $\lim_{t \rightarrow \infty} (w(t) - z(t))^2 = 0$.

3. Application to renewal equations

We apply this result to a renewal equation (see [2,3,7,8,14,12,17])

$$\begin{cases} \frac{\partial}{\partial t} n + \frac{\partial}{\partial a} n + \frac{\partial}{\partial x} (Vn) = 0, \\ n(t, 0, x) = 2n(t, 1, x). \end{cases} \tag{20}$$

We notice that the monotone decay assumption (9) is satisfied in [7] and the main equation (20) is similar to the Vlasov equation with friction [9,10]. In order to extend the result of Proposition 2.1, we define z solution to the transport equation

$$\frac{\partial}{\partial t} z + \frac{\partial}{\partial a} z = V(t, z), \tag{21}$$

with periodic boundary condition

$$z(t, 0) = z(t, 1). \tag{22}$$

Here, two difficulties appear. The first one is that n is no longer conservative and according to ideas developed in [12,13,16] we have to use ψ solution to the backward equation ($n\psi$ is conservative)

$$\begin{cases} \frac{\partial}{\partial t} \psi + \frac{\partial}{\partial a} \psi + V \frac{\partial}{\partial x} \psi = 0, \\ \psi(t, 0) = \frac{1}{2} \psi(t, 1). \end{cases} \tag{23}$$

The second difficulty is that we can only conclude that $n\psi$ concentrate its mass in $x = z(t, a)$ and $n \sim m(t, a) \delta_{x=z(t,a)}$, thus we have to study $m(t, a)$ separately.

Proposition 3.1. *Assume (9) and n positive solution to (20) where $\int n(0, a, x) 2^{a-t} da dx = 1$ then we have*

$$\begin{aligned} \int n(t, a, x) 2^{a-t} da dx &= 1, \quad \forall t \geq 0, \\ \frac{d}{dt} \int n(t, a, x) 2^{a-t} \|x - z(t, a)\|_2^2 da dx &= 2 \int n 2^{a-t} (V(t, x) - V(t, z), x - z) da dx < 0, \end{aligned} \tag{24}$$

and

$$\begin{cases} \frac{\partial}{\partial t} \int n dx + \frac{\partial}{\partial a} \int n dx = 0, \\ \int n(t, 0, x) dx = 2 \int n(t, 1, x) dx. \end{cases} \tag{25}$$

Proof. It suffices to notice that $\psi(t, a) = 2^{a-t}$ is solution to (23). Indeed, in this case, we have

$$\begin{cases} \frac{\partial}{\partial t} n\psi + \frac{\partial}{\partial a} n\psi + \frac{\partial}{\partial x} Vn\psi = 0, \\ n(t, 0, x)\psi(t, 0, x) = n(t, 1, x)\psi(t, 1, x), \end{cases} \tag{26}$$

and using (21) and (22), we find

$$\begin{cases} \frac{\partial}{\partial t} n\psi(x - z)^2 + \frac{\partial}{\partial a} n\psi(x - z)^2 + \frac{\partial}{\partial x} Vn(x - z)^2\psi = 2n(V(t, x) - V(t, z), x - z), \\ n(t, 0, x)\psi(t, 0, x)(x - z(t, 0))^2 = n(t, 1, x)\psi(t, 1)(x - z(t, 1))^2. \end{cases} \tag{27}$$

Then, integrating (26) with respect to a and x , we find that $\int n(t, a, x)\psi da dx$ is a constant. Now, integrating (27) with respect to a and x , we find (24) and we conclude to the convergence of n to a Dirac mass. \square

Thus, we can reduce the PDE (20) to two equations (21) and (25) with less variables. Indeed, the numerical simulation of (20) requires the computation of P^2 discretization points, while the numerical simulation of (21) and (25) need at most $2P$ points. An example of application is given by [7]. There a system of N renewal PDEs (similar to (20)) is considered and the gain due to the reduction of the model in $2N$ equations with less variables is important. This system modelizes the growth of follicles structured in age and maturity with the maturity speed rate which satisfies the condition (9). The simplified model we obtain using Proposition 3.1 is easier to compute numerically and can be compared to the classical model of Lacker [11].

References

- [1] F. Bolley, C. Villani, Weighted Csiszar–Kullback–Pinsker inequalities and application to transport equation inequalities, *Ann. Fac. Sci. Toulouse Ser. 6* 14 (3) (2005) 331–352.
- [2] J.M. Cushing, *An Introduction to Structured Population Dynamics*, SIAM, Philadelphia, 1998.
- [3] O. Diekmann, M. Gyllenberg, J.A.J. Metz, Steady state analysis of structured population models, *Theor. Pop. Biol.* 63 (2003) 309–338.
- [4] R.J. DiPerna, Measure valued solutions to conservation laws, *Arch. Rational Mech. Anal.* 88 (1985) 223–270.
- [5] R.J. DiPerna, P.L. Lions, Ordinary differential equations, transport theory and Sobolev spaces, *Invent. Math.* 98 (1989) 511–547.
- [6] R.J. DiPerna, P.L. Lions, Solutions globales d'équations du type Vlasov–Poisson, *C. R. Acad. Sci. Paris, Ser. I* 307 (1988) 655–658.
- [7] N. Echenim, D. Monniaux, M. Sorine, F. Clement, Multi-scale modeling of the follicle selection process in the ovary, *Math. Biosci.* 198 (1) (2005) 57–79.
- [8] M. Iannelli, Age-structured population, in: M. Hazewinkel (Ed.), *Encyclopedia of Mathematics*, supplement II, Kluwer Academics, 2000, pp. 21–23.
- [9] P.E. Jabin, Macroscopic limit of Vlasov type equations with friction, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 17 (2000) 651–672.
- [10] P.E. Jabin, Large time concentration for solutions to kinetic equations with energy dissipation, *Comm. Partial Differential Equations* 25 (2000) 541–557.
- [11] H.M. Lacker, A. Percus, How do ovarian follicles interact? A many body problem with unusual symmetry and symmetry-breaking properties, *J. Stat. Phys.* 63 (1991) 1133–1161.
- [12] P. Michel, S. Mischler, B. Perthame, General relative entropy inequality: an illustration on growth models, *J. Math. Pures Appl.* 84 (9) (2005) 1235–1260.
- [13] S. Mischler, B. Perthame, L. Ryzhik, Stability in a nonlinear population maturation model, *Math. Models Meth. Appl. Sci.* 12 (12) (2002) 1751–1772.
- [14] J.A.J. Metz, O. Diekmann, *The Dynamics of Physiologically Structured Populations*, Lecture Notes in Biomathematics, vol. 68, Springer-Verlag, 1986.
- [15] F. Otto, M. Westdickenberg, Eulerian calculus for the contraction in the Wasserstein distance, *SIAM J. Math. Anal.* 37 (2005) 1227–1255.
- [16] B. Perthame, *Transport Equations in Biology*, *Frontiers in Mathematics*, Birkhäuser, 2007.
- [17] M. Rotenberg, Transport theory for growing cell populations, *J. Theor. Biol.* 103 (1983) 181–199.