

Stationary distributions for jump processes with memory

K. Burdzy^{a,1}, T. Kulczycki^{b,c,2} and R. L. Schilling^{d,3}

^a*Department of Mathematics, Box 354350, University of Washington, Seattle, WA 98195, USA. E-mail: burdzy@math.washington.edu*

^b*Institute of Mathematics, Polish Academy of Sciences, ul. Kopernika 18, 51-617 Wrocław, Poland. E-mail: t.kulczycki@impan.pl*

^c*Institute of Mathematics and Computer Science, Wrocław University of Technology, Wybrzeże Wyspińskiego 27, 50-370 Wrocław, Poland*

^d*Institut für Mathematische Stochastik, TU Dresden, D-01062 Dresden, Germany. E-mail: rene.schilling@tu-dresden.de*

Received 13 September 2010; revised 1 April 2011; accepted 7 April 2011

Abstract. We analyze a jump processes Z with a jump measure determined by a “memory” process S . The state space of (Z, S) is the Cartesian product of the unit circle and the real line. We prove that the stationary distribution of (Z, S) is the product of the uniform probability measure and a Gaussian distribution.

Résumé. Nous proposons d'étudier un processus à sauts Z avec une mesure de sauts déterminée par un processus S représentant une “mémoire”. L'espace d'états de (Z, S) est le produit Cartésien du cercle trigonométrique et de l'axe réel. Nous démontrons que la distribution stationnaire de (Z, S) est la mesure produit d'une loi uniforme et d'une loi Gaussienne.

MSC: Primary 60J35; secondary 60H10; 60G51; 60J75; 60J55

Keywords: Stationary distribution; Stable Lévy process; Process with memory

1. Introduction

We are going to find stationary distributions for processes with jumps influenced by “memory.” This paper is a companion to [3]. The introduction to that paper contains a review of various sources of inspiration for this project, related models and results.

We will analyze a pair of real-valued processes (Y, S) such that S is a “memory” in the sense that $dS_t = W(Y_t) dt$ where W is a C^3 function. The process Y is a jump process “mostly” driven by a stable process but the process S affects the rate of jumps of Y . We refer the reader to Section 2 for a formal presentation of this model as it is too long for the introduction. The present article illustrates advantages of semi-discrete models introduced in [5] since the form of the stationary distribution for (Y, S) was conjectured in [5], Example 3.8. We would not find it easy to conjecture the stationary distribution for this process in a direct way.

The main result of this paper, i.e. Theorem 3.7, is concerned with the stationary distribution of a transformation of (Y, S) . In order to obtain non-trivial results, we “wrap” Y on the unit circle, so that the state space for the transformed process is compact. In other words, we consider $(Z_t, S_t) = (e^{iY_t}, S_t)$. The stationary distribution for (Z_t, S_t) is the product of the uniform distribution on the circle and the normal distribution.

The Gaussian distribution of the “memory” process appeared in models discussed in [2,3]. In each of those papers, memory processes similar to S effectively represented “inert drift.” A heuristic argument given in the introduction

¹Supported in part by NSF Grant DMS-09-06743 and by grant N N201 397137, MNiSW, Poland.

²Supported in part by grant N N201 373136, MNiSW, Poland.

³Supported in part by DFG grant Schi 419/5-1.

Table 1
Frequently used symbols

$a \vee b, a \wedge b$	$\max(a, b), \min(a, b)$
a_+, a_-	$\max(a, 0), -\min(a, 0)$
$ x _{\ell^1}$	$\sum_{j=1}^m x_j $ where $x = (x_1, \dots, x_m) \in \mathbb{R}^m$
e_k	the k th unit base vector in the usual orthonormal basis for \mathbb{R}^n
\mathcal{A}_α	$\alpha \Gamma(\frac{1+\alpha}{2}) \frac{2^{\alpha-1}}{\sqrt{\pi} \Gamma(1-\alpha/2)}, \alpha \in (0, 2)$
D^α	$\frac{\partial^{ \alpha }}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}, \alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}_0^d$
C^k	k -times continuously differentiable functions
C_b^k, C_c^k, C_0^k	functions in C^k which, together with all their derivatives up to order k , are “bounded,” are “compactly supported,” and “vanish at infinity,” respectively
$C_*(\mathbb{R}^2)$	all bounded and uniformly continuous functions $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $\text{supp}(f) \subset \mathbb{R} \times [-N, N]$ for some $N > 0$
$C_*(\mathbb{R}^2)$	$C_*(\mathbb{R}^2) \cap C_b^2(\mathbb{R}^2)$
\mathbb{S}	$\{z \in \mathbb{C}: z = 1\}$ unit circle in \mathbb{C}

to [3] provides a justification for the Gaussian distribution, using the concepts of kinetic energy associated to drift and Gibbs measure. The conceptual novelty of the present paper is that the Gaussian distribution of S in the stationary regime cannot be explained by kinetic energy because S affects the jump distribution and not the drift of Z .

The product form of the stationary distribution for a two-component Markov process is obvious if the two components are independent Markov processes. The product form is far from obvious if the components are not independent but it does appear in a number of contexts, from queuing theory to mathematical physics. The paper [5] was an attempt to understand this phenomenon for a class of models. The unexpected appearance of the Gaussian distribution in some stationary measures was noticed in [4] before it was explored more deeply in [2,5].

We turn to the technical aspects of the paper. The main effort is directed at determining the domain and a core of the generator of the process. A part of the argument is based on an estimate of the smoothness of the stochastic flow of solutions to (2.3).

1.1. Notation

Since the paper uses a large amount of notation, we collect some of the most frequently used symbols in Table 1, for easy reference. Constants c without sub- or superscript are generic and may change their value from line to line.

2. The construction of the process and its generator

Let $\mathbb{S} = \{z \in \mathbb{C}: |z| = 1\}$ be the unit circle in \mathbb{C} . Consider a C^3 function $V: \mathbb{S} \rightarrow \mathbb{R}$ such that $\int_{\mathbb{S}} V(z) dz = 0$ and set $W(x) = V(e^{ix}), x \in \mathbb{R}$. Assume that V is not identically constant. In this paper we will be interested in the Markov process (Y_t, S_t) with state space \mathbb{R}^2 and generator $\mathcal{G}^{(Y,S)}$ of the following form

$$\mathcal{G}^{(Y,S)} f(y, s) = -(-\Delta_y)^{\alpha/2} f(y, s) + Rf(y, s) + W(y) f_s(y, s), \quad (2.1)$$

with a domain that will be specified later. Here, $(y, s) \in \mathbb{R}^2, \alpha \in (0, 2)$ and

$$\begin{aligned} -(-\Delta_y)^{\alpha/2} f(y, s) &= \mathcal{A}_\alpha \lim_{\varepsilon \rightarrow 0^+} \int_{|y-x| > \varepsilon} \frac{f(x, s) - f(y, s)}{|y-x|^{1+\alpha}} dx, \\ Rf(y, s) &= \int_{-\pi+y}^{\pi+y} (f(x, s) - f(y, s)) ((W(y) - W(x))s)_+ dx. \end{aligned} \quad (2.2)$$

Since $-(-\Delta)^{\alpha/2}, \alpha \in (0, 2)$, is the generator of the symmetric α -stable process on \mathbb{R} , we may think of the process Y_t as the perturbed symmetric α -stable process and S_t as the memory which changes the jumping measure of the process Y_t .

The definition of (Y, S) is informal. Below we will construct this process in a direct way and we will show that this process has the generator (2.1); see Proposition 2.4. Our construction is based on the so-called *construction of Meyer*; see, e.g., [7,8] or [1], Section 3.1.

For any $(y, s) \in \mathbb{R}^2$ let

$$g(y, s, x) = ((W(y) - W(y+x))s)_+ \mathbb{1}_{(-\pi, \pi)}(x), \quad x \in \mathbb{R},$$

and

$$\|g(y, s, \cdot)\|_1 = \int_{-\pi}^{\pi} ((W(y) - W(y+x))s)_+ dx.$$

Let $\bar{g}(y, s, x) := g(y, s, x)/\|g(y, s, \cdot)\|_1$ if $\|g(y, s, \cdot)\|_1 \neq 0$. We let $\bar{g}(y, s, \cdot)$ be the delta function at 0 when $\|g(y, s, \cdot)\|_1 = 0$. If $\|g(y, s, \cdot)\|_1 \neq 0$, we let $F_{y,s}(\cdot)$ denote the cumulative distribution function of a random variable with density $\bar{g}(y, s, \cdot)$. If $\|g(y, s, \cdot)\|_1 = 0$, we let $F_{y,s}(\cdot)$ denote the cumulative distribution function of a random variable that is identically equal to 0. We have

$$F_{y,s}^{-1}(v) = \inf \left\{ x \in \mathbb{R} : \int_{-\infty}^x \frac{g(y, s, z)}{\|g(y, s, \cdot)\|_1} dz \geq v \right\}$$

so for any v , the function $(y, s) \rightarrow F_{y,s}^{-1}(v)$ is measurable. If \mathcal{U} is a uniformly distributed random variable on $(0, 1)$, then $F_{y,s}^{-1}(\mathcal{U})$ has the density $\bar{g}(y, s, \cdot)$. Let $(\mathcal{U}_n)_{n \in \mathbb{N}}$ be countably many independent copies of \mathcal{U} and set $\eta_n(y, s) = F_{y,s}^{-1}(\mathcal{U}_n)$.

Let $X(t)$ be a symmetric α -stable process on \mathbb{R} , $\alpha \in (0, 2)$, starting from 0 and $N(t)$ a Poisson process with intensity 1. We assume that $(\mathcal{U}_n)_{n \in \mathbb{N}}$, $X(\cdot)$ and $N(\cdot)$ are independent.

Let $0 < \sigma_1 < \sigma_2 < \dots$ be the times of jumps of $N(t)$. Consider any $y, s \in \mathbb{R}$ and for $t \geq 0$ let

$$\begin{aligned} Y_t^1 &= y + X_t, \\ S_t^1 &= s + \int_0^t W(Y_r^1) dr, \\ \widehat{\sigma}_1(t) &= \int_0^t \|g(Y_r^1, S_r^1, \cdot)\|_1 dr, \\ \tau_1 &= \inf_{t \geq 0} \{\widehat{\sigma}_1(t) = \sigma_1\} \quad (\inf \emptyset = \infty). \end{aligned}$$

Now we proceed recursively. If $Y_t^j, S_t^j, \widehat{\sigma}_j(t)$ are well defined on $[0, \tau_j)$ and $\tau_j < \infty$ then we define for $t \geq \tau_j$,

$$\begin{aligned} Y_t^{j+1} &= y + X_t + \sum_{n=1}^j \eta_n(Y^n(\tau_n-), S^n(\tau_n-)), \\ S_t^{j+1} &= s + S^j(\tau_j-) + \int_{\tau_j-}^t W(Y_r^{j+1}) dr, \\ \widehat{\sigma}_{j+1}(t) &= \tau_j + \int_{\tau_j}^t \|g(Y_r^{j+1}, S_r^{j+1}, \cdot)\|_1 dr, \\ \tau_{j+1} &= \inf_{t \geq \tau_j} \{\widehat{\sigma}_{j+1}(t) = \sigma_{j+1}\}. \end{aligned}$$

Let $\tau_0 = 0$ $(Y_t, S_t) = (Y_t^j, S_t^j)$ for $\tau_{j-1} \leq t < \tau_j$, $j \geq 1$. It is easy to see that (Y_t, S_t) is defined for all $t \geq 0$, a.s. If we put $\sigma(t) = \int_0^t \|g(Y_r, S_r, \cdot)\|_1 dr$ then we can represent (Y_t, S_t) by the following closed-form expression,

$$\begin{cases} Y_t = y + X_t + \sum_{n=1}^{N(\sigma(t))} \eta_n(Y(\tau_n-), S(\tau_n-)), \\ S_t = s + \int_0^t W(Y_r) dr. \end{cases} \quad (2.3)$$

We define the semigroup $\{T_t\}_{t \geq 0}$ of the process (Y_t, S_t) for $f \in C_b(\mathbb{R}^2)$ by

$$T_t f(y, s) = \mathbb{E}^{(y, s)} f(Y_t, S_t), \quad (y, s) \in \mathbb{R}^2.$$

By $\mathcal{G}^{(Y, S)}$ we denote the generator of $\{T_t\}_{t \geq 0}$ and its domain by $\mathcal{D}(\mathcal{G}^{(Y, S)})$. We will show in Proposition 2.4 that $C_*^2(\mathbb{R}^2) \subset \mathcal{D}(\mathcal{G}^{(Y, S)})$ and that $\mathcal{G}^{(Y, S)} f$ is given by (2.1) for $f \in C_*^2(\mathbb{R}^2)$, see Section 1.1 for the definition of $C_*^2(\mathbb{R}^2)$.

Our construction of (Y_t, S_t) is a deterministic map

$$\{(\mathcal{U}_n)_{n \in \mathbb{N}}, (N(t))_{t \geq 0}, (X(t))_{t \geq 0}\} \rightarrow \{(Y(t))_{t \geq 0}, (S(t))_{t \geq 0}\}.$$

This easily implies the strong Markov property for (Y, S) . We will verify that $(Z_t, S_t) := (e^{iY_t}, S_t)$ is also a strong Markov process. We first show that the transition function of (Y_t, S_t) is periodic.

Lemma 2.1. *Let (Y_t, S_t) be the Markov process defined by (2.3). Then*

$$\mathbb{P}^{(y+2\pi, s)}(Y_t \in A + 2\pi, S_t \in B) = \mathbb{P}^{(y, s)}(Y_t \in A, S_t \in B)$$

for all $(y, s) \in \mathbb{R}^2$ and all Borel sets $A, B \subset \mathbb{R}$.

Proof. Let X_t be a symmetric α -stable process, starting from 0, $\alpha \in (0, 2)$, and let $N(t)$ be a Poisson process with intensity 1. By (Y_t^y, S_t^s) we denote the process given by (2.3) with initial value $(Y_0^y, S_0^s) = (y, s)$. The process $(\tilde{Y}_t, \tilde{S}_t) := (Y_t^{y+2\pi}, S_t^s)$ has the following representation

$$\begin{aligned} \tilde{Y}_t &= y + 2\pi + X_t + \sum_{n=1}^{N(\tilde{\sigma}(t))} \eta_n(\tilde{Y}(\tilde{\tau}_n -), \tilde{S}(\tilde{\tau}_n -)), \\ \tilde{S}_t &= s + \int_0^t W(\tilde{Y}_r) dr, \end{aligned}$$

where $\tilde{\sigma}(t) = \int_0^t \|g(\tilde{Y}_r, \tilde{S}_r, \cdot)\|_1 dr$ and $\tilde{\tau}_k = \inf_{t \geq 0} \{\tilde{\sigma}(t) = \sigma_k\}$.

Note that for all $x \in \mathbb{R}$,

$$g(y - 2\pi, s, x) = g(y, s, x) \quad \text{and, therefore,} \quad \|g(y - 2\pi, s, \cdot)\|_1 = \|g(y, s, \cdot)\|_1.$$

It follows that $\eta_n(y - 2\pi, s)$ has the same distribution as $\eta_n(y, s)$. Since the function W is periodic with period 2π , we have $W(\tilde{Y}_r) = W(\tilde{Y}_r - 2\pi)$. Moreover, $\|g(\tilde{Y}_r, \tilde{S}_r, \cdot)\|_1 = \|g(\tilde{Y}_r - 2\pi, \tilde{S}_r, \cdot)\|_1$ and $\eta_n(\tilde{Y}(\tilde{\tau}_n -), \tilde{S}(\tilde{\tau}_n -))$ has the same distribution as $\eta_n(\tilde{Y}(\tilde{\tau}_n -) - 2\pi, \tilde{S}(\tilde{\tau}_n -))$. This means that we can rewrite the representation of $(Y_t^{y+2\pi}, S_t^s)$ in the following way:

$$\begin{aligned} \tilde{Y}_t &= y + 2\pi + X_t + \sum_{n=1}^{N(\tilde{\sigma}(t))} \eta_n(\tilde{Y}(\tilde{\tau}_n -) - 2\pi, \tilde{S}(\tilde{\tau}_n -)), \\ \tilde{S}_t &= s + \int_0^t W(\tilde{Y}_r - 2\pi) dr, \end{aligned}$$

where $\tilde{\sigma}(t) = \int_0^t \|g(\tilde{Y}_r - 2\pi, \tilde{S}_r, \cdot)\|_1 dr$ and $\tilde{\tau}_k = \inf_{t \geq 0} \{\tilde{\sigma}(t) = \sigma_k\}$.

By subtracting 2π from both sides of the first equation we get

$$\begin{aligned} \tilde{Y}_t - 2\pi &= y + X_t + \sum_{n=1}^{N(\tilde{\sigma}(t))} \eta_n(\tilde{Y}(\tilde{\tau}_n -) - 2\pi, \tilde{S}(\tilde{\tau}_n -)), \\ \tilde{S}_t &= s + \int_0^t W(\tilde{Y}_r - 2\pi) dr, \end{aligned}$$

Table 2
Notation for processes and their generators

Process	Semigroup	Generator and domain
(Y_t, S_t)	$T_t, t \geq 0$	$(\mathcal{G}^{(Y,S)}, \mathcal{D}(\mathcal{G}^{(Y,S)}))$
$(Z_t, S_t) = (e^{iY_t}, S_t)$	$T_t^{\mathbb{S}}, t \geq 0$	$(\mathcal{G}, \mathcal{D}(\mathcal{G}))$
$(\hat{Y}_t, \hat{S}_t) = (\hat{Y}_0 + X_t, \hat{S}_0 + \int_0^t W(\hat{Y}_r) dr)$	$\hat{T}_t, t \geq 0$	$(\mathcal{G}^{(\hat{Y}, \hat{S})}, \mathcal{D}(\mathcal{G}^{(\hat{Y}, \hat{S})}))$
$(\hat{Z}_t, \hat{S}_t) = (e^{i\hat{Y}_t}, \hat{S}_t)$	$\hat{T}_t^{\mathbb{S}}, t \geq 0$	$(\hat{\mathcal{G}}, \mathcal{D}(\hat{\mathcal{G}}))$

with $\tilde{\sigma}(t)$ and $\tilde{\tau}_k$ as before. Substituting $\hat{Y}_t := \tilde{Y}_t - 2\pi$ we see that this is the defining system of equations for the process (Y_t^y, S_t^s) . Therefore, the processes (Y_t^y, S_t^s) and $(Y_t^{y+2\pi}, S_t^s)$ have the same law. \square

We can now argue exactly as in [3], Corollary 2.3, to see that $(Z_t, S_t) = (e^{iY_t}, S_t)$ is indeed a strong Markov process. We define the transition semigroup of (Z_t, S_t) for $f \in C_0(\mathbb{S} \times \mathbb{R})$ by

$$T_t^{\mathbb{S}} f(z, s) = \mathbb{E}^{(z,s)} f(Z_t, S_t), \quad (z, s) \in \mathbb{S} \times \mathbb{R}. \tag{2.4}$$

The generator of $\{T_t^{\mathbb{S}}\}_{t \geq 0}$ and its domain will be denoted \mathcal{G} and $\mathcal{D}(\mathcal{G})$.

In the sequel we will need the following auxiliary processes

$$\begin{aligned} \hat{Y}_t &= \hat{Y}_0 + X_t, \\ \hat{S}_t &= \hat{S}_0 + \int_0^t W(\hat{Y}_r) dr, \\ \hat{Z}_t &= e^{i\hat{Y}_t}, \end{aligned}$$

where X_t is a symmetric α -stable Lévy process on \mathbb{R} , $\alpha \in (0, 2)$, starting from 0. We will use notation from Table 2.

We will now identify the generators of the processes (Y_t, S_t) and (Z_t, S_t) and link them with the generators of the processes (\hat{Y}_t, \hat{S}_t) and (\hat{Z}_t, \hat{S}_t) .

Proposition 2.2. *Let (Y_t, S_t) be the process defined by (2.3) and let $f \in C_*(\mathbb{R}^2)$. Then*

$$\lim_{t \rightarrow 0^+} \frac{T_t f - f}{t} \text{ exists} \iff \lim_{t \rightarrow 0^+} \frac{\hat{T}_t f - f}{t} \text{ exists,}$$

in the norm $\|\cdot\|_{\infty}$. If one, hence both, limits exist, then

$$\lim_{t \rightarrow 0^+} \frac{T_t f - f}{t} = \lim_{t \rightarrow 0^+} \frac{\hat{T}_t f - f}{t} + Rf, \tag{2.5}$$

where Rf is given by (2.2).

Corollary 2.3. *We have*

$$f \in \mathcal{D}(\mathcal{G}) \cap C_c(\mathbb{S} \times \mathbb{R}) \iff f \in \mathcal{D}(\hat{\mathcal{G}}) \cap C_c(\mathbb{S} \times \mathbb{R}).$$

If $f \in \mathcal{D}(\mathcal{G}) \cap C_c(\mathbb{S} \times \mathbb{R})$ then

$$\mathcal{G}f = \hat{\mathcal{G}}f + R^{\mathbb{S}}f,$$

where

$$R^{\mathbb{S}}f(z, s) = \int_{\mathbb{S}} (f(w, s) - f(z, s))((V(z) - V(w))s)_+ dw.$$

Proposition 2.4. *Let (Y_t, S_t) be the process defined by (2.3). Then $C_*^2(\mathbb{R}^2) \subset \mathcal{D}(\mathcal{G}^{(Y,S)})$ and for $f \in C_*^2(\mathbb{R}^2)$ we have*

$$\mathcal{G}^{(Y,S)} f(y, s) = -(-\Delta_y)^{\alpha/2} f(y, s) + Rf(y, s) + W(y) f_s(y, s) \quad (2.6)$$

for all $(y, s) \in \mathbb{R}^2$ with Rf given by (2.2).

Moreover, $C_*^2(\mathbb{R}^2) \subset \mathcal{D}(\mathcal{G}^{(\hat{Y}, \hat{S})})$ and for $f \in C_*^2(\mathbb{R}^2)$ we have

$$\mathcal{G}^{(\hat{Y}, \hat{S})} f(y, s) = -(-\Delta_y)^{\alpha/2} f(y, s) + W(y) f_s(y, s) \quad (2.7)$$

for all $(y, s) \in \mathbb{R}^2$.

By $\text{Arg}(z)$ we denote the argument of $z \in \mathbb{C}$ contained in $(-\pi, \pi]$. For $g \in C^2(\mathbb{S})$ let us put

$$\begin{aligned} Lg(z) &= \mathcal{A}_\alpha \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{S} \cap \{|\text{Arg}(w/z)| > \varepsilon\}} \frac{g(w) - g(z)}{|\text{Arg}(w/z)|^{1+\alpha}} dw \\ &\quad + \mathcal{A}_\alpha \sum_{n \in \mathbb{Z} \setminus \{0\}} \int_{\mathbb{S}} \frac{g(w) - g(z)}{|\text{Arg}(w/z) + 2n\pi|^{1+\alpha}} dw, \end{aligned} \quad (2.8)$$

where dw denotes the arc length measure on \mathbb{S} ; note that $\int_{\mathbb{S}} dw = 2\pi$. It is clear that for $f \in C_c^2(\mathbb{S} \times \mathbb{R})$, $z = e^{iy}$, $y, s \in \mathbb{R}$ we have

$$-(-\Delta_y)^{\alpha/2} \tilde{f}(y, s) = L_z f(z, s). \quad (2.9)$$

Corollary 2.5. *We have $C_c^2(\mathbb{S} \times \mathbb{R}) \subset \mathcal{D}(\mathcal{G})$ and for $f \in C_c^2(\mathbb{S} \times \mathbb{R})$ we have*

$$\mathcal{G}f(z, s) = L_z f(z, s) + R^{\mathbb{S}} f(z, s) + V(z) f_s(z, s)$$

for all $(z, s) \in \mathbb{S} \times \mathbb{R}$, where L is given by (2.8).

We also have $C_c^2(\mathbb{S} \times \mathbb{R}) \subset \mathcal{D}(\hat{\mathcal{G}})$ and for $f \in C_c^2(\mathbb{S} \times \mathbb{R})$ we have

$$\hat{\mathcal{G}}f(z, s) = L_z f(z, s) + V(z) f_s(z, s)$$

for all $(z, s) \in \mathbb{S} \times \mathbb{R}$.

Remark 2.6. *Proposition 2.4 shows that for $f \in C_*^2(\mathbb{R}^2)$ the generator of the process (Y_t, S_t) defined by (2.3) is of the form (2.1). This is a standard result, the so-called “construction of Meyer,” but we include our own proof of this result so that the paper is self-contained. Moreover, Proposition 2.2, Corollaries 2.3 and 2.5 are needed to identify a core for \mathcal{G} . Corollary 2.5 is also needed to find the stationary measure for (Z_t, S_t) .*

We will need two auxiliary results.

Lemma 2.7. *There exists a constant $c = c(M) > 0$ such that for any $x \in [-\pi, \pi]$ and any $u_1 = (y_1, s_1) \in \mathbb{R}^2$, $u_2 = (y_2, s_2) \in \mathbb{R}^2$ with $s_1, s_2 \in [-M, M]$ we have*

$$|g(u_1, x) - g(u_2, x)| \leq c(|u_2 - u_1| \wedge 1).$$

Proof. From $a_+ = (a + |a|)/2$ we conclude that $|a_+ - b_+| \leq |a - b|$ for all $a, b \in \mathbb{R}$.

Let $x \in [-\pi, \pi]$, $u_1 = (y_1, s_1) \in \mathbb{R}^2$, $u_2 = (y_2, s_2) \in \mathbb{R}^2$ and $s_1, s_2 \in [-M, M]$. We have

$$\begin{aligned} &|g(u_1, x) - g(u_2, x)| \\ &\leq \left| \left((W(y_1) - W(y_1 + x))s_1 \right)_+ - \left((W(y_2) - W(y_2 + x))s_2 \right)_+ \right| \end{aligned}$$

$$\begin{aligned}
 &\leq |(W(y_1) - W(y_1 + x))s_1 - (W(y_2) - W(y_2 + x))s_2| \\
 &\leq |(W(y_1) - W(y_1 + x))s_1 - (W(y_1) - W(y_1 + x))s_2| \\
 &\quad + |(W(y_1) - W(y_1 + x))s_2 - (W(y_2) - W(y_2 + x))s_2| \\
 &\leq |W(y_1) - W(y_1 + x)||s_1 - s_2| + |W(y_1) - W(y_2)||s_2| \\
 &\quad + |W(y_1 + x) - W(y_2 + x)||s_2| \\
 &\leq 2\|W\|_\infty|s_1 - s_2| + 2M\|W'\|_\infty|y_1 - y_2|.
 \end{aligned}$$

Since, trivially, $|g(u_1, x) - g(u_2, x)| \leq 4\|W\|_\infty M$, the claim follows with $c = 4(\|W\|_\infty + \|W'\|_\infty)(M + 1)$. \square

As an easy corollary of Lemma 2.7 we get

Lemma 2.8. *There exists a constant $c = c(M) > 0$ such that for any $u_1 = (y_1, s_1) \in \mathbb{R}^2$, $u_2 = (y_2, s_2) \in \mathbb{R}^2$ with $s_1, s_2 \in [-M, M]$ we have*

$$\left| \|g(u_1, \cdot)\|_1 - \|g(u_2, \cdot)\|_1 \right| \leq c(|u_2 - u_1| \wedge 1).$$

Proof of Proposition 2.2. Let $f \in C_*(\mathbb{R}^2)$. Throughout the proof we will assume that $\text{supp}(f) \subset \mathbb{R} \times (-M_0, M_0)$ for some $M_0 > 0$. Note that

$$|S_t| = \left| S_0 + \int_0^t W(Y_r) dr \right| \leq |S_0| + \|W\|_\infty \leq M_0 + \|W\|_\infty$$

for all starting points $(Y_0, S_0) = (y, s) \in \mathbb{R} \times [-M_0, M_0]$ and all $0 \leq t \leq 1$. Put

$$M_1 = M_0 + \|W\|_\infty.$$

If $(Y_0, S_0) = (y, s) \notin \mathbb{R} \times [-M_1, M_1]$, then

$$|S_t| = \left| S_0 + \int_0^t W(Y_r) dr \right| > M_1 - \|W\|_\infty = M_0, \quad 0 \leq t \leq 1,$$

so $f(Y_t, S_t) = 0$. It follows that for any $(y, s) \notin \mathbb{R} \times [-M_1, M_1]$ and $0 < h \leq 1$ we have

$$\frac{\mathbb{E}^{(y,s)} f(Y_h, S_h) - f(y, s)}{h} = 0.$$

By the same argument,

$$\frac{\mathbb{E}^{(y,s)} f(\hat{Y}_h, \hat{S}_h) - f(y, s)}{h} = 0.$$

It now follows from the definition of $Rf(y, s)$ that $Rf(y, s) = 0$ for $(y, s) \notin \mathbb{R} \times [-M_1, M_1]$. It is, therefore, enough to consider $(y, s) \in \mathbb{R} \times [-M_1, M_1]$.

The arguments above tell us that for all starting points $(Y_0, S_0) = (y, s) \in \mathbb{R} \times [-M_1, M_1]$ and all $0 \leq t \leq 1$, $|S_t| \leq |S_0| + \|W\|_\infty \leq M_1 + \|W\|_\infty$. Setting

$$M = M_1 + \|W\|_\infty,$$

we get from the definition of the function g that

$$\|g(Y_r, S_r, \cdot)\|_1 \leq 2\pi 2\|W\|_\infty M, \quad 0 \leq r \leq 1,$$

and so

$$\sigma(t) = \int_0^t \|g(Y_r, S_r, \cdot)\|_1 dr \leq 4\pi \|W\|_\infty M t = c_0 t, \quad 0 \leq t \leq 1,$$

with the constant $c_0 = 4\pi \|W\|_\infty M$.

From now on we will assume that $(y, s) \in \mathbb{R} \times [-M_1, M_1]$ and $0 < h \leq 1$. We have

$$\begin{aligned} \frac{T_h f(y, s) - f(y, s)}{h} &= \frac{\mathbb{E}^{(y,s)} f(Y_h, S_h) - f(y, s)}{h} \\ &= \frac{1}{h} \mathbb{E}^{(y,s)} [f(Y_h, S_h) - f(y, s); N(\sigma(h)) = 0] \\ &\quad + \frac{1}{h} \mathbb{E}^{(y,s)} [f(Y_h, S_h) - f(y, s); N(\sigma(h)) = 1] \\ &\quad + \frac{1}{h} \mathbb{E}^{(y,s)} [f(Y_h, S_h) - f(y, s); N(\sigma(h)) \geq 2] \\ &= \text{I} + \text{II} + \text{III}. \end{aligned}$$

Since $\sigma(h) \leq c_0 h$ we obtain

$$\begin{aligned} |\text{III}| &\leq \frac{2\|f\|_\infty}{h} \mathbb{P}^{(y,s)} [N(\sigma(h)) \geq 2] \leq \frac{2\|f\|_\infty}{h} \mathbb{P}^{(y,s)} [N(c_0 h) \geq 2] \\ &= 2\|f\|_\infty \frac{1 - e^{-c_0 h} - c_0 h e^{-c_0 h}}{h} \xrightarrow{h \rightarrow 0^+} 0 \end{aligned}$$

uniformly for all $(y, s) \in \mathbb{R} \times [-M_1, M_1]$.

Now we will consider the expression I. We have

$$\begin{aligned} \text{I} &= \frac{1}{h} \mathbb{E}^{(y,s)} \left[f \left(y + X_h, s + \int_0^h W(y + X_r) dr \right) - f(y, s); N(\sigma(h)) = 0 \right] \\ &= \frac{1}{h} \mathbb{E}^{(y,s)} [f(\hat{Y}_h, \hat{S}_h) - f(y, s); N(\sigma(h)) = 0] \\ &= \frac{1}{h} \mathbb{E}^{(y,s)} [f(\hat{Y}_h, \hat{S}_h) - f(y, s)] - \frac{1}{h} \mathbb{E}^{(y,s)} [f(\hat{Y}_h, \hat{S}_h) - f(y, s); N(\sigma(h)) \geq 1] \\ &= \text{I}_1 + \text{I}_2. \end{aligned}$$

Note that

$$\text{I}_1 = \frac{\hat{T}_h f(y, s) - f(y, s)}{h}.$$

It will suffice to prove that $\text{I}_2 \rightarrow 0$ and $\text{II} \rightarrow Rf$. We have

$$\begin{aligned} |\text{I}_2| &\leq \frac{1}{h} \mathbb{E}^{(y,s)} [|f(\hat{Y}_h, \hat{S}_h) - f(y, s)|; N(c_0 h) \geq 1] \\ &= \frac{1 - e^{-c_0 h}}{h} \mathbb{E}^{(y,s)} [|f(\hat{Y}_h, \hat{S}_h) - f(y, s)|]. \end{aligned}$$

Recall that $f \in C_*(\mathbb{R}^2)$ is bounded and uniformly continuous. We will use the following modulus of continuity

$$\varepsilon(f; \delta) = \varepsilon(\delta) = \sup_{(y,s) \in \mathbb{R}^2} \sup_{|y_1| \vee |s_1| \leq \delta} |f(y + y_1, s + s_1) - f(y, s)|.$$

Clearly, $\varepsilon(\delta) \leq 2\|f\|_\infty$ and $\lim_{\delta \rightarrow 0^+} \varepsilon(\delta) = 0$.

Note that for $\hat{Y}_0 = y$, $\hat{S}_0 = s$ we have $\hat{Y}_h - y = X_h$, $\hat{S}_h - s = \int_0^h W(\hat{Y}_r) dr$ which gives $|\hat{S}_t - s| \leq h\|W\|_\infty$ for all $t \leq h$. It follows that

$$\mathbb{E}^{(y,s)}[|f(\hat{Y}_h, \hat{S}_h) - f(y, s)|] \leq \mathbb{E}^{(y,s)}\left[\varepsilon\left(\sup_{0 < t \leq h} (|X_t| \vee h\|W\|_\infty)\right)\right].$$

Since $t \mapsto X_t$ is right-continuous and $X_0 \equiv 0$ we have, a.s.,

$$\sup_{0 < t \leq h} |X_t| \xrightarrow{h \rightarrow 0^+} 0 \quad \text{and, therefore,} \quad \varepsilon\left(\sup_{0 < t \leq h} (|X_t| \vee h\|W\|_\infty)\right) \xrightarrow{h \rightarrow 0^+} 0.$$

By the bounded convergence theorem

$$\mathbb{E}^{(y,s)}\left[\varepsilon\left(\sup_{0 < t \leq h} (|X_t| \vee h\|W\|_\infty)\right)\right] \xrightarrow{h \rightarrow 0^+} 0$$

uniformly for all $(y, s) \in \mathbb{R} \times [-M_1, M_1]$ because the expression $\varepsilon(\sup_{0 < t \leq h} |X_t| \vee h\|W\|_\infty)$ does not depend on (y, s) . It follows that

$$|\mathbb{I}_2| \xrightarrow{h \rightarrow 0^+} 0$$

uniformly for all $(y, s) \in \mathbb{R} \times [-M_1, M_1]$.

Now we turn to II. We have

$$\begin{aligned} \mathbb{II} &= \frac{1}{h} \mathbb{E}^{(y,s)}[f(Y_h, S_h) - f(Y_h, s); N(\sigma(h)) = 1] \\ &\quad + \frac{1}{h} \mathbb{E}^{(y,s)}[f(Y_h, s) - f(y, s); N(\sigma(h)) = 1] \\ &= \mathbb{II}_1 + \mathbb{II}_2. \end{aligned}$$

Since $\sigma(h) \leq c_0 h$

$$\begin{aligned} |\mathbb{II}_1| &\leq \frac{1}{h} \mathbb{E}^{(y,s)}\left[\left|f\left(Y_h, s + \int_0^h W(Y_r) dr\right) - f(Y_h, s)\right|; N(c_0 h) \geq 1\right] \\ &\leq \frac{1}{h} \mathbb{E}^{(y,s)}\left[\varepsilon\left(\left|\int_0^h W(Y_r) dr\right|\right); N(c_0 h) \geq 1\right] \\ &\leq \frac{1}{h} \mathbb{E}^{(y,s)}[\varepsilon(h\|W\|_\infty); N(c_0 h) \geq 1] \\ &= \frac{1 - e^{-c_0 h}}{h} \mathbb{E}^{(y,s)}[\varepsilon(h\|W\|_\infty)] \xrightarrow{h \rightarrow 0^+} 0 \end{aligned}$$

uniformly for all $(y, s) \in \mathbb{R} \times [-M_1, M_1]$. It will suffice to show that $\mathbb{II}_2 \rightarrow Rf$.

From now on we will use the following shorthand notation

$$U_t := (Y_t, S_t), \quad \hat{U}_t := (\hat{Y}_t, \hat{S}_t), \quad u := (y, s).$$

We have

$$\begin{aligned} \mathbb{II}_2 &= \frac{1}{h} \mathbb{E}^{(y,s)}[f(y + X_h + \eta_1(U_{\tau_1-}), s) - f(y + \eta_1(U_{\tau_1-}), s); N(\sigma(h)) \geq 1] \\ &\quad + \frac{1}{h} \mathbb{E}^{(y,s)}[f(y + \eta_1(U_{\tau_1-}), s) - f(y, s); N(\sigma(h)) \geq 1] \end{aligned}$$

$$\begin{aligned}
& - \frac{1}{h} \mathbb{E}^{(y,s)} [f(y + X_h + \eta_1(U_{\tau_1-}), s) - f(y, s); N(\sigma(h)) \geq 2] \\
& = \mathbb{I}_{2a} + \mathbb{I}_{2b} + \mathbb{I}_{2c}.
\end{aligned}$$

Observe that

$$|\mathbb{I}_{2c}| \leq 2 \|f\|_\infty \frac{1 - e^{-c_0 h} - c_0 h e^{-c_0 h}}{h} \xrightarrow{h \rightarrow 0^+} 0$$

and that the convergence is uniform in $(y, s) \in \mathbb{R} \times [-M_1, M_1]$.

Moreover,

$$\begin{aligned}
|\mathbb{I}_{2a}| & \leq \frac{1}{h} \mathbb{E}^{(y,s)} [|f(y + X_h + \eta_1(U_{\tau_1-}), s) - f(y + \eta_1(U_{\tau_1-}), s)|; N(c_0 h) \geq 1] \\
& \leq \frac{1}{h} \mathbb{E}^{(y,s)} \left[\varepsilon \left(\sup_{0 \leq t \leq h} |X_h| \right); N(c_0 h) \geq 1 \right] \\
& = \frac{1 - e^{-c_0 h}}{h} \mathbb{E}^{(y,s)} \left[\varepsilon \left(\sup_{0 \leq t \leq h} |X_h| \right) \right] \\
& \xrightarrow{h \rightarrow 0^+} 0
\end{aligned}$$

uniformly for all $(y, s) \in \mathbb{R} \times [-M_1, M_1]$. It will suffice to show that $\mathbb{I}_{2b} \rightarrow Rf$.

Note that

$$N(\sigma(h)) \geq 1 \iff \tau_1 \leq h \iff \int_0^h \|g(U_r, \cdot)\|_1 dr \geq \sigma_1. \quad (2.10)$$

We claim that

$$\int_0^h \|g(U_r, \cdot)\|_1 dr \geq \sigma_1 \iff \int_0^h \|g(\hat{U}_r, \cdot)\|_1 dr \geq \sigma_1. \quad (2.11)$$

First, we assume that $\int_0^h \|g(U_r, \cdot)\|_1 dr \geq \sigma_1$. This implies that $\tau_1 \leq h$. Recall that $U_r = \hat{U}_r$ for $r < \tau_1$. Hence

$$\int_0^h \|g(\hat{U}_r, \cdot)\|_1 dr \geq \int_0^{\tau_1} \|g(\hat{U}_r, \cdot)\|_1 dr = \int_0^{\tau_1} \|g(U_r, \cdot)\|_1 dr = \sigma_1,$$

where the last equality follows from the definition of τ_1 .

Now let us assume that $\int_0^h \|g(U_r, \cdot)\|_1 dr < \sigma_1$. This implies that $\tau_1 > h$. Using again $U_r = \hat{U}_r$ for $r < h < \tau_1$, we obtain

$$\sigma_1 > \int_0^h \|g(U_r, \cdot)\|_1 dr = \int_0^h \|g(\hat{U}_r, \cdot)\|_1 dr,$$

which finishes the proof of (2.11).

By (2.10) and (2.11) we obtain

$$\begin{aligned}
\mathbb{I}_{2b} & = \frac{1}{h} \mathbb{E}^{(y,s)} [f(y + \eta_1(U_{\tau_1-}), s) - f(y, s); \tau_1 \leq h] \\
& = \frac{1}{h} \mathbb{E}^{(y,s)} [f(y + \eta_1(U_{(\tau_1 \wedge h)-}), s) - f(y, s); \tau_1 \leq h] \\
& = \frac{1}{h} \mathbb{E}^{(y,s)} \left[f(y + \eta_1(\hat{U}_{(\tau_1 \wedge h)-}), s) - f(y, s); \int_0^h \|g(\hat{U}_r, \cdot)\|_1 dr \geq \sigma_1 \right].
\end{aligned}$$

We will use the following abbreviations:

$$\begin{aligned} u &= (y, s), \\ A &= \left\{ \int_0^h \|g(\hat{U}_r, \cdot)\|_1 \, dr \geq \sigma_1 \right\}, \\ B &= \left\{ \int_0^h \|g(u, \cdot)\|_1 \, dr \geq \sigma_1 \right\}, \\ F &= \frac{1}{h} (f(y + \eta_1(\hat{U}_{(\tau_1 \wedge h)^-}), s) - f(y, s)). \end{aligned}$$

This allows us to rewrite Π_{2b} as

$$\Pi_{2b} = \mathbb{E}^u[F; A] = \mathbb{E}^u[F; B] + \mathbb{E}^u[F, A \setminus B] - \mathbb{E}^u[F; B \setminus A].$$

Recall that $X = (X_t)_{t \geq 0}$, $N = (N(t))_{t \geq 0}$ and $\mathcal{U} = (\mathcal{U}_n)_{n \in \mathbb{N}}$ are independent. Therefore the probability measure \mathbb{P} can be written in the form $\mathbb{P} = \mathbb{P}_X \otimes \mathbb{P}_N \otimes \mathbb{P}_{\mathcal{U}}$; the conditional probability, given N or \mathcal{U} , is \mathbb{P}_X and the corresponding expectation is denoted by \mathbb{E}_X . In a similar way $\mathbb{P}_{(X, N)} = \mathbb{P}_X \otimes \mathbb{P}_N$ and $\mathbb{E}_{(X, N)}$ denote conditional probability and conditional expectation if \mathcal{U} is given. As usual, the initial (time-zero) value of the process under consideration is given as a superscript. Note that $\hat{U}_t = (\hat{Y}_t, \hat{S}_t)$ is a function of X and does not depend on N or \mathcal{U} . In particular, \hat{U}_t and σ_1 are independent. Since σ_1 is the time of the first jump of the Poisson process $N(t)$, it is exponentially distributed with parameter 1. It follows that

$$\begin{aligned} & |\mathbb{E}^u[F, A \setminus B]| \\ & \leq \frac{2\|f\|_\infty}{h} \mathbb{P}^u \left[\int_0^h \|g(\hat{U}_r, \cdot)\|_1 \, dr \geq \sigma_1 > \int_0^h \|g(u, \cdot)\|_1 \, dr \right] \\ & = \frac{2\|f\|_\infty}{h} \mathbb{E}_X^u \left[e^{-\int_0^h \|g(u, \cdot)\|_1 \, dr} - e^{-\int_0^h \|g(\hat{U}_r, \cdot)\|_1 \, dr}; \int_0^h \|g(\hat{U}_r, \cdot)\|_1 \, dr > \int_0^h \|g(u, \cdot)\|_1 \, dr \right] \\ & \leq \frac{2\|f\|_\infty}{h} \mathbb{E}_X^u \left| e^{-\int_0^h \|g(u, \cdot)\|_1 \, dr} - e^{-\int_0^h \|g(\hat{U}_r, \cdot)\|_1 \, dr} \right| \\ & \leq \frac{2\|f\|_\infty}{h} \mathbb{E}_X^u \left| \int_0^h (\|g(u, \cdot)\|_1 - \|g(\hat{U}_r, \cdot)\|_1) \, dr \right| \\ & \leq 2\|f\|_\infty \mathbb{E}_X^u \sup_{0 \leq r \leq h} \left| \|g(u, \cdot)\|_1 - \|g(\hat{U}_r, \cdot)\|_1 \right|. \end{aligned}$$

For the penultimate inequality we used the elementary estimate $|e^{-a} - e^{-b}| \leq |a - b|$, $a, b \geq 0$. From Lemma 2.8 we infer that the last expression is bounded by

$$\begin{aligned} & 2\|f\|_\infty c \mathbb{E}_X^u \left[\sup_{0 \leq r \leq h} (|\hat{U}_r - u| \wedge 1) \right] \\ & = 2\|f\|_\infty c \mathbb{E}_X^u \left[\sup_{0 \leq r \leq h} \left(\left| \left(X_r, \int_0^r W(\hat{Y}_t) \, dt \right) \right| \wedge 1 \right) \right] \\ & \xrightarrow{h \rightarrow 0^+} 0 \end{aligned}$$

uniformly for all $(y, s) \in \mathbb{R} \times [-M_1, M_1]$. This convergence follows from the right-continuity of X_r and the fact that $|\int_0^r W(\hat{Y}_t) \, dt| \leq h \|W\|_\infty$.

A similar argument shows that $|\mathbb{E}^u[F; B \setminus A]| \xrightarrow{h \rightarrow 0^+} 0$ uniformly in $(y, s) \in \mathbb{R} \times [-M_1, M_1]$. It will suffice to show that $\mathbb{E}^u[F; B] \rightarrow Rf$.

We have

$$\begin{aligned} \mathbb{E}^u[F; B] &= \frac{1}{h} \mathbb{E}^{(y,s)} \left[f(y + \eta_1(\hat{U}_{(\tau_1 \wedge h)-}), s) - f(y, s); \int_0^h \|g(u, \cdot)\|_1 dr \geq \sigma_1 \right] \\ &= \frac{1}{h} \mathbb{E}^{(y,s)} \left[f(y + \eta_1(\hat{U}_{(\tau_1 \wedge h)-}), s) - f(y + \eta_1(u), s); \int_0^h \|g(u, \cdot)\|_1 dr \geq \sigma_1 \right] \\ &\quad + \frac{1}{h} \mathbb{E}^{(y,s)} \left[f(y + \eta_1(u), s) - f(y, s); \int_0^h \|g(u, \cdot)\|_1 dr \geq \sigma_1 \right] \\ &= A + B. \end{aligned}$$

In order to deal with A and B we introduce the following auxiliary notation.

Recall that X , N and \mathcal{U} are independent. As before let $\mathbb{E}_{(X,N)}^{(y,s)}$ be the conditional expectation given \mathcal{U} ; the superscript (y, s) indicates that $Y_0 = y$ and $S_0 = s$. Moreover, $\mathbb{E}_{\mathcal{U}}$ denotes conditional expectation given X and N .

Lemma 2.9. *Let $u_1 = (y_1, s_1) \in \mathbb{R}^2$, $u_2 = (y_2, s_2) \in \mathbb{R}^2$ be such that $s_1, s_2 \in [-M, M]$ and $\|g(u_2, \cdot)\|_1 > 0$. Then we have*

$$|\mathbb{E}_{\mathcal{U}}(f(y + \eta_1(u_1), s) - f(y + \eta_1(u_2), s))| \leq c \left(\frac{|u_1 - u_2|}{\|g(u_2, \cdot)\|_1} \wedge 1 \right)$$

for some $c = c(f, M) > 0$.

Proof. We will distinguish two cases: $\|g(u_1, \cdot)\|_1 = 0$ and $\|g(u_1, \cdot)\|_1 > 0$.

Assume that $\|g(u_1, \cdot)\|_1 = 0$. Then by Lemma 2.8 we have

$$\|g(u_2, \cdot)\|_1 = \left| \|g(u_2, \cdot)\|_1 - \|g(u_1, \cdot)\|_1 \right| \leq c|u_2 - u_1|.$$

Hence,

$$|\mathbb{E}_{\mathcal{U}}(f(y + \eta_1(u_1), s) - f(y + \eta_1(u_2), s))| \leq 2\|f\|_{\infty} \leq \frac{2\|f\|_{\infty} c|u_1 - u_2|}{\|g(u_2, \cdot)\|_1}.$$

Now we will consider the second case: $\|g(u_1, \cdot)\|_1 > 0$. We have

$$\begin{aligned} &|\mathbb{E}_{\mathcal{U}}(f(y + \eta_1(u_1), s) - f(y + \eta_1(u_2), s))| \\ &= \left| \int_{-\pi}^{\pi} \frac{f(y+x, s)}{\|g(u_1, \cdot)\|_1} g(u_1, x) dx - \int_{-\pi}^{\pi} \frac{f(y+x, s)}{\|g(u_2, \cdot)\|_1} g(u_2, x) dx \right| \\ &\leq \frac{|\int_{-\pi}^{\pi} f(y+x, s)[g(u_1, x)\|g(u_2, \cdot)\|_1 - g(u_2, x)\|g(u_1, \cdot)\|_1] dx|}{\|g(u_1, \cdot)\|_1 \|g(u_2, \cdot)\|_1} \\ &\leq \frac{\|f\|_{\infty}}{\|g(u_1, \cdot)\|_1 \|g(u_2, \cdot)\|_1} \left[\int_{-\pi}^{\pi} |g(u_1, x)\|g(u_2, \cdot)\|_1 - g(u_1, x)\|g(u_1, \cdot)\|_1| dx \right. \\ &\quad \left. + \int_{-\pi}^{\pi} |g(u_1, x)\|g(u_1, \cdot)\|_1 - g(u_2, x)\|g(u_1, \cdot)\|_1| dx \right]. \end{aligned}$$

By Lemmas 2.7 and 2.8 this is bounded from above by

$$\begin{aligned} &\frac{\|f\|_{\infty}}{\|g(u_1, \cdot)\|_1 \|g(u_2, \cdot)\|_1} \left[\int_{-\pi}^{\pi} g(u_1, x) dx c'|u_2 - u_1| + 2\pi c''|u_1 - u_2| \|g(u_1, \cdot)\|_1 \right] \\ &\leq \frac{(c' + 2\pi c'')\|f\|_{\infty}|u_2 - u_1|}{\|g(u_2, \cdot)\|_1}. \end{aligned}$$

The lemma follows now from the observation that

$$|\mathbb{E}_{\mathcal{U}}(f(y + \eta_1(u_1), s) - f(y + \eta_1(u_2), s))| \leq 2\|f\|_{\infty}. \quad \square$$

Proof of Proposition 2.2 (continued). We go back to $A + B$. If $\|g(u, \cdot)\|_1 = 0$ then $A + B = 0 = Rf(y, s)$. The proof of the proposition is complete in this case.

We will consider the case $\|g(u, \cdot)\|_1 > 0$. Because of the independence of σ_1 , X_t and $(\eta_1(u))_{u \in \mathbb{R}^2}$ we get

$$|A| = \left| \frac{1}{h} \mathbb{E}_{(X,N)}^u \left[\mathbb{E}_{\mathcal{U}} [f(y + \eta_1(\hat{U}_{(\tau_1 \wedge h)-}), s) - f(y + \eta_1(u), s)]; h\|g(u, \cdot)\|_1 \geq \sigma_1 \right] \right|.$$

By Lemma 2.9 this is bounded from above by

$$\begin{aligned} & \left| \frac{1}{h} \mathbb{E}_{(X,N)}^u \left[c \left(\frac{|\hat{U}_{((\tau_1 \wedge h)-)} - u|}{\|g(u, \cdot)\|_1} \wedge 1 \right); h\|g(u, \cdot)\|_1 \geq \sigma_1 \right] \right| \\ & \leq \left| \frac{c}{h\|g(u, \cdot)\|_1} \mathbb{E}_{(X,N)}^u \left[\sup_{0 \leq r \leq h} |\hat{U}_r - u| \wedge \|g(u, \cdot)\|_1; h\|g(u, \cdot)\|_1 \geq \sigma_1 \right] \right| \\ & = \left| \frac{c}{h\|g(u, \cdot)\|_1} \mathbb{E}_{(X,N)}^u \left[\sup_{0 \leq r \leq h} \left| \left(X_r, \int_0^r W(\hat{Y}_t) dt \right) \right| \wedge \|g(u, \cdot)\|_1; h\|g(u, \cdot)\|_1 \geq \sigma_1 \right] \right| \\ & \leq \left| \frac{c}{h\|g(u, \cdot)\|_1} \mathbb{E}_{(X,N)}^u \left[\sup_{0 \leq r \leq h} |(X_r, h\|W\|_{\infty})| \wedge \|g(u, \cdot)\|_1; h\|g(u, \cdot)\|_1 \geq \sigma_1 \right] \right|. \end{aligned}$$

Using the independence of X and σ_1 this is equal to

$$\begin{aligned} & \frac{c}{h\|g(u, \cdot)\|_1} (1 - e^{-h\|g(u, \cdot)\|_1}) \mathbb{E}_X^u \left[\sup_{0 \leq r \leq h} |(X_r, h\|W\|_{\infty})| \wedge \|g(u, \cdot)\|_1 \right] \\ & \leq c \mathbb{E}_X^u \left[\sup_{0 \leq r \leq h} |(X_r, h\|W\|_{\infty})| \wedge \|g(u, \cdot)\|_1 \right] \xrightarrow{h \rightarrow 0^+} 0 \end{aligned}$$

uniformly for all $u = (y, s) \in \mathbb{R} \times [-M_1, M_1]$.

It will suffice to show that $B \xrightarrow{h \rightarrow 0^+} Rf$. Because of the independence of η_1 and σ_1 we get

$$\begin{aligned} B &= \mathbb{E}_{\mathcal{U}} [f(y + \eta_1(u), s) - f(y, s)] \frac{1}{h} (1 - e^{-h\|g(u, \cdot)\|_1}) \\ &= \int_{-\pi}^{\pi} (f(y + x, s) - f(y, s)) ((W(y) - W(y + x))s)_+ dx \frac{1 - e^{-h\|g(u, \cdot)\|_1}}{h\|g(u, \cdot)\|_1} \\ &= Rf(y, s) \frac{1 - e^{-h\|g(u, \cdot)\|_1}}{h\|g(u, \cdot)\|_1} \\ &= Rf(y, s) + Rf(y, s) \left(\frac{1 - e^{-h\|g(u, \cdot)\|_1}}{h\|g(u, \cdot)\|_1} - 1 \right). \end{aligned} \quad (2.12)$$

For $u = (y, s) \in \mathbb{R} \times [-M_1, M_1]$ we have

$$\begin{aligned} |Rf(y, s)| &\leq 2\|f\|_{\infty} 2\pi 2\|W\|_{\infty} M_1 = 8\pi\|f\|_{\infty}\|W\|_{\infty} M_1, \\ \|g(u, \cdot)\|_1 &\leq 2\pi 2\|W\|_{\infty} M_1 = 4\pi\|W\|_{\infty} M_1. \end{aligned}$$

Note that for any $h, c > 0$ we have

$$-\frac{hc}{2} \leq \frac{1 - e^{-hc} - hc}{hc} \leq 0.$$

Therefore,

$$\left| \frac{1 - e^{-h\|g(u, \cdot)\|_1}}{h\|g(u, \cdot)\|_1} - 1 \right| \leq \frac{h\|g(u, \cdot)\|_1}{2} \leq \frac{4\pi\|W\|_\infty M_1}{2} h.$$

It follows that the expression in (2.12) tends to $Rf(y, s)$ when $h \rightarrow 0^+$ uniformly for all $u = (y, s) \in \mathbb{R} \times [-M_1, M_1]$. We have shown that $\mathbf{B} \xrightarrow{h \rightarrow 0^+} Rf$. This was the last step in the proof. \square

We will now introduce some further notation. Let \mathbb{N} be the positive integers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For any $f : \mathbb{S} \rightarrow \mathbb{R}$ we set

$$\tilde{f}(x) := f(e^{ix}), \quad x \in \mathbb{R}.$$

We say that $f : \mathbb{S} \rightarrow \mathbb{R}$ is *differentiable* at $z = e^{ix}$, $x \in \mathbb{R}$, if and only if \tilde{f} is differentiable at x and we put

$$f'(z) := (\tilde{f})'(x), \quad \text{where } z = e^{ix}, x \in \mathbb{R}.$$

Analogously, we say that $f : \mathbb{S} \rightarrow \mathbb{R}$ is *n times differentiable* at $z = e^{ix}$, $x \in \mathbb{R}$, if and only if \tilde{f} is *n times differentiable* at x and we write

$$f^{(n)}(z) = (\tilde{f})^{(n)}(x), \quad \text{where } z = e^{ix}, x \in \mathbb{R}.$$

In a similar way we define for $f : \mathbb{S} \times \mathbb{R} \rightarrow \mathbb{R}$

$$\tilde{f}(y, s) = f(e^{iy}, s), \quad y, s \in \mathbb{R}. \quad (2.13)$$

We say that $D^\alpha f(z, s)$, $z = e^{iy}$, $y, s \in \mathbb{R}$, $\alpha \in \mathbb{N}_0^2$, exists if and only if $D^\alpha \tilde{f}(y, s)$ exists and we set

$$D^\alpha f(z, s) = D^\alpha \tilde{f}(y, s), \quad \text{where } z = e^{iy}, y, s \in \mathbb{R}.$$

When writing $C^2(\mathbb{S})$, $C_c^2(\mathbb{S} \times \mathbb{R})$, etc., we are referring to the derivatives defined above.

Proof of Corollary 2.3. We will use the notation \tilde{f} introduced in (2.13). Let $f \in C_c(\mathbb{S} \times \mathbb{R})$. Then $\tilde{f} \in C_*(\mathbb{R}^2)$. Let $z = e^{iy}$, $z \in \mathbb{S}$, $s \in \mathbb{R}$. We have, cf. [3], Eq. (2.9),

$$\frac{T_t^{\mathbb{S}} f(z, s) - f(z, s)}{t} = \frac{T_t \tilde{f}(y, s) - \tilde{f}(y, s)}{t}, \quad (2.14)$$

$$\frac{\hat{T}_t^{\mathbb{S}} f(z, s) - f(z, s)}{t} = \frac{\hat{T}_t \tilde{f}(y, s) - \tilde{f}(y, s)}{t}. \quad (2.15)$$

Using this and Proposition 2.2 we get that $\lim_{t \rightarrow 0^+} (T_t^{\mathbb{S}} f - f)/t$ exists if and only if $\lim_{t \rightarrow 0^+} (\hat{T}_t^{\mathbb{S}} f - f)/t$ exists, where both limits are in $\|\cdot\|_\infty$ norm. Consequently,

$$f \in \mathcal{D}(\mathcal{G}) \cap C_c(\mathbb{S} \times \mathbb{R}) \iff f \in \mathcal{D}(\hat{\mathcal{G}}) \cap C_c(\mathbb{S} \times \mathbb{R}).$$

The second assertion of the proposition follows from (2.5), the definition of the infinitesimal generator and from the fact that for $z \in \mathbb{S}$ and $s \in \mathbb{R}$

$$R\tilde{f}(y, s) = R^{\mathbb{S}} f(z, s), \quad z = e^{iy}. \quad (2.16)$$

\square

Proof of Proposition 2.4. Note that (2.6) follows from (2.7) by Proposition 2.2. So it is sufficient to show (2.7).

Pick $f \in C_*^2(\mathbb{R}^2)$. Throughout this proof we assume that $\text{supp}(f) \subset \mathbb{R} \times (-M_0, M_0)$ for some $M_0 > 0$. With exactly the same argument as at the beginning of the proof of Proposition 2.2, we can restrict our attention to $(y, s) \in \mathbb{R} \times [-M_1, M_1]$ where $M_1 := M_0 + \|W\|_\infty$. We have for $0 < h < 1$,

$$\begin{aligned} & \frac{\hat{T}_h f(y, s) - f(y, s)}{h} \\ &= \frac{\mathbb{E}^{(y,s)} f(\hat{Y}_h, \hat{S}_h) - \mathbb{E}^{(y,s)} f(\hat{Y}_h, s)}{h} + \frac{\mathbb{E}^{(y,s)} f(\hat{Y}_h, s) - \mathbb{E}^{(y,s)} f(y, s)}{h} \\ &= \text{I} + \text{II}. \end{aligned}$$

We get

$$\begin{aligned} \text{I} &= \frac{1}{h} \mathbb{E}^{(y,s)} \left[\frac{\partial f}{\partial s}(\hat{Y}_h, \xi) (\hat{S}_h - s) \right] \\ &= \frac{1}{h} \mathbb{E}^{(y,s)} \left[\frac{\partial f}{\partial s}(\hat{Y}_h, \xi) \int_0^h W(\hat{Y}_t) dt \right] \\ &= \mathbb{E}^{(y,s)} \left[\frac{1}{h} \frac{\partial f}{\partial s}(y, s) \int_0^h W(y) dt \right] + \mathbb{E}^{(y,s)} \left[\frac{1}{h} \left[\frac{\partial f}{\partial s}(\hat{Y}_h, \xi) - \frac{\partial f}{\partial s}(y, s) \right] \int_0^h W(y) dt \right] \\ &\quad + \mathbb{E}^{(y,s)} \left[\frac{1}{h} \frac{\partial f}{\partial s}(\hat{Y}_h, \xi) \int_0^h (W(\hat{Y}_t) - W(y)) dt \right] \\ &= \text{I}_1 + \text{I}_2 + \text{I}_3, \end{aligned}$$

where ξ is a point between s and \hat{S}_h . Note that $|\hat{Y}_h - y| = |X_h|$ and $|\xi - s| \leq |\hat{S}_h - s| \leq h \|W\|_\infty$. Moreover,

$$\begin{aligned} |W(\hat{Y}_h) - W(y)| &\leq (2\|W\|_\infty) \wedge (\|W'\|_\infty |\hat{Y}_h - y|) \\ &\leq c(|X_h| \wedge 1) \\ &\leq c\left(\sup_{0 \leq t \leq h} |X_t| \wedge 1\right) \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{\partial f}{\partial s}(\hat{Y}_h, \xi) - \frac{\partial f}{\partial s}(y, s) \right| \\ &\leq 2 \left\| \frac{\partial f}{\partial s} \right\|_\infty \wedge \left[\left(\left\| \frac{\partial^2 f}{\partial s^2} \right\|_\infty + \left\| \frac{\partial^2 f}{\partial s \partial y} \right\|_\infty \right) (|\hat{Y}_h - y| + |\xi - s|) \right] \\ &\leq c((|X_h| + h) \wedge 1), \end{aligned}$$

where $c = c(W, f)$. It follows that

$$|\text{I}_2| \leq c \|W\|_\infty \mathbb{E}^{(y,s)} \left((|X_h| + h) \wedge 1 \right) \xrightarrow{h \rightarrow 0^+} 0,$$

uniformly for all $u = (y, s) \in \mathbb{R} \times [-M_1, M_1]$. In a similar way

$$|\text{I}_3| \leq c \left\| \frac{\partial f}{\partial s} \right\|_\infty \mathbb{E}^{(y,s)} \left[\sup_{0 \leq t \leq h} |X_t| \wedge 1 \right] \xrightarrow{h \rightarrow 0^+} 0,$$

uniformly for all $u = (y, s) \in \mathbb{R} \times [-M_1, M_1]$. So

$$\text{I} \xrightarrow{h \rightarrow 0^+} \frac{\partial f}{\partial s}(y, s) W(y)$$

uniformly for all $u = (y, s) \in \mathbb{R} \times [-M_1, M_1]$.

It is well known that

$$\text{II} = \frac{\mathbb{E}^{(y,s)}(f(y + X_h, s) - f(y, s))}{h} \xrightarrow{h \rightarrow 0^+} -(-\Delta_y)^{\alpha/2} f(y, s)$$

uniformly in $u = (y, s)$.

Combining the estimates for I and II shows that $f \in \mathcal{D}(\mathcal{G}^{(\hat{Y}, \hat{S})})$ and that (2.7) holds. \square

Proof of Corollary 2.5. Let $f \in C_c^2(D \times \mathbb{R})$. Then $\tilde{f} \in C_*^2(\mathbb{R}^2) \subset \mathcal{D}(\mathcal{G}^{(Y,S)})$. By (2.14) $f \in \mathcal{D}(\mathcal{G})$. Now let $z = e^{iy}$, $z \in D$, $s \in \mathbb{R}$. By (2.14), Proposition 2.4, (2.9) and (2.16) we get

$$\begin{aligned} \mathcal{G}f(z, s) &= \mathcal{G}^{(Y,S)}\tilde{f}(y, s) \\ &= -(-\Delta_y)^{\alpha/2}\tilde{f}(y, s) + R\tilde{f}(y, s) + W(y)\tilde{f}_s(y, s) \\ &= L_z f(z, s) + R^D f(z, s) + V(z)f_s(z, s). \end{aligned}$$

The proof for $\hat{\mathcal{G}}$ is the same. \square

3. Stationary measure

The aim of this section is to show that the process (Z_t, S_t) has a unique stationary measure. First we will show that $C_c^2(\mathbb{S} \times \mathbb{R})$ is a core for $(\mathcal{G}, \mathcal{D}(\mathcal{G}))$. For this we will need two auxiliary lemmas.

Lemma 3.1. $C_c^2(\mathbb{S} \times \mathbb{R})$ is a core for $\hat{\mathcal{G}}$.

Proof. Here we will use the results from [3]. Note that (\hat{Y}_t, \hat{S}_t) is the solution of a SDE of the form (3.1) in [3]. Since $V: \mathbb{S} \rightarrow \mathbb{R}$ is a C^3 function, [3], Theorem 3.1, see also [3], Proposition 3.6, guarantees that $\hat{T}_t f \in C_*^2(\mathbb{R}^2)$ for all $f \in C_*^2(\mathbb{R}^2)$.

Now let $f \in C_c^2(\mathbb{S} \times \mathbb{R})$. Then $\tilde{f} \in C_*^2(\mathbb{R}^2)$ and $\hat{T}_t \tilde{f} \in C_*^2(\mathbb{R}^2)$. For $z = e^{iy}$, $z \in \mathbb{S}$, $s \in \mathbb{R}$ we get as in [3], Eq. (2.9), $\hat{T}_t^{\mathbb{S}} f(z, s) = \hat{T}_t \tilde{f}(y, s)$. Hence, $\hat{T}_t^{\mathbb{S}} f \in C_c^2(\mathbb{S} \times \mathbb{R})$. This means that $\hat{T}_t^{\mathbb{S}}: C_c^2(\mathbb{S} \times \mathbb{R}) \rightarrow C_c^2(\mathbb{S} \times \mathbb{R})$. Since $C_c^2(\mathbb{S} \times \mathbb{R})$ is dense in $C_0(\mathbb{S} \times \mathbb{R})$ – the Banach space where the semigroup $\{\hat{T}_t^{\mathbb{S}}\}_{t \geq 0}$ is defined – [6], Proposition 1.3.3, applies and shows that $C_c^2(\mathbb{S} \times \mathbb{R})$ is a core for $(\hat{\mathcal{G}}, \mathcal{D}(\hat{\mathcal{G}}))$. \square

Lemma 3.2. $C_c(\mathbb{S} \times \mathbb{R}) \cap \mathcal{D}(\mathcal{G}) = C_c(\mathbb{S} \times \mathbb{R}) \cap \mathcal{D}(\hat{\mathcal{G}})$ is a core for $(\mathcal{G}, \mathcal{D}(\mathcal{G}))$ and $(\hat{\mathcal{G}}, \mathcal{D}(\hat{\mathcal{G}}))$.

Proof. The equality of the two families of functions follows from Corollary 2.3.

By Corollary 2.5, $C_c^2(\mathbb{S} \times \mathbb{R}) \subset C_c(\mathbb{S} \times \mathbb{R}) \cap \mathcal{D}(\mathcal{G})$ and $C_c^2(\mathbb{S} \times \mathbb{R})$ is dense in $C_0(\mathbb{S} \times \mathbb{R})$ where the semigroups $\{T_t^{\mathbb{S}}\}_{t \geq 0}$, $\{\hat{T}_t^{\mathbb{S}}\}_{t \geq 0}$ are defined; so $C_c(\mathbb{S} \times \mathbb{R}) \cap \mathcal{D}(\mathcal{G})$ is dense in $C_0(\mathbb{S} \times \mathbb{R})$.

By the definition of the processes S_t and \hat{S}_t and the boundedness of W it is easy to see that $T_t^{\mathbb{S}}: C_c(\mathbb{S} \times \mathbb{R}) \rightarrow C_c(\mathbb{S} \times \mathbb{R})$ and $\hat{T}_t^{\mathbb{S}}: C_c(\mathbb{S} \times \mathbb{R}) \rightarrow C_c(\mathbb{S} \times \mathbb{R})$. It follows that $T_t^{\mathbb{S}}$ and $\hat{T}_t^{\mathbb{S}}$ map $C_c(\mathbb{S} \times \mathbb{R}) \cap \mathcal{D}(\mathcal{G})$ into itself. Now [6], Proposition 1.3.3, gives that $C_c(\mathbb{S} \times \mathbb{R}) \cap \mathcal{D}(\mathcal{G})$ is a core for \mathcal{G} and $\hat{\mathcal{G}}$. \square

Proposition 3.3. $C_c^2(\mathbb{S} \times \mathbb{R})$ is a core for \mathcal{G} .

Proof. Pick $f \in \mathcal{D}(\mathcal{G}) \cap C_c(\mathbb{S} \times \mathbb{R})$. We have $f \in \mathcal{D}(\hat{\mathcal{G}}) \cap C_c(\mathbb{S} \times \mathbb{R})$ and $C_c^2(\mathbb{S} \times \mathbb{R})$ is a core for $\hat{\mathcal{G}}$ so there exists a sequence $(f_n)_{n=1}^{\infty}$, where $f_n \in C_c^2(\mathbb{S} \times \mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} (\|f_n - f\|_{\infty} + \|\hat{\mathcal{G}}f_n - \hat{\mathcal{G}}f\|_{\infty}) = 0.$$

Since $f \in C_c(\mathbb{S} \times \mathbb{R})$, there exists some $M > 0$ such that $\text{supp}(f) \subset \mathbb{S} \times [-M, M]$. Let $g \in C_c^\infty(\mathbb{R})$ be such that $0 \leq g \leq 1$, $\|g'\|_\infty \leq 1$, $g \equiv 1$ on $[-M-1, M+1]$ and $g \equiv 0$ on $(-\infty, -M-3] \cup [M+3, \infty)$. Put

$$g_n(z, s) := g(s)f_n(z, s), \quad (z, s) \in \mathbb{S} \times \mathbb{R},$$

and note that $f(z, s) = g(s)f(z, s)$. Therefore

$$|g_n(z, s) - f(z, s)| = |g(s)f_n(z, s) - g(s)f(z, s)| \leq |f_n(z, s) - f(z, s)|$$

and

$$\|g_n - f\|_\infty \leq \|f_n - f\|_\infty.$$

Since $g_n \in C_c^2(\mathbb{S} \times \mathbb{R}) \subset \mathcal{D}(\hat{\mathcal{G}})$, we find for $(z, s) \in \mathbb{S} \times [-M, M]$,

$$\begin{aligned} |\hat{\mathcal{G}}g_n(z, s) - \hat{\mathcal{G}}f(z, s)| &= \left| V(z) \frac{\partial g_n}{\partial s}(z, s) + L_z g_n(z, s) - \hat{\mathcal{G}}f(z, s) \right| \\ &= \left| V(z) \frac{\partial f_n}{\partial s}(z, s) + L_z f_n(z, s) - \hat{\mathcal{G}}f(z, s) \right| \\ &= |\hat{\mathcal{G}}f_n(z, s) - \hat{\mathcal{G}}f(z, s)| \\ &\leq \|\hat{\mathcal{G}}f_n - \hat{\mathcal{G}}f\|_\infty, \end{aligned}$$

whereas for $(z, s) \notin \mathbb{S} \times [-M, M]$,

$$\hat{\mathcal{G}}f(z, s) = V(z) \frac{\partial f}{\partial s}(z, s) + L_z f(z, s) = 0,$$

and

$$\begin{aligned} |\hat{\mathcal{G}}g_n(z, s) - \hat{\mathcal{G}}f(z, s)| &= \left| V(z) \frac{\partial g_n}{\partial s}(z, s) + L_z g_n(z, s) \right| \\ &= \left| V(z)g'(s)f_n(z, s) + V(z) \frac{\partial f_n}{\partial s}(z, s)g(s) + g(s)L_z f_n(z, s) \right| \\ &\leq |V(z)| \cdot |g'(s)| \cdot |f_n(z, s)| + |g(s)| \left| V(z) \frac{\partial f_n}{\partial s}(z, s) + L_z f_n(z, s) \right| \\ &\leq \|V\|_\infty |f_n(z, s)| + |\hat{\mathcal{G}}f_n(z, s)| \\ &= \|V\|_\infty |f_n(z, s) - f(z, s)| + |\hat{\mathcal{G}}f_n(z, s) - \hat{\mathcal{G}}f(z, s)| \\ &\leq \|V\|_\infty \|f_n - f\|_\infty + \|\hat{\mathcal{G}}f_n - \hat{\mathcal{G}}f\|_\infty. \end{aligned}$$

Hence

$$\|g_n - f\|_\infty + \|\hat{\mathcal{G}}g_n - \hat{\mathcal{G}}f\|_\infty \leq (1 + \|V\|_\infty) \|f_n - f\|_\infty + \|\hat{\mathcal{G}}f_n - \hat{\mathcal{G}}f\|_\infty$$

and we see that

$$\lim_{n \rightarrow \infty} (\|g_n - f\|_\infty + \|\hat{\mathcal{G}}g_n - \hat{\mathcal{G}}f\|_\infty) = 0.$$

Note that for every $M > 0$ there exists a constant $C_{M,V} > 0$ such that

$$\|R^{\mathbb{S}}h\|_\infty \leq C_{M,V} \|h\|_\infty$$

for all $h \in C_c(\mathbb{S} \times \mathbb{R})$ such that $\text{supp}(h) \subset \mathbb{S} \times [-M + 3, M + 3]$. Hence

$$\begin{aligned} & \|g_n - f\|_\infty + \|\mathcal{G}g_n - \mathcal{G}f\|_\infty \\ &= \|g_n - f\|_\infty + \|\hat{\mathcal{G}}g_n - \hat{\mathcal{G}}f + R^{\mathbb{S}}g_n - R^{\mathbb{S}}f\|_\infty \\ &\leq \|g_n - f\|_\infty + \|\hat{\mathcal{G}}g_n - \hat{\mathcal{G}}f\|_\infty + \|R^{\mathbb{S}}g_n - R^{\mathbb{S}}f\|_\infty \\ &\leq (1 + \|V\|_\infty)\|f_n - f\|_\infty + \|\hat{\mathcal{G}}f_n - \hat{\mathcal{G}}f\|_\infty + C_{M,V}\|g_n - f\|_\infty \\ &\leq (1 + \|V\|_\infty + C_{M,V})\|f_n - f\|_\infty + \|\hat{\mathcal{G}}f_n - \hat{\mathcal{G}}f\|_\infty \\ &\xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

This shows that for every $f \in \mathcal{D}(\mathcal{G}) \cap C_c(\mathbb{S} \times \mathbb{R})$ there exists a sequence $(g_n)_{n=1}^\infty$, such that $g_n \in C_c^2(\mathbb{S} \times \mathbb{R})$ and

$$\|g_n - f\|_\infty + \|\mathcal{G}g_n - \mathcal{G}f\|_\infty \xrightarrow{n \rightarrow \infty} 0.$$

Since we know that $\mathcal{D}(\mathcal{G}) \cap C_c(\mathbb{S} \times \mathbb{R})$ is a core for $(\mathcal{G}, \mathcal{D}(\mathcal{G}))$, we conclude that $C_c^2(\mathbb{S} \times \mathbb{R})$ is also a core for $(\mathcal{G}, \mathcal{D}(\mathcal{G}))$. \square

We will now identify the form of the stationary distribution of the process (Z_t, S_t) . For this we need two auxiliary results, Lemma 3.4 and Proposition 3.5.

Lemma 3.4. *For any $f \in C^2(\mathbb{S})$ we have*

$$\int_{\mathbb{S}} Lf(z) \, dz = 0.$$

The proof of this lemma is the same as the proof of [3], Lemma 2.8, and is omitted.

Proposition 3.5. *Let*

$$\pi(dz, ds) = \frac{1}{2\pi} e^{-\pi s^2} \, dz \, ds.$$

Then for any $f \in C_c^2(\mathbb{S} \times \mathbb{R})$ we have

$$\int_{\mathbb{S}} \int_{\mathbb{R}} \mathcal{G}f(z, s) \pi(dz, ds) = 0.$$

Proof. Let $f \in C_c^2(\mathbb{S} \times \mathbb{R})$. By Corollary 2.5 we have

$$\begin{aligned} & 2\pi \int_{\mathbb{S}} \int_{\mathbb{R}} \mathcal{G}f(z, s) \pi(dz, ds) \\ &= \int_{\mathbb{R}} \int_{\mathbb{S}} L_z f(z, s) \, dz e^{-\pi s^2} \, ds + \int_{\mathbb{S}} V(z) \int_{\mathbb{R}} f_s(z, s) e^{-\pi s^2} \, ds \, dz \\ &\quad + \int_{\mathbb{R}} \int_{\mathbb{S}} \int_{\mathbb{S}} (f(w, s) - f(z, s)) ((V(z) - V(w))s)_+ \, dw \, dz e^{-\pi s^2} \, ds \\ &= \text{I} + \text{II} + \text{III}. \end{aligned}$$

From Lemma 3.4 we know that $\text{I} = 0$. Integrating by parts we obtain

$$\text{II} = 2\pi \int_{\mathbb{S}} \int_{\mathbb{R}} V(z) f(z, s) e^{-\pi s^2} \, ds \, dz.$$

Now we will simplify III. Note that $a_+ = (a + |a|)/2$, $a \in \mathbb{R}$. Hence

$$\begin{aligned} & \int_{\mathbb{S}} \int_{\mathbb{S}} (f(w, s) - f(z, s)) ((V(z) - V(w))s)_+ dw dz \\ &= \frac{s}{2} \int_{\mathbb{S}} \int_{\mathbb{S}} (f(w, s) - f(z, s)) (V(z) - V(w)) dw dz \\ & \quad + \frac{|s|}{2} \int_{\mathbb{S}} \int_{\mathbb{S}} (f(w, s) - f(z, s)) |V(z) - V(w)| dw dz \\ &= \text{III}_1 + \text{III}_2. \end{aligned}$$

By interchanging w and z in III_2 we get

$$\text{III}_2 = \frac{|s|}{2} \int_{\mathbb{S}} \int_{\mathbb{S}} (f(z, s) - f(w, s)) |V(w) - V(z)| dw dz = -\text{III}_2,$$

which means that $\text{III}_2 = 0$.

By assumption, $\int_{\mathbb{S}} V(z) dz = 0$. Therefore

$$\begin{aligned} \text{III}_1 &= \frac{s}{2} \int_{\mathbb{S}} f(w, s) dw \int_{\mathbb{S}} V(z) dz - \frac{s}{2} \int_{\mathbb{S}} f(w, s) V(w) dw \int_{\mathbb{S}} dz \\ & \quad - \frac{s}{2} \int_{\mathbb{S}} f(z, s) V(z) dz \int_{\mathbb{S}} dw + \frac{s}{2} \int_{\mathbb{S}} f(z, s) dz \int_{\mathbb{S}} V(w) dw \\ &= -2\pi s \int_{\mathbb{S}} f(z, s) V(z) dz. \end{aligned}$$

Informally, $\text{III} = \int ((\text{III}_1)e^{-\pi s^2}) ds$, so

$$\text{III} = -2\pi \int_{\mathbb{S}} \int_{\mathbb{R}} V(z) f(z, s) e^{-\pi s^2} s ds dz.$$

Consequently $\text{I} + \text{II} + \text{III} = 0$. □

Theorem 3.6. *The measure*

$$\pi(dz, ds) = \frac{1}{2\pi} e^{-\pi s^2} dz ds \tag{3.1}$$

is a stationary distribution of the process (Z_t, S_t) .

Proof. Let (Y_t, S_t) be the Markov process given by (2.3) and let (Z_t, S_t) be the Markov process where $Z_t = e^{iY_t}$. By $\{T_t^{\mathbb{S}}\}_{t \geq 0}$ we denote the transition semigroup of (Z_t, S_t) on the Banach space $C_0(\mathbb{S} \times \mathbb{R})$, cf. (2.4), and by \mathcal{G} we denote its generator. Let $\mathcal{P}(\mathbb{R} \times \mathbb{R})$ and $\mathcal{P}(\mathbb{S} \times \mathbb{R})$ denote the sets of all probability measures on $\mathbb{R} \times \mathbb{R}$ and $\mathbb{S} \times \mathbb{R}$ respectively. In this proof, for any $\tilde{\mu} \in \mathcal{P}(\mathbb{S} \times \mathbb{R})$ we define $\mu \in \mathcal{P}(\mathbb{R} \times \mathbb{R})$ by $\mu([0, 2\pi) \times \mathbb{R}) = 1$ and $\mu(A \times B) = \tilde{\mu}(e^{iA} \times B)$ for Borel sets $A \subset [0, 2\pi)$, $B \subset \mathbb{R}$.

Consider any $\tilde{\mu} \in \mathcal{P}(\mathbb{S} \times \mathbb{R})$ and the corresponding $\mu \in \mathcal{P}(\mathbb{R} \times \mathbb{R})$. For this μ there exists a Markov process (Y_t, S_t) given by (2.3) such that (Y_0, S_0) has the distribution μ . It follows that for any $\tilde{\mu} \in \mathcal{P}(\mathbb{S} \times \mathbb{R})$ there exists a Markov process (Z_t, S_t) with $Z_t = e^{iY_t}$ and with initial distribution $\tilde{\mu}$. By [6], Proposition 4.1.7, (Z_t, S_t) is a solution of the martingale problem for $(\mathcal{G}, \tilde{\mu})$. By [6], Theorem 4.4.1, for any $\tilde{\mu} \in \mathcal{P}(\mathbb{S} \times \mathbb{R})$, uniqueness holds for the martingale problem for $(\mathcal{G}, \tilde{\mu})$. Hence the martingale problem for \mathcal{G} is well posed.

Proposition 3.3 gives that $C_c^2(\mathbb{S} \times \mathbb{R})$ is a core for \mathcal{G} . By Proposition 3.5 and [6], Proposition 4.9.2, we get that π is a stationary measure for \mathcal{G} . This means that π is a stationary distribution for (Z_t, S_t) . □

Theorem 3.7. *The measure π defined in (3.1) is the unique stationary distribution of the process (Z_t, S_t) .*

Proof. The proof is similar to the proof of [3], Theorem 2.1.2.

Step 1. Suppose that (Y_t, S_t) satisfies

$$Y_t = y + X_t, \quad S_t = s + \int_0^t W(Y_r) dr,$$

where $X_0 = 0$. Suppose that X_t is a stable Lévy process with $X_0 = 0$. The following Lévy inequality for symmetric Lévy processes is well known

$$\mathbb{P}\left(\sup_{0 \leq r \leq \tau} |X_r| > \varepsilon\right) \leq 2\mathbb{P}(|X_\tau| > \varepsilon) \leq 1 - \delta.$$

It follows that for every $\tau < \infty$, $y, s \in \mathbb{R}$ and $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\mathbb{P}^{y,s}\left(\sup_{0 \leq r \leq \tau} |Y_r - y| \leq \varepsilon\right) = \mathbb{P}\left(\sup_{0 \leq r \leq \tau} |X_r| \leq \varepsilon\right) \geq \delta. \quad (3.2)$$

Step 2. Recall that $V \in C^3$ and it is not identically constant. This and the fact that $\int_{\mathbb{S}} V(z) dz = 0$ imply that W is strictly positive on some interval and strictly negative on some other interval. We fix some $a_1, a_2 \in (-\pi, \pi)$, $b_1 > 0$, $b_2 < 0$ and $\varepsilon_0 \in (0, \pi/100)$, such that $V(z) > b_1$ for $z \in \mathbb{S}$, $\text{Arg}(z) \in [a_1 - 4\varepsilon_0, a_1 + 4\varepsilon_0]$, and $V(z) < b_2$ for $z \in \mathbb{S}$, $\text{Arg}(z) \in [a_2 - 4\varepsilon_0, a_2 + 4\varepsilon_0]$.

Suppose that there exist two stationary probability distributions π and $\hat{\pi}$ for (Z, S) . Let $((Z_t, S_t))_{t \geq 0}$ and $((\hat{Z}_t, \hat{S}_t))_{t \geq 0}$ be processes with (Z_0, S_0) and (\hat{Z}_0, \hat{S}_0) distributed according to π and $\hat{\pi}$, respectively. The transition probabilities for these processes are the same as for the processes which are solutions to (2.3). Recall that X denotes the driving stable Lévy process for Z and τ_1 is the time of the first “extra jump” in the representation (2.3).

We will show that $S_t \neq 0$ for some $t > 0$, a.s. Suppose that the event $A = \{S_t = 0 \text{ for all } t \geq 0\}$ has strictly positive probability. On A we have $Y_t = X_t + y$ for all $t \geq 0$, according to (2.3). Recall that $W(x) > 0$ for all x in the set $\Gamma := \bigcup_{k \in \mathbb{Z}} (a_1 - 4\varepsilon_0 + 2\pi k, a_1 + 4\varepsilon_0 + 2\pi k)$. It is easy to see that X enters $\Gamma - y$ at a finite time s_0 , a.s. Hence, Y enters Γ at a finite time s_0 , on the event A . Since Y is right-continuous, $Y_t \in \Gamma$ for all $t \in (s_0, s_1)$ for some random $s_1 > s_0$. This and (2.3) imply that $S_t \neq 0$ for some $t \in (s_0, s_1)$, on the event A . This contradicts the definition of A and hence it proves that $S_t \neq 0$ for some $t > 0$, a.s.

Assume without loss of generality that $S_t > 0$ for some $t > 0$, with positive probability. Then there exist $\varepsilon_1 > 0$, $t_1 > 0$ and $p_1 > 0$ such that

$$\mathbb{P}^\pi(S_{t_1} > \varepsilon_1, \tau_1 > t_1) > p_1.$$

Let $F_1 = \{S_{t_1} > \varepsilon_1, \tau_1 > t_1\}$ and $t_2 = \varepsilon_1/(2\|W\|_\infty)$. Clearly, for some $p_2 > 0$ we have

$$\mathbb{P}^\pi(\exists t \in [t_1, t_1 + t_2]: \text{Arg}(Z_t) \in [a_2 - \varepsilon_0, a_2 + \varepsilon_0], \tau_1 > t_1 + t_2 | F_1) > p_2.$$

Since $\text{Arg}(Z_t)$ has right-continuous paths, this implies that there exist $\varepsilon_1 > 0$, $t_1 > 0$, $t_3 \in [t_1, t_1 + t_2]$ and $p_3 > 0$ such that

$$\mathbb{P}^\pi(S_{t_1} > \varepsilon_1, \text{Arg}(Z_{t_3}) \in [a_2 - 2\varepsilon_0, a_2 + 2\varepsilon_0], \tau_1 > t_3) > p_3.$$

Note that $|S_{t_3} - S_{t_1}| \leq \|W\|_\infty t_2 < \varepsilon_1/2$. Hence,

$$\mathbb{P}^\pi(S_{t_3} > \varepsilon_1/2, \text{Arg}(Z_{t_3}) \in [a_2 - 2\varepsilon_0, a_2 + 2\varepsilon_0], \tau_1 > t_3) > p_3.$$

Let $\varepsilon_2 \in (\varepsilon_1/2, \infty)$ be such that

$$\mathbb{P}^\pi(S_{t_3} \in [\varepsilon_1/2, \varepsilon_2], \text{Arg}(Z_{t_3}) \in [a_2 - 2\varepsilon_0, a_2 + 2\varepsilon_0], \tau_1 > t_3) > p_3/2.$$

Set $t_4 = 2\varepsilon_2/|b_2|$ and $t_5 = t_3 + t_4$. By (3.2), for any $\varepsilon_3 > 0$ and some $p_4 > 0$,

$$\mathbb{P}^\pi \left(\sup_{t_3 \leq r \leq t_5} |X_r - X_{t_3}| \leq \varepsilon_3, S_{t_3} \in [\varepsilon_1/2, \varepsilon_2], \right. \\ \left. \text{Arg}(Z_t) \in [a_2 - 3\varepsilon_0, a_2 + 3\varepsilon_0] \text{ for all } t \in [t_3, t_5], \tau_1 > t_5 \right) > p_4.$$

Since $V(x) < b_2 < 0$ for $x \in [a_2 - 3\varepsilon_0, a_2 + 3\varepsilon_0]$, if the event in the last formula holds then

$$S_{t_5} = S_{t_3} + \int_{t_3}^{t_5} V(Z_s) ds \leq \varepsilon_2 + b_2 t_4 \leq -\varepsilon_2.$$

This implies that,

$$\mathbb{P}^\pi \left(\sup_{t_3 \leq r \leq t_5} |X_r - X_{t_3}| \leq \varepsilon_3, S_{t_3} \geq \varepsilon_1/2, S_{t_5} \leq -\varepsilon_2, \tau_1 > t_5 \right) > p_4. \quad (3.3)$$

Step 3. By the Lévy–Itô representation we can write the stable Lévy process X in the form $X_t = J_t + \tilde{X}_t$, where J is a compound Poisson process comprising all jumps of X which are greater than ε_0 and $\tilde{X} = X - J$ is an independent Lévy process (accounting for all small jumps of X). Denote by $\lambda = \lambda(\alpha, \varepsilon_0)$ the rate of the compound Poisson process J and let (\tilde{Y}, \tilde{S}) be the solution to (2.3), with X_t replaced by \tilde{X}_t for $t \geq t_3$. Similarly $\tilde{\tau}_1$ denotes the first “extra jump” in the representation (2.3) for the process (\tilde{Y}, \tilde{S}) . Moreover, we take $\varepsilon_3 < \varepsilon_0/2$. By our construction $\sup_{t_3 \leq r \leq t_5} |X_r - X_{t_3}| \leq \varepsilon_3$ entails that $\sup_{t_3 \leq r \leq t_5} |J_r - J_{t_3}| = 0$; therefore, (3.3) becomes

$$\mathbb{P}^\pi \left(\sup_{t_3 \leq r \leq t_5} |\tilde{X}_r - \tilde{X}_{t_3}| \leq \varepsilon_3, \tilde{S}_{t_3} \geq \frac{\varepsilon_1}{2}, \tilde{S}_{t_5} \leq -\varepsilon_2, \tilde{\tau}_1 > t_5 \right) \\ \geq \mathbb{P}^\pi \left(\sup_{t_3 \leq r \leq t_5} |\tilde{X}_r - \tilde{X}_{t_3}| \leq \varepsilon_3, \sup_{t_3 \leq r \leq t_5} |J_r - J_{t_3}| = 0, \tilde{S}_{t_3} \geq \frac{\varepsilon_1}{2}, \tilde{S}_{t_5} \leq -\varepsilon_2, \tilde{\tau}_1 > t_5 \right) \\ > p_4 > 0.$$

Let τ be the time of the first jump of J in the interval $[t_3, t_5]$; we set $\tau = t_5$ if there is no such jump. We can represent $\{(Y_t, S_t), 0 \leq t \leq \tau\}$ in the following way: $(Y_t, S_t) = (\tilde{Y}_t, \tilde{S}_t)$ for $0 \leq t < \tau$, $S_\tau = \tilde{S}_\tau$, and $Y_\tau = \tilde{Y}_{\tau-} + J_\tau - J_{\tau-}$. Note that $\tilde{Y}_t = y + \tilde{X}_t$ if $t < \tau_1$.

We say that a non-negative measure μ_1 is a component of a non-negative measure μ_2 if $\mu_2 = \mu_1 + \mu_3$ for some non-negative measure μ_3 . Let $\mu(dz, ds) = \mathbb{P}^\pi(Z_\tau \in dz, S_\tau \in ds)$. We will argue that $\mu(dz, ds)$ has a component with a density bounded below by $c_2 > 0$ on $\mathbb{S} \times (-\varepsilon_2, \varepsilon_1/2)$. We find for every Borel set $A \subset \mathbb{S}$ of arc length $|A|$ and every interval $(s_1, s_2) \subset (-\varepsilon_2, \varepsilon_1/2)$

$$\mu(A \times (s_1, s_2)) \\ = \mathbb{P}^\pi(Z_\tau \in A, S_\tau \in (s_1, s_2)) \\ \geq \mathbb{P}^\pi \left(Z_\tau \in A, S_\tau \in (s_1, s_2), \sup_{t_3 \leq r \leq t_5} |\tilde{X}_r - \tilde{X}_{t_3}| \leq \varepsilon_3, \tilde{S}_{t_3} \geq \frac{\varepsilon_1}{2}, \tilde{S}_{t_5} \leq -\varepsilon_2, \tilde{\tau}_1 > t_5 \right) \\ \geq \mathbb{P}^\pi \left(e^{i(J_\tau - J_{\tau-})} \in e^{-i\tilde{X}_{\tau-}} A, \tilde{S}_\tau \in (s_1, s_2), \right. \\ \left. \sup_{t_3 \leq r \leq t_5} |\tilde{X}_r - \tilde{X}_{t_3}| \leq \varepsilon_3, \tilde{S}_{t_3} \geq \varepsilon_1/2, \tilde{S}_{t_5} \leq -\varepsilon_2, \tilde{\tau}_1 > t_5, N^J = 1 \right).$$

Here N^J counts the number of jumps of the process J occurring during the interval $[t_3, t_5]$. Without loss of generality we can assume that $\varepsilon_0 < 2\pi$. In this case the density of the jump measure of J is bounded below by $c_3 > 0$ on $(2\pi, 4\pi)$. Observe that the processes (\tilde{X}, \tilde{S}) and J are independent. Conditional on $\{N^J = 1\}$, τ is uniformly distributed on

$[t_3, t_5]$, and the probability of the event $\{N^J = 1\}$ is $\lambda(t_5 - t_3)e^{-\lambda(t_5 - t_3)}$. Thus,

$$\begin{aligned} & \mu(A \times (s_1, s_2)) \\ & \geq c_3 |A| \mathbb{P}^\pi \left(\tilde{X}_\tau \in (s_1, s_2) \mid \sup_{t_3 \leq r \leq t_5} |\tilde{X}_r - \tilde{X}_{t_3}| \leq \varepsilon_3, \tilde{S}_{t_3} \geq \varepsilon_1/2, \tilde{S}_{t_5} \leq -\varepsilon_2, \tilde{\tau}_1 > t_5, N^J = 1 \right) \\ & \quad \times p_4 \cdot \lambda(t_5 - t_3) e^{-\lambda(t_5 - t_3)}. \end{aligned}$$

Since the process \tilde{S} spends at least $(s_2 - s_1)/\|W\|_\infty$ units of time in (s_1, s_2) we finally arrive at

$$\mu(A, (s_1, s_2)) \geq p_4 \lambda e^{-\lambda(t_5 - t_3)} c_3 |A| (s_2 - s_1) / \|W\|_\infty.$$

This proves that $\mu(dz, ds)$ has a component with a density bounded below by $c_2 = p_4 \lambda e^{-\lambda(t_5 - t_3)} c_3 / \|W\|_\infty$ on $\mathbb{S} \times (-\varepsilon_2, \varepsilon_1/2)$.

Step 4. Let $\varepsilon_4 = \frac{\varepsilon_1}{2} \wedge \varepsilon_2 > 0$. We have shown that for some stopping time τ , $\mathbb{P}^\pi(Z_\tau \in dz, S_\tau \in ds)$ has a component with a density bounded below by $c_2 > 0$ on $\mathbb{S} \times (-\varepsilon_4, \varepsilon_4)$. We can prove in an analogous way that for some stopping time $\hat{\tau}$ and $\hat{\varepsilon}_4 > 0$, $\mathbb{P}^{\hat{\pi}}(\hat{Z}_{\hat{\tau}} \in dz, \hat{S}_{\hat{\tau}} \in ds)$ has a component with a density bounded below by $\hat{c}_2 > 0$ on $\mathbb{S} \times (-\hat{\varepsilon}_4, \hat{\varepsilon}_4)$.

Since $\pi \neq \hat{\pi}$, there exists a Borel set $A \subset \mathbb{S} \times \mathbb{R}$ such that $\pi(A) \neq \hat{\pi}(A)$. Moreover, since any two stationary probability measures are either mutually singular or identical, cf. [9], Chapter 2, Theorem 4, we have $\pi(A) > 0$ and $\hat{\pi}(A) = 0$ for some A . By the strong Markov property applied at τ and the ergodic theorem, see [9], Chapter 1, page 12, we have \mathbb{P}^π -a.s.

$$\lim_{t \rightarrow \infty} (1/t) \int_\tau^t \mathbb{1}_{\{(Z_s, S_s) \in A\}} ds = \pi(A) > 0.$$

Similarly, we see that $\mathbb{P}^{\hat{\pi}}$ -a.s.

$$\lim_{t \rightarrow \infty} (1/t) \int_{\hat{\tau}}^t \mathbb{1}_{\{(\hat{Z}_s, \hat{S}_s) \in A\}} ds = \hat{\pi}(A) = 0.$$

Since the distributions of (Z_τ, S_τ) and $(\hat{Z}_{\hat{\tau}}, \hat{S}_{\hat{\tau}})$ have mutually absolutely continuous components, the last two statements contradict each other. This shows that we must have $\pi = \hat{\pi}$. \square

Acknowledgement

We are grateful to an anonymous referee for suggestions for improvement.

References

- [1] M. Barlow, A. Grigor'yan and T. Kumagai. Heat kernel upper bounds for jump processes and the first exit time. *J. Reine Angew. Math.* **626** (2009) 135–157. [MR2492992](#)
- [2] R. Bass, K. Burdzy, Z. Chen and M. Hairer. Stationary distributions for diffusions with inert drift. *Probab. Theory Related Fields* **146** (2010) 1–47. [MR2550357](#)
- [3] K. Burdzy, T. Kulczycki and R. Schilling. Stationary distributions for jump processes with inert drift. Preprint, 2010. Available at [arXiv:1009.2347](#).
- [4] K. Burdzy and D. White. A Gaussian oscillator. *Electron. Commun. Probab.* **9** (2004) 92–95. [MR2108855](#)
- [5] K. Burdzy and D. White. Markov processes with product-form stationary distribution. *Electron. Commun. Probab.* **13** (2008) 614–627. [MR2461535](#)
- [6] S. N. Ethier and T. G. Kurtz. *Markov Processes: Characterization and Convergence*. Wiley, New York, 1986. [MR0838085](#)
- [7] N. Ikeda, N. Nagasawa and S. Watanabe. A construction of Markov processes by piecing out. *Proc. Japan Acad. Ser. A Math. Sci.* **42** (1966) 370–375. [MR0202197](#)
- [8] P.-A. Meyer. Renaissance, recollements, mélanges, ralentissement de processus de Markov. *Ann. Inst. Fourier (Grenoble)* **25** (1975) 464–497. [MR0415784](#)
- [9] Ya. G. Sinai. *Topics in Ergodic Theory*. Princeton Univ. Press, Princeton, NJ, 1994. [MR1258087](#)