

A branching-selection process related to censored Galton–Watson processes

Olivier Couronné and Lucas Gerin

Université Paris Ouest Nanterre La Défense, Equipe Modal'X, 200 avenue de la République, 92000 Nanterre, France.
E-mail: ocouronn@u-paris10.fr

Received 2 November 2011; revised 4 June 2012; accepted 19 June 2012

Abstract. We obtain the asymptotics for the speed of a particular case of a particle system with branching and selection introduced by Bérard and Gouéré [*Comm. Math. Phys.* **298** (2010) 323–342]. The proof is based on a connection with a supercritical Galton–Watson process *censored* at a certain level.

Résumé. Nous étudions un cas particulier de système de particules avec branchement et sélection introduit par Bérard et Gouéré [*Comm. Math. Phys.* **298** (2010) 323–342]. Nous obtenons l'asymptotique pour la vitesse, en remarquant un lien avec un processus de Galton–Watson surcritique censuré à un certain niveau.

MSC: 60J80

Keywords: Galton–Watson process; Censored branching process; Branching and selection

1. Models and results

1.1. The censored Galton–Watson process

For a probability distribution \mathcal{X} on non-negative integers, the Galton–Watson process with *offspring* \mathcal{X} is the process defined by $Z_0^a = a$ and

$$Z_{k+1}^a = \sum_{i=1}^{Z_k^a} X_{k,i},$$

where the $X_{k,i}$'s are i.i.d. copies of \mathcal{X} ($Z_{k+1}^a = 0$ if $Z_k^a = 0$). We write $(Z_k) = (Z_k^1)$. We deal here with supercritical Galton–Watson processes (i.e. when $\mathbb{E}(\mathcal{X}) > 1$) and to avoid trivialities we impose $\mathcal{X}(0) > 0$. In this case, it is well-known (we refer to [1] for basic facts on branching processes) that Z_k dies with a certain probability $0 < q < 1$ which is the unique solution in $(0, 1)$ of

$$f(x) := \sum_{i \geq 0} \mathcal{X}(i)x^i = x.$$

The first step of this article will be to study a kind of constrained Galton–Watson process, in which the constraint is a “roof” at a given height that prevents the process from exploding.

Definition 1. Given N a positive integer and a probability distribution \mathcal{X} on integers, the Galton–Watson process censored at level $N \geq 2$ with offspring \mathcal{X} is the process $(X_k^N)_{k \geq 0}$ defined by $X_0^N = N$ and, for $k \geq 0$,

$$X_{k+1}^N = \begin{cases} \min(N, \sum_{i=1}^{X_k^N} X_{k,i}) & \text{if } X_k^N > 0, \\ 0 & \text{otherwise,} \end{cases}$$

where the $X_{k,i}$'s are i.i.d. copies of \mathcal{X} .

The process $(X_k^N)_{k \geq 0}$ is a finite state Markov process which is typically stuck for a long time on N , but eventually dies. Let U_N be the survival time of the Galton–Watson process:

$$U_N = \min\{k \geq 0, X_k^N = 0\}.$$

Let us describe heuristically the asymptotic behavior of U_N . When the censored process dies, it happens very suddenly: in the uncensored underlying process, the progenies of the N particles pass away almost simultaneously in a few generations. This latter event occurs with a probability close to q^N , and therefore the censored process is expected to survive a time close to a geometric random variable with parameter q^N .

We state this in the following theorem, which may be derived from results of [5].

Theorem 1. *The following convergence holds in law:*

$$U_N q^N \xrightarrow{N \rightarrow \infty} \mathcal{E}(1), \quad (1)$$

where $\mathcal{E}(1)$ stands for the exponential distribution with parameter one. The convergence also holds in mean:

$$\mathbb{E}(U_N) \sim (1/q)^N \quad \text{as } N \rightarrow \infty. \quad (2)$$

The convergence in law (1) is a consequence of ([5], Th. 1 (3), (4)) and the asymptotic (2) follows from ([5], Th. 1 (2)). The results of [5] are actually much accurate. As we will only need these simple estimates, and for the sake of completeness, we provide in the two following sections an elementary and more concise proof of Theorem 1. We also mention [6], in which the Galton–Watson process is censored by a function depending on time; the author obtained a criterion for the degeneracy of the process.

1.2. A connection with a branching-selection process

The present authors considered Theorem 1 when trying to find an asymptotic of the speed of a very particular case of a certain branching particle system with selection, studied by Bérard and Guéré ([2], Section 7, Theorem 5). The *branching-selection* process we will study is a generalization of their Bernoulli branching-selection process and can be described as follows. For a distribution $(\mathcal{X}, \mathcal{X}')$ on \mathbb{N}^2 such that the expectation of \mathcal{X} is strictly greater than 1 and $\mathcal{X} + \mathcal{X}' \geq 1$ a.s., the particle system is the discrete-time particle system of N particles moving on \mathbb{Z} and starting from the origin such that, at each time unit,

1. Each of the N particles is replaced by \mathcal{X} particles just on the right of that particle and by \mathcal{X}' particles on the same position, independently of each other;
2. Among all these new particles, we keep only the N rightmost particles.

It is convenient to see the process of the locations of the N particles as a finite point measure. We set $Y_0^N = N\delta_0$ and write

$$Y_k^N = \sum_{\ell \geq 0} \delta_\ell Y_k^N(\ell),$$

where $Y_k^N(\ell)$ is the number of particles at time k at the position ℓ ; by construction we have at any time $\sum_{\ell \geq 0} Y_k^N(\ell) = N$. We write

$$\max Y_k^N := \max\{\ell \geq 0; Y_k^N(\ell) > 0\}$$

for the position of the rightmost particle at time k . We have $\max Y_k^N \leq k$ by construction. Since $\mathbb{E}(\mathcal{X}) > 1$, and since $Y_{k+1}^N \geq Y_k^N$ by the property of \mathcal{X}' , it is likely for N large that $\max Y_k^N$ is close to k . Remark also that in this process, all the particles at time k are on $\{\max Y_k^N - 1, \max Y_k^N\}$ after the selection.

Bérard and Gouéré in fact studied the particle system defined as follows. At every time unit each particle is duplicated and moves one step forward with probability $\alpha \in (\frac{1}{2}, 1)$ and stays put with probability $1 - \alpha$. With our notations, this corresponds to the case where \mathcal{X} is a Bernoulli variable $B(2, \alpha)$ with $\alpha > \frac{1}{2}$, and $\mathcal{X}' = 2 - \mathcal{X}$ (we will denote their model as the *Bernoulli branching-selection process*). They proved that $(\max Y_k^N)/k$ converges almost surely to a constant $v_N(\alpha)$, and noticed that $v_N(\alpha)$ lies between $(1 - \exp(-c_1(\alpha)N))$ and $(1 - \exp(-c_2(\alpha)N))$ for some positive constants $c_1(\alpha)$ and $c_2(\alpha)$. The stress of their paper was a large class of distributions for the walk performed by the particles, and the case of the Bernoulli random walk with $\alpha > 1/2$ was in fact just mentioned as a degenerate case. The existence of the asymptotic speed $v_N(\alpha)$ would apply as well in our case, but not their proof for the bounds on this speed.

The key point for our result is that the process of the number of particles that are at the rightmost possible position $(Y_k^N(k))_k$ has the same law as the censored Galton–Watson process $(X_k^N)_k$, when the offspring distribution is \mathcal{X} .

Theorem 2. *For the branching-selection process with the distribution $(\mathcal{X}, \mathcal{X}')$ on \mathbb{N}^2 such that $\mathbb{E}(\mathcal{X}) > 1$ and $\mathcal{X} + \mathcal{X}' \geq 1$ a.s., we have*

$$\lim_{N \rightarrow \infty} \frac{1 - v_N(\alpha)}{q^N} = 1,$$

where q is the extinction probability of a Galton–Watson process whose offspring is \mathcal{X} .

In particular, for the Bernoulli branching-selection model studied in [2] with $\alpha > \frac{1}{2}$,

$$\lim_{N \rightarrow \infty} \frac{1 - v_N(\alpha)}{(q_\alpha)^N} = 1,$$

where

$$q_\alpha = \frac{1 - 2\alpha(1 - \alpha) - \sqrt{1 - 4\alpha(1 - \alpha)}}{2\alpha^2}$$

is the extinction probability of a Galton–Watson process whose offspring is the Binomial distribution with parameters $(2, \alpha)$.

Remark that the exact law of \mathcal{X}' has no influence, and we could have always taken

$$\mathcal{X}' = 1_{\mathcal{X}=0}.$$

Using \mathcal{X}' allows us essentially to have a true generalization of the Bernoulli branching-selection process.

The article is organized as follows. The proof of Theorem 1 is decomposed over Sections 2 and 3. A key ingredient in Section 3 is to compare our censored process at the last time it reaches the level N , and the classical Galton–Watson process starting at N and conditioned to die. The main contribution is the application to the branching-selection particle system and is given in Section 4.

2. Preliminaries

We will repeatedly use the following left-tail bound for a sum of copies of \mathcal{X} :

Lemma 1. *Let X_1, \dots, X_N be i.i.d. copies of law \mathcal{X} . There exist $b, c > 0$ such that, when N is large enough,*

$$P(X_1 + \dots + X_N \leq N(1 + b)) \leq \exp(-cN).$$

To prove this, we choose M such that $\mathbb{E}[\min\{X_1, M\}] > 1$. We then can apply ([4], III, Chap. 4) to the bounded random variables $\min\{X_i, M\}$ and we get the desired bound. This implies in particular that

$$\mathbb{P}(X_{k+1}^N < N | X_k^N = N) \leq \exp(-cN). \quad (3)$$

Note that throughout the paper, c stands for a generic positive constant, which might differ at each appearance. We introduce the last time for which our process is equal to N . This variable will turn out to be equivalent to U_N and more simple to approximate by a geometric variable.

Definition 2. *Let*

$$V_N = \max\{k \geq 0, X_k^N = N\},$$

and let q_N be the probability that $(X_k^N)_k$ does not ever hit N after time zero:

$$q_N = \mathbb{P}(X_k^N < N, \forall k > 0) = \mathbb{P}(V_N = 0).$$

We deduce from (3) that

$$q_N \leq \mathbb{P}(X_1^N < N) \leq \exp(-cN). \quad (4)$$

We will compare q_N with the probability that the classical Galton–Watson process (Z_k^N) starting from N particles dies out. By independence of the progeny of the N particles, we have $\mathbb{P}((Z_k^N) \text{ dies out}) = q^N$. By conditioning over the first passage over some level n , we get

$$\mathbb{P}((Z_k) \text{ dies out} | \exists k \geq 0 \text{ such that } Z_k \geq n) \leq q^n. \quad (5)$$

3. Proof of Theorem 1

The main tool of the proof is the following lemma, which says that the Galton–Watson process (Z_k^N) , conditioned to die, roughly behaves like X_k^N after its last passage in N . Recall that q stands for the probability that Z dies while q_N is the probability that X^N does not ever hit N .

Lemma 2. *There exists $c > 0$ such that for N large enough*

$$\left| \frac{q_N}{q^N} - 1 \right| \leq \exp(-cN).$$

Proof. By Lemma 1, we can take $b > 0$ and $c > 0$ such that

$$\mathbb{P}(Z_1^N < (1+b)N) \leq \exp(-cN).$$

The point is that if $(Z_k^N)_k$ dies out, then with high probability it does not hit N after time $k = 0$. Let $R = \inf\{k \geq 1, Z_k^N \geq N\}$ (with the convention that $R = +\infty$ if Z^N does not hit N after time zero),

$$\begin{aligned} & \mathbb{P}(R < \infty; Z^N \text{ dies out}) \\ & \leq \mathbb{P}(Z_1^N < (1+b)N; R < \infty; Z^N \text{ dies out}) + \mathbb{P}(Z_1^N \geq (1+b)N; Z^N \text{ dies out}) \\ & \leq \mathbb{P}(Z_1^N < (1+b)N; R < \infty; Z^N \text{ dies out}) + q^{(1+b)N} \\ & \leq \sum_{\ell \geq N} \mathbb{P}(Z_R^N = \ell; R < +\infty; Z_1^N < (1+b)N; Z^N \text{ dies out}) + q^{(1+b)N} \end{aligned}$$

$$\begin{aligned}
&\leq \sum_{\ell \geq N} \mathbb{P}(Z^N \text{ d.o.} | Z_R^N = \ell; R < +\infty; Z_1^N < (1+b)N) \mathbb{P}(Z_R = \ell; R < +\infty; Z_1^N < (1+b)N) + q^{(1+b)N} \\
&\leq \sum_{\ell \geq N} \mathbb{P}(Z^N \text{ d.o.} | Z_R^N = \ell; R < +\infty) \mathbb{P}(Z_1^N < (1+b)N) + q^{(1+b)N} \\
&\leq \sum_{\ell \geq N} q^\ell e^{-cN} + q^{(1+b)N} \leq \frac{1}{1-q} \exp(-cN) q^N + q^{(1+b)N}.
\end{aligned}$$

Finally, for N large enough

$$\begin{aligned}
\mathbb{P}(Z^N \text{ dies out}) - \mathbb{P}(V_N = 0) &= \mathbb{P}(Z^N \text{ dies out}, R < \infty) \\
&\leq q^N \exp(-c'N),
\end{aligned} \tag{6}$$

which finishes the proof. \square

By construction $U_N \geq V_N$, we first prove that U_N is close to V_N , in the following sense.

Lemma 3. *The sequence $(\frac{U_N}{1+V_N})_{N \geq 1}$ converges to 1 in probability.*

Proof. By definition we have $U_N \geq V_N + 1$. For $\varepsilon > 0$,

$$\mathbb{P}\left(\frac{U_N}{1+V_N} - 1 \leq -\varepsilon\right) \leq \mathbb{P}(U_N - (V_N + 1) > \sqrt{\varepsilon}N) + \mathbb{P}\left(U_N < \frac{1}{\sqrt{\varepsilon}}N\right).$$

We study the first term using a standard technique in branching processes theory:

$$\mathbb{P}((Z_k^1) \text{ dies}; Z_K > 0) = q - f_K(0),$$

where f_K is both the K th iterate of f and the generating function of Z_K . Our assumptions imply that $|q - f_K(0)| \leq \exp(-cK)$ for some $c > 0$ ([1], Th. 1, Chap. I.11).

By the Markov property, the process $(X_{k+\ell}^N)_{\ell \geq 0}$ conditioned to $X_k^N = N$ has the same law as $(X_\ell^N)_{\ell \geq 0}$, it follows that

$$\mathbb{P}(U_N - (V_N + 1) > K) = \mathbb{P}(U_N > K + 1 | V_N = 0).$$

Recall (6):

$$\mathbb{P}(V_N = 0) \geq \mathbb{P}(Z^N \text{ dies out}) - 2q^N \exp(-c'N) \geq q^N/2$$

for N large enough. We then write

$$\begin{aligned}
\mathbb{P}(U_N > K + 1 | V_N = 0) &= \mathbb{P}(Z_{K+1}^N > 0 | Z_k^N < N \text{ for each } k \geq 1) \\
&\leq \mathbb{P}(Z_{K+1}^N > 0 | (Z_k^N) \text{ dies out}) \times \frac{\mathbb{P}((Z_k^N) \text{ dies out})}{\mathbb{P}(Z_k^N < N \text{ for each } k \geq 1)} \\
&\leq N \mathbb{P}(Z_{K+1}^1 > 0 | (Z_k^N) \text{ dies out}) \times \frac{q^N}{q^N/2} \\
&\leq 2N \frac{\exp(-cK)}{q},
\end{aligned} \tag{7}$$

where we finally used ([1], Sec. I.11). We obtain

$$\mathbb{P}(U_N - V_N - 1 > \sqrt{\varepsilon}N) \leq CN \exp(-c\sqrt{\varepsilon}N)$$

for some C and $c > 0$ and for any $N \geq 1$ and $\varepsilon > 0$. The second term is handled thanks to (4) as follows. For $N > 1/\varepsilon$,

$$\begin{aligned} \mathbb{P}\left(U_N < \left\lfloor \frac{1}{\sqrt{\varepsilon}} N \right\rfloor\right) &\leq \mathbb{P}\left(V_N < \left\lfloor \frac{1}{\sqrt{\varepsilon}} N \right\rfloor\right) \\ &\leq \mathbb{P}\left(\exists v \leq \left\lfloor \frac{1}{\sqrt{\varepsilon}} N \right\rfloor, V_N = v\right) \\ &\leq \mathbb{P}\left(\exists v \leq \left\lfloor \frac{1}{\sqrt{\varepsilon}} N \right\rfloor, X_v^N = N \text{ and } X_{v+1}^N < N\right) \\ &\leq \frac{1}{\sqrt{\varepsilon}} N \exp(-cN) \end{aligned}$$

for a certain $c > 0$. Letting N go to infinity finishes the proof. \square

We introduce another process and an associated time whose properly renormalized law will converge and which is equivalent to V_N . Let A_k be the k th passage in N of the censored process: $A_0 = 0$ and for $k \geq 0$, $A_{k+1} = \inf\{\ell > A_k, X_\ell^N = N\}$, with the usual convention that $\inf(\emptyset) = +\infty$. Let T be the survival time of A :

$$T = \sup\{k \geq 0, A_k < \infty\}.$$

It is clear by construction that T has the law of $\mathcal{G}(q_N) - 1$, where $\mathcal{G}(r)$ is a geometric of parameter r . By the Markov property of our branching process, the variables $A_{k+1} - A_k$ are independent and identically distributed. With high probability, $A_{k+1} = A_k + 1$, we can be more precise:

Lemma 4. *There exists $c > 0$ such that for all N*

$$\mathbb{P}(A_{k+1} > A_k + 1) \leq \exp(-cN).$$

There exists $c, C > 0$ such that for all $K \geq 1$, for all N large enough,

$$\mathbb{P}(A_k + K < A_{k+1} < \infty) \leq C \exp\left(-c \frac{K}{\log N} - cN\right).$$

Proof. The first assertion is just (3). The second one is a consequence of the following inequality: for any $0 < i < N$

$$\mathbb{P}(0 < Z_k^i < N \text{ for all } k \leq K) \leq C \exp\left(-c \frac{K}{\log N}\right). \quad (8)$$

To prove so, take $1 < \mu_\star < \mu \leq +\infty$. The sequence Z_k/μ_\star^k tends to infinity with positive probability (see [1], Th. 3, Chap. I.10). Hence we have positive constants δ, β such that, for any integers k, n ,

$$\mathbb{P}(Z_k^n > \delta \mu_\star^k) \geq \mathbb{P}(Z_k^1 > \delta \mu_\star^k) \geq \beta. \quad (9)$$

Now, set $m = \lceil \log(N/\delta) / \log(\mu_\star) \rceil$. We assume that N is large enough, such that $m \geq 1$. We first assume that $K > m$.

$$\begin{aligned} \{0 < Z_k < N \text{ for all } k \leq K\} &\subset \{Z_m < N\} \cap \{Z_m > 0 \text{ and } Z_{2m} < N\} \\ &\quad \cap \{Z_{2m} > 0 \text{ and } Z_{3m} < N\} \cap \cdots \cap \{Z_{\lfloor K/m \rfloor m} > 0 \text{ and } Z_{\lfloor K/m \rfloor m} < N\}. \end{aligned}$$

Thanks to (9), the probability of the right-hand side is smaller than $(1 - \beta)^{\lfloor K/m \rfloor}$. If $K \leq m$ then $K/\log N \leq 2/\log \mu_\star$ for N large enough. By choosing C large enough, the right-hand side in (8) is greater than one, and the claimed inequality also holds.

To finish the proof of the lemma, it suffices to remark that for $K \geq 1$,

$$\begin{aligned} \mathbb{P}(A_k + K < A_{k+1} < \infty) &\leq \mathbb{P}(A_k + 1 < A_{k+1} < \infty) \mathbb{P}(0 < Z_\ell^N < N \forall \ell \leq K) \\ &\leq C \exp(-cN) \exp\left(-c' \frac{K}{\log N}\right). \end{aligned} \quad \square$$

Lemma 5. *The sequence $(\frac{1+V_N}{1+T})_{N \geq 1}$ converges to one in probability.*

Proof. By construction we have $T \leq V_N$. In order to bound the difference, we first notice that, T being a $\mathcal{G}(q_N) - 1$, we have for large enough N the bound

$$\mathbb{P}(T < 1/\sqrt{q_N}) \leq (1/\sqrt{q_N} + 1) \mathbb{P}(T = 0) \leq (1/\sqrt{q_N} + 1) q_N \leq 2\sqrt{q_N} \leq 3q^{N/2}, \quad (10)$$

where we finally used Lemma 2. We now write

$$V_N \stackrel{(\text{law})}{=} \sum_{k=0}^T A_k \stackrel{(\text{law})}{=} T + \sum_{k=0}^T \tilde{A}_k,$$

where (\tilde{A}_k) is a sequence of i.i.d. random variables with the same law as $A_1 - 1$ conditioned by $A_1 < \infty$, and the sequence is independent from T .

Now take $\varepsilon > 0$. By Lemma 4, for N large enough we have $\mathbb{E}(\tilde{A}_1) \leq \varepsilon/2$.

$$\begin{aligned} \mathbb{P}(V_N + 1 - (T + 1) \geq \varepsilon(T + 1)) &= \mathbb{P}\left(\sum_{k=0}^T \tilde{A}_k \geq \varepsilon(T + 1)\right) \\ &\leq \mathbb{P}\left(\sum_{k=0}^T \tilde{A}_k - \mathbb{E}(\tilde{A}_k) \geq (T + 1)\varepsilon/2\right) \\ &\leq \mathbb{P}\left(\sum_{k=0}^T \tilde{A}_k - \mathbb{E}(\tilde{A}_k) \geq \varepsilon(T + 1)/2; T \geq 1/\sqrt{q_N}\right) + \mathbb{P}(T < 1/\sqrt{q_N}) \\ &\leq \sum_{t \geq 1/\sqrt{q_N}} \mathbb{P}\left(\sum_{k=0}^t \tilde{A}_k - \mathbb{E}(\tilde{A}_k) \geq \varepsilon(t + 1)/2\right) + \mathbb{P}(T < 1/\sqrt{q_N}) \\ &\leq \sum_{t \geq 1/2\sqrt{q_N}} \mathbb{P}\left(\sum_{k=0}^t \tilde{A}_k - \mathbb{E}(\tilde{A}_k) \geq \varepsilon(t + 1)/2\right) + \mathbb{P}(T < 1/\sqrt{q_N}), \end{aligned}$$

where we finally used Lemma 2. The second term on the right-hand side goes to zero thanks to (10). Let us now handle the sum. Using Lemma 4 we have

$$\mathbb{P}(\tilde{A}_k \geq K) = \mathbb{P}(K \leq A_1 < \infty) \mathbb{P}(A_1 < \infty)^{-1} \leq 2 \exp(-cK/\log N)$$

for a certain $c > 0$. From such a tail, we deduce (see [4], Sec. III.4) that there exists $c > 0$ such that for all t, ε and N ,

$$\mathbb{P}\left(\sum_{k=0}^t \tilde{A}_k - \mathbb{E}(\tilde{A}_k) \geq \varepsilon t/2\right) \leq \exp\left(-c\varepsilon \frac{t}{\log N}\right),$$

and the sum goes to zero. □

End of proof of Theorem 1. We now write

$$U_N q^N = \frac{U_N}{V_N + 1} \frac{q^N}{q_N} q_N (T + 1) \frac{V_N + 1}{T + 1},$$

where, when N goes to infinity,

$$U_N / (V_N + 1) \rightarrow 1 \quad (\text{in prob.}) \text{ by Lemma 3,}$$

$$q^N / q_N \rightarrow 1 \quad \text{by Lemma 2,}$$

$$q_N / (T + 1) \rightarrow \mathcal{E}(1) \quad (\text{in law}) \text{ since } T \text{ is geometric with parameter } q_N,$$

$$(V_N + 1) / (T + 1) \rightarrow 1 \quad (\text{in prob.}) \text{ by Lemma 5.}$$

This proves the convergence in distribution. For the convergence in mean, recall that one may write

$$\mathbb{E}[V_N] = \mathbb{E}\left[T + \sum_{k=0}^T \tilde{A}_k\right] = \mathbb{E}[T](1 + \mathbb{E}[\tilde{A}_1]) = q^N (1 + o(1)),$$

thanks to Lemmas 2 and 4. Since $\mathbb{E}[U_N] = \mathbb{E}[V_N] + \mathbb{E}[U_N - V_N]$, we finally obtain with (7)

$$\mathbb{E}[U_N] \sim \mathbb{E}[V_N] \sim (1/q)^N, \tag{11}$$

which will be useful in the next section. \square

4. Application: Branching-selection particle system

We now describe more precisely the connection with the branching-selection process defined in the Introduction, in order to exploit the results of the previous section. Recall that $Y_k^N(\ell)$ is the number of particles at time k at the position ℓ and that $v_N := \lim_{k \rightarrow \infty} (\max Y_k^N) / k$ (a.s.).

The connection with the censored Galton–Watson process is the following lemma, whose proof is immediate by construction of the process.

Lemma 6. *In the branching-selection process with a pair $(\mathcal{X}, \mathcal{X}')$ of laws on \mathbb{N} , the number of particles that are at the rightmost possible position has the law of a censored Galton–Watson process, when the offspring distribution is \mathcal{X} :*

$$(Y_k^N(k))_{k \geq 0} \stackrel{\text{law}}{=} (X_k^N)_{k \geq 0}.$$

In particular, if we define V^1 as the last time at which the N particles are at the rightmost possible location:

$$V^1 = \max\{k; Y_k^N(k) = N\},$$

then $V^1 \stackrel{\text{law}}{=} V_N$, where V_N , defined in the previous section, is the last time at which the censored Galton–Watson hits N .

We now are ready to estimate v_N .

Lemma 7. *For all integer $N \geq 1$,*

$$v_N \geq 1 - \frac{1}{E(V_N) + 1},$$

where the progeny distribution defining V_N is \mathcal{X} .

Proof. The proof is inspired by that of Proposition 4 in [2]. The main idea is to design a *dominated* process \tilde{Y}_k moving slower than the *true* process Y_k .

Let us skip the exponent N in order to lighten the notations, and consider an i.i.d. family $(\mathcal{X}_{\ell,k,i}, \mathcal{X}'_{\ell,k,i})_{\ell \geq 0, k \geq 0, i \in [1, N]}$ of integer variables with the same law as $(\mathcal{X}, \mathcal{X}')$.

We sample the particle system (Y_k) by means of these variables: the population Y_k at time k being given, we let

$$T_{k+1} = \sum_{\ell=0}^{\infty} \sum_{i=1}^{Y_k(\ell)} (\mathcal{X}_{\ell,k,i} \delta_{\ell+1} + \mathcal{X}'_{\ell,k,i} \delta_{\ell})$$

be the locations of the particles after branching. The population Y_{k+1} is then obtained by keeping only the N rightmost particles among T_{k+1} .

We modify Y in the following way. At time $V^1 + 1$, let continue the process as if all the particles were at position V^1 at time $V^1 + 1$. More precisely, set $(Y_k^{(1)}) = (Y_k)$ and introduce a process $(Y_k^{(2)})_{k \geq V^1}$ starting from

$$Y_{V^1+1}^{(2)} = N \delta_{V^1}.$$

Then, for $k \geq V^1 + 1$, $Y_{k+1}^{(2)}$ is sampled from $Y_k^{(2)}$ exactly as Y_{k+1} is sampled from Y_k^N (with the same $(\mathcal{X}_{\ell,k,i}, \mathcal{X}'_{\ell,k,i})$'s). The main point is that, at time $V^1 + 1$, the N particles of $Y_{V^1+1}^{(1)}$ are at positions V^1 or $V^1 + 1$ while the N particles of $Y_{V^1+1}^{(2)}$ are at position V^1 . Hence, the point measure $Y_{V^1+1}^{(1)}$ dominates $Y_{V^1+1}^{(2)}$, and this domination will continue throughout the process, since the same $(\mathcal{X}, \mathcal{X}')$'s are used to generate $Y^{(1)}$ and $Y^{(2)}$.

Now, let V^2 be such that $V^1 + 1 + V^2$ is the last time k at which the N particles are at the rightmost possible location (which is position $k - 1$) for $Y^{(2)}$:

$$V^2 = \max\{k; Y_k^{(2)}(k - 1) = N\} - (V^1 + 1).$$

The random time V^2 has the same law as V^1 and, as a function of $\{\mathcal{X}_{\ell,k,i}, \mathcal{X}'_{\ell,k,i}; \ell > V^1\}$, is independent of V^1 . We define recursively similar processes $Y^{(3)}, Y^{(4)}, \dots$ and random variables V^3, V^4, \dots . Setting also

$$\Gamma^0 = 0, \quad \Gamma^{i+1} = \Gamma^i + V^i + 1,$$

the Γ^i 's are renewals.

Let us introduce a last process $(\tilde{Y}_k)_{k \geq 0}$ as follows: if $k \in [\Gamma^{i-1}, \Gamma^i)$, then

$$\tilde{Y}_k = Y_k^{(i)}.$$

One can see in Fig. 1 the first steps of a realization of the process (Y_k) , and in Fig. 2 the corresponding modified process (\tilde{Y}_k) . By construction, Y_k dominates \tilde{Y}_k for each k , and in particular $\max Y_k \geq \max \tilde{Y}_k$. Let us also note that at each renewal Γ^i the quantity $\max \tilde{Y}_k$ is decreased by one. Set I_k be the renewal process associated to the renewals Γ^i , that is I_k is the only integer i such that $\Gamma^i \leq k < \Gamma^{i+1}$. On the one hand we have

$$\Gamma^{i+1} - \Gamma^i \stackrel{\text{law}}{=} V^i + 1,$$

so by applying the renewal theorem (see for instance [3], Chap. 3.4) to the renewals $\Gamma^1, \Gamma^2, \dots$ we get

$$\lim_{k \rightarrow \infty} \frac{1}{k} \mathbb{E}(I_k) \rightarrow \frac{1}{\mathbb{E}[V^1] + 1}.$$

On the second hand we have

$$\max \tilde{Y}_k = k - I_k.$$




Fig. 1. A realization of the *true* process Y with $N = 3$ particles, for the particular case of the Bernoulli branching-selection process. At each step are represented the 6 particles (after replication) and the 3 leftmost particles are struck. Here, we see that $V_1 = 3$, since from time $k = 5$ there can be no more particles at the rightmost possible position, and because at time 4 only one particle is on the rightmost possible position.




Fig. 2. A realization of the associated *dominated* process: at time $V_N^1 + 1 = 4$, we let the process restart with *all* particles at position 3.

Hence we get

$$\frac{1}{k} \mathbb{E}[\max Y_k] \geq \frac{1}{k} \mathbb{E}[\max \tilde{Y}_k] \xrightarrow{k \rightarrow \infty} 1 - \frac{1}{\mathbb{E}[V^1] + 1}.$$

To conclude, recall that thanks to the connection with the censored Galton–Watson process $\mathbb{E}[V^1] = \mathbb{E}[V_N]$. \square

Lemma 8. For all integer $N \geq 1$,

$$v_N \leq 1 - \frac{1}{E(U_N)}.$$

Proof. The proof is an adaptation of the previous one, replacing V^1 by U^1 , which is the first time at which there is no more particle at the rightmost possible position:

$$U^1 = \min\{k; Y_k^N(k) = 0\}.$$

(On the example drawn in Fig. 1, one has $U^1 = 5$.) Then, U^1 has the same law as the U_N defined in Section 2. In a similar manner to the previous proof, we let the process restart at time U^1 as if all the particles were at position $U^1 - 1$. The end of the proof is similar. \square

We now combine these two estimates of v_N with the results of the previous section in order to prove Theorem 2:

$$\frac{1}{E(U_N)} \leq 1 - v_N \leq \frac{1}{E(V_N) + 1}$$

and both sides, thanks to (11), are equivalent to q^N .

Acknowledgments

We warmly thank J.-B. Gouéré for many helpful comments on this work. We are also very thankful to the two referees and the associate editor for recommending numerous improvements.

References

- [1] K. B. Athreya and P. E. Ney. *Branching Processes*. Dover Publications Inc., Mineola, NY, 2004. [MR2047480](#)
- [2] J. Bérard and J.-B. Gouéré. Brunet–Derrida behavior of branching-selection particle systems on the line. *Comm. Math. Phys.* **298** (2010) 323–342. [MR2669438](#)
- [3] R. Durrett. *Probability: Theory and Examples*, 2nd edition. Duxbury Press, Belmont, CA, 1996. [MR1609153](#)
- [4] V. V. Petrov. *Sums of Independent Random Variables*. Springer, New York, 1975. [MR0388499](#)
- [5] V. A. Topchiĭ. The time to extinction for Galton–Watson processes censored at a high level. *Teor. Veroyatn. Primen.* **54** (2009) 271–287. [MR2761556](#)
- [6] A. N. Zubkov. A degeneracy condition for a bounded branching process. *Mat. Zametki* **8** (1970) 9–18. [MR0267649](#)