

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

RADIAL MAXIMAL FUNCTION CHARACTERIZATIONS FOR HARDY SPACES ON RD-SPACES

Loukas Grafakos & Liguang Liu & Dachun Yang (corresponding author)

**Tome 137
Fascicule 2**

2009

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique
pages 225-251

RADIAL MAXIMAL FUNCTION CHARACTERIZATIONS FOR HARDY SPACES ON RD-SPACES

BY LOUKAS GRAFAKOS, LIGUANG LIU
& DACHUN YANG (CORRESPONDING AUTHOR)

ABSTRACT. — An RD-space \mathcal{X} is a space of homogeneous type in the sense of Coifman and Weiss with the additional property that a reverse doubling property holds. The authors prove that for a space of homogeneous type \mathcal{X} having “dimension” n , there exists a $p_0 \in (n/(n+1), 1)$ such that for certain classes of distributions, the $L^p(\mathcal{X})$ quasi-norms of their radial maximal functions and grand maximal functions are equivalent when $p \in (p_0, \infty]$. This result yields a radial maximal function characterization for Hardy spaces on \mathcal{X} .

Texte reçu le 17 septembre 2007, révisé le 29 juillet 2008, accepté le 23 septembre 2008

LOUKAS GRAFAKOS, Department of Mathematics, University of Missouri, Columbia, MO 65211, USA • *E-mail* : loukas@math.missouri.edu

LIGUANG LIU, School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems, Ministry of Education, Beijing 100875, People's Republic of China • *E-mail* : liuliguang@mail.bnu.edu.cn

DACHUN YANG (CORRESPONDING AUTHOR), School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems, Ministry of Education, Beijing 100875, People's Republic of China • *E-mail* : dcyang@bnu.edu.cn

2000 Mathematics Subject Classification. — 42B25; 42B30, 47B38, 47A30.

Key words and phrases. — Space of homogeneous type, approximation of the identity, space of test function, grand maximal function, radial maximal function, Hardy space.

The first author was supported by grant DMS 0400387 of the National Science Foundation of the USA and the University of Missouri Research Council. The third (corresponding) author was supported by the National Science Foundation for Distinguished Young Scholars (Grant No. 10425106) of China.

RÉSUMÉ (*Caractérisations de fonctions radiales maximales pour les espaces de Hardy sur les RD-espaces*)

Un RD-espace X est un espace de type homogène au sens de Coifman et Weiss, possédant en outre une propriété de doublement inverse. Les auteurs prouvent que pour un espace de type homogène X de « dimension » n , il existe un $p_0 \in (n/(n+1), 1)$ tel que les quasi-normes $L^p(X)$ des fonctions radiales maximales et grand-maximales d'une certaine classe de distributions soient équivalentes lorsque $p \in (p_0, \infty]$. Ce résultat fournit une caractérisation des espaces de Hardy sur X en termes de fonctions radiales maximales.

1. Introduction

The theory of Hardy spaces on Euclidean spaces plays an important role in harmonic analysis and partial differential equations and has been systematically studied and developed; see, for example, [7, 19, 8, 21]. It is well known that spaces of homogeneous type, in the sense of Coifman and Weiss [3], are a natural setting of the Calderón-Zygmund theory of singular integrals; see also [4].

A space of homogeneous type is a set \mathcal{X} equipped with a metric d and a regular Borel measure μ having the doubling property. Coifman and Weiss [4] introduced the atomic Hardy space $H_{\text{at}}^p(\mathcal{X})$ for $p \in (0, 1]$ and further established a molecular characterization for $H_{\text{at}}^1(\mathcal{X})$. Moreover, under the assumption that the measure of any ball in \mathcal{X} is equivalent to its radius (i. e., \mathcal{X} is an Ahlfors 1-regular metric measure space), when $p \in (1/2, 1]$, Macías and Segovia [14] used distributions acting on certain spaces of Lipschitz functions to obtain a grand maximal function characterization for $H_{\text{at}}^p(\mathcal{X})$; Han [10] further established a Lusin-area characterization for $H_{\text{at}}^p(\mathcal{X})$, and Duong and Yan [6] characterized these atomic Hardy spaces in terms of Lusin-area functions associated with certain Poisson semigroups. Also in this setting, a deep result of Uchiyama [22] states that if $p \in (p_0, 1]$ for some p_0 near 1, for functions in $L^1(\mathcal{X})$, the $L^p(\mathcal{X})$ quasi-norms of the grand maximal functions as in [14] are equivalent to the $L^p(\mathcal{X})$ quasi-norms of the radial maximal functions defined via some kernels in [4].

An important special class of spaces of homogeneous type is called RD-spaces, which is introduced in [12] (see also [11, 15]) and modeled on Euclidean spaces with A_∞ -weights (Muckenhoupt's class), Ahlfors n -regular metric measure spaces (see, for example, [13]), Lie groups of polynomial growth (see, for example, [23, 24, 1]) and Carnot-Carathéodory spaces with doubling measure (see, for example, [16, 17, 5, 20, 18]). A Littlewood-Paley theory of Hardy spaces on RD-spaces was established in [11], and these Hardy spaces are shown

to coincide with some of Triebel-Lizorkin spaces in [12]. The grand, nontangential and dyadic maximal function characterizations of these Hardy spaces have recently been established in [9].

The main purpose of this paper is twofold: first to generalize the results of Uchiyama [22] to the setting of RD-spaces and second to replace the space $L^1(\mathcal{X})$ used by Uchiyama in [22] by certain spaces of distributions developed in [11, 12]. In other words, we build on the work of Uchiyama [22] to establish a radial maximal function characterization for the Hardy spaces in [11].

To state our main results, we need to recall some definitions and notation. We begin with the classical notions of spaces of homogeneous type ([3], [4]) and RD-spaces ([12]).

DEFINITION 1.1. — *Let (\mathcal{X}, d) be a metric space with a regular Borel measure μ such that all balls defined by d have finite and positive measures. For any $x \in \mathcal{X}$ and $r > 0$, set $B(x, r) \equiv \{y \in \mathcal{X} : d(x, y) < r\}$.*

(i) *The triple (\mathcal{X}, d, μ) is called a space of homogeneous type if there exists a constant $C_0 \geq 1$ such that for all $x \in \mathcal{X}$ and $r > 0$,*

$$(1.1) \quad \mu(B(x, 2r)) \leq C_0 \mu(B(x, r)) \text{ (doubling property).}$$

(ii) *Let $0 < \kappa \leq n$. The triple (\mathcal{X}, d, μ) is called a (κ, n) -space if there exist constants $0 < C_1 \leq 1$ and $C_2 \geq 1$ such that for all $x \in \mathcal{X}$, $0 < r < \text{diam}(\mathcal{X})/2$ and $1 \leq \lambda < \text{diam}(\mathcal{X})/(2r)$,*

$$(1.2) \quad C_1 \lambda^\kappa \mu(B(x, r)) \leq \mu(B(x, \lambda r)) \leq C_2 \lambda^n \mu(B(x, r)),$$

where $\text{diam}(\mathcal{X}) \equiv \sup_{x, y \in \mathcal{X}} d(x, y)$.

A space of homogeneous type is called an RD-space, if it is a (κ, n) -space for some $0 < \kappa \leq n$, i. e., some “reverse” doubling condition holds.

REMARK 1.2. — (i) *A regular Borel measure μ has the property that open sets are measurable and every set is contained in a Borel set with the same measure; see [13].*

(ii) *The number n in some sense measures the “dimension” of \mathcal{X} . Obviously any (κ, n) space is a space of homogeneous type with $C_0 = C_2 2^n$. Conversely, any space of homogeneous type satisfies the second inequality of (1.2) with $C_2 = C_0$ and $n = \log_2 C_0$.*

(iii) *If μ is doubling, then μ satisfies (1.2) if and only if there exist constants $a_0 > 1$ and $\tilde{C}_0 > 1$ such that for all $x \in \mathcal{X}$ and $0 < r < \text{diam}(\mathcal{X})/a_0$,*

$$\mu(B(x, a_0 r)) \geq \tilde{C}_0 \mu(B(x, r)) \text{ (reverse doubling property)}$$

(If $a_0 = 2$, this is the classical reverse doubling condition), and equivalently, for all $x \in \mathcal{X}$ and $0 < r < \text{diam}(\mathcal{X})/a_0$,

$$B(x, a_0 r) \setminus B(x, r) \neq \emptyset;$$

see [12]. From this, it follows that if \mathcal{X} is an RD-space, then $\mu(\{x\}) = 0$ for all $x \in \mathcal{X}$.

Throughout the whole paper, we always assume that \mathcal{X} is an RD-space and $\mu(\mathcal{X}) = \infty$. For any $x, y \in \mathcal{X}$ and $\delta > 0$, set $V_\delta(x) \equiv \mu(B(x, \delta))$ and $V(x, y) \equiv \mu(B(x, d(x, y)))$. It follows from (1.1) that $V(x, y) \sim V(y, x)$. The following notion of approximations of the identity on RD-spaces is a variant of that in [12, Definitions 2.1, 2.2]; see also [11].

DEFINITION 1.3. — Let $\epsilon_1 \in (0, 1]$, $\epsilon_2 > 0$ and $\epsilon_3 > 0$. A sequence $\{S_k\}_{k \in \mathbb{Z}}$ of bounded linear integral operators on $L^2(\mathcal{X})$ is said to be a special approximation of the identity of order $(\epsilon_1, \epsilon_2, \epsilon_3)$ (for short, $(\epsilon_1, \epsilon_2, \epsilon_3)$ -SAOTI), if there exists a constant $C_3 > \sqrt{2}$ such that for all $k \in \mathbb{Z}$ and all x, x', y and $y' \in \mathcal{X}$, $S_k(x, y)$, the integral kernel of S_k is a function from $\mathcal{X} \times \mathcal{X}$ into $[0, \infty)$ satisfying

- (i) $S_k(x, y) \leq C_3 \frac{1}{V_{2^{-k}}(x)+V_{2^{-k}}(y)+V(x,y)} \frac{2^{-k\epsilon_2}}{(2^{-k}+d(x,y))^{\epsilon_2}}$;
- (ii) $|S_k(x, y) - S_k(x', y)| \leq C_3 \frac{d(x,x')^{\epsilon_1}}{(2^{-k}+d(x,y))^{\epsilon_1}} \frac{1}{V_{2^{-k}}(x)+V_{2^{-k}}(y)+V(x,y)} \frac{2^{-k\epsilon_2}}{(2^{-k}+d(x,y))^{\epsilon_2}}$ for $d(x, x') \leq (2^{-k} + d(x, y))/2$;
- (iii) Property (ii) holds with x and y interchanged;
- (iv) $|[S_k(x, y) - S_k(x, y')] - [S_k(x', y) - S_k(x', y')]| \leq C_3 \frac{d(x,x')^{\epsilon_1}}{(2^{-k}+d(x,y))^{\epsilon_1}} \frac{d(y,y')^{\epsilon_1}}{(2^{-k}+d(x,y))^{\epsilon_1}} \times \frac{1}{V_{2^{-k}}(x)+V_{2^{-k}}(y)+V(x,y)} \frac{2^{-k\epsilon_3}}{(2^{-k}+d(x,y))^{\epsilon_3}}$ for $d(x, x') \leq (2^{-k} + d(x, y))/3$ and $d(y, y') \leq (2^{-k} + d(x, y))/3$;
- (v) $C_3 V_{2^{-k}}(x) S_k(x, x) > 1$ for all $x \in \mathcal{X}$ and $k \in \mathbb{Z}$;
- (vi) $\int_{\mathcal{X}} S_k(x, y) d\mu(y) = 1 = \int_{\mathcal{X}} S_k(x, y) d\mu(x)$.

We remark that (i) and (v) of Definition 1.3 imply that $C_3 > \sqrt{2}$. The existence of $(\epsilon_1, \epsilon_2, \epsilon_3)$ -SAOTI's was proved in [12, Theorem 2.1].

The following spaces of test functions play a key role in the theory of function spaces on RD-spaces; see [12, 11].

DEFINITION 1.4. — Let $x_1 \in \mathcal{X}$, $r \in (0, \infty)$, $\beta \in (0, 1]$ and $\gamma \in (0, \infty)$. A function φ on \mathcal{X} is said to be a test function of type (x_1, r, β, γ) if there exists a nonnegative constant C such that

- (i) $|\varphi(x)| \leq C \frac{1}{V_r(x_1)+V(x_1,x)} \left(\frac{r}{r+d(x_1,x)}\right)^\gamma$ for all $x \in \mathcal{X}$;
- (ii) $|\varphi(x) - \varphi(y)| \leq C \left(\frac{d(x,y)}{r+d(x_1,x)}\right)^\beta \frac{1}{V_r(x_1)+V(x_1,x)} \left(\frac{r}{r+d(x_1,x)}\right)^\gamma$ for all $x, y \in \mathcal{X}$ satisfying $d(x, y) \leq (r + d(x_1, x))/2$.

We denote by $\mathcal{G}(x_1, r, \beta, \gamma)$ the set of all test functions of type (x_1, r, β, γ) . If $\varphi \in \mathcal{G}(x_1, r, \beta, \gamma)$, we define its norm by $\|\varphi\|_{\mathcal{G}(x_1, r, \beta, \gamma)} \equiv \inf\{C : (i) \text{ and } (ii) \text{ hold}\}$. The space $\mathcal{G}(x_1, r, \beta, \gamma)$ is called the space of test functions.

Throughout the whole paper, we fix $x_1 \in \mathcal{X}$. Let $\mathcal{G}(\beta, \gamma) \equiv \mathcal{G}(x_1, 1, \beta, \gamma)$. It is easy to see that for any $x_2 \in \mathcal{X}$ and $r > 0$, we have $\mathcal{G}(x_2, r, \beta, \gamma) = \mathcal{G}(\beta, \gamma)$ with equivalent norms (but with constants depending on x_1, x_2 and r). Moreover, $\mathcal{G}(\beta, \gamma)$ is a Banach space.

For any given $\epsilon \in (0, 1]$, let $\mathcal{G}_0^\epsilon(\beta, \gamma)$ be the completion of the space $\mathcal{G}(\epsilon, \epsilon)$ in $\mathcal{G}(\beta, \gamma)$ when $\beta, \gamma \in (0, \epsilon]$. Obviously $\mathcal{G}_0^\epsilon(\epsilon, \epsilon) = \mathcal{G}(\epsilon, \epsilon)$. Moreover, $\varphi \in \mathcal{G}_0^\epsilon(\beta, \gamma)$ if and only if $\varphi \in \mathcal{G}(\beta, \gamma)$ and there exists $\{\phi_i\}_{i \in \mathbb{N}} \subset \mathcal{G}(\epsilon, \epsilon)$ such that $\|\varphi - \phi_i\|_{\mathcal{G}(\beta, \gamma)} \rightarrow 0$ as $i \rightarrow \infty$. If $\varphi \in \mathcal{G}_0^\epsilon(\beta, \gamma)$, define $\|\varphi\|_{\mathcal{G}_0^\epsilon(\beta, \gamma)} \equiv \|\varphi\|_{\mathcal{G}(\beta, \gamma)}$. Obviously $\mathcal{G}_0^\epsilon(\beta, \gamma)$ is a Banach space and $\|\varphi\|_{\mathcal{G}_0^\epsilon(\beta, \gamma)} = \lim_{i \rightarrow \infty} \|\phi_i\|_{\mathcal{G}(\beta, \gamma)}$ for the above chosen $\{\phi_i\}_{i \in \mathbb{N}}$. It is known that $\mathcal{G}_0^\epsilon(\beta, \gamma)$ is dense in $L^p(\mathcal{X})$ for $p \in [1, \infty)$; see [12, Corollary 2.1]. Let $(\mathcal{G}_0^\epsilon(\beta, \gamma))'$ be the set of all bounded linear functionals f from $\mathcal{G}_0^\epsilon(\beta, \gamma)$ to \mathbb{C} . Denote by $\langle f, \varphi \rangle$ the natural pairing of elements $f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))'$ and $\varphi \in \mathcal{G}_0^\epsilon(\beta, \gamma)$.

Let $\epsilon \in (0, 1)$, $\beta, \gamma \in (0, \epsilon)$ and $p \in (0, \infty]$. If $f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))'$, then for all $x \in \mathcal{X}$, we define the grand maximal function of f to be

$$f^*(x) \equiv \sup \{ |\langle f, \varphi \rangle| : \varphi \in \mathcal{G}(\epsilon, \epsilon), \|\varphi\|_{\mathcal{G}(x, r, \epsilon, \epsilon)} \leq 1 \text{ for some } r > 0 \}.$$

Define the corresponding Hardy space by

$$H^{*,p}(\mathcal{X}) \equiv \{ f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))' : \|f^*\|_{L^p(\mathcal{X})} < \infty \}.$$

For any $f \in H^{*,p}(\mathcal{X})$, set $\|f\|_{H^{*,p}(\mathcal{X})} \equiv \|f^*\|_{L^p(\mathcal{X})}$. Let $\{Q_\alpha^k : k \in \mathbb{Z}, \alpha \in I_k\}$ be the Christ dyadic cube collection of \mathcal{X} , where I_k is some index set; see [2]. For any $f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))'$, we define the dyadic maximal function $\mathcal{M}_d(f)$ of f by setting, for all $x \in \mathcal{X}$,

$$\mathcal{M}_d(f)(x) \equiv \sup_{k \in \mathbb{Z}, \alpha \in I_k} \left\{ \frac{1}{\mu(Q_\alpha^k)} \int_{Q_\alpha^k} |S_k(f)(y)| d\mu(y) \right\} \chi_{Q_\alpha^k}(x),$$

and define $H_d^p(\mathcal{X})$ to be the corresponding Hardy space; see [9, Definition 2.9]. When $p \in (1, \infty]$, it was proved in [9, Corollary 3.12] that $H^{*,p}(\mathcal{X}) = H_d^p(\mathcal{X}) = L^p(\mathcal{X})$ with equivalent norms.

DEFINITION 1.5. — Let $\epsilon_1 \in (0, 1]$, $\epsilon_2 > 0$, $\epsilon_3 > 0$, $\epsilon \in (0, \epsilon_1 \wedge \epsilon_2)$ and $\{S_k\}_{k \in \mathbb{Z}}$ be an $(\epsilon_1, \epsilon_2, \epsilon_3)$ -SAOTI. Let $p \in (0, \infty]$ and $f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))'$ with $\beta, \gamma \in (0, \epsilon)$. Define the radial maximal function of f to be $\mathcal{M}_0(f)(x) \equiv \sup_{k \in \mathbb{Z}} |S_k(f)(x)|$ for all $x \in \mathcal{X}$. The corresponding Hardy spaces are defined by

$$H_0^p(\mathcal{X}) \equiv \{ f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))' : \|\mathcal{M}_0(f)\|_{L^p(\mathcal{X})} < \infty \},$$

and moreover, we define $\|f\|_{H_0^p(\mathcal{X})} \equiv \|\mathcal{M}_0(f)\|_{L^p(\mathcal{X})}$.

The properties (i) and (ii) in Definition 1.3 imply that $\mathcal{M}_0(f)(x) \lesssim f^*(x)$ for all $x \in \mathcal{X}$. In what follows, for simplicity of presentation, for any $t > 0$, we use the notation

$$(1.3) \quad S_t(x, y) \equiv \sum_{k \in \mathbb{Z}} S_k(x, y) \chi_{(2^{-k-1}, 2^{-k}]}(t).$$

By (1.3) and Definition 1.5, it is easy to see that for all $x \in \mathcal{X}$,

$$(1.4) \quad \mathcal{M}_0(f)(x) = \sup_{t > 0} |S_t(f)(x)|.$$

Obviously, S_t satisfies (i) through (vi) in Definition 1.3 with 2^{-k} replaced by t . From (iv) and (v) in Definition 1.3, it follows easily that there exist constants $C_4 \in (0, (C_3)^{-2/\epsilon_1})$ and $C_5 > 1$ such that for all $t > 0$ and all $x, y \in \mathcal{X}$ satisfying $d(x, y) < C_4 t$,

$$(1.5) \quad C_5 V_t(x) S_t(x, y) > 1.$$

This observation is used in applications below.

Denote by \mathcal{M} the centered Hardy-Littlewood maximal operator. To be precise, for any $f \in L^1_{\text{loc}}(\mathcal{X})$ and $x \in \mathcal{X}$, set

$$\mathcal{M}(f)(x) \equiv \sup_{r > 0} \frac{1}{\mu(B(x, r))} \int_{B(x, r)} |f(y)| d\mu(y).$$

Then \mathcal{M} is weak-type $(1, 1)$ and bounded on $L^p(\mathcal{X})$ for $p \in (1, \infty]$ in [4, 3]. It is not so difficult to show that for all $x \in \mathcal{X}$, $\mathcal{M}_0(f)(x) \lesssim \mathcal{M}(f)(x)$ and $\mathcal{M}_d(f)(x) \lesssim \mathcal{M}(\mathcal{M}_0(f))(x)$ by their definitions and Lemma 2.1 (iv) below. Therefore, we have $H^p_0(\mathcal{X}) = L^p(\mathcal{X})$ with equivalent norms when $p \in (1, \infty]$.

The main result of this paper concerns the spaces $H^p_0(\mathcal{X})$ and $H^{*,p}(\mathcal{X})$, and is as follows.

THEOREM 1.6. — *Let $\epsilon_1 \in (0, 1]$, $\epsilon_2 > 0$, $\epsilon_3 > 0$ and $\epsilon \in (0, \epsilon_1 \wedge \epsilon_2)$. Let $\{S_k\}_{k \in \mathbb{Z}}$ be an $(\epsilon_1, \epsilon_2, \epsilon_3)$ -SAOTI and \mathcal{M}_0 be as in (1.4). Then there exist $\sigma \in (0, 1/2)$ and $\eta \in (0, (1 - \sigma)^{1/\epsilon} \wedge (1/2))$, both depending only on \mathcal{X} and ϵ , such that for any given $p \in (n/(n + \log_\eta(1 - \sigma)), \infty]$ and all $f \in (\mathcal{G}^\epsilon_0(\beta, \gamma))'$ with $\beta \in (0, \log_\eta(1 - \sigma))$ and $\gamma \in (0, \epsilon)$,*

$$\|f^*\|_{L^p(\mathcal{X})} \leq C \|\mathcal{M}_0(f)\|_{L^p(\mathcal{X})},$$

where C is a positive constant independent of f .

Theorem 1.6 will be a consequence of the following key proposition.

PROPOSITION 1.7. — *With the notation of Theorem 1.6, for any $\delta_0 \in (0, \log_\eta(1 - \sigma))$, there exists a positive constant C , depending on \mathcal{X} , ϵ and δ_0 ,*

such that for all $f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))'$ with $\beta \in (0, \log_\eta(1 - \sigma))$ and $\gamma \in (0, \epsilon)$, and all $\varphi \in \mathcal{G}(x_0, r_0, \epsilon, \epsilon)$ satisfying $\|\varphi\|_{\mathcal{G}(x_0, r_0, \epsilon, \epsilon)} \leq 1$ for some $x_0 \in \mathcal{X}$ and $r_0 > 0$,

$$|\langle f, \varphi \rangle| \leq C \left[\mathcal{M}([\mathcal{M}_0(f)]^{n/(n+\delta_0)})(x_0) \right]^{(n+\delta_0)/n}.$$

We remark that in Theorem 1.6 and Proposition 1.7, it is not necessary to assume that $\{S_k\}_{k \in \mathbb{Z}}$ has the property (vi) of Definition 1.3. Moreover, Theorem 1.6 follows easily from Proposition 1.7; see Section 3 below. The main ingredient in the proof of Proposition 1.7 is to expand φ as in Proposition 1.7 into a sum of certain S_t as in (1.3); see (3.11) below. When \mathcal{X} is an Ahlfors 1-regular metric measure space, for any Lipschitz function with bounded support, Uchiyama in [22] established an expansion similar to (3.11), which holds pointwise. Unlike [22], we prove that for any φ as in Proposition 1.7, (3.11) also holds in $\mathcal{G}_0^\epsilon(\beta, \gamma)$ with β and γ as in Proposition 1.7. This allows us to relax the assumption $f \in L^1(\mathcal{X})$ to $f \in (\mathcal{G}_0^\epsilon(\beta, \gamma))'$.

From the fact $\mathcal{M}_0(f) \lesssim f^*$ and Theorem 1.6, it follows that for p in a certain range of $(0, 1]$, $H_0^p(\mathcal{X})$ coincides with $H^{*,p}(\mathcal{X})$ as a subspace of certain distribution spaces $(\mathcal{G}_0^\epsilon(\beta, \gamma))'$. Recall that when $p \in (n/(n + 1), 1]$, [9, Remark 3.16] and [9, Corollary 4.19] tell us that the definition of $H^{*,p}(\mathcal{X})$ is independent of the choices of $(\mathcal{G}_0^\epsilon(\beta, \gamma))'$ with $\beta, \gamma \in (n(1/p - 1), \epsilon)$. Therefore, we deduce the following conclusion.

COROLLARY 1.8. — *Let $\epsilon_1, \epsilon_2, \epsilon_3, \sigma$ and η be as in Theorem 1.6 and $\epsilon \in (0, \epsilon_1 \wedge \epsilon_2)$. Let $p_0 \equiv n/(n + \log_\eta(1 - \sigma))$ and $p \in (p_0, 1]$. Then $H_0^p(\mathcal{X}) = H^{*,p}(\mathcal{X})$ with equivalent quasi-norms, where $H_0^p(\mathcal{X})$ and $H^{*,p}(\mathcal{X})$ are defined via $(\mathcal{G}_0^\epsilon(\beta, \gamma))'$ with some $\beta \in (n(1/p - 1), n(1/p_0 - 1))$ and $\gamma \in (n(1/p - 1), \epsilon)$. Consequently, the definition of $H_0^p(\mathcal{X})$ is independent of the choices of $(\epsilon_1, \epsilon_2, \epsilon_3)$ -SAOTI and $(\mathcal{G}_0^\epsilon(\beta, \gamma))'$ with β and γ as above.*

REMARK 1.9. — *We point out that in Theorem 1.6 and Proposition 1.7, it is not necessary to assume that \mathcal{X} satisfies the reverse doubling condition determined by the first inequality of (1.2). However, the assertion $H_0^p(\mathcal{X}) = L^p(\mathcal{X})$ when $p \in (1, \infty]$ and Corollary 1.8 do need this assumption, since, to obtain these conclusions, we need to use the Calderón reproducing formulae in [12], which depend on the reverse doubling condition.*

The organization of this paper is as follows. In Section 2, we give some technical lemmas which will be used in the proof of Proposition 1.7. Section 3 is the main part of this paper, which contains a proof of Proposition 1.7 and also of Theorem 1.6.

In this paper we use the following notation: $\mathbb{N} \equiv \{1, 2, \dots\}$, $\mathbb{Z}_+ \equiv \mathbb{N} \cup \{0\}$ and $\mathbb{R}_+ \equiv [0, \infty)$. For any $p \in [1, \infty]$, we denote by p' the conjugate index, namely, $1/p + 1/p' = 1$. We also denote by C a positive constant independent of the

main parameters involved, which may vary at different occurrences. Constants with subscripts do not change through the whole paper. We use $f \lesssim g$ and $f \gtrsim g$ to denote $f \leq Cg$ and $f \geq Cg$, respectively. If $f \lesssim g \lesssim f$, we then write $f \sim g$. For any $a, b \in \mathbb{R}$, set $a \wedge b \equiv \min\{a, b\}$ and $a \vee b \equiv \max\{a, b\}$. For any set E , we denote by $\sharp E$ the cardinality of E .

2. Some technical lemmas

In this section, we establish several technical lemmas which will be used in the proof of Proposition 1.7. The following lemma includes some basic properties on RD-spaces, which are used throughout the paper; see, for example, [12, 9, 11].

LEMMA 2.1. — *Let $\delta > 0$, $a > 0$, $r > 0$ and $\theta \in (0, 1)$. Then,*

- (i) *For all $x, y \in \mathcal{X}$ and $r > 0$, $V_r(x)+V(x, y) \sim V_r(y)+V(y, x) \sim \mu(B(y, r+d(y, x))) \sim \mu(B(x, r+d(x, y)))$.*
- (ii) *If $x, x', x_1 \in \mathcal{X}$ satisfy $d(x, x') \leq \theta(r+d(x, x_1))$, then $r+d(x, x_1) \sim r+d(x', x_1)$ and $\mu(B(x, r+d(x, x_1))) \sim \mu(B(x', r+d(x', x_1)))$.*
- (iii) *$\int_{\mathcal{X}} \frac{1}{V_r(x)+V(x, y)} \left(\frac{r}{r+d(x, y)}\right)^a d(x, y)^\eta d\mu(x) \leq Cr^\eta$ uniformly in $x \in \mathcal{X}$ and $r > 0$ if $a > \eta \geq 0$.*
- (iv) *For all $f \in L^1_{loc}(\mathcal{X})$ and $x \in \mathcal{X}$, $\int_{d(x, y) > \delta} \frac{1}{V(x, y)} \frac{\delta^a}{d(x, y)^a} |f(y)| d\mu(y) \leq CM(f)(x)$ uniformly in $\delta > 0$, $f \in L^1_{loc}(\mathcal{X})$ and $x \in \mathcal{X}$.*

When $\delta = 0$, the following lemma provides a property of Carleson measures on RD-spaces; see [12, Proposition 5.14].

LEMMA 2.2. — *Let $p \in (1, \infty]$ and $\delta \geq 0$. Let ν be a non-negative measure on $\mathcal{X} \times \mathbb{R}_+$ such that for all $x \in \mathcal{X}$ and $r > 0$,*

$$(2.1) \quad \nu(B(x, r) \times (0, r)) \leq [\mu(B(x, r))]^{1+\delta}.$$

Then there exists a positive constant C such that for all $f \in L^p(\mathcal{X})$,

$$\left\{ \int_{\mathcal{X} \times \mathbb{R}_+} |F(r, y, f)|^{p(1+\delta)} d\nu(y, r) \right\}^{1/(p(1+\delta))} \leq C \|f\|_{L^p(\mathcal{X})},$$

where and in what follows, $F(r, x, f) \equiv S_r(f)(x)$ for all $r > 0$ and $x \in \mathcal{X}$.

Proof. — Fix $\lambda > 0$ and let $W_\lambda \equiv \{(x, r) \in \mathcal{X} \times \mathbb{R}_+ : |F(r, x, f)| > \lambda\}$. For any $\ell \in \mathbb{Z}$, set

$$W_{\ell, \lambda} \equiv \left\{ x \in \mathcal{X} : \sup_{2^{\ell-1} < r \leq 2^\ell} |F(r, x, f)| > \lambda \right\}.$$

For each $N \in \mathbb{N}$, let $E_N \equiv \{x \in \mathcal{X} : \sup_{r > 2^N} |F(r, x, f)| > \lambda\}$. It is easy to deduce that

$$(2.2) \quad \lim_{N \rightarrow \infty} E_N = \emptyset.$$

To prove (2.2), notice that $\lim_{N \rightarrow \infty} E_N = \bigcap_{N \in \mathbb{N}} E_N$ since $E_{N+1} \subset E_N$ for any $N \in \mathbb{N}$. Suppose that (2.2) fails, that is, there exists an $x \in \bigcap_{N \in \mathbb{N}} E_N$. Thus for any $N \in \mathbb{N}$, there exists $r_N > 2^N$ satisfying that $|F(r_N, x, f)| > \lambda$. By this, (1.3), Hölder’s inequality and Lemma 2.1 (iii), we obtain

$$\lambda < |F(r_N, x, f)| = \left| \int_{\mathcal{X}} S_{r_N}(x, y) f(y) d\mu(y) \right| \lesssim \frac{1}{[V_{2^N}(x)]^{1/p}} \left\{ \int_{\mathcal{X}} |f(y)|^p d\mu(y) \right\}^{1/p},$$

which implies that $V_{2^N}(x) \lesssim \lambda^{-p} \|f\|_{L^p(\mathcal{X})}^p < \infty$ for all $N \in \mathbb{N}$, and hence $\mu(\mathcal{X}) < \infty$. This contradicts the assumption $\mu(\mathcal{X}) = \infty$. Thus, (2.2) holds.

It is not so difficult to prove that for any given $N \in \mathbb{N}$, there exist $L_N < N$ with $L_N \in \mathbb{Z}$ or $L_N = -\infty$, a set of indices $I_{N,\ell}$ with $\ell \in \{L_N + 1, \dots, N\}$, and disjoint balls $\{B(y_{\ell,j}^N, 2^\ell)\}_{L_N < \ell \leq N, j \in I_{N,\ell}}$ satisfying

- (i) $y_{\ell,j}^N \in W_{\ell,\lambda}$;
- (ii) $B(y_{\ell,j}^N, 2^\ell) \cap (\bigcup_{m=\ell+1}^N \bigcup_{i \in I_{N,m}} B(y_{m,i}^N, 2^m)) = \emptyset$;
- (iii) for any $x \in W_{\ell,\lambda}$, $B(x, 2^\ell) \cap (\bigcup_{m=\ell}^N \bigcup_{i \in I_{N,m}} B(y_{m,i}^N, 2^m)) \neq \emptyset$.

In fact, we start with $\ell = N$ and choose an arbitrary point in $W_{N,\lambda}$ as $y_{N,1}^N$. Then we find a point $y_{N,2}^N \in W_{N,\lambda} \setminus B(y_{N,1}^N, 2^N)$ such that $B(y_{N,2}^N, 2^N) \cap B(y_{N,1}^N, 2^N) = \emptyset$. Continuing in this way, by Zorn’s lemma and the doubling property of the measure μ , we arrive at $I_{N,N} = \mathbb{N}$ or $I_{N,N}$ will be a finite set. We then consider $\ell = N - 1$. In this way, one finds the desired balls.

From (i), (ii) and (iii), it follows that for each $N \in \mathbb{N}$,

$$(2.3) \quad W_\lambda \subset \left(\bigcup_{\ell=L_N+1}^N \bigcup_{j \in I_{N,\ell}} [B(y_{\ell,j}^N, 2^{\ell+1}) \times (0, 2^\ell)] \right) \cup (E_N \times (2^N, \infty)).$$

To see (2.3), notice that for any $(x, r) \in W_\lambda$, $|F(r, x, f)| > \lambda$. If $r > 2^N$, then $(x, r) \in E_N \times (2^N, \infty)$; otherwise there exists $\ell \leq N$ such that $2^{\ell-1} < r \leq 2^\ell$, which implies that $x \in W_{\ell,\lambda}$. By Property (iii) above, there exist integers $\ell \leq m \leq N$ and $j \in I_{N,m}$ such that $B(x, 2^\ell) \cap B(y_{m,j}^N, 2^m) \neq \emptyset$. Noticing that $\ell \leq m$, we have $x \in B(y_{m,j}^N, 2^{m+1})$ and thus $(x, r) \in B(y_{m,j}^N, 2^{m+1}) \times (0, 2^m)$, which yields (2.3) then.

For any $N \in \mathbb{N}$, by (2.3),

$$\nu(W_\lambda) \leq \sum_{\ell=L_N+1}^N \sum_{j \in I_{N,\ell}} \nu(B(y_{\ell,j}^N, 2^{\ell+1}) \times (0, 2^\ell)) + \nu(E_N \times (2^N, \infty)).$$

Letting $N \rightarrow \infty$ in the formula above, then using (2.1) and (2.2), we obtain

$$\begin{aligned}
 \nu(W_\lambda) &\leq \lim_{N \rightarrow \infty} \sum_{\ell=L_N+1}^N \sum_{j \in I_{N,\ell}} [\mu(B(y_{\ell,j}^N, 2^{\ell+1}))]^{1+\delta} \\
 (2.4) \quad &\leq \lim_{N \rightarrow \infty} \left\{ \sum_{\ell=L_N+1}^N \sum_{j \in I_{N,\ell}} \mu(B(y_{\ell,j}^N, 2^{\ell+1})) \right\}^{1+\delta},
 \end{aligned}$$

where in the second step we use the fact $(\sum_{j \in \mathbb{N}} |a_j|)^\kappa \leq \sum_{j \in \mathbb{N}} |a_j|^\kappa$ for any $\kappa \in (0, 1]$. Choose $\tilde{p} \in (1, p)$. For any $N, \ell \in (L_N, N]$ and any given $j \in I_{N,\ell}$, by Property (i), the size condition of $S_{-\ell}$, (1.2) and Hölder’s inequality, we have

$$\begin{aligned}
 \lambda &< \sup_{2^{\ell-1} < r \leq 2^\ell} |F(r, y_{\ell,j}^N, f)| \\
 &\lesssim \frac{1}{V_{2^\ell}(y_{\ell,j}^N)} \int_{B(y_{\ell,j}^N, 2^\ell)} |f(z)| d\mu(z) \\
 &\quad + \sum_{k=1}^\infty \int_{2^{\ell+k-1} \leq d(y_{\ell,j}^N, z) < 2^{\ell+k}} \frac{1}{V_{2^\ell}(y_{\ell,j}^N) + V(y_{\ell,j}^N, z)} \left(\frac{2^\ell}{2^\ell + d(y_{\ell,j}^N, z)} \right)^{\epsilon_2} |f(z)| d\mu(z) \\
 &\lesssim \sum_{k=0}^\infty 2^{-k\epsilon_2} \left\{ \frac{1}{V_{2^{\ell+k}}(y_{\ell,j}^N)} \int_{B(y_{\ell,j}^N, 2^{\ell+k})} |f(z)|^{p/\tilde{p}} d\mu(z) \right\}^{\tilde{p}/p} \\
 &\lesssim \inf \left\{ \left[\mathcal{M}(|f|^{p/\tilde{p}})(z_{\ell,j}^N) \right]^{\tilde{p}/p} : z_{\ell,j}^N \in B(y_{\ell,j}^N, 2^\ell) \right\},
 \end{aligned}$$

which together with the pairwise disjointness of the balls $\{B(y_{\ell,j}^N, 2^\ell)\}_{L_N < \ell \leq N, j \in I_{N,\ell}}$ and $\tilde{L}^p(\mathcal{X})$ -boundedness of \mathcal{M} yields that for all $N \in \mathbb{N}$,

$$\sum_{\ell=L_N+1}^N \sum_{j \in I_{N,\ell}} \mu(B(y_{\ell,j}^N, 2^{\ell+1})) \lambda^p \lesssim \int_{\mathcal{X}} \left[\mathcal{M}(|f|^{p/\tilde{p}})(z) \right]^{\tilde{p}} d\mu(z) \lesssim \|f\|_{L^p(\mathcal{X})}^p.$$

Combining this with (2.4) shows that $\lambda^{p(1+\delta)} \nu(W_\lambda) \lesssim \|f\|_{L^p(\mathcal{X})}^{p(1+\delta)}$. Then the desired conclusion follows from the Marcinkiewicz interpolation theorem, which completes the proof of Lemma 2.2. \square

LEMMA 2.3. — *Let $x_0 \in \mathcal{X}$, $r_0 > 0$ and g be a non-negative function on \mathcal{X} . Then for any $t \in (0, 1]$, there exist $\{x_j\}_j \subset \mathcal{X}$ with $x_j \equiv x_j(g, t, x_0, r_0)$ and positive constants C_6 and C_7 depending only on \mathcal{X} such that for all $x \in \mathcal{X}$,*

$$(2.5) \quad 1 \leq \sum_j \chi_{B(x_j, C_4 t r_j)}(x) \leq C_6$$

and

$$(2.6) \quad g(x_j)^{1/2} \leq C_7 F(tr_j, x_j, g^{1/2} \chi_{B(x_j, C_4 tr_j)}),$$

where $r_j \equiv r_0 + d(x_j, x_0)$ and F is as in Lemma 2.2. In particular, there exists a constant $C_8 > 1$ such that for all j and all $x \in B(x_j, C_4 tr_j)$,

$$(2.7) \quad \begin{aligned} & \frac{2^a C_8}{V_{r_j}(x_j)} \left(\frac{r_0}{r_j}\right)^a V_{tr_j}(x_j) S_{tr_j}(x_j, x) \chi_{B(x_j, C_4 tr_j)}(x) \\ & \geq \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^a, \end{aligned}$$

where S_{tr_j} is as in (1.3). Moreover, C_6 through C_8 are independent of x_0, r_0 and g .

Proof. — For any $x \in \mathcal{X}$, set $r_x \equiv r_0 + d(x_0, x)$. By Zorn’s lemma, there exists a set of points $\{y_j\}_j \subset \mathcal{X}$ satisfying that $y_j \notin \cup_{i=1}^{j-1} B(y_i, C_4 tr_{y_i}/4)$ and $\mathcal{X} = \cup_j B(y_j, C_4 tr_{y_j}/4)$. From the selection of $\{y_j\}_j$, it follows that for any $i \neq j$,

$$(2.8) \quad d(y_i, y_j) \geq \frac{1}{4} C_4 t \min\{r_{y_i}, r_{y_j}\},$$

and that for all $x \in \mathcal{X}$,

$$(2.9) \quad \sum_j \chi_{B(y_j, C_4 tr_{y_j}/4)}(x) \geq 1.$$

By (1.2) and the disjointness of $\{B(y_i, C_4 tr_0/8)\}_i$, we know that $\{y_j\}_j$ is at most countable. For every y_j , choose $x_j \equiv x_j(g, t, x_0, r_0)$ satisfying that

$$(2.10) \quad d(x_j, y_j) < C_4 tr_{y_j}/4$$

and

$$(2.11) \quad g(x_j)^{1/2} \leq \frac{1}{\mu(B(y_j, C_4 tr_{y_j}/4))} \int_{B(y_j, C_4 tr_{y_j}/4)} g(z)^{1/2} d\mu(z).$$

Let $r_j \equiv r_{x_j}$. By (2.10) and the triangle inequality for d together with $C_4 < 1/2$ and $t \in (0, 1]$, we have

$$(2.12) \quad r_{y_j}/2 \leq r_j \leq 2r_{y_j},$$

which together with (2.9) implies the left-hand side inequality of (2.5).

For any $x \in \mathcal{X}$, set $J(x) \equiv \{j : d(x_j, x) < C_4 tr_j\}$. Notice that for any $j \in J(x)$, by (2.12), $r_j/2 \leq r_x \leq 2r_j$. This together with (2.10) and (2.12) yields that $J(x) \subset \tilde{J}(x)$, where $\tilde{J}(x) \equiv \{j : d(y_j, x) < 3C_4 tr_x\}$. It follows from (1.2) and the pairwise disjointness of $\{B(y_i, C_4 tr_0/8)\}_i$ that $\#\tilde{J}(x) < \infty$. Thus, we may assume that $r_{y_1} = \min\{r_{y_j} : j \in J(x)\}$. Therefore, by (2.8),

$\{B(y_j, C_4tr_{y_1}/8)\}_{j \in J(x)}$ are mutually disjoint. Furthermore, for any $j \in J(x)$, by (2.12) and $r_j/2 \leq r_x \leq 2r_j$, we have

$$B(y_j, C_4tr_x/32) \subset B(y_j, C_4tr_{y_1}/8) \subset B(x, 4C_4tr_x) \subset B(y_j, 7C_4tr_x).$$

From this and (1.2), it follows that $\#J(x)$ is bounded by a positive constant which depends only on \mathcal{X} . This implies the validity of (2.5).

The fact that $B(y_j, C_4tr_{y_j}/4) \subset B(x_j, C_4tr_j)$ together with (2.11) and (1.5) implies (2.6). Since for any $x \in B(x_j, C_4tr_j)$, we have $r_j/2 \leq r_x \leq 2r_j$. Using this fact, (1.5), (2.5) and Lemma 2.1 (i), we obtain (2.7), which completes the proof of Lemma 2.3. \square

LEMMA 2.4. — *Let $t \in (0, 1]$, $a \in [0, \infty)$, $b \in (a, \infty)$, $M \in [0, \infty)$ and $\{x_j\}_j \subset \mathcal{X}$ satisfying*

$$(2.13) \quad \sum_j \chi_{B(x_j, C_4tr_j)}(x) \leq C_6,$$

where $r_j \equiv r_0 + d(x_j, x_0)$ with $r_0 > 0$ and $x_0 \in \mathcal{X}$ and C_6 is as in (2.5). For any j and $x \in \mathcal{X}$, set

$$u_j(x) \equiv \frac{1}{V_{r_j}(x_j)} \left(\frac{1}{r_j}\right)^a \frac{V_{C_4tr_j}(x_j)}{V_{tr_j}(x_j) + V_{tr_j}(x) + V(x_j, x)} \\ \times \left(\frac{tr_j}{tr_j + d(x_j, x)}\right)^b \chi_{[M, \infty)}\left(\frac{d(x_j, x)}{tr_j}\right),$$

where $\chi_{[M, \infty)}$ is the characteristic function of the interval $[M, \infty)$. Then there exists $C_9 > 1$ independent of x_0, r_0 and M such that for all $x \in \mathcal{X}$,

$$\sum_j u_j(x) \leq C_9 \max\{t^b, (1 + M)^{-b}\} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{1}{r_0 + d(x_0, x)}\right)^a.$$

Proof. — For any $k \in \mathbb{Z}$, set $J(k) \equiv \{j : 2^{k-1} \leq r_j < 2^k\}$ and $v_k \equiv \sum_{j \in J(k)} u_j$. For any fixed $x \in \mathcal{X}$, let

$$W_1 \equiv \{k \in \mathbb{Z} : (r_0 + d(x_0, x))/2 \leq 2^k < 4(r_0 + d(x_0, x))\},$$

$$W_2 \equiv \{k \in \mathbb{Z} : 2^k < (r_0 + d(x_0, x))/2\},$$

and

$$W_3 \equiv \{k \in \mathbb{Z} : 2^k \geq 4(r_0 + d(x_0, x))\}.$$

We then write,

$$\sum_j u_j(x) = \sum_{k \in W_1} \sum_{j \in J(k)} u_j(x) + \sum_{k \in W_2} \sum_{j \in J(k)} u_j(x) + \sum_{k \in W_3} \sum_{j \in J(k)} u_j(x) \equiv Z_1 + Z_2 + Z_3.$$

To estimate Z_1 , notice that for any $k \in W_1$ and $j \in J(k)$, we have

$$(2.14) \quad \mu(B(x_j, r_0 + d(x_j, x_0))) \sim \mu(B(x_0, r_j)) \sim \mu(B(x_0, 2^k)) \sim V_{r_0}(x_0) + V(x_0, x).$$

For any $j \in J(k)$ and $z \in B(x_j, C_4tr_j)$, we have $d(z, x_j) < C_4tr_j < C_4t2^k$, which together with Lemma 2.1 further implies that for all $z \in B(x_j, C_4tr_j)$,

$$(2.15) \quad t2^k + d(x_j, x) \sim t2^k + d(z, x), \quad V_{t2^k}(x_j) + V(x_j, x) \sim V_{t2^k}(z) + V(z, x)$$

and

$$(2.16) \quad \chi_{[M, \infty)} \left(\frac{d(x_j, x)}{tr_j} \right) \leq \chi_{[M, \infty)} \left(\frac{d(x_j, x)}{t2^{k-1}} \right) \leq \chi_{[M-2C_4, \infty)} \left(\frac{d(z, x)}{t2^{k-1}} \right).$$

From (2.13) through (2.16), it follows that

$$\begin{aligned} v_k(x) &\lesssim \frac{1}{2^{a(k-1)}} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \\ &\quad \times \sum_{j \in J(k)} \frac{V_{C_4tr_j}(x_j)}{V_{t2^k}(x_j) + V_{t2^k}(x) + V(x_j, x)} \left(\frac{t2^k}{t2^k + d(x_j, x)} \right)^b \chi_{[M, \infty)} \left(\frac{d(x_j, x)}{t2^{k-1}} \right) \\ &\lesssim \frac{1}{2^{ak}} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \\ &\quad \times \int_{\mathcal{X}} \frac{1}{V_{t2^k}(z) + V_{t2^k}(x) + V(z, x)} \left(\frac{t2^k}{t2^k + d(z, x)} \right)^b \chi_{[M-2C_4, \infty)} \left(\frac{d(z, x)}{t2^{k-1}} \right) d\mu(z). \end{aligned}$$

Denote by J the integral in the last formula. When $0 \leq M \leq 4C_4 + 1$, by Lemma 2.1,

$$J \leq \int_{\mathcal{X}} \frac{1}{V_{t2^k}(z) + V_{t2^k}(x) + V(z, x)} \left(\frac{t2^k}{t2^k + d(z, x)} \right)^b d\mu(z) \lesssim 1 \lesssim (1 + M)^{-b}.$$

When $M > 1 + 4C_4$, we have $M - 2C_4 > (1 + M)/2$. For any $i \in \mathbb{N}$ and $x \in \mathcal{X}$, set

$$R_i \equiv \{z \in \mathcal{X} : 2^{i+k-2}(M - 2C_4)t \leq d(z, x) < 2^{i+k-1}(M - 2C_4)t\}.$$

We then obtain

$$\begin{aligned} J &= \sum_{i=1}^{\infty} \int_{R_i} \frac{1}{V_{t2^k}(z) + V_{t2^k}(x) + V(z, x)} \left(\frac{t2^k}{t2^k + d(z, x)} \right)^b d\mu(z) \\ &\lesssim \sum_{i=1}^{\infty} [1 + 2^{i-1}(M - 2C_4)]^{-b} \lesssim (M - 2C_4)^{-b} \lesssim (1 + M)^{-b}. \end{aligned}$$

Therefore, for all $k \in W_1$ and $x \in \mathcal{X}$, we have

$$v_k(x) \lesssim \frac{1}{2^{ak}} \frac{(1 + M)^{-b}}{V_{r_0}(x_0) + V(x_0, x)},$$

which together with the fact $\#W_1 \leq 5$ yields that

$$Z_1 \leq \sum_{k \in W_1} \frac{1}{2^{ak}} \frac{(1 + M)^{-b}}{V_{r_0}(x_0) + V(x_0, x)} \lesssim \frac{1}{(r_0 + d(x_0, x))^a} \frac{(1 + M)^{-b}}{V_{r_0}(x_0) + V(x_0, x)}.$$

To estimate Z_2 , notice that for any $k \in W_2$ and $j \in J(k)$, we have $r_0 + d(x_0, x) \leq r_0 + d(x_0, x_j) + d(x_j, x) < 2^k + d(x, x_j) \leq (r_0 + d(x_0, x))/2 + d(x, x_j)$.

From this, it follows that for any $j \in J(k)$,

$$(2.17) \quad d(x, x_j) > (r_0 + d(x_0, x))/2 > 2^k$$

and thus

$$(2.18) \quad V(x_j, x) \gtrsim \mu(B(x, r_0 + d(x_0, x))) \sim V_{r_0}(x_0) + V(x_0, x).$$

For $k \in \mathbb{Z}$ and $j \in J(k)$, we have $V_{2^k}(x_j) \sim V_{2^k}(x_0)$ and $B(x_j, C_4 t r_j) \subset B(x_0, (1 + C_4 t)2^k)$, which together with (1.2) and (2.13) yields that

$$(2.19) \quad \sum_{j \in J(k)} \frac{V_{C_4 t 2^k}(x_j)}{V_{2^k}(x_j)} \lesssim \int_{B(x_0, (1+C_4 t)2^k)} \frac{1}{V_{2^k}(x_0)} \sum_{j \in J(k)} \chi_{B(x_j, C_4 t r_j)}(x) d\mu(x) \lesssim 1.$$

For any $k \in \mathbb{Z}$ and $j \in J(k)$, we have $V_{r_j}(x_j) \gtrsim V_{2^k}(x_j)$, which together with (2.17), (2.18), and (2.19) implies that

$$\begin{aligned} Z_2 &\lesssim \sum_{k \in W_2} \sum_{j \in J(k)} \frac{1}{2^{ak}} \frac{1}{V_{2^k}(x_j)} \frac{V_{C_4 t 2^k}(x_j)}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{t2^k}{t2^k + r_0 + d(x_0, x)} \right)^b \\ &\lesssim t^b \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{1}{r_0 + d(x_0, x)} \right)^a. \end{aligned}$$

To estimate Z_3 , notice that for any $k \in W_3$ and $j \in J(k)$,

$$(2.20) \quad V_{r_j}(x_j) \sim \mu(B(x_j, r_0 + d(x_j, x_0))) \gtrsim \mu(B(x_0, 2^k)) \gtrsim V_{r_0}(x_0) + V(x_0, x).$$

Moreover, since

$$2^{k-1} \leq r_0 + d(x_j, x_0) \leq r_0 + d(x_j, x) + d(x, x_0) \leq 2^{k-2} + d(x_j, x),$$

we have $d(x_j, x) \geq 2^{k-2} \geq r_j/4$. Consequently, for any $k \in W_3$ and $j \in J(k)$,

$$(2.21) \quad V_{t r_j}(x_j) + V_{t r_j}(x) + V(x_j, x) \geq V(x_j, x) \gtrsim V_{r_j/2}(x_j).$$

An argument similar to (2.19) yields that

$$(2.22) \quad \sum_{j \in J(k)} \frac{V_{C_4 t r_j}(x_j)}{V_{r_j/2}(x_j)} \lesssim 1.$$

Applying (2.20) through (2.22) and the fact that $d(x_j, x) \geq r_j/4$, we obtain

$$\begin{aligned} Z_3 &\lesssim \sum_{k \in W_3} \sum_{j \in J(k)} \frac{1}{2^{ka}} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \frac{V_{C_4 t r_j}(x_j)}{V_{r_j/2}(x_j)} \left(\frac{t}{1+t} \right)^b \\ &\lesssim t^b \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{1}{r_0 + d(x_0, x)} \right)^a, \end{aligned}$$

which completes the proof of Lemma 2.4. □

3. Proofs of Theorem 1.6 and Proposition 1.7

Assuming Proposition 1.7 for the moment, we now prove Theorem 1.6.

Proof of Theorem 1.6. — Choose $\delta_0 \in (0, \log_\eta(1-\sigma))$ satisfying $p > n/(n+\delta_0)$. By Proposition 1.7 and the $L^{p(n+\delta_0)/n}(\mathcal{X})$ -boundedness of \mathcal{M} , we obtain

$$\|f^*\|_{L^p(\mathcal{X})} \lesssim \left\| \left[\mathcal{M}([\mathcal{M}_0(f)]^{n/(n+\delta_0)}) \right]^{(n+\delta_0)/n} \right\|_{L^p(\mathcal{X})} \lesssim \|\mathcal{M}_0(f)\|_{L^p(\mathcal{X})},$$

which completes the proof of Theorem 1.6. □

The rest of this section is devoted to the proof of Proposition 1.7. The key for the proof of Proposition 1.7 is to obtain a desired expansion (3.11) for any $\varphi \in \mathcal{G}(x_0, r_0, \epsilon, \epsilon)$ in terms of the given $(\epsilon_1, \epsilon_2, \epsilon_3)$ -SAOTI and to show this expansion converges in $\mathcal{G}_0^\epsilon(\beta, \gamma)$. To this end, for a given $\varphi \in \mathcal{G}(x_0, r_0, \epsilon, \epsilon)$, we construct a sequence of functions $\{\varphi_s\}_{s \in \mathbb{Z}_+}$ and obtain some desired estimates for these functions.

Proof of Proposition 1.7. — Let $\varphi \in \mathcal{G}(x_0, r_0, \epsilon, \epsilon)$ satisfy $\|\varphi\|_{\mathcal{G}(x_0, r_0, \epsilon, \epsilon)} \leq 1$ for some $x_0 \in \mathcal{X}$ and $r_0 > 0$. Write φ as

$$\varphi = (\Re(\varphi) \vee 0) - (\Re(\varphi) \wedge 0) + i[(\Im(\varphi) \vee 0) - (\Im(\varphi) \wedge 0)],$$

where $\Re(\varphi)$ and $\Im(\varphi)$ represent the real part and the imaginary part of the function φ , respectively. Since $\|\varphi\|_{\mathcal{G}(x_0, r_0, \epsilon, \epsilon)} \leq 1$, it follows easily that each term of the decomposition above has a norm in $\mathcal{G}(x_0, r_0, \epsilon, \epsilon)$ at most 1. Thus, we may assume that φ is non-negative.

Fix $A \equiv 2^{\epsilon+1}C_8$ and some $\sigma \in (0, 1/(1+AC_3C_9))$. Choose positive numbers H and η such that

$$H \leq \min \left\{ \frac{1}{2} \left(\frac{1-\sigma-AC_3C_9\sigma}{2} \right)^{1/\epsilon}, \frac{1}{2} \left(\frac{(1-\sigma-AC_3C_9\sigma)(1-\sigma)}{4AC_3C_9\sigma} \right)^{1/\epsilon_1} \right\}$$

and

$$\eta < \min \left\{ (1-\sigma)^{1/\epsilon}, \frac{H}{C_4}, \frac{1}{AC_3C_9}, \left(\frac{1-\sigma}{2} \right)^{1/\epsilon_1}, H \left(\frac{1}{AC_3C_9} \right)^{1/\epsilon_2} \right\}.$$

Choose $\delta_0 \in (0, \log_\eta(1-\sigma))$. For any $s \in \mathbb{N}$, applying Lemma 2.3 with $t = \eta^s$ and $g = \mathcal{M}_0(f)$, we obtain $\{x_{s,j} : s \in \mathbb{N}, j = 1, \dots, j(s)\} \subset \mathcal{X}$, where $j(s)$ can be finite or ∞ . For each $s \in \mathbb{N}$ and $j = 1, \dots, j(s)$, let $r_{s,j} \equiv r_0 + d(x_{s,j}, x_0)$. Lemma 2.3 further implies that

(A) for any $x \in \mathcal{X}$ and $s \in \mathbb{N}$, $1 \leq \sum_{j=1}^{j(s)} \chi_{B_{s,j}}(x) \leq C_6$, where $B_{s,j} \equiv B(x_{s,j}, C_4\eta^s r_{s,j})$;

(B) $[\mathcal{M}_0(f)(x_{s,j})]^{1/2} \leq C_7 F(\eta^s r_{s,j}, x_{s,j}, [\mathcal{M}_0(f)]^{1/2} \chi_{B_{s,j}})$.

Let us now inductively construct $\{\epsilon_{s,j} : s \in \mathbb{N}, j = 1, \dots, j(s)\} \subset \{-1, 0, 1\}$ and functions $\{\varphi_s\}_{s \in \mathbb{Z}_+}$ satisfying that for all $x \in \mathcal{X}$,

- (C) for all $s \in \mathbb{Z}_+$, $|\varphi_s(x)| \leq (1 - \sigma)^s \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon$;
- (D) $\varphi_0(x) \equiv \varphi(x)$, and for any $s \in \mathbb{N}$,

$$(3.1) \quad \varphi_s(x) \equiv \varphi(x) - A\sigma \sum_{i=1}^s \sum_{j=1}^{j(i)} (1 - \sigma)^{i-1} \epsilon_{i,j} \times \frac{V_{C_4 \eta^s r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}} \right)^\epsilon S_{\eta^i r_{i,j}}(x_{i,j}, x).$$

In fact, obviously, φ_0 satisfies (C). Suppose that $\{\epsilon_{i,j} : i = 1, \dots, s - 1; j = 1, \dots, j(i)\}$ and $\{\varphi_i\}_{i=0}^{s-1}$ satisfying (C) have been constructed. Then for each $j = 1, \dots, j(s)$, let $\epsilon_{s,j} \equiv \text{sign}(\varphi_{s-1}(x_{s,j}))$, and φ_s be as in (D). Now it remains to verify that φ_s satisfies (C). To this end, for all $x \in \mathcal{X}$, set

$$(3.2) \quad \omega_s(x) \equiv A\sigma \sum_{j=1}^{j(s)} (1 - \sigma)^{s-1} \epsilon_{s,j} \frac{V_{C_4 \eta^s r_{s,j}}(x_{s,j})}{V_{r_{s,j}}(x_{s,j})} \left(\frac{r_0}{r_{s,j}} \right)^\epsilon S_{\eta^s r_{s,j}}(x_{s,j}, x).$$

By the size condition of S_t , $\epsilon_2 > \epsilon$ and Lemma 2.4 with $t = \eta^s$, $a = \epsilon$, $b = \epsilon_2$ and $M = 0$, we obtain

$$(3.3) \quad \begin{aligned} |\omega_s(x)| &\leq AC_3\sigma \sum_{j=1}^{j(s)} (1 - \sigma)^{s-1} \frac{V_{C_4 \eta^s r_{s,j}}(x_{s,j})}{V_{r_{s,j}}(x_{s,j})} \left(\frac{r_0}{r_{s,j}} \right)^\epsilon \\ &\quad \times \frac{1}{V_{\eta^s r_{s,j}}(x_{s,j}) + V_{\eta^s r_{s,j}}(x) + V(x_{s,j}, x)} \left(\frac{\eta^s r_{s,j}}{\eta^s r_{s,j} + d(x_{s,j}, x)} \right)^{\epsilon_2} \\ &\leq AC_3C_9\sigma(1 - \sigma)^{s-1} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon. \end{aligned}$$

If $d(x, y) \leq \eta^{s-1}(r_0 + d(x_0, x))/2$, then for any $i = 1, \dots, s - 1$, we have that $d(x, y) \leq (\eta^i r_{i,j} + d(x, x_{i,j}))/2$. This combined with the regularity of $S_{\eta^i r_{i,j}}$ and Lemma 2.4 with $t = \eta^i$, $a = \epsilon + \epsilon_1$, $b = \epsilon_1 + \epsilon_2$ and $M = 0$ yields that

$$\begin{aligned} &|\varphi_{s-1}(x) - \varphi_{s-1}(y)| \\ &\leq |\varphi(x) - \varphi(y)| + AC_3\sigma \sum_{i=1}^{s-1} \sum_{j=1}^{j(i)} (1 - \sigma)^{i-1} \\ &\quad \times \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}} \right)^\epsilon |S_{\eta^i r_{i,j}}(x_{i,j}, x) - S_{\eta^i r_{i,j}}(x_{i,j}, y)| \\ &\leq |\varphi(x) - \varphi(y)| + AC_3\sigma \sum_{i=1}^{s-1} \sum_{j=1}^{j(i)} \frac{(1 - \sigma)^{i-1}}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}} \right)^\epsilon \left(\frac{\eta^i r_{i,j}}{\eta^i r_{i,j} + d(x_{i,j}, x)} \right)^{\epsilon_2} \\ &\quad \times \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{\eta^i r_{i,j}}(x_{i,j}) + V_{\eta^i r_{i,j}}(x) + V(x_{i,j}, x)} \left(\frac{d(x, y)}{\eta^i r_{i,j} + d(x_{i,j}, x)} \right)^{\epsilon_1} \end{aligned}$$

$$(3.4) \quad \leq |\varphi(x) - \varphi(y)| + \frac{AC_3C_9\sigma}{1 - \sigma - \eta^{\epsilon_1}} \left[\left(\frac{1 - \sigma}{\eta^{\epsilon_1}} \right)^{s-1} - 1 \right] \\ \times \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)} \right)^{\epsilon_1} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon.$$

When

$$(3.5) \quad d(x, y) < H\eta^{s-1}(r_0 + d(y, x_0)),$$

the assumption that $H < 1/4$ gives $d(x, y) < 2H\eta^{s-1}(r_0 + d(x_0, x)) < \eta^{s-1}(r_0 + d(x_0, x))/2$, which together with (3.4) and the regularity of φ yields that

$$(3.6) \quad |\varphi_{s-1}(x) - \varphi_{s-1}(y)| \leq \left[(2H)^\epsilon \left(\frac{\eta^\epsilon}{1 - \sigma} \right)^{s-1} + \frac{(2H)^{\epsilon_1} AC_3C_9\sigma}{1 - \sigma - \eta^{\epsilon_1}} \left(1 - \left(\frac{\eta^{\epsilon_1}}{1 - \sigma} \right)^{s-1} \right) \right] \\ \times \frac{(1 - \sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon \\ \leq \lambda_H \frac{(1 - \sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon,$$

where

$$\lambda_H \equiv (2H)^\epsilon + \frac{(2H)^{\epsilon_1} AC_3C_9\sigma}{1 - \sigma - \eta^{\epsilon_1}}.$$

For any $\lambda \in \mathbb{R}$ and $s \in \mathbb{N}$, set

$$\Omega_{s,\lambda} \equiv \left\{ x \in \mathcal{X} : \varphi_{s-1}(x) > \lambda(1 - \sigma)^{s-1} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon \right\}.$$

If $\varphi_{s-1}(x_{s,j}) < 0$, then by (3.6) and the definition of $\Omega_{s,\lambda}$, we obtain

$$(3.7) \quad B(x_{s,j}, \eta^{s-1}r_{s,j}/3) \cap \Omega_{s,\lambda_H} = \emptyset.$$

In fact, for any $x \in B(x_{s,j}, \eta^{s-1}r_{s,j}/3)$, by (3.6),

$$\varphi_{s-1}(x) \leq \varphi_{s-1}(x_{s,j}) + \lambda_H \frac{(1 - \sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon \\ < \lambda_H \frac{(1 - \sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon,$$

which implies that $x \notin \Omega_{s,\lambda_H}$ and thus (3.7) holds.

For any $x \in \mathcal{X}$, by (3.2), we write

$$\omega_s(x) = \sum_{\{j: \varphi_{s-1}(x_{s,j}) > 0\}} A\sigma(1 - \sigma)^{s-1} \frac{VC_4\eta^{sr_{s,j}}(x_{s,j})}{V_{r_{s,j}}(x_{s,j})} \left(\frac{r_0}{r_{s,j}} \right)^\epsilon S_{\eta^{sr_{s,j}}}(x_{s,j}, x) \\ - \sum_{\{j: \varphi_{s-1}(x_{s,j}) < 0\}} \dots \equiv \text{I} - \text{II}.$$

We now turn to obtain a desired upper bound for φ_s by considering two cases: $x \in \Omega_{s,\lambda_H}$ and $x \notin \Omega_{s,\lambda_H}$. For any $x \in \Omega_{s,\lambda_H}$, Property (A) implies that $x \in B(x_{s,j_0}, C_4\eta^s r_{s,j_0})$ for some j_0 . By this and the assumption $\eta < H/C_4$, we have $d(x, x_{s,j_0}) < C_4\eta^s r_{s,j_0} \leq H\eta^{s-1} r_{s,j_0}$, which together with (3.7) implies that $\varphi_{s-1}(x_{s,j_0}) > 0$. Thus, by (2.7),

$$\begin{aligned} \text{I} &\geq A\sigma(1-\sigma)^{s-1} \frac{V_{C_4\eta^s r_{s,j_0}}(x_{s,j_0})}{V_{r_{s,j_0}}(x_{s,j_0})} \left(\frac{r_0}{r_{s,j_0}}\right)^\epsilon S_{\eta^s r_{s,j_0}}(x_{s,j_0}, x) \\ &\geq \frac{A\sigma(1-\sigma)^{s-1}}{C_8 2^\epsilon} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon. \end{aligned}$$

On the other hand, by (3.7) and Lemma 2.4,

$$\text{II} \leq \frac{AC_3 C_9 \sigma(1-\sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon \max\{\eta^{s\epsilon_2}, (1 + H\eta^{-1})^{-\epsilon_2}\}.$$

From the assumption on η , it follows easily that

$$\frac{A}{C_8 2^\epsilon} - AC_3 C_9 \max\{\eta^{s\epsilon_2}, (1 + H\eta^{-1})^{-\epsilon_2}\} \geq 1,$$

which together with the estimates for I and II yields that when $x \in \Omega_{s,\lambda_H}$,

$$(3.8) \quad \omega_s(x) \geq \frac{\sigma(1-\sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon.$$

Thus, by (2.1) and the fact that (C) holds for $s - 1$, we obtain that when $x \in \Omega_{s,\lambda_H}$,

$$\varphi_s(x) = \varphi_{s-1}(x) - \omega_s(x) \leq \frac{(1-\sigma)^s}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon.$$

Let now $x \notin \Omega_{s,\lambda_H}$. Notice that (C) holds for $s - 1$. From this and (3.3), we deduce that

$$\begin{aligned} \varphi_s(x) &= \varphi_{s-1}(x) - \omega_s(x) \\ &\leq \lambda_H \frac{(1-\sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon \\ &\quad + AC_3 C_9 \sigma(1-\sigma)^{s-1} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon \\ &\leq \frac{(1-\sigma)^s}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon, \end{aligned}$$

where in the last inequality, we used the fact $\lambda_H + AC_3 C_9 \sigma \leq 1 - \sigma$. Thus, we obtain a desired upper bound of φ_s .

Let us now show that φ_s has the desired lower bound also by considering two cases: $x \notin \Omega_{s,-\lambda_H}$ and $x \in \Omega_{s,-\lambda_H}$. For every $x_{s,j}$ satisfying $\varphi_{s-1}(x_{s,j}) > 0$, by (3.6) and an argument similar to the proof of (3.7), we have $B(x_{s,j}, H\eta^{s-1} r_{s,j}) \cap$

$(\mathcal{X} \setminus \Omega_{s,-\lambda_H}) = \emptyset$. From this, Lemmas 2.3 and 2.4 together with an argument similar to the estimates for I and II, it follows that when $x \notin \Omega_{s,-\lambda_H}$,

$$(3.9) \quad \omega_s(x) \leq -\frac{\sigma(1-\sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon.$$

Therefore, (3.9) and the lower bound of φ_{s-1} yields that when $x \notin \Omega_{s,-\lambda_H}$,

$$(3.10) \quad \varphi_s(x) = \varphi_{s-1}(x) - \omega_s(x) \geq -\frac{(1-\sigma)^s}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon;$$

and the validity of (C) for $s - 1$ together with (3.3) and the assumption $\lambda_H + AC_3C_9\sigma \leq 1 - \sigma$ implies that when $x \in \Omega_{s,-\lambda_H}$,

$$\begin{aligned} \varphi_s(x) &\geq \varphi_{s-1}(x) - \omega_s(x) \\ &\geq -\lambda_H \frac{(1-\sigma)^{s-1}}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon \\ &\quad - AC_3C_9\sigma(1-\sigma)^{s-1} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon \\ &\geq -\frac{(1-\sigma)^s}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)} \right)^\epsilon, \end{aligned}$$

which together with (3.10) further yields the desired lower bound of φ_s . This finishes the proofs of (C) and (D).

It follows from (C) and (D) that for all $x \in \mathcal{X}$,

$$(3.11) \quad \varphi(x) = A\sigma \sum_{i=1}^{\infty} \sum_{j=1}^{j(i)} (1-\sigma)^{i-1} \epsilon_{i,j} \frac{V_{C_4\eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}} \right)^\epsilon S_{\eta^i r_{i,j}}(x_{i,j}, x).$$

Set

$$\Phi_L(x) \equiv A\sigma \sum_{i=1}^L \sum_{\{j: \eta^i r_{i,j} \leq L^i\}} (1-\sigma)^{i-1} \epsilon_{i,j} \frac{V_{C_4\eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}} \right)^\epsilon S_{\eta^i r_{i,j}}(x_{i,j}, x).$$

By (A) and (1.2), it is easy to see that the series in Φ_L has only finitely many terms. To verify that (3.11) holds in $\mathcal{G}_0^\epsilon(\beta, \gamma)$, it suffices to show that Φ_L converges to φ in $\mathcal{G}_0^\epsilon(\beta, \gamma)$ as $L \rightarrow \infty$. Notice that for any i and j , $S_{\eta^i r_{i,j}}(x_{i,j}, \cdot) \in \mathcal{G}(x_{i,j}, \eta^i r_{i,j}, \epsilon, \epsilon)$. Thus $\Phi_L \in \mathcal{G}(\epsilon, \epsilon)$ since it has only finite terms. Now it remains to show

$$\lim_{L \rightarrow \infty} \|\varphi - \Phi_L\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)} = 0.$$

To this end, we write

$$|\varphi(x) - \Phi_L(x)| \leq \left| A\sigma \sum_{i=1}^{\infty} \sum_{\{j: \eta^i r_{i,j} > L^i\}} (1-\sigma)^{i-1} \epsilon_{i,j} \frac{V_{C_4\eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}} \right)^\epsilon S_{\eta^i r_{i,j}}(x_{i,j}, x) \right|$$

$$\begin{aligned}
 & + \left| A\sigma \sum_{i=L}^{\infty} \sum_{\{j: \eta^i r_{i,j} \leq L^i\}} (1-\sigma)^{i-1} \epsilon_{i,j} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon S_{\eta^i r_{i,j}}(x_{i,j}, x) \right| \\
 & \equiv \Phi_L^1(x) + \Phi_L^2(x).
 \end{aligned}$$

Let us first prove that Φ_L^1 converges to zero in $\mathcal{G}(x_0, r_0, \beta, \gamma)$ as $L \rightarrow \infty$. Notice that $\epsilon > \gamma$ and $r_{i,j} = r_0 + d(x_{i,j}, x_0)$. By the size condition of S_t and Lemma 2.4, we obtain that for all $x \in \mathcal{X}$,

$$\begin{aligned}
 |\Phi_L^1(x)| & \lesssim \sum_{i=1}^{\infty} \sum_{\{j: \eta^i r_{i,j} > L^i\}} (1-\sigma)^{i-1} \left(\frac{r_0 \eta^i}{L^i}\right)^{\epsilon-\gamma} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\gamma S_{\eta^i r_{i,j}}(x_{i,j}, x) \\
 (3.12) \quad & \lesssim \frac{(1-\sigma)\eta^{\epsilon-\gamma}}{L^{\epsilon-\gamma} - (1-\sigma)\eta^{\epsilon-\gamma}} \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma.
 \end{aligned}$$

For any $x, y \in \mathcal{X}$ satisfying $d(x, y) \leq (r_0 + d(x_0, x))/2$, we write

$$\begin{aligned}
 & |\Phi_L^1(x) - \Phi_L^1(y)| \\
 & \leq A\sigma \sum_{i=1}^{\infty} \sum_{j \in W_i^1} (1-\sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon |S_{\eta^i r_{i,j}}(x_{i,j}, x) - S_{\eta^i r_{i,j}}(x_{i,j}, y)| \\
 & \quad + \sum_{i=1}^{\infty} \sum_{j \in W_i^2} \dots + \sum_{i=1}^{\infty} \sum_{j \in W_i^3} \dots \equiv \sum_{i=1}^{\infty} (Z_i^1 + Z_i^2 + Z_i^3),
 \end{aligned}$$

where

$$W_i^1 \equiv \{j \in \mathbb{N} : \eta^i r_{i,j} > L^i, d(x, y) \leq (\eta^i r_{i,j} + d(x, x_{i,j}))/2\},$$

$$W_i^2 \equiv \{j \in \mathbb{N} : \eta^i r_{i,j} > L^i, d(x, y) > (\eta^i r_{i,j} + d(x, x_{i,j}))/2, d(y, x_{i,j}) \geq d(x, x_{i,j})\}$$

and

$$W_i^3 \equiv \{j \in \mathbb{N} : \eta^i r_{i,j} > L^i, d(x, y) > (\eta^i r_{i,j} + d(x, x_{i,j}))/2, d(y, x_{i,j}) < d(x, x_{i,j})\}.$$

By the regularity of $S_{\eta^i r_{i,j}}$, Lemma 2.4 and $\eta^i(r_0 + d(x_{i,j}, x_0)) > L^i$, we obtain

$$\begin{aligned}
 Z_i^1 & \lesssim \sum_{j \in W_i^1} (1-\sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0 \eta^i}{L^i}\right)^{\epsilon-\gamma} \left(\frac{r_0}{r_{i,j}}\right)^\gamma \\
 & \quad \times \frac{1}{V_{\eta^i r_{i,j}}(x_{i,j}) + V_{\eta^i r_{i,j}}(x) + V(x_{i,j}, x)} \left(\frac{d(x, y)}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_1} \left(\frac{\eta^i r_{i,j}}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_2} \\
 & \lesssim \left(\frac{(1-\sigma)\eta^{\epsilon-\gamma-\epsilon_1}}{L^{\epsilon-\gamma}}\right)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x, x_0)}\right)^{\epsilon_1} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma.
 \end{aligned}$$

By the size condition of $S_{\eta^i r_{i,j}}$, $d(x, y) > (\eta^i r_{i,j} + d(x, x_{i,j}))/2$, $d(y, x_{i,j}) \geq d(x, x_{i,j})$, $r_{i,j} \geq r_0$ and Lemma 2.4,

$$\begin{aligned} Z_i^2 &\lesssim \sum_{j \in W_i^2} (1 - \sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon [S_{\eta^i r_{i,j}}(x_{i,j}, x) + S_{\eta^i r_{i,j}}(x_{i,j}, y)] \\ &\lesssim \sum_{j \in W_i^2} (1 - \sigma)^{i-1} \left(\frac{r_0 \eta^i}{L^i}\right)^{\epsilon-\gamma} \left(\frac{r_0}{r_{i,j}}\right)^\gamma \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{d(x, y)}{\eta^i r_0}\right)^{\epsilon_1} \\ &\quad \times \frac{1}{V_{\eta^i r_{i,j}}(x_{i,j}) + V_{\eta^i r_{i,j}}(x) + V(x_{i,j}, x)} \left(\frac{\eta^i r_{i,j}}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_2 + \epsilon_1} \\ &\lesssim \left(\frac{(1 - \sigma)\eta^{\epsilon-\gamma-\epsilon_1}}{L^{\epsilon-\gamma}}\right)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^{\epsilon_1} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma. \end{aligned}$$

From the assumption $d(x, y) \leq (r_0 + d(x_0, x))/2$, it follows that

(3.13)

$$r_0 + d(x_0, x) \sim r_0 + d(y, x_0) \quad \text{and} \quad V_{r_0}(x_0) + V(x_0, x) \sim V_{r_0}(x_0) + V(x_0, y),$$

which together with the definition of W_i^3 and an argument similar to the estimation of Z_i^2 yields that

$$Z_i^3 \lesssim \left(\frac{(1 - \sigma)\eta^{\epsilon-\gamma-\epsilon_1}}{L^{\epsilon-\gamma}}\right)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^{\epsilon_1} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma.$$

By the estimates of Z_i^1 , Z_i^2 and Z_i^3 , we obtain that when L is large enough and satisfies $L^{\epsilon-\gamma} > (1 - \sigma)\eta^{\epsilon-\gamma-\epsilon_1}$,

$$\begin{aligned} |\Phi_L^1(x) - \Phi_L^1(y)| &\lesssim \frac{(1 - \sigma)\eta^{\epsilon-\gamma-\epsilon_1}}{L^{\epsilon-\gamma} - (1 - \sigma)\eta^{\epsilon-\gamma-\epsilon_1}} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^{\epsilon_1} \\ &\quad \times \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma, \end{aligned}$$

which together with (3.12) implies that $\lim_{L \rightarrow \infty} \|\Phi_L^1\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)} = 0$.

Now we consider $\|\Phi_L^2\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)}$. By the size condition of $S_{\eta^i r_{i,j}}$ and Lemma 2.4, we obtain that for all $x \in \mathcal{X}$,

$$\begin{aligned} |\Phi_L^2(x)| &\lesssim \sum_{i=L}^\infty \sum_j (1 - \sigma)^{i-1} \frac{1}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\gamma \\ &\quad \times \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{\eta^i r_{i,j}}(x_{i,j}) + V_{\eta^i r_{i,j}}(x) + V(x_{i,j}, x)} \left(\frac{\eta^i r_{i,j}}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_2} \\ (3.14) \quad &\lesssim (1 - \sigma)^L \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma. \end{aligned}$$

For any $x, y \in \mathcal{X}$ satisfying $d(x, y) \leq (r_0 + d(x_0, x))/2$, we write

$$\begin{aligned} & |\Phi_L^2(x) - \Phi_L^2(y)| \\ & \leq A\sigma \sum_{i=L}^{\infty} \sum_{j \in W_i^4} (1 - \sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon |S_{\eta^i r_{i,j}}(x_{i,j}, x) - S_{\eta^i r_{i,j}}(x_{i,j}, y)| \\ & \quad + \sum_{i=L}^{\infty} \sum_{j \in W_i^5} \dots + \sum_{i=L}^{\infty} \sum_{j \in W_i^6} \dots \equiv \sum_{i=L}^{\infty} (Z_i^4 + Z_i^5 + Z_i^6), \end{aligned}$$

where

$$W_i^4 \equiv \{j \in \mathbb{N} : d(x, y) \leq (\eta^i r_{i,j} + d(x, x_{i,j}))/2\},$$

$$W_i^5 \equiv \{j \in \mathbb{N} : d(x, y) > (\eta^i r_{i,j} + d(x, x_{i,j}))/2, d(y, x_{i,j}) \geq d(x, x_{i,j})\}$$

and

$$W_i^6 \equiv \{j \in \mathbb{N} : d(x, y) > (\eta^i r_{i,j} + d(x, x_{i,j}))/2, d(y, x_{i,j}) < d(x, x_{i,j})\}.$$

For any $i \in \mathbb{N}$ and $k = 4, 5, 6$, by $d(x, y) \leq (r_0 + d(x_0, x))/2$, (3.2), (3.3) and (3.13),

$$(3.15) \quad Z_i^k \leq |\omega_i(x)| + |\omega_i(y)| \lesssim (1 - \sigma)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon.$$

The regularity of $S_{\eta^i r_{i,j}}$ and Lemma 2.4 show that

$$\begin{aligned} (3.16) \quad Z_i^4 & \lesssim \sum_{j \in W_i^4} (1 - \sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon \left(\frac{d(x, y)}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_1} \\ & \quad \times \frac{1}{V_{\eta^i r_{i,j}}(x_{i,j}) + V_{\eta^i r_{i,j}}(x) + V(x_{i,j}, x)} \left(\frac{\eta^i r_{i,j}}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_2} \\ & \lesssim \left(\frac{1 - \sigma}{\eta^{\epsilon_1}}\right)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^{\epsilon_1} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon. \end{aligned}$$

For any $j \in W_i^5$, we have $d(y, x_{i,j}) \geq d(x, x_{i,j})$. From this, the size condition of $S_{\eta^i r_{i,j}}$ and Lemma 2.4, we deduce that

$$\begin{aligned} (3.17) \quad Z_i^5 & \lesssim \sum_{j \in W_i^5} (1 - \sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon \left(\frac{d(x, y)}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_1} \\ & \quad \times \frac{1}{V_{\eta^i r_{i,j}}(x_{i,j}) + V_{\eta^i r_{i,j}}(x) + V(x_{i,j}, x)} \left(\frac{\eta^i r_{i,j}}{\eta^i r_{i,j} + d(x_{i,j}, x)}\right)^{\epsilon_2} \\ & \lesssim \left(\frac{1 - \sigma}{\eta^{\epsilon_1}}\right)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^{\epsilon_1} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon, \end{aligned}$$

By an argument similar to the estimate of Z_i^3 , we obtain

$$(3.18) \quad Z_i^6 \lesssim \left(\frac{1-\sigma}{\eta^{\epsilon_1}}\right)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^{\epsilon_1} \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\epsilon.$$

The geometric means of (3.16) and (3.15), (3.17) and (3.15), (3.18) and (3.15), respectively, give that for all $i \in \mathbb{N}$ and $k = 4, 5, 6$,

$$Z_i^k \lesssim \left(\frac{1-\sigma}{\eta^\beta}\right)^i \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^\beta \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma.$$

Using this and the assumption $\eta^\beta > 1 - \sigma$, we obtain, for all $x, y \in \mathcal{X}$ satisfying $d(x, y) \leq (r_0 + d(x_0, x))/2$, the following estimate:

$$|\Phi_L^2(x) - \Phi_L^2(y)| \lesssim \left(\frac{1-\sigma}{\eta^\beta}\right)^L \frac{1}{V_{r_0}(x_0) + V(x_0, x)} \left(\frac{d(x, y)}{r_0 + d(x_0, x)}\right)^\beta \left(\frac{r_0}{r_0 + d(x_0, x)}\right)^\gamma.$$

This together with (3.14) yields that $\lim_{L \rightarrow \infty} \|\Phi_L^2\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)} = 0$. Therefore, we obtain $\lim_{L \rightarrow \infty} \|\Phi_L\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)} = 0$ and thus $\lim_{L \rightarrow \infty} \|\Phi_L\|_{\mathcal{G}(x_1, 1, \beta, \gamma)} = 0$, which implies that (3.11) holds in $\mathcal{G}_0^\epsilon(\beta, \gamma)$.

Set

$$\nu \equiv \sum_{i=1}^\infty \sum_{j=1}^{j(i)} (1-\sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon \delta_{(x_{i,j}, \eta^i r_{i,j})},$$

where $\delta_{(x,r)}$ is the Dirac measure of point $(x, r) \in \mathcal{X} \times \mathbb{R}_+$. Notice that for any given i and j , by Property (B) above, we have

$$|\langle S_{\eta^i r_{i,j}}(x_{i,j}, \cdot), f \rangle| \leq \mathcal{M}_0(f)(x_{i,j}) \lesssim [F(\eta^i r_{i,j}, x_{i,j}, [\mathcal{M}_0(f)]^{1/2} \chi_{B_{i,j}})]^2.$$

Then, by this and (3.11),

$$\begin{aligned} |\langle f, \varphi \rangle| &\lesssim \sum_{i=1}^\infty \sum_{j=1}^{j(i)} (1-\sigma)^{i-1} \frac{V_{C_4 \eta^i r_{i,j}}(x_{i,j})}{V_{r_{i,j}}(x_{i,j})} \left(\frac{r_0}{r_{i,j}}\right)^\epsilon [F(\eta^i r_{i,j}, x_{i,j}, [\mathcal{M}_0(f)]^{1/2} \chi_{B_{i,j}})]^2 \\ &= \int_{\mathcal{X} \times \mathbb{R}_+} [F(r, x, [\mathcal{M}_0(f)]^{1/2} \chi_{B(x, C_4 r)})]^2 d\nu(x, r). \end{aligned}$$

Denote by $\lfloor \log_2 r_0 \rfloor$ the largest integer no more than $\log_2 r_0$. Since $r_{i,j} \geq r_0$, we then have

$$(3.19) \quad \begin{aligned} \nu &\lesssim (r_0)^\epsilon \sum_{t=\lfloor \log_2 r_0 \rfloor + 1}^\infty 2^{-t\epsilon} V_{2^t}(x_0)^{-1} \sum_{\{i, j: 2^{t-1} \leq r_{i,j} < 2^t\}} (1-\sigma)^i V_{C_4 \eta^i r_{i,j}}(x_{i,j}) \delta_{(x_{i,j}, \eta^i r_{i,j})} \\ &\equiv (r_0)^\epsilon \sum_{t=\lfloor \log_2 r_0 \rfloor + 1}^\infty 2^{-t\epsilon} \nu_t. \end{aligned}$$

If $2^{t-1} \leq r_{i,j} < 2^t$, then for any $z \in B(x_{i,j}, C_4\eta^i r_{i,j})$, we have

$$d(z, x_0) \leq d(z, x_{i,j}) + d(x_{i,j}, x_0) \leq 2r_{i,j} < 2^{t+1},$$

which implies that $B(x_{i,j}, C_4\eta^i r_{i,j}) \subset B(x_0, 2^{t+1})$. From this, it follows that for any $(x, r) \in \text{supp } \nu_t$,

$$(3.20) \quad F(r, x, [\mathcal{M}_0(f)]^{1/2} \chi_{B(x, C_4 r)}) \leq F(r, x, [\mathcal{M}_0(f)]^{1/2} \chi_{B(x_0, 2^{t+1})}).$$

By (3.19) and (3.20),

$$(3.21)$$

$$|\langle f, \varphi \rangle| \lesssim (r_0)^\epsilon \sum_{t=\lfloor \log_2 r_0 \rfloor}^\infty 2^{-t\epsilon} \int_{\mathcal{X} \times \mathbb{R}_+} \left[F(r, x, [\mathcal{M}_0(f)]^{1/2} \chi_{B(x, 2^{t+1})}) \right]^2 d\nu_t(x, r).$$

We claim that for any fixed $\delta_0 \in (0, \log_\eta(1-\sigma))$, there exists a positive constant C_{δ_0} independent of t such that for all $x \in \mathcal{X}$ and $r > 0$,

$$(3.22) \quad \nu_t(B(x, r) \times (0, r)) \leq C_{\delta_0} \left(\frac{V_r(x)}{V_{2^t}(x_0)} \right)^{(n+\delta_0)/n}.$$

Assume this claim for the moment. By this claim, (3.21) and Lemma 2.2, we have

$$\begin{aligned} |\langle f, \varphi \rangle| &\lesssim (r_0)^\epsilon \sum_{t=\lfloor \log_2 r_0 \rfloor + 1}^\infty 2^{-t\epsilon} \left\{ \frac{1}{V_{2^t}(x_0)} \int_{B(x_0, C_4 2^{t+1})} [\mathcal{M}_0(f)(x)]^{n/(n+\delta_0)} d\mu(x) \right\}^{(n+\delta_0)/n} \\ &\lesssim \left[\mathcal{M}((\mathcal{M}_0(f))^{n/(n+\delta_0)})(x_0) \right]^{(n+\delta_0)/n}, \end{aligned}$$

where in the last step, we use $(r_0)^\epsilon \sum_{t=\lfloor \log_2 r_0 \rfloor + 1}^\infty 2^{-t\epsilon} \lesssim 1$. Thus the desired conclusion of Proposition 1.7 holds.

To finish the proof of Proposition 1.7, we still need to verify the validity of (3.22). For any $x \in \mathcal{X}$ and $r > 0$, set

$$W \equiv \{(i, j) : 2^{t-1} \leq r_{i,j} < 2^t, \eta^i r_{i,j} < r, d(x, x_{i,j}) < r\}.$$

For any $(i, j) \in W$, since $C_4 < 1$, we have $B(x_{i,j}, C_4\eta^i r_{i,j}) \subset B(x, 2r)$, which further implies that

$$(3.23) \quad \sum_{\{j: (i, j) \in W\}} V_{C_4\eta^i r_{i,j}}(x_{i,j}) \lesssim V_r(x).$$

If $(i, j) \in W$, then $2^t < 2r_{i,j} < 2r\eta^{-i}$. From this, we deduce that for any $z \in B(x_0, 2^t)$,

$$d(z, x) \leq d(z, x_0) + d(x_0, x_{i,j}) + d(x_{i,j}, x) \leq 2^t + 2^t + r \leq (4\eta^{-i} + 1)r,$$

and thus $B(x_0, 2^t) \subset B(x, (4\eta^{-i} + 1)r)$. Moreover, by (1.2),

$$(3.24) \quad V_{2^t}(x_0) \lesssim (4\eta^{-i} + 1)^n V_r(x) \lesssim \eta^{-in} V_r(x).$$

Combining (3.19), (3.23), (3.24) and the assumption $\delta_0 < \log_\eta(1 - \sigma)$ yields that

$$\begin{aligned} & \nu_t(B(x, r) \times (0, r)) \\ &= [V_{2^t}(x_0)]^{-1} \int_{B(x,r) \times (0,r)} \sum_{\{i, j: 2^{t-1} \leq r_{i,j} < 2^t\}} (1 - \sigma)^i V_{C_4 \eta^i r_{i,j}}(x_{i,j}) d\delta_{(x_{i,j}, \eta^i r_{i,j})} \\ &\lesssim [V_{2^t}(x_0)]^{-1} \sum_{(i, j) \in W} (1 - \sigma)^i V_{C_4 \eta^i r_{i,j}}(x_{i,j}) \left(\frac{\eta^{-in} V_r(x)}{V_{2^t}(x_0)} \right)^{\delta_0/n} \\ &\lesssim \left(\frac{V_r(x)}{V_{2^t}(x_0)} \right)^{(n+\delta_0)/n} \sum_{i \in \mathbb{N}} \left(\frac{1 - \sigma}{\eta^{\delta_0}} \right)^i \lesssim \left(\frac{V_r(x)}{V_{2^t}(x_0)} \right)^{(n+\delta_0)/n}, \end{aligned}$$

which implies the claim, and hence completes the proof of Proposition 1.7. \square

REMARK 3.1. — *By the proof of Theorem 1.6, we need the assumption $p > n/(n + \delta_0)$, while in the proof of Proposition 1.7, we also need $\delta_0 < \log_\eta(1 - \sigma)$. Thus, Theorem 1.6 holds for any $p \in (n/(n + \log_\eta(1 - \sigma)), \infty]$. In fact, Theorem 1.6 and Proposition 1.7 hold for all $\sigma > 0$ and $\eta > 0$, if $\sigma, \eta, A > 0$ in (3.1) and $H > 0$ in (3.5) satisfy the following five inequalities: $H \leq 1/3$; $C_4 \eta < H$; $\eta^{\epsilon_1} < 1 - \sigma < \eta^\beta$; $\frac{A}{C_8 2^\epsilon} - AC_3 C_9 \max\{\eta^{\epsilon_2}, (1 + H\eta^{-1})^{-\epsilon_2}\} \geq 1$; and*

$$\left(\frac{H}{1 - H} \right)^\epsilon + \left(\frac{H}{1 - H} \right)^{\epsilon_1} \frac{AC_3 C_9 \sigma}{1 - \sigma - \eta^{\epsilon_1}} \leq 1 - \sigma - AC_3 C_9 \sigma.$$

Acknowledgments. — The authors are greatly indebted to the referees for their very valuable remarks which made this article more readable.

BIBLIOGRAPHY

- [1] G. ALEXOPOULOS – “Spectral multipliers on Lie groups of polynomial growth”, *Proc. Amer. Math. Soc.* **120** (1994), p. 973–979.
- [2] M. CHRIST – “A $T(b)$ theorem with remarks on analytic capacity and the Cauchy integral”, *Colloq. Math.* **60/61** (1990), p. 601–628.
- [3] R. R. COIFMAN & G. WEISS – *Analyse harmonique non-commutative sur certains espaces homogènes*, Lecture Notes in Math., Vol. 242, Springer, 1971.
- [4] ———, “Extensions of Hardy spaces and their use in analysis”, *Bull. Amer. Math. Soc.* **83** (1977), p. 569–645.
- [5] D. DANIELLI, N. GAROFALO & D.-M. NHIEU – “Non-doubling Ahlfors measures, perimeter measures, and the characterization of the trace spaces of Sobolev functions in Carnot-Carathéodory spaces”, *Mem. Amer. Math. Soc.* **182** (2006), p. 119.

- [6] X. T. DUONG & L. YAN – “Hardy spaces of spaces of homogeneous type”, *Proc. Amer. Math. Soc.* **131** (2003), p. 3181–3189.
- [7] C. FEFFERMAN & E. M. STEIN – “ H^p spaces of several variables”, *Acta Math.* **129** (1972), p. 137–193.
- [8] L. GRAFAKOS – *Classical Fourier analysis*, second ed., Graduate Texts in Math., vol. 249, Springer, 2008.
- [9] L. GRAFAKOS, L. LIU & D. YANG – “Maximal function characterizations of Hardy spaces on RD-spaces and their applications”, *Sci. China (Ser. A)* **51** (2008), p. 2253–2284.
- [10] Y. HAN – “Triebel-Lizorkin spaces on spaces of homogeneous type”, *Studia Math.* **108** (1994), p. 247–273.
- [11] Y. HAN, D. MÜLLER & D. YANG – “Littlewood-Paley characterizations for Hardy spaces on spaces of homogeneous type”, *Math. Nachr.* **279** (2006), p. 1505–1537.
- [12] ———, “A theory of Besov and Triebel-Lizorkin spaces on metric measure spaces modeled on Carnot-Carathéodory spaces”, to appear in *Abstr. Appl. Anal.*, Art. ID 893409, 2008.
- [13] J. HEINONEN – *Lectures on analysis on metric spaces*, Universitext, Springer, 2001.
- [14] R. A. MACÍAS & C. SEGOVIA – “A decomposition into atoms of distributions on spaces of homogeneous type”, *Adv. in Math.* **33** (1979), p. 271–309.
- [15] D. MÜLLER & D. YANG – “A difference characterization of Besov and Triebel-Lizorkin spaces on RD-spaces”, to appear in *Forum Math.*
- [16] A. NAGEL & E. M. STEIN – “Differentiable control metrics and scaled bump functions”, *J. Differential Geom.* **57** (2001), p. 465–492.
- [17] ———, “On the product theory of singular integrals”, *Rev. Mat. Iberoamericana* **20** (2004), p. 531–561.
- [18] A. NAGEL, E. M. STEIN & S. WAINGER – “Balls and metrics defined by vector fields. I. Basic properties”, *Acta Math.* **155** (1985), p. 103–147.
- [19] E. M. STEIN – *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, 1993.
- [20] ———, “Some geometrical concepts arising in harmonic analysis”, *Geom. Funct. Anal.*, Special Volume, Part I (2000), p. 434–453, GAFA 2000 (Tel Aviv, 1999).
- [21] E. M. STEIN & G. WEISS – “On the theory of harmonic functions of several variables. I. The theory of H^p -spaces”, *Acta Math.* **103** (1960), p. 25–62.
- [22] A. UCHIYAMA – “A maximal function characterization of H^p on the space of homogeneous type”, *Trans. Amer. Math. Soc.* **262** (1980), p. 579–592.

- [23] N. T. VAROPOULOS – “Analysis on Lie groups”, *J. Funct. Anal.* **76** (1988), p. 346–410.
- [24] N. T. VAROPOULOS, L. SALOFF-COSTE & T. COULHON – *Analysis and geometry on groups*, Cambridge Tracts in Mathematics, vol. 100, Cambridge University Press, 1992.