



ANNALES

DE

L'INSTITUT FOURIER

Xianghong GONG

Levi-flat invariant sets of holomorphic symplectic mappings

Tome 51, n° 1 (2001), p. 151-208.

http://aif.cedram.org/item?id=AIF_2001__51_1_151_0

© Association des Annales de l'institut Fourier, 2001, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

LEVI-FLAT INVARIANT SETS OF HOLOMORPHIC SYMPLECTIC MAPPINGS

by Xianghong GONG

1. Introduction.

Let φ be a biholomorphic mapping defined near the origin of \mathbb{C}^{2n} with $\varphi(0) = 0$. Recall that φ is *symplectic* if it preserves the holomorphic symplectic 2-form $\sum_{j=1}^n d\xi_j \wedge d\eta_j$. Assume that the linear part of φ is diagonalizable. In suitable local holomorphic symplectic coordinates, φ is then given by

$$(1.1) \quad \xi_j \rightarrow \lambda_j \xi_j + u_j(\xi, \eta), \quad \eta_j \rightarrow \lambda_j^{-1} \eta_j + v_j(\xi, \eta), \quad j = 1, \dots, n,$$

where u_j, v_j are holomorphic functions starting with terms of order at least two. One says that $\lambda = (\lambda_1, \dots, \lambda_n)$ is *non-resonant*, if

$$(1.2) \quad \lambda_1^{\alpha_1} \lambda_2^{\alpha_2} \dots \lambda_n^{\alpha_n} \neq 1$$

for all integers $\alpha_1, \dots, \alpha_n$ with $|\alpha_1| + \dots + |\alpha_n| > 0$. G.D. Birkhoff [4] showed that under formal symplectic transformations, φ is equivalent to the formal symplectic mapping

$$(1.3) \quad \xi_j \rightarrow \lambda_j \xi_j e^{\omega_j}, \quad \eta_j \rightarrow \lambda_j^{-1} \eta_j e^{-\omega_j}, \quad \omega_j = \partial_{\zeta_j} H$$

Supported in part by NSF grants DMS-9704835 and DMS-0096047.

Keywords: Levi-flat set – Segre variety – Holomorphic symplectic map – Birkhoff normal form.

Math. classification: 37G05 – 32V40 – 70H06.

with H a formal power series in the products $\zeta_1 = \xi_1\eta_1, \dots, \zeta_n = \xi_n\eta_n$ without constant term.

Analogous to work of C.L. Siegel [17] on Hamiltonian systems, a result of H. Rüssmann [12] says that as a rule the Birkhoff normal form (1.3) of φ is not realizable by convergent symplectic transformations. In fact, one aspect of divergence for Birkhoff's normalization for Hamiltonian systems was understood by Siegel [16] much earlier. Recall that holomorphic functions invariant under a holomorphic symplectic mapping or the flow of a holomorphic Hamiltonian system are called *first-integrals*. Siegel [16] showed that there are Hamiltonian systems of degree of freedom larger than one that admit no first-integrals other than functions of their Hamiltonians. On the other hand, when admitting a certain system of first-integrals, a holomorphic symplectic mapping or Hamiltonian system can be transformed into the Birkhoff normal form by convergent transformations. This is the so-called integrability of Hamiltonian systems (or symplectic mappings) via first-integrals, which was already observed by Birkhoff [3] for the case of one degree of freedom. In general cases, such integrability results were obtained by Rüssmann [13] for analytic Hamiltonian systems with two degrees of freedom and by J. Vey [18] for analytic Hamiltonian systems and symplectic mappings of any degrees of freedom, for which the quadratic forms of the first-integrals satisfy a certain non-degeneracy condition. Most recently H. Ito [10] studied the convergence of Birkhoff's normalization for analytic Hamiltonian systems and symplectic mappings which admit certain first-integrals with degenerate quadratic forms. The reader is also referred to results of L.H. Eliasson [9] on smooth real Hamiltonian systems. We should mention that the results in [13], [18], and [10], based on the first-integral method, hold for both real and complex cases.

In case the holomorphic symplectic map is in its normal form (1.3), it is clear that for each j the quantity $\xi_j\eta_j$ is invariant under the map. In particular the set defined by $\Re\xi_j\eta_j = 0$ ($j = 1, \dots, n$) is invariant. More generally, if H_j are holomorphic functions of n variables, the set defined by $\Re\{H_j(\xi\eta)\} = 0$ is invariant.

In Vey's theorem and in Ito's one has of course the existence of a (singular) foliation of \mathbb{C}^{2n} by (n -codimensional) invariant sets. In the present work, we show the convergence of (the reduction to) the normal form under the sole hypothesis of just one appropriate invariant real analytic set containing the origin. The invariant set that we assume to exist is of co-dimension n in \mathbb{C}^{2n} , and Levi-flat, as above. A generic n -dimensional

real analytic set in \mathbb{C}^{2n} is said to be *Levi-flat* if its regular set (away from singular points and *CR* singularities) is foliated by n -dimensional complex manifolds. The main result of this paper is the convergence of the normalization under the assumption of the existence of an invariant set in two special cases (see below): (1.4) and (1.5).

We now state the following.

THEOREM 1.1. — *Let φ be a holomorphic symplectic mapping of \mathbb{C}^{2n} given by (1.1)–(1.2), and let $M \subset \mathbb{C}^{2n}$ be the real analytic set*

$$(1.4) \quad \Re \left\{ \sum a_{jk} \xi_k \eta_k \right\} + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, n,$$

where $\text{rank}(a_{jk}) = n$ and $R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = O(3)$ are real-valued analytic functions. Assume that M is Levi-flat and $\varphi(M) = M$. Then φ can be transformed into the Birkhoff normal form (1.3) by convergent symplectic transformations.

It turns out that the integrability of φ in terms of Levi-flat invariant sets of the form (1.4) is well connected to its first-integrals; namely, we shall prove that φ has n first-integrals if it has a Levi-flat invariant set (1.4). Thus Theorem 1.1 follows eventually from the above-mentioned result of Vey. Our next result is the following.

THEOREM 1.2. — *Let φ, M be as in Theorem 1.1 except that M is given by*

$$(1.5) \quad |\xi_j|^2 - |\eta_{\sigma(j)}|^2 + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, n,$$

where σ is a permutation of $1, \dots, n$ and $R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = O(3)$ are real-valued analytic functions. Assume that $\varphi(M) = M$. Then $|\lambda_{\sigma(j)}| \cdot |\lambda_j| = 1$ for $j = 1, \dots, n$. Moreover, φ has n formal first-integrals $\delta_j \kappa$ such that φ can be transformed into (1.3) by convergent symplectic transformations, provided $\sigma^2 = \text{Id}$ and all $\delta_j \kappa \equiv 1$.

For the definition of $\delta_j \kappa$ ($\delta_j \kappa \equiv 1$ if $\sigma = \text{Id}$), see (4.37) below.

In contrary to case (1.4), the Levi-flat invariant set (1.5), however, leads to meromorphic eigenfunctions of the holomorphic symplectic mappings. We shall show that such a mapping φ has n meromorphic functions m_j satisfying $m_j \circ \varphi = \lambda_j \lambda_{\sigma(j)} m_j$. An example (end of Section 2) shows that it is not enough to assume the existence of an invariant Levi-flat n codimensional set in \mathbb{C}^{2n} for $n \geq 2$, but in case of (1.5) we have no counterexample if one drops the restrictive hypotheses on the permutation σ

and on $\delta_j\kappa$. It also remains open if the Birkhoff normalization converges for the holomorphic symplectic mappings of \mathbb{C}^2 that have a Levi-flat real hypersurface. Only for a special case shall we prove the following.

THEOREM 1.3. — *Let φ be a holomorphic symplectic mapping of \mathbb{C}^2 given by (1.1)–(1.2). Let $M \subset \mathbb{C}^2: r = 0$ be a real analytic Levi-flat hypersurface invariant under φ . Assume that the quadratic form of r defines a real hypersurface in \mathbb{C}^2 . Then φ can be transformed into its Birkhoff normal form by convergent symplectic transformations.*

Note that Theorem 3 is meant for the case $|\lambda| = 1$, since Moser [11] already showed that a holomorphic symplectic mapping of \mathbb{C}^2 is always normalizable by convergent transformations if $|\lambda| \neq 1$.

The present paper relies on some techniques used in very recent joint work of D. Burns and the author [6], where singular Levi-flat real analytic hypersurfaces of \mathbb{C}^n are studied. These techniques will allow us to construct holomorphic first-integrals or meromorphic eigenfunctions for the holomorphic symplectic mappings. Motivated by applications to holomorphic symplectic mappings, we shall also extend some results in [6] to Levi-flat sets of higher codimensions. One of main ingredients used in [6] is that of Segre varieties, which is a family of invariant complex varieties associated to a real analytic hypersurface in complex space [15]. In dealing with singular Levi-flat structure one encounters with difficulties of constructing formal normal forms. In [6] and in this work, the Segre varieties serve as an essential tool to avoid the formal normal forms of Levi-flat sets. As another application of Segre varieties we shall also use Segre varieties to prove part of Theorem 1.3 directly (see Proposition 4.2). The reader is also referred to work of E. Bedford [2] on the domain of holomorphy of complements of singular Levi-flat hypersurfaces. For other applications of Segre varieties, see work of S.M. Webster [19] on algebraic real hypersurfaces with non-degenerate Levi-form and work of Diederich-Fornaess [8] on complex varieties in real analytic sets.

The paper is organized as follows. Sections 2 and 3 consist in a preliminary study of Levi-flat invariant sets of codimension n . In Section 2, we study those Levi-flat sets defined by real quadratic forms. Their classification happens to be already non-trivial, and in fact we succeed only in classifying Levi-flat sets of special types. In Section 3, the Levi-flat sets that have been studied serve as models for more general Levi-flat sets. At the expense of having the hypotheses looking more artificial, the reader can

read Section 3 without having read Section 2. Also, the results but not the proofs in Section 3 are to be used in Section 4.

In Section 4, the simplified equations that we found for the Levi-flat invariant sets allow one to study the defining functions of those sets, i.e., functions whose at least the common zero level set is invariant. From this study one is able to get either enough first-integrals, or meromorphic eigenfunctions.

In case enough first-integrals are obtained, the convergence of normal forms follows from Vey's theorem. This however does not establish Theorem 1.2, which correspond to a case for which in Section 4 one obtains only meromorphic eigenfunctions. This latter case is treated in Section 5, using a KAM method.

In Section 6 we shall formulate analogous results for holomorphic Hamiltonian systems, based on an observation regarding normal forms of the time-one mappings of Hamiltonian systems. The paper is concluded with an appendix about two equivalent Birkhoff normal forms of holomorphic symplectic mappings.

Acknowledgment. The author is indebted to Dan Burns for the insights in Segre varieties through the collaboration [6]. The author would also like to thank Jean-Pierre Rosay for helpful suggestions.

2. Invariant Levi-flat sets defined by real quadratic forms.

In this section we shall study certain classes of Levi-flat sets of high codimension in \mathbb{C}^{2n} defined by real quadratic forms. Here the classification is not complete. In fact, we shall only consider 4 families of Levi-flat sets of high codimension under an additional condition that the sets are invariant under an elliptic complex linear symplectic transformation. A complete classification of Levi-flat quadratic hypersurfaces of \mathbb{C}^n is in [6].

Recall that a germ M of real analytic set of dimension k at $0 \in \mathbb{C}^n$ admits a decomposition $M = M^* \cup M_s$, where M^* consists of points x near which M is a k -dimensional submanifold of \mathbb{C}^n . By a result of Bruhat and Cartan [5], if M is irreducible, M^* is the union of a finite number of topological components M_j^* with the origin of \mathbb{C}^n in the closure of each M_j^* . We define the *CR* dimension of M to be

$$\liminf_{M^* \ni p \rightarrow 0} \dim_{\mathbb{C}} T_p M \cap \sqrt{-1} T_p M \equiv \dim_{CR} M.$$

Let M_{cs} be the set of points $x \in M^*$ at which the complex dimension of $T_x M \cap \sqrt{-1}T_x M$ is larger than $\dim_{CR} M$. Then M_{cs} is a real analytic subset of M^* . We say that M is *Levi-flat*, if $M_c \equiv M^* \setminus M_{cs}$ is foliated by complex submanifolds of dimension $\dim_{CR} M$. Notice that we allow M_{cs} to contain some components of M^* . In general, one has $\dim_{CR} M \geq \max\{\dim M - n, 0\}$. We say that M is *generic*, or of *minimal CR dimension*, when $\dim_{CR} M = \max\{\dim M - n, 0\}$, i.e., when $T_x M_c + \sqrt{-1}T_x M_c = \mathbb{C}^n$ for all $x \in M_c$, or $T_x M_c \cap \sqrt{-1}T_x M_c = 0$ for all $x \in M_c$. In the latter case M_c is said to be *totally real*. We shall also use a standard fact that the real analytic submanifold M_c is Levi-flat, if and only if the Lie brackets $[L_i, \bar{L}_j]$ remain in the span of $L_1, \dots, L_m, \bar{L}_1, \dots, \bar{L}_m$ for any local basis L_1, \dots, L_m ($m = \dim_{CR} M$) of $(1, 0)$ tangent vectors of M_c , i.e., tangent vectors of M_c of the form $\sum_{j=1}^n a_j(z, \bar{z}) \frac{\partial}{\partial z_j}$ with a_j being analytic.

We shall need some basic properties about the complexification of real analytic varieties. Recall that a germ M of real analytic variety of \mathbb{R}^n at 0 is contained in a unique germ M^c of complex variety at $0 \in \mathbb{C}^n$ such that $M = M^c \cap \mathbb{R}^n$, and such that germs of holomorphic function at $0 \in \mathbb{C}^n$ vanishing on M^c are precisely the germs of complex-valued real analytic function at $0 \in \mathbb{R}^n$ that vanish on M . We shall refer the germ M^c as the Bruhat-Whitney-Cartan *complexification* of M . A result of H. Cartan [7] says that M is irreducible if and only if M^c is irreducible; consequently, a germ of real analytic function vanishing on an open subset of M^* , of which the closure contains the origin, vanishes entirely on the germ M .

The main purpose of this section is to study Levi-flat sets that are defined by quadratic forms and invariant under the \mathbb{C} -linear symplectic mapping

$$(2.1) \quad \Lambda: \xi_j \rightarrow \lambda_j \xi_j, \quad \eta_j \rightarrow \lambda_j^{-1} \eta_j, \quad j = 1, \dots, n,$$

where $\lambda_1, \dots, \lambda_n$ satisfy the non-resonance condition (1.2). Throughout the section, we assume that Λ is *elliptic*, that is that $|\lambda_j| = 1, j = 1, \dots, n$.

A complex-valued quadratic form on \mathbb{C}^{2n} is a quadratic homogeneous polynomial in $\xi, \eta, \bar{\xi}, \bar{\eta}$; the space of all complex-valued quadratic forms will be denoted by E . Let E^{psh} be the set of complex-valued quadratic forms containing no terms of the form $\xi^\alpha \eta^\beta \bar{\xi}^\gamma \bar{\eta}^\delta$ with $|\alpha| + |\beta| = 1$, and E^{hrrm} the set of quadratic forms containing only the terms $\xi^\alpha \eta^\beta \bar{\xi}^\gamma \bar{\eta}^\delta$ with $|\alpha| + |\beta| = 1$. Sending f to $f \circ \Lambda$, the transformation $\Lambda: E \rightarrow E$ is \mathbb{C} -linear. Put

$$\lambda_{-j} = \bar{\lambda}_j, \quad \xi_{-j} = \bar{\xi}_j, \quad \eta_{-j} = \bar{\eta}_j.$$

Notice that (2.1) holds for $j = \pm 1, \dots, \pm n$. The eigenvalues of the linear transformation $\Lambda: E \rightarrow E$ are 1 and $\lambda_i \lambda_j$ ($i + j \neq 0$), to which the corresponding eigen-spaces are

$$E_1 = \bigoplus_{i=1}^n \mathbb{C} \cdot \xi_i \eta_i \oplus \mathbb{C} \cdot \bar{\xi}_i \bar{\eta}_i \oplus \mathbb{C} \cdot \xi_i \bar{\xi}_i \oplus \mathbb{C} \cdot \eta_i \bar{\eta}_i,$$

$$E_{ij} = \mathbb{C} \cdot \xi_i \xi_j \oplus \mathbb{C} \cdot \bar{\eta}_i \bar{\eta}_j \oplus \mathbb{C} \cdot \xi_i \bar{\eta}_j \oplus \mathbb{C} \cdot \bar{\eta}_i \xi_j$$

with

$$i, j = \pm 1, \dots, \pm n, \quad j \geq i, \quad i + j \neq 0.$$

Let \mathcal{Q} be the vanishing set of finitely many real quadratic forms on \mathbb{C}^{2n} , and let $\mathcal{I} = \mathcal{I}_{\mathcal{Q}}$ be the set of complex-valued quadratic forms vanishing on \mathcal{Q} . Assume that \mathcal{Q} is invariant under Λ , so \mathcal{I} is invariant under Λ also. From the theory of linear algebra one knows that

$$\mathcal{I} = \mathcal{I} \cap E_1 \oplus \bigoplus \mathcal{I} \cap E_{ij} \equiv \mathcal{I} \cap E_1 \oplus \mathcal{I} \cap E_0.$$

We shall only classify the Levi-flat sets \mathcal{Q} for which \mathcal{I} is contained in one of the subspaces

$$E_1 \cap E^{psh}, \quad E_1 \cap E^{hrm}, \quad E_0 \cap E^{psh}, \quad E_0 \cap E^{hrm}.$$

We start with the following.

PROPOSITION 2.1. — *Let $\mathcal{Q} \subset \mathbb{C}^{2n}$ be a generic real analytic set defined by real quadratic forms. Assume that \mathcal{Q} has codimension $m \leq 2n$ and $\mathcal{I}_{\mathcal{Q}}$ is contained in $E^{psh} \cap E_1$. Then there exists an $m \times n$ matrix (a_{ij}) of rank m such that*

$$(2.2) \quad \mathcal{Q}: \Re \left\{ \sum a_{ij} \xi_j \eta_j \right\} = 0, \quad i = 1, \dots, m.$$

Conversely, (2.2) is a Levi-flat real analytic set in \mathbb{C}^{2n} of minimal CR dimension $2n - m$, provided the rank of (a_{ij}) is m .

Proof. — Take k quadratic forms $q_i(\xi\eta) = \sum_{j=1}^n a_{ij} \xi_j \eta_j$ so that $\Re q_1, \dots, \Re q_k$ form a basis for real quadratic forms vanishing on \mathcal{Q} . Since \mathcal{Q} is generic and of codimension at most $2n$, then \mathcal{Q} is not contained in any proper complex subvariety of \mathbb{C}^{2n} . This implies that q_1, \dots, q_k are linearly independent over \mathbb{C} . For the proof, we assume for the sake of contradiction that $q_k = c_1 q_1 + \dots + c_k q_{k-1}$. Then

$$\sum_{j=1}^{k-1} \Im c_j \cdot \Im q_j = -\Re q_k + \sum_{j=1}^{k-1} \Re c_j \Re q_j$$

vanishes on \mathcal{Q} . Hence, $\sum_{j=1}^{k-1} \Im c_j \cdot q_j$ vanishes on \mathcal{Q} . Since $\Re q_1, \dots, \Re q_k$ are linearly independent over \mathbb{R} , then at least one of $\Im c_j$ is nonzero. Therefore, \mathcal{Q} is contained in the complex variety $\sum_{j=1}^{k-1} \Im c_j \cdot q_j = 0$, which is a contradiction.

To show that $k = m$, we need only to verify the last statement in the proposition. Without loss of generality, one may assume that $(a_{ij})_{1 \leq i, j \leq m}$ is nonsingular. Then $\mathcal{Q} \setminus \cup_{j=1}^m \{\xi_j = 0\}$ is parameterized by

$$\eta_i = \sqrt{-1} \left\{ \sum_{j \leq m} b_{ij} t_j + \sum_{j > m} c_{ij} \xi_j \eta_j \right\} / \xi_i, \quad i = 1, \dots, m$$

with $(b_{ij}) = (a_{ij})_{1 \leq i, j \leq m}^{-1}$ and $t_i \in \mathbb{R}$. This shows that $\mathcal{Q} \setminus \cup_{j=1}^m \{\xi_j = 0\}$ is a generic Levi-flat CR manifold of CR dimension $2n - m$. Next, we want to show that $\mathcal{Q} \cap \cup_{j=1}^m \{\xi_j = 0\}$ has dimension less than $4n - m$. It suffices to verify that $\mathcal{Q}' = \mathcal{Q} \cap \{\xi_1 = 0\}$ has dimension less than $4n - m$. Note that the rank of $(a_{ij})_{1 \leq i \leq m, 2 \leq j \leq m}$ is $m - 1$. Without loss of generality, we may further assume that $(a_{ij})_{2 \leq i, j \leq m}$ has rank $m - 1$. Obviously, \mathcal{Q}' is contained in $\mathbb{C} \times \mathcal{Q}''$ for $\mathcal{Q}'' \subset \mathbb{C}^{2n-2} : \Re \left\{ \sum_{j=2}^m a_{ij} \xi_j \eta_j \right\} = 0, i = 2, \dots, m$. Applying induction on n and m , one can verify that $\mathcal{Q} \cap \{\xi_1 = \eta_1 = 0\}$ is a real analytic set of dimension at most $4(n - 1) - (m - 1)$; hence $\dim \mathcal{Q}' < 4n - m$. The proof of the proposition is complete. \square

PROPOSITION 2.2. — *Let $\mathcal{Q} \subset \mathbb{C}^{2n}$ be an irreducible generic real analytic set defined by real quadratic forms and of dimension at least $2n$. Assume that $\Lambda(\mathcal{Q}) = \mathcal{Q}$, and that \mathcal{I} is contained in $E^{psh} \cap E_0$. There is a \mathbb{C} -linear symplectic transformation which leaves Λ in the diagonal form (2.1) and transforms \mathcal{Q} into the set*

$$(2.3) \quad \xi_j \eta_{\sigma(j)} + \mu_j \bar{\xi}_{\sigma(j)} \bar{\eta}_j = 0, \quad j = 1, \dots, 2L$$

with $\sigma(2j - 1) = 2j = \sigma^{-1}(2j - 1)$ for $j = 1, \dots, L$, and

$$(2.4) \quad |\mu_j| = 1, \quad \mu_{\sigma(j)} = \mu_j.$$

Proof. — Since \mathcal{I} is invariant under Λ , we have $\mathcal{I} = \oplus I_{ij}^{psh}$ for

$$I_{ij}^{psh} \equiv \mathcal{I} \cap E^{psh} \cap E_{ij}.$$

Assume that $I_{ij}^{psh} \neq \{0\}$. We first want to show that $\dim_{\mathbb{C}} I_{ij}^{psh} = 1$. Without loss of generality, one may assume that $i, j > 0$. Assume for the sake of contradiction that $\dim_{\mathbb{C}} I_{ij}^{psh} > 1$. Since $\dim_{\mathbb{C}} E_{ij}^{psh} = 2$ then

$\mathcal{I}_{ij}^{ps h} = E_{ij}^{ps h}$; hence, \mathcal{Q} is contained in $\xi_i \xi_j = 0$, which contradicts that \mathcal{Q} is generic and of dimension at least $2n$. Next, we want to show that

$$(2.5) \quad \dim_{\mathbb{C}} \mathcal{I}_{ij}^{ps h} + \dim_{\mathbb{C}} \mathcal{I}_{i'j'}^{ps h} \leq 1$$

for $\{i, j\} \neq \{i', j'\}$ and $\{i, j\} \cap \{i', j'\} \neq \emptyset$. Let us focus on the case $i, j > 0$, and other cases can be checked similarly. Assume for the sake of contradiction that on \mathcal{Q}

$$a\xi_i \xi_j + b\bar{\eta}_i \bar{\eta}_j = 0, \quad c\xi_i \xi_{j'} + d\bar{\eta}_i \bar{\eta}_{j'} = 0$$

for some $a, b, c, d \neq 0$. Eliminating ξ_i and $\bar{\eta}_i$ yields

$$(2.6) \quad ad\xi_j \bar{\eta}_{j'} - bc\xi_{j'} \bar{\eta}_j = 0$$

on $\mathcal{Q} \setminus (\{\xi_i = 0\} \cup \{\xi_{j'} = 0\})$. Since \mathcal{Q} is irreducible and generic, then (2.6) holds on \mathcal{Q} also, which contradicts that $\mathcal{I} \cap E^{hrm} = \{0\}$.

Next, we want to show that all $I_{ii}^{ps h} = 0$. Otherwise, one may assume that $I_{1,1}^{ps h} \neq 0$, so \mathcal{Q} is defined by the vanishing of $\xi_1^2 - a\bar{\eta}_1^2$ and other quadratic forms independent of ξ_1, η_1 . Denote by $\mathcal{Q}' \subset \mathbb{C}^{2n-2}$ the vanishing set of the complex-valued quadratic forms in \mathcal{I} that are independent of ξ_1, η_1 . Obviously, the decomposition

$$\mathcal{Q} = (\{\xi_1 + \sqrt{a}\bar{\eta}_1 = 0\} \cup \{\xi_1 - \sqrt{a}\bar{\eta}_1 = 0\}) \times \mathcal{Q}'$$

contradicts that \mathcal{Q} is irreducible.

To achieve (2.3), let \mathcal{I} contain one of quadrics

$$\xi_i \eta_j + a\bar{\eta}_i \bar{\xi}_j, \quad \xi_i \xi_j + a\bar{\eta}_i \bar{\eta}_j, \quad \eta_i \eta_j + a\bar{\xi}_i \bar{\xi}_j$$

with $i \neq j$ and $a \neq 0$. By substituting $(\eta_j, -\xi_j)$ for (ξ_j, η_j) in the second quadratic form and replacing the third quadratic form by its conjugate divided by \bar{a} , one may assume that the first quadratic form is in \mathcal{I} . Next, interchange (ξ_i, η_i) with (ξ_1, η_1) and (ξ_j, η_j) with (ξ_2, η_2) , respectively. Thus \mathcal{I} contains

$$\xi_1 \eta_2 + \mu_1 \bar{\eta}_1 \bar{\xi}_2, \quad \mu_1 \neq 0.$$

By the transformation $(\xi_1, \eta_1) \rightarrow (\sqrt{|\mu_1|} \xi_1, \eta_1 / \sqrt{|\mu_1|})$, one gets $|\mu_1| = 1$. Inductively, one finds the remaining coordinates $(\xi_3, \eta_3, \dots, \xi_n, \eta_n)$ so that \mathcal{Q} is given by (2.3)–(2.4). The proof of the proposition is complete. \square

Analogous to Proposition 2.2, we have the following.

PROPOSITION 2.3. — *Let $\mathcal{Q} \subset \mathbb{C}^{2n}$ be an irreducible generic real analytic set defined by real quadratic forms and of dimension at least $2n$.*

Assume that \mathcal{Q} is invariant under Λ , and that \mathcal{I} is contained in $E^{hrm} \cap E_0$ and satisfies the non-degeneracy conditions

$$(2.7) \quad \dim_{\mathbb{C}} \mathcal{I}_{ij}^{hrm} \cdot \dim_{\mathbb{C}} \mathcal{I}_{jk}^{hrm} = 0$$

for all distinct i, j, k . Then there is a \mathbb{C} -linear symplectic transformation which leaves Λ in the diagonal form (2.1) and transforms \mathcal{Q} into

$$(2.8) \quad \xi_j \bar{\xi}_{\sigma(j)} - \mu_j \bar{\eta}_j \eta_{\sigma(j)} = 0, \quad j = 1, \dots, 2L$$

with $\mu_{\sigma(j)} = \bar{\mu}_j$, $|\mu_j| = 1$, and $\sigma(2j-1) = 2j = \sigma^{-1}(2j-1)$ for $j = 1, \dots, L$.

The proof is almost the same as that of Proposition 2.2, except (2.7), replacing (2.5), is part of the assumptions of the proposition. One can show that \mathcal{Q} is equivalent to $\xi_j \bar{\eta}_{\sigma(j)} + \mu_j \bar{\eta}_j \xi_{\sigma(j)} = 0, j = 1, \dots, 2L$. However, the latter is equivalent to (2.8) by a symplectic transformation. The details are left to the reader.

We depart from the symplectic coordinates for a moment.

PROPOSITION 2.4. — Let $\mathcal{Q} \subset \mathbb{C}^n$ be defined by the vanishing of some linear combinations of $z_1 \bar{z}_1, \dots, z_n \bar{z}_n$. Assume that \mathcal{Q} is generic, Levi-flat, and of codimension $m \leq n$. Then m is less than n and rearranging z_1, \dots, z_n gives

$$(2.9) \quad \mathcal{Q}: z_j \bar{z}_j - c_j z_{k_j} \bar{z}_{k_j} = 0, \quad j = 1, \dots, m$$

for some $c_j > 0$ and $k_j > m$.

Proof. — For the space of \mathbb{R} -linear combinations of $z_1 \bar{z}_1, \dots, z_n \bar{z}_n$ that vanish on \mathcal{Q} , choose a basis q_1, \dots, q_k so that for a possible rearrangement of z_1, \dots, z_n

$$(2.10) \quad q_j(z, \bar{z}) = z_j \bar{z}_j - \sum_{\alpha > k} a_{j\alpha} z_{\alpha} \bar{z}_{\alpha}, \quad j = 1, \dots, k.$$

Since \mathcal{Q} is not contained in $\cup_{j=1}^n z_j = 0$, then

$$D = \left\{ (z_{k+1}, \dots, z_n) \mid \sum_{l > k} a_{jl} z_l \bar{z}_l > 0, j = 1, \dots, k \right\}$$

is a nonempty open set in \mathbb{C}^{n-k} , of which the boundary contains the origin. Obviously, $\{q_1 = \dots = q_k = 0\} \setminus \cup_{j=1}^k \{z_j = 0\}$ is a smooth submanifold in \mathbb{C}^n of codimension k ; in particular, $k \geq m$. On the other hand, if $k > m$, then \mathcal{Q}^* is contained in $\cup_{j=1}^k \{z_j = 0\}$, which contradicts that \mathcal{Q} is generic. Therefore, $k = m$.

To determine $a_{j\alpha}$ we consider $(1, 0)$ tangent vectors of $\mathcal{Q}_c \setminus \cup_{j=1}^k \{z_j = 0\}$, spanned by

$$(2.11) \quad L_\alpha = \sum_{j=1}^m a_{j\alpha} \frac{\bar{z}_\alpha}{\bar{z}_j} \frac{\partial}{\partial z_j} + \frac{\partial}{\partial z_\alpha}, \quad \alpha > m.$$

One has

$$[L_\alpha, \bar{L}_\beta] = \sum_{j=1}^m a_{j\alpha} a_{j\beta} \frac{z_\alpha \bar{z}_\beta}{z_j \bar{z}_j^2} \frac{\partial}{\partial \bar{z}_j} - \sum_{j=1}^m a_{j\alpha} a_{j\beta} \frac{z_\alpha \bar{z}_\beta}{\bar{z}_j z_j^2} \frac{\partial}{\partial z_j}, \quad \alpha \neq \beta.$$

Since \mathcal{Q} is Levi-flat, the Lie bracket $[L_\alpha, \bar{L}_\beta]$, restricted to $\mathcal{Q}_c \setminus \cup \{z_j = 0\}$, is locally in the span of vector fields (2.11) and their conjugates. This implies that $[L_\alpha, \bar{L}_\beta]$ vanishes on $\mathcal{Q}_c \setminus \cup \{z_j = 0\}$, i.e.,

$$a_{j\alpha} a_{j\beta} \frac{z_\alpha \bar{z}_\beta}{z_j \bar{z}_j^2} = 0, \quad \text{on } \mathcal{Q} \setminus \cup_{j=1}^m \{z_j = 0\}.$$

Since $\mathcal{Q} \setminus \cup_{j=1}^m \{z_j = 0\}$ is not contained in $z_\alpha z_\beta = 0$, then $a_{j\alpha} a_{j\beta} = 0$ for $\alpha \neq \beta$. On the other hand, from (2.10) one knows that at least one of $a_{j\alpha} (\alpha > m)$ is positive. This shows that \mathcal{Q} has the form (2.9). The proof of the proposition is complete. \square

Returning to symplectic coordinates, we now want to prove the following.

PROPOSITION 2.5. — *Let $\mathcal{Q} \subset \mathbb{C}^{2n}$ be a Levi-flat real analytic set defined by elements in $E^{hm} \cap E_1$. Assume that \mathcal{Q} is generic and of codimension $m \leq 2n$, and that \mathcal{Q} contains no \mathbb{C} -linear coordinate subspaces (i.e., vanishing sets of some of $\xi_1, \eta_1, \dots, \xi_n, \eta_n$) of dimension greater than $2n - 2m$. Then under a change of symplectic coordinates which leaves Λ in the form (2.1), \mathcal{Q} is given by*

$$(2.12) \quad |\xi_j|^2 - \mu_j |\eta_{\sigma(j)}|^2 = 0, \quad j = 1, \dots, m,$$

where $\sigma: \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ is injective, and μ_j satisfy

$$\mu_j = \begin{cases} 1/\mu_{\sigma(j)} > 0 & \text{if } \tau_j \text{ is even and positive,} \\ 1 & \text{otherwise} \end{cases}$$

with τ_j being the smallest positive integer satisfying $\sigma^{\tau_j}(j) = j$ ($\tau_j = 0$ when $\sigma^k(j) > m$ for some k).

Proof. — In view of Proposition 2.4, such a set takes the form (2.9) with z_1, \dots, z_{2n} being a permutation of $\xi_1, \eta_1, \dots, \xi_n, \eta_n$. Since \mathcal{Q} does not contain any \mathbb{C} -linear coordinate subspace of dimension greater than

$2n - 2m$, all k_1, \dots, k_m in (2.9) are distinct; consequently, $m \leq n$ and \mathcal{Q} is given by m equations

$$(2.14) \quad |\xi_{i_1}|^2 = c'_{i_1} |\xi_{\alpha(i_1)}|^2, \quad |\xi_{i_2}|^2 = c''_{i_2} |\eta_{\beta(i_2)}|^2, \quad |\eta_{i_3}|^2 = c''_{i_3} |\eta_{\gamma(i_3)}|^2,$$

in which $c'_{i_1}, c''_{i_2}, c''_{i_3}$ are positive, and indices i_1, i_2, i_3 run over subsets I_1, I_2, I_3 of $\{1, \dots, n\}$, respectively. Moreover, I_1, I_2 and $\alpha(I_1)$ are mutually disjoint, and so are $I_3, \beta(I_2)$, and $\gamma(I_3)$. We shall first permute the symplectic coordinates to eliminate the first and last groups of equations in (2.14). For each $j \in \alpha(I_1)$, we change the coordinates (ξ_j, η_j) to $(-\eta_j, \xi_j)$, and for each $k \in \beta^{-1}(\alpha(I_1) \cap \beta(I_2))$, we change the coordinates (ξ_k, η_k) to $(-\eta_k, \xi_k)$ and replace c''_k with $1/c''_k$. Thus, equations in the first group are eliminated. Next, we apply the coordinate change (ξ_j, η_j) to $(-\eta_j, \xi_j)$ for each $j \in I_3$ and $j \in \beta^{-1}(I_3 \cap \beta(I_2))$. After the above coordinate changes, only the second group of equations in (2.14) remains. By a permutation of symplectic coordinates, we achieve (2.12) for an injective σ .

To obtain the normalization (2.13), we apply induction on the number of equations in (2.12). If $m = 1$, one readily obtains (2.13) by applying the transformation $(\xi_j, \eta_j) \rightarrow (\sqrt{\mu_1}\xi_j, \eta_j/\sqrt{\mu_1})$ for $j = 1, \sigma(1)$.

For induction, we assume that (2.13) can be achieved if the number of equations in (2.12) is less than m . We first consider the case there exists the smallest integer τ_1 with $\sigma^{\tau_1}(1) = 1$. Note that such τ_1 always exists if $\sigma^k(1) \leq m$ for all integer k . In this case, the equations in (2.12) break into two groups: the equations involving only $(\xi_{\sigma^k(1)}, \eta_{\sigma^k(1)})$ for $k = 1, \dots, \tau_1$ and the ones not involving any of those variables. If both groups are non-empty, (2.12) follows from the induction assumption. Thus, one may assume, without loss of generality, that $\tau_1 = m$, $\sigma^j(1) = j + 1$ for $j = 1, \dots, m - 1$.

Consider a symplectic transformation

$$(2.15) \quad \xi_j \rightarrow c_j \xi_j, \quad \eta_j \rightarrow \eta_j / c_j, \quad c_j > 0, \quad j = 1, \dots, n.$$

The transformation (2.15) sends (2.12) to $|\xi_j|^2 - \tilde{\mu}_j |\eta_{\sigma(j)}|^2 = 0$ with $\tilde{\mu}_j = c_j^2 c_{j+1}^2 \mu_j$. Put $c_1 = 1$ and $c_j = 1/(c_{j-1} \sqrt{\mu_{j-1}})$ for $j = 2, \dots, m$. Then $\tilde{\mu}_j = 1$ for $j = 1, \dots, m - 1$. To normalize $\tilde{\mu}_m$, consider first the case that m is even. In this case

$$\delta_1 \mu \equiv \mu_1 \mu_2^{-1} \mu_3 \cdots \mu_m^{-1}$$

is invariant under (2.15). Therefore, (2.12) are equivalent to $|\xi_j|^2 - \mu'_j |\eta_{j+1}|^2 = 0, j = 1, \dots, m$ for $\mu'_1 = (\delta, \mu)^{1/m}$ and $\mu'_{\sigma(j)} = 1/\mu'_j$, since $\delta_1 \mu' = \delta_1 \mu$. Consider now the case that m is odd. In this case we apply

another transformation (2.15) with $c_1 = (\mu'_m)^{-1/4}$ and $c_{j+1} = 1/c_j$ for $j = 1, \dots, m - 1$, which gives us (2.12) with $\mu_j = 1$ for $j = 1, \dots, m$.

We now consider the case that $\sigma^k(1) = m' > m$ for some k . In this case $1 \leq \tau^{-1}(m') \leq m$. Let τ be the largest integer such that $\sigma^{-\tau}(m')$ is well-defined. The equations in (2.12) are divided into two groups: the ones involving $\eta_{m'}, \xi_{\sigma^{-1}(m')}, \eta_{\sigma^{-1}(m')}, \dots, \xi_{\sigma^{-\tau}(m')}$ only and the ones not involving $\xi_{\sigma^{-k}(m')}, \eta_{\sigma^{-k}(m')}$ for $k = 0, \dots, \tau$. Thus by the induction assumption, one may assume that $\tau = m - 1$. Put $j_0 = \sigma^{-m+1}(m')$ and $j_k = \sigma^k(j_0)$ for $k = 1, \dots, m - 1$. Thus, (2.12) becomes

$$|\xi_{k_j}|^2 - \mu_{k_j} |\eta_{k_{j+1}}|^2 = 0, \quad j = 0, \dots, m - 1$$

with $j_0, \dots, j_{m-1} \leq m$ being distinct and $j_m = m' > m$. Take $c_{k_{m-1}} = 1$, and $c_{k_j} = 1/(c_{k_{j+1}} \sqrt{\mu_{k_j}})$ for $k = m - 2, \dots, 0$. Then applying the transformation (2.15) one gets $\mu_{k_j} = 1$ for $j = 0, \dots, m - 1$. The proposition is proved by induction. □

To conclude this section we remark that, for holomorphic symplectic mappings, having a Levi-flat invariant set of minimal CR dimension is not a sufficient condition for the convergence of their normalizations. In fact, the sets M_2, \dots, M_5 below are combinations of 4 types of Levi-flat sets (2.2), (2.3), (2.8) and (2.12).

Examples. — The following are Levi-flat sets of minimal CR dimension:

$$\begin{aligned} M_1 &\subset \mathbb{C}^4: \xi_2^2 - \bar{\eta}_2^2 = 0, \\ M_2 &\subset \mathbb{C}^6: \xi_2 \eta_2 + \overline{\xi_2 \eta_2} = \xi_3 \xi_3 + \overline{\eta_3 \eta_3} = 0, \\ M_3 &\subset \mathbb{C}^6: |\xi_2|^2 - |\eta_2|^2 = \xi_2 \bar{\eta}_3 + \xi_3 \bar{\eta}_2 = 0, \\ M_4 &\subset \mathbb{C}^6: |\xi_2|^2 - |\eta_2|^2 = |\xi_3|^2 - |\eta_3|^2 = |\xi_2|^2 - |\xi_3|^2 = 0, \\ M_5 &\subset \mathbb{C}^8: \xi_2 \bar{\xi}_3 - \bar{\eta}_2 \eta_3 = \xi_3 \bar{\xi}_4 - \bar{\eta}_3 \eta_4 = 0. \end{aligned}$$

Note that all these sets contain the (ξ_1, η_1) -subspace. Let ψ be an elliptic holomorphic symplectic transformation of \mathbb{C}^2 which is not normalizable by any convergent symplectic transformation. For each M_j , one can find a map φ which is the product of ψ with a suitable elliptic \mathbb{C} -linear symplectic transformation of \mathbb{C}^{2n-2} ($n = 2, 3$, or 4) such that M_j is invariant under φ . However, φ is not normalizable by any convergent symplectic transformation.

3. Normalization of Levi-flat invariant sets.

Throughout the paper we denote by $[p]_k$ the sum of homogeneous terms of order k , whenever p is a multivariable power series.

In Section 2, four special Levi-flat classes of quadrics were introduced: the sets defined by (2.2), (2.3), (2.8) and (2.12), respectively. We now study Levi-flat sets defined by equations whose lowest order terms correspond to these quadratic polynomials, and we try to simplify their equations by changing holomorphic coordinates. For this purpose, we shall combine two singularities (2.2) and (2.3) since such combination results in no extra difficulties. One can also treat two singularities (2.8) and (2.12) simultaneously. Notice that quantities μ_j in Section 2 are symplectic invariants, but are not the holomorphic ones. Since holomorphic coordinates, not the symplectic ones, are used throughout this section, we can restrict ourselves to the case all $\mu_j = 1$. We shall also use a simpler permutation σ to simplify notations.

To combine two types of singularities (2.2) and (2.3), we let σ be a permutation of $1, \dots, m$ ($m \leq n$) satisfying

$$(3.1) \quad \sigma(j) = \begin{cases} j, & \text{if } 1 \leq j \leq K, \\ \sigma^{-1}(j) \neq j, & \text{if } K < j \leq K + 2L = m. \end{cases}$$

In particular, $\sigma^2 = \text{Id}$. Put

$$(3.2) \quad q_j(\xi, \eta) = \begin{cases} \sum_{k=1}^n a_{jk} \xi_k \eta_k, & 1 \leq j \leq K, \\ \xi_j \eta_{\sigma(j)}, & K < j \leq m \end{cases}$$

with

$$(3.3) \quad \det(a_{ij})_{1 \leq i, j \leq K} \neq 0.$$

Define the set

$$(3.4) \quad \mathcal{Q} \subset \mathbb{C}^{2n}: q_j(\xi, \eta) + \bar{q}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, m.$$

Note that we allow $K = 0$ or m , so that sets (2.2) and (2.3) (all $\mu_j = 1$) are indeed special cases of (3.4). In Proposition 3.5 below, we shall prove that (3.4) is generic, irreducible and Levi-flat.

One of main results of this section is the following.

THEOREM 3.1. — *Let $M \subset \mathbb{C}^{2n}$ be a real analytic variety defined by*

$$(3.5) \quad q_j(\xi, \eta) + \bar{q}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, m,$$

where σ, q_j , and a_{ij} are given by (3.1)–(3.3) and R_j are real analytic and satisfy the reality condition

$$(3.6) \quad R_{\sigma(j)}(\xi, \eta, \bar{\xi}, \bar{\eta}) = \overline{R_j(\bar{\xi}, \bar{\eta}, \xi, \eta)}.$$

Assume that M is Levi-flat. Then there are m holomorphic functions $H_j(\xi, \eta) = O(3)$ such that M , as a germ of real analytic set at the origin, is given by

$$(3.7) \quad q_j(\xi, \eta) + \bar{q}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) + H_j(\xi, \eta) + \overline{H_{\sigma(j)}(\bar{\xi}, \bar{\eta})} = 0, \quad j = 1, \dots, m.$$

To combine two singularities (2.8) and (2.12), let σ be still given by (3.1), and put

$$(3.8) \quad q_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = \begin{cases} |\xi_j|^2 - |\eta_j|^2, & 1 \leq j \leq K, \\ \xi_j \bar{\xi}_{\sigma(j)} - \bar{\eta}_j \eta_{\sigma(j)}, & K < j \leq m. \end{cases}$$

We now state the following rigidity result.

THEOREM 3.2. — *Let $M \subset \mathbb{C}^{2n}$ be a Levi-flat real analytic variety defined by*

$$(3.9) \quad q_j(\xi, \eta, \bar{\xi}, \bar{\eta}) + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad 1 \leq j \leq m,$$

where σ, q_j are given (3.1) and (3.8), and $R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = O(3)$ are real analytic and satisfy the reality condition (3.6). Then near the origin, M is holomorphically equivalent to the set

$$(3.10) \quad \mathcal{Q} \subset \mathbb{C}^{2n}: q_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, m.$$

Note that (3.10) becomes (2.12) (all $\mu_j = 1$ and $\sigma = \text{Id}$) when $K = 0$, or (2.8) (all $\mu_j = 1$) when $K = m$. As a consequence of Proposition 3.7 we shall see that the set (3.10) is generic, irreducible, and Levi-flat.

The main purpose of this section is to prove Theorem 3.1 and Theorem 3.2. We start with the following.

LEMMA 3.3. — *Let $V \subset \mathbb{C}^{2m} \times \mathbb{C}^n$ be a complex variety defined by*

$$f_j = x_j^2 + y_j^2 + q_j(x, y, z) + h_j(x, y, z) = 0, \quad j = 1, \dots, m,$$

where q_j is a quadratic polynomial in $x_{j+1}, y_{j+1}, \dots, x_m, y_m, z$ for $j = 1, \dots, m$, and h_j are holomorphic functions of order at least 3. Let f be a holomorphic function vanishing on V . Then, as a germ of holomorphic function at the origin, f admits a decomposition

$$f = a_1 f_1 + a_2 f_2 + \dots + a_n f_n$$

for some holomorphic functions a_j with $\min_j \{\text{ord } a_j\} = \text{ord } f - 2$.

Proof. — Without loss of generality one may assume that f_j is a quadratic Weierstrass polynomial in x_j . Put $f = g_0$. Using the Weierstrass division theorem, one gets

$$g_{j-1} = a_j f_j + g_j, \quad j = 1, \dots, m$$

with $\min\{\text{ord } a_j + 2, \text{ord } g_j\} = \text{ord}\{g_{j-1}\} \geq \text{ord } f$. Hence

$$f = a_1 f_1 + \dots + a_m f_m + g_m.$$

Assuming that $f \neq 0$, we would like to show that $\text{ord } g_m > \text{ord } f$ and $\min\{\text{ord } a_j\} = \text{ord } f - 2$.

Consider the dilation $(x, y, z) \rightarrow (\epsilon x, \epsilon y, \epsilon z)$. Let V^ϵ be defined by

$$x_j^2 + y_j^2 + q_j(x, y, z) + h_j(x, y, z, \epsilon) \equiv f_j(\epsilon x, \epsilon y, \epsilon z)/\epsilon^2 = 0$$

for $j = 1, \dots, m$. Obviously, $h_j = 0$ for $\epsilon = 0$. Put

$$\Delta_r = \{t \in \mathbb{C}: |t| < r\}$$

and let Δ_r^k be the products of k copies of Δ_r . We shall also drop the superscript in Δ_r^k whenever the dimension k is clear from context. Let π be the projection $(x, y, z) \rightarrow (y, z)$. Choose δ, δ' so small that the restriction $\pi: V^\epsilon \equiv V^\epsilon \cap (\Delta_\delta^m \times \Delta_{\delta'}^{m+n}) \rightarrow \Delta_{\delta'}^{m+n}$ is a 2^m -to-1 branched covering, for which the branch locus is denoted by B^ϵ . Obviously,

$$B^0 = V^0 \cap \{(y_m^2 + q_m(x, y, z)) \cdots (y_1^2 + q_1(x, y, z)) = 0\}$$

is a proper subvariety of V^0 ; hence, B^ϵ is also a proper subvariety of V^ϵ for small ϵ . Off the branch locus B^ϵ , V^ϵ is locally given by $x = d_J(y, z, \epsilon)$, $J \in \mathbb{Z}_2^m = \{0, 1\}^m$ with $d_J(\cdot, \epsilon) \rightarrow d_J(\cdot, 0)$ as $\epsilon \rightarrow 0$. Note that for $\epsilon = 0$ the k -th coordinate of d_J is obtained by choosing a square root of $-y_k^2 - q_k(x, y, z)$, starting with $k = m$. This implies that for $J = (j_1, \dots, j_m) \in \mathbb{Z}_2^m$ and $J_k = (j_k, \dots, j_m)$, one has $d_J(y, z, 0) = (X_{J_1}(y, z), X_{J_2}(y, z), \dots, X_{J_m}(y, z))$. Moreover,

$$(3.11) \quad X_{(0, J_{k+1})}(y, z) \neq X_{(1, J_{k+1})}(y, z), \quad J_{k+1} \in \mathbb{Z}_2^{m-k-1}$$

for $(y, z) \notin B^0$ and $1 \leq k \leq m$.

Assume for the sake of contradiction that $\text{ord } g_m = \text{ord } f \equiv k$. Note that

$$(3.12) \quad g_m(\epsilon x, \epsilon y, \epsilon z)/\epsilon^k = [g_m]_k(x, y, z) + \epsilon \tilde{g}_m(x, y, z, \epsilon)$$

vanishes on V^ϵ , and that $\text{ord } \tilde{g}_m > k$. Since q_j is independent of x_1, \dots, x_j , one gets from $[g_{j-1}]_k = [a_j]_{k-2}[f_j]_2 + [g_j]_k$ that

$$[g_m]_k(x, y, z) = \sum_{\alpha \in \mathbb{Z}_2^m} c_\alpha(y, z)x^\alpha.$$

Now (3.12) yields

$$\sum_{\alpha \in \mathbb{Z}_2^m} c_\alpha(y, z)d_J^\alpha(y, z, \epsilon) + \epsilon \tilde{g}_m(d_J(y, z, \epsilon), y, z, \epsilon) = 0, \quad J \in \mathbb{Z}_2^m$$

for $(y, z) \notin \pi(B^\epsilon)$. Letting $\epsilon \rightarrow 0$, one gets

$$\sum_{\alpha \in \mathbb{Z}_2^m} c_\alpha(y, z)d_J^\alpha(y, z, 0) = 0, \quad J \in \mathbb{Z}_2^m$$

for $(y, z) \notin \pi(B^0)$. The above are 2^m linear equations in 2^m unknowns $c_\alpha(y, z)$, for which the coefficients $d_J^\alpha(y, z), \alpha, J \in \mathbb{Z}_2^m$ form a block matrix

$$D = \begin{pmatrix} D' & D' \\ X_0 D' & X_1 D' \end{pmatrix},$$

where D' is the matrix formed by $d_{(0, J')}^{(0, \alpha')}$ with $\alpha', J' \in \mathbb{Z}_2^{m-1}$, and X_0, X_1 are diagonal $2^{m-1} \times 2^{m-1}$ matrices of which the J' -th diagonal entry is $X_{(0, J')}(y, z)$ and $X_{(1, J')}(y, z)$, respectively. From (3.11) it follows that $\det(X_0 - X_1) \neq 0$. Applying induction on m , one gets $\det D = \det(X_0 - X_1) \det D' \neq 0$. Therefore, $c_\alpha = 0$. This shows that $\text{ord } g_m > \text{ord } f$ and $\text{ord } f = \min\{\text{ord } a_j\} + 2$.

However, g_m still vanishes on V , so one can find another decomposition for g_m . Inductively, one can at least achieve

$$(3.13) \quad f = a_1 f_1 + \dots + a_m f_m, \quad \min\{\text{ord } a_m\} = \text{ord } f - 2$$

for some formal power series a_j . By a theorem of M. Artin [1], there are convergent power series $\tilde{a}_j(x, y, z) = a_j(x, y, z) + O(k)$ satisfying (3.13). This completes the proof of the lemma. \square

LEMMA 3.4. — *Let V, q_1, \dots, q_m be as in Lemma 3.3, and let $q_j(0, 0, z) = \sum_{k=1}^n a_{jk} z_k^2$. Assume that the rank of (a_{jk}) is m . Then V is irreducible.*

Proof. — As in the proof of Lemma 3.3, let π be the projection $(x, y, z) \rightarrow (y, z)$. Choose $\delta, \delta' > 0$ such that the restriction $\pi: V \equiv V \cap (\Delta_\delta^m \times \Delta_{\delta'}^{m+n}) \rightarrow \Delta_{\delta'}^{m+n}$ is a 2^m -to-1 branched covering, for which the

set of points of *branch order* 2^m , i.e., the points $p \in V$ with $\pi^{-1}\pi(p) = \{p\}$, contains a subset B satisfying additional equations

$$(3.14) \quad 2x_j + \partial_{x_j} h_j = \partial_{x_j} f_j = 0, \quad j = 1, \dots, m.$$

For a possible smaller δ' , the above equations define a one-to-one covering over $\Delta_{\delta'}^{m+n}$. Solving for x from (3.14) and inserting x into $f_j(x, y, z)$ yields new functions

$$(3.15) \quad \tilde{f}_j(y, z) = y_j^2 + q_j(0, y, z) + \tilde{h}_j(y, z) = 0, \quad j = 1, \dots, m$$

on B . Since (a_{jk}) is of rank m , one may assume that $(a_{jk})_{1 \leq j, k \leq m}$ is nonsingular. Fix (y^0, z^0) on $y_j^2 + q_j(0, y, z) = 0, j = 1, \dots, m$, so that $z_{m+1}^0 = \dots = z_n^0 = 0$, while none of $y_1^0, z_1^0, \dots, y_m^0, z_m^0$ is zero. Consider the dilation $g_j(y, z, \epsilon) \equiv \tilde{f}_j(\epsilon y, \epsilon z)/\epsilon^2$. Then $\partial g/\partial(z_1, \dots, z_m)$ is nonsingular for $(y, z) = (y^0, z^0)$ and $\epsilon = 0$. By the implicit function theorem, one can verify that for small ϵ , there exist $x(\epsilon), z(\epsilon)$ such that $x(\epsilon), \epsilon y^0, z(\epsilon)$ satisfy (3.14)–(3.15) and $|x(\epsilon)|, |z(\epsilon)| \leq c\epsilon$ for some constant c ($z_j(\epsilon) = 0$ for $j > m$).

On the other hand, the singular locus V_s of V is contained in

$$\det \partial f/\partial y = 2^m y_1 y_2 \dots y_m + O(m+1) = 0,$$

which obviously does not contain $(x(\epsilon), \epsilon y^0, z(\epsilon)) \in B$ for small $\epsilon \neq 0$. This shows that V contains a smooth point of branch order 2^m . Hence, V^* is connected, i.e., V is irreducible. This completes the proof of the lemma. \square

Note that the above proof is based on branched coverings and the existence of smooth branch points of maximal branch order. Applying the same argument to the family of complex varieties

$$V^\epsilon: f_j^\epsilon(x, y, z) \equiv f_j(\epsilon x, \epsilon y, \epsilon z)/\epsilon^2, \quad j = 1, \dots, m$$

yields that any neighborhood of the origin contains another neighborhood D , independent of ϵ , such that each $V^\epsilon \cap D$ is connected for small ϵ .

We now apply the above lemmas to the real analytic variety given by (3.5) and (3.6).

PROPOSITION 3.5. — *Let $M \subset \mathbb{C}^{2n}$ be a real analytic set given by (3.5)–(3.6). Then M is a generic and irreducible real analytic variety of codimension m . Also, a germ of any real analytic function f vanishing on an open subset of M^* , of which the closure contains the origin, can be decomposed into $a_1 r_1 + \dots + a_m r_m$ for germs of some complex-valued real analytic functions a_j with $\min_j \{\text{ord } a_j\} = \text{ord } f - 2$. Moreover, the set \mathcal{Q} , defined by (3.4), is Levi-flat.*

Proof. — Put $r_j^\epsilon(\xi, \eta, \bar{\xi}, \bar{\eta}) = r_j(\epsilon\xi, \eta, \epsilon\bar{\xi}, \bar{\eta})/\epsilon^2$ for $j = 1, \dots, m$. Note that \mathcal{Q} is defined by $r_1^0 = \dots = r_m^0 = 0$. At $\xi = (0, \dots, 0), \eta = (1, \dots, 1)$, one has

$$\det \partial r^0 / \partial(\xi_1, \dots, \xi_m) = \det A \neq 0, \quad A = (a_{ij})_{1 \leq i, j \leq K}.$$

This shows that $\dim \mathcal{Q} \geq 4n - m$, and that \mathcal{Q} is generic if $\dim \mathcal{Q} = 4n - m$. Now the implicit function theorem implies that M enjoys the same property.

To show that $\dim M = 4n - m$, we consider the complex variety

$$\mathcal{M} \subset \mathbb{C}^{2n} \times \mathbb{C}^{2n} : r_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = \bar{r}_j(\bar{\xi}, \bar{\eta}, \xi, \eta) = 0, \quad j = 1, \dots, m,$$

where $\xi, \eta, \bar{\xi}, \bar{\eta}$ are now independent complex variables. By the reality conditions (3.6), \mathcal{M} is actually defined by $r_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0$ for $j = 1, \dots, m$; more explicitly, \mathcal{M} is given by

$$\begin{aligned} \xi_j \eta_j + \sum_{k=K+1}^n b_{jk} \xi_k \eta_k + \sum_{k=1}^n c_{jk} \bar{\xi}_k \bar{\eta}_k + \tilde{R}_j(\xi, \eta, \bar{\xi}, \bar{\eta}) &= 0, \quad 1 \leq j \leq K, \\ \xi_j \eta_{\sigma(j)} + \bar{\xi}_{\sigma(j)} \bar{\eta}_j + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) &= 0, \quad K < j \leq m, \end{aligned}$$

where $(c_{jk})_{1 \leq j, k \leq K}$ is nonsingular. Choose complex linear coordinates x, y, z so that

$$(3.16) \quad \begin{aligned} \xi_j \eta_j &= x_j^2 + y_j^2, & \bar{\xi}_j \bar{\eta}_j &= z_j^2 + z_{m+j}^2, & 1 \leq j \leq K, \\ \xi_j \eta_{\sigma(j)} &= x_j^2 + y_j^2, & \bar{\xi}_{\sigma(j)} \bar{\eta}_j &= z_j^2 + z_{m+K+j}^2, & K < j \leq m. \end{aligned}$$

Substitute the rest of z -variables for $\xi, \eta, \bar{\xi}, \bar{\eta}$ that do not appear in (3.16). Obviously, in the new coordinates, \mathcal{M} has defining functions satisfying all the assumptions in Lemma 3.4; hence, \mathcal{M} is irreducible and of codimension m . Since $M \equiv \{(\xi, \eta, \bar{\xi}, \bar{\eta}) : (\xi, \eta) \in M\} \subset \mathcal{M}$ is of real dimension at least $4n - m$, then \mathcal{M} is the Bruhat-Whitney-Cartan complexification of M . A result of H. Cartan [7] implies that the complexification of f vanishes on \mathcal{M} . In view of Lemma 3.3, the complexification of f , and hence f , admits a decomposition $a_1 r_1 + \dots + a_m r_m$.

The above argument shows especially that \mathcal{Q} is irreducible, generic, and of codimension m . Finally, we want to show that \mathcal{Q} is Levi-flat. Note that $\mathcal{Q} \setminus \cup_{j=1}^m \{\eta_j = 0\}$ is the disjoint union of graphs over the $(\xi_{m+1}, \dots, \xi_n, \eta)$ -space, which are given implicitly by

$$\begin{aligned} \sum_{k=1}^n a_{jk} \xi_k \eta_k &= it_j, & \sigma(j) &= j, \\ \xi_j \eta_{\sigma(j)} &= t_j + it_{\sigma(j)}, & \xi_{\sigma(j)} \eta_j &= -(t_j - it_{\sigma(j)}), \quad \sigma(j) > j \end{aligned}$$

with $t_j \in \mathbb{R}$. Hence, $\mathcal{Q} \setminus \cup_{j=1}^m \{\eta_j = 0\}$ is smooth and Levi-flat. Since Q_c is generic, it is contained in the closure of $\mathcal{Q} \setminus \cup_{j=1}^m \{\xi_j = 0\}$. Therefore, Q_c is Levi-flat. This completes the proof of the proposition. \square

Before we prove the first main result of this section, we should recall the Segre varieties associated to a real variety [15]. Let M be a germ of real analytic variety at $0 \in \mathbb{C}^n$ defined by $r_1(z, \bar{z}) = \dots = r_m(z, \bar{z}) = 0$, where r_j are real power series convergent on a polydisc $\Delta_\epsilon^n \times \Delta_\epsilon^n \subset \mathbb{C}^n \times \mathbb{C}^n$. Then the Segre varieties are the family of complex varieties

$$Q_w = \{z \in \Delta_\epsilon^n : r_1(z, \bar{w}) = \dots = r_m(z, \bar{w}) = 0\}, \quad w \in \Delta_\epsilon^n.$$

For a fixed $w \in \Delta_\epsilon^n$, the Segre variety Q_w could be an empty set, or the whole polydisc Δ_ϵ^n , or a proper subvariety of dimension between 0 and $n - 1$. For our applications of Segre varieties, we shall only deal with real analytic sets M that admit defining functions r_1, \dots, r_m such that germs of real analytic function vanishing on M are always generated by r_1, \dots, r_m . This implies that the Segre varieties Q_w are independent of the choice of such defining functions r_j , except that a possible smaller polydisc $\Delta_\epsilon^n \times \Delta_\epsilon^n$ has to be chosen for a given set of such defining functions. Also, note that, given two real analytic varieties M, M' admitting such defining functions and a biholomorphic mapping ϕ sending M onto M' , ϕ must send Segre varieties of M into Segre varieties of M' (by shrinking the polydiscs on which M, M' are defined if necessary). Finally, we should mention that if a germ of complex variety V at w is contained in M , then V is also contained in the Segre variety Q_w of M .

Proof of Theorem 3.1. — Following [6], we start with the dilation M^ϵ defined by

$$r_j^\epsilon(\xi, \eta, \bar{\xi}, \bar{\eta}) \equiv r_j(\epsilon\xi, \epsilon\eta, \epsilon\bar{\xi}, \epsilon\bar{\eta})/\epsilon^2 = 0, \quad j = 1, \dots, m.$$

Write

$$r_j^\epsilon(\xi, \eta, \bar{\xi}, \bar{\eta}) = q_j(\xi, \eta, \bar{\xi}, \bar{\eta}) + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}, \epsilon).$$

Parameterize

$$\gamma^\epsilon = M^\epsilon \cap \{\eta_j = 1, \quad j \leq m; \quad \xi_j = \eta_j = 0, \quad j > m\}$$

by $z = \gamma^\epsilon(t) = (\xi(t), \eta(t))$ ($t \in \mathbb{R}^m$) with

$$\left. \begin{aligned} \sum_{k=1}^m a_{jk} \xi_k(t) &= it_j + \alpha_j(t, \epsilon), \quad \alpha_j = \bar{\alpha}_j \quad \text{if } \sigma(j) = j, \\ \xi_j(t) &= t_j + it_{\sigma(j)} + \alpha_j(t, \epsilon) \\ \xi_{\sigma(j)}(t) &= -(t_j - it_{\sigma(j)} - \bar{\alpha}_j(t, \epsilon)) \end{aligned} \right\} \text{if } \sigma(j) > j.$$

Then α_j should satisfy

$$2\alpha_j + R_j(\xi(t), \eta(t), \bar{\xi}(t), \bar{\eta}(t), \epsilon) = 0, \quad \sigma(j) \geq j.$$

The implicit function theorem implies that the above equations have a unique solution

$$\alpha_j = \alpha_j(t, \epsilon), \quad \sigma(j) \geq j$$

with $\alpha_j|_{\epsilon=0} \equiv 0$.

The Segre variety $Q_{\gamma^\epsilon(t)}$ of M^ϵ is defined by

$$(3.17) \quad \left. \begin{aligned} -it_j + \alpha_j(t, \epsilon) + \sum_{k=1}^n a_{jk} \xi_k \eta_k + R_j &= 0 \quad \text{if } \sigma(j) = j, \\ -t_j - it_{\sigma(j)} + \alpha_j(t, \epsilon) + \xi_j \eta_{\sigma(j)} + R_j &= 0 \\ t_j - it_{\sigma(j)} + \bar{\alpha}_j(t, \epsilon) + \xi_{\sigma(j)} \eta_j + R_{\sigma(j)} &= 0 \end{aligned} \right\} \quad \text{if } \sigma(j) > j$$

for $R_j = R_j(\xi, \eta, \bar{\gamma}^\epsilon(t), \epsilon)$. By applying the fixed-point theorem, one can solve (3.17) to get

$$t_j = ih_j(\xi, \eta, \epsilon), \quad j = 1, \dots, m,$$

where h_j are holomorphic functions of ξ, η, ϵ . It is easy to see that

$$(3.18) \quad h_j(\xi, \eta, 0) = \sum_{k=1}^m c_{jk} q_k(\xi, \eta), \quad \det(c_{jk}) \neq 0.$$

This implies that for small ϵ , all $h_j(\xi, \eta, \epsilon)$ are well-defined on the domain

$$D_s = \Delta_2^{2n} \cap \{|q_j(\xi, \eta)| < s, j = 1, \dots, m\},$$

where $s > 0$ is small and independent of ϵ . Clearly, $\gamma^\epsilon(t)$ is contained in D_s when ϵ, t are small. As mentioned after the proof of Lemma 3.4, D_s contains a neighborhood D' of the origin such that for each small ϵ the smooth locus of $M^\epsilon \cap D'$ is connected.

Note that $\partial(r_1^\epsilon, \dots, r_m^\epsilon)/\partial(\xi_1, \dots, \xi_m)$ is nonsingular near γ^ϵ . Hence, M^ϵ is smooth and of CR -dimension $2n - m$ near γ^ϵ . This implies that M^ϵ is Levi-flat near γ^ϵ ; in particular, the branch $Q'_{\gamma^\epsilon(t)}$ of $Q_{\gamma^\epsilon(t)}$, which passes through $\gamma^\epsilon(t)$, is contained in M^ϵ .

Next, we would like to show that $Q'_{\gamma^\epsilon(t)} \subset M^\epsilon$ sweep out an open subset of $(M^\epsilon)^* \cap D'$. To this end we take a polydisc $\Delta_{s'}^{2n} \subset D'$. Rewrite (3.17) as

$$\xi_j = \eta_j^{-1} \left(\sum_{k=1}^K b_{jk} (it_k - \alpha_k - R_k) + \sum_{k>K} b_{jk} \xi_k \eta_k \right), \quad \sigma(j) = j,$$

$$\left. \begin{aligned} \xi_j &= \eta_{\sigma(j)}^{-1}(t_j + it_{\sigma(j)} - \alpha_j - R_j) \\ \xi_{\sigma(j)} &= \eta_j^{-1}((it_{\sigma(j)} - t_j - \bar{\alpha}_j) - R_{\sigma(j)}) \end{aligned} \right\} \sigma(j) > j.$$

Applying the fixed-point theorem to the above equations, one can get a unique solution $\xi_j = p_j^\epsilon(\eta, \xi'', t)$ defined on $E^m \times \Delta_{s''}^{2n-2m}$ for $|t| < s''$, where $s'' < s'$ is a positive constant depending only on s' and

$$E = \{w \in \mathbb{C}: s'/2 < |w| < 3/2, |\arg w| < \pi/2\}.$$

This shows that when ϵ, t are sufficiently small, the branch $Q'_{\gamma^\epsilon(t)}$ contains the graph

$$G_t^\epsilon \subset \mathbb{C}^{2n}: \xi_j = p_j^\epsilon(\xi'', \eta, t), \quad (\eta, \xi'') \in E^m \times \Delta_{s''}^{2n-2m}.$$

Obviously, $G_t^\epsilon \cap \Delta_{s''}^{2n}$ is nonempty if ϵ, t are small, and $(\partial p^\epsilon / \partial t)(\eta, \xi'', t)$ is nonsingular on $E^m \times \Delta_{s''}^{2n-2m+1}$ for $\epsilon = 0$. This shows that $G^\epsilon \equiv \cup G_t^\epsilon \subset M^\epsilon$ is an immersed real manifold of dimension $2n - m$. Thus, $G^\epsilon \cap (M^\epsilon)^* \cap \Delta_{s''}^{2n}$ is nonempty. On the other hand, G_t^ϵ is contained in the Segre variety $Q'_{\gamma^\epsilon(t)}$ on which the holomorphic functions $h_j(\xi, \eta, \epsilon)$ are pure imaginary. Therefore, all $\Re\{h_j(\xi, \eta, \epsilon)\}$ vanish on a nonempty open subset of $(M^\epsilon)^* \cap \Delta_{s''}^{2n}$. The latter is, however, a totally real subspace of $(M^\epsilon)^* \cap D$ of maximal dimension. Therefore, the complexification of each $\Re\{h_j(\xi, \eta, \epsilon)\}$ vanishes on $M^\epsilon \cap D$. In return, $\Re\{h_j(\xi, \eta, \epsilon)\}$ vanishes on M^ϵ .

We now fix a small ϵ . In view of Proposition 3.5, we have

$$(3.19) \quad \Re\{h_j(\xi, \eta, \epsilon)\} = \sum d_{jk}(\xi, \eta, \bar{\xi}, \bar{\eta}) r_k^\epsilon(\xi, \eta, \bar{\xi}, \bar{\eta}), \quad j = 1, \dots, m.$$

In particular, $h_j(\xi, \eta, \epsilon) - h_j(0, 0, \epsilon) = O(|(\xi, \eta)|^2)$. From (3.18), we know that $(d_{jk})(0)$ is invertible. Let (d'_{jk}) be the inverse of the constant matrix $(d_{jk})(0)$. Put

$$h_j^*(\xi, \eta) = \sum_{k=1}^m d'_{jk} h_k(\xi, \eta, \epsilon).$$

Then q_j is the quadratic form of h_j^* . Notice that the right-hand side of (3.19) is real; in particular, its quadratic form is real also. Hence, $d_{jk}(0) = \bar{d}_{j\sigma(k)}(0)$. From $\sigma^2 = \text{Id}$, it follows that $d'_{jk} = \bar{d}'_{\sigma(j)k}$. On M^ϵ , we have $\bar{h}_j(\bar{\xi}, \bar{\eta}, \epsilon) = -h_j(\xi, \eta, \epsilon)$. Now a straightforward computation shows that $h_j^*(\xi, \eta, \epsilon) = -\bar{h}_{\sigma(j)}^*(\bar{\xi}, \bar{\eta}, \epsilon)$ on M^ϵ . This shows that M^ϵ , and hence M , is of the form (3.7). The proof of the theorem is complete. \square

We now turn to the proof of Theorem 3.2. The rest of the section is to show that the Levi-flat analytic set defined by (3.9) is holomorphically equivalent to the set (3.10).

We shall use general holomorphic coordinates for the rest of section. For convenience, we set

$$z = (x, y), \quad x' = (x_1, \dots, x_m), \quad x'' = (x_{m+1}, \dots, x_n), \text{ etc,}$$

where $(x, y) = (x_1, \dots, x_n, y_1, \dots, y_n)$ are coordinates of \mathbb{C}^{2n} . In particular, the last $m - K$ equations in (3.9) are equivalent to the last $m - K$ ones of

$$(3.20) \quad r_j = x_j \bar{y}_{\sigma(j)} + \bar{x}_{\sigma(j)} y_j + R_j(z, \bar{z}) = 0, \quad j = 1, \dots, m.$$

The first K equations in (3.9) are transformed into the first K ones of (3.20) by the transformation

$$\xi_j = (1 + i)x_j + y_j, \quad \eta_j = (1 + i)x_j + iy_j, \quad j = 1, \dots, K.$$

Thus, M is given by (3.20) with σ of the form (3.1), and the reality condition (3.6) now reads

$$(3.21) \quad R_{\sigma(j)}(z, \bar{z}) = \bar{R}_j(\bar{z}, z).$$

Thus, Theorem 3.2 is reduced to the following.

THEOREM 3.6. — *Let $M \subset \mathbb{C}^{2n}$ be a Levi-flat analytic set given by (3.20)–(3.21). Then M is holomorphically equivalent to the set*

$$(3.22) \quad \mathcal{Q} \subset \mathbb{C}^{2n}: x_j \bar{y}_{\sigma(j)} + \bar{x}_{\sigma(j)} y_j = 0, \quad j = 1, 2, \dots, m.$$

We need the following.

PROPOSITION 3.7. — *Let $M \subset \mathbb{C}^{2n}$ be a real analytic set defined by (3.20)–(3.21) with σ of the form (3.1). Then M is generic, irreducible and of codimension m . Also, a germ of any real analytic function f vanishing on a topological component of M , of which the closure contains the origin, admits a decomposition $a_1 r_1 + \dots + a_m r_m$ for some complex-valued real analytic functions a_j with $\min\{\text{ord } a_j\} = \text{ord } f - 2$. Moreover, the set \mathcal{Q} , defined by (3.22), is Levi-flat.*

Proof. — The proof is almost the same as that of Proposition 3.5. Take a point $z^0 = (x^0, y^0)$ on \mathcal{Q} such that $x_j^0 \neq 0 \neq y_j^0$ for $j = 1, \dots, m$. Then $\partial(q_1, \dots, q_m)/\partial(x_1, \dots, x_m) \neq 0$ at z^0 . This shows that \mathcal{Q} has codimension at most m , and that \mathcal{Q} is generic if $\text{codim } \mathcal{Q} = m$. By the implicit function theorem, one then knows that M has codimension at most m , and it is generic if the codimension is m . Using Lemma 3.4, one shows that the complexification of M is the irreducible complex variety of codimension m given by $r_j(z, \bar{z}) = 0, j = 1, \dots, m$. Since M^* is a totally

real subspace in $(\mathcal{M})^*$ of maximal dimension, the complexification of f vanishes on \mathcal{M} . Lemma 3.3 says that the complexification, and hence f , can be decomposed into $a_1 r_1 + \dots + a_m r_m$ for some real analytic functions a_j .

Finally, $\mathcal{Q} \setminus \cup_{j=1}^m \{x_j = 0\}$ is the disjoint union of complex manifolds

$$y_j = it_j x_j, \quad 1 \leq j \leq m$$

with parameters $t_j \in \mathbb{C}$ satisfying $t_{\sigma(j)} = \bar{t}_j$. Therefore, \mathcal{Q} is Levi-flat. The proof of Proposition 3.7 is complete. \square

LEMMA 3.8. — *Let σ be a permutation of $1, \dots, m$, and let l_1, \dots, l_m be \mathbb{C} -linear functions on \mathbb{C}^{2n} satisfying*

$$(3.23) \quad x_j \overline{l_{\sigma(j)}(x, y)} + \bar{x}_{\sigma(j)} l_j(x, y) = 0.$$

Then $l_j(x, y) = c_j x_j$ and

$$(3.24) \quad c_{\sigma(j)} + \bar{c}_j = 0.$$

If $c, d \in \mathbb{C}^m$ satisfy (3.24) with $c - d \in (\mathbb{C}^*)^m$, equations

$$(3.25) \quad x_j^* = y_j - c_j x_j, \quad y_j^* = (y_j - d_j x_j)/(c_j - d_j), \quad 1 \leq j \leq m$$

define a nonsingular linear transformation with

$$(3.26) \quad x_j^* \bar{y}_{\sigma(j)}^* + \bar{x}_{\sigma(j)}^* y_j^* = x_j \bar{y}_{\sigma(j)} + \bar{x}_{\sigma(j)} y_j.$$

Proof. — By (3.23) it is clear that $l(x, y) = (c_1 x_1, \dots, c_m x_m)$ with c_j satisfying (3.24). Now, $c - d \in (\mathbb{C}^*)^m$ implies that the transformation (3.25) is nonsingular. In view of (3.24), the computation for (3.26) is straightforward. The details are left to the reader. \square

PROPOSITION 3.9. — *Let $M \subset \mathbb{C}^{2n}$ be a Levi-flat real analytic set defined by (3.20)–(3.21). Then in suitable holomorphic coordinates, M is given by*

$$(3.27) \quad x_j \bar{y}_{\sigma(j)} + \bar{x}_{\sigma(j)} y_j + R_j(z, \bar{z}) = 0, \quad j = 1, \dots, m$$

with

$$(3.28) \quad R_j(z, \bar{z}) = \sum_{k,l=1}^m x_k \bar{y}_l a_{jkl}(z, \bar{z}) + \sum_{k,l=1}^m \bar{x}_k y_l \bar{a}_{\sigma(j)kl}(\bar{z}, z) \\ + \sum_{k,l=1}^m x_k y_l b_{jkl}(z, \bar{z}'') + \sum_{k,l=1}^m \bar{x}_k \bar{y}_l \bar{b}_{\sigma(j)kl}(\bar{z}, z''),$$

$$(3.29) \quad b_{jkl}(z, 0'') \equiv b_{\sigma(j)kl}(\bar{z}, 0'') \equiv 0,$$

$$(3.30) \quad a_{j k \sigma(j)}(z, 0) \equiv a_{j j l}(0, \bar{z}) \equiv 0.$$

Proof. — Put $r_j^\epsilon(z, \bar{z}) = r_j(\epsilon z, \epsilon \bar{z})/\epsilon^2$ for $\epsilon > 0$. Let M^ϵ be the dilation of M defined by $r_1^\epsilon = \dots = r_m^\epsilon = 0$. Set

$$E^\epsilon: \partial r_1^\epsilon \wedge \dots \wedge \partial r_m^\epsilon \neq 0.$$

Obviously, $M^0 \cap E^0$ contains an m -dimensional totally real submanifold parameterized by

$$(3.31) \quad x' = (1, \dots, 1), \quad x'' = y'' = (0, \dots, 0),$$

$$(3.32) \quad y_j = y_j^0(t) = \begin{cases} it_j, & \sigma(j) = j, \\ t_j + it_{\sigma(j)} \equiv -\bar{y}_{\sigma(j)}, & \sigma(j) > j \end{cases}$$

for $t_j \in \mathbb{R}$. By the implicit function theorem, $M^\epsilon \cap E^\epsilon$ also contains an m -dimensional totally real submanifold $\gamma^\epsilon(t)$ parameterized by (3.31) and

$$y_j = y_j^\epsilon(t) = y^0(t) + \alpha_j(t, \epsilon), \quad -1 < t_j < 1, \quad j = 1, \dots, m,$$

where $\alpha_j(t)$ are real analytic functions in t and ϵ , satisfying $\alpha_{\sigma(j)}(t, \epsilon) = \bar{\alpha}_j(t, \epsilon)$. The Segre variety $Q_{\gamma^\epsilon(t)}$ is defined by

$$(3.33) \quad y_j + x_j \bar{y}_{\sigma(j)}(t) + R_j(z, \gamma^\epsilon(t), \epsilon) = 0, \quad j = 1, \dots, m.$$

Note that $R_j \equiv 0$ when $\epsilon = 0$. By the fixed-point theorem, $Q_{\gamma^\epsilon(t)} \equiv Q_{\gamma^\epsilon(t)} \cap \{|x_j| < 2, |y_j| < 1/2\}$ is a graph of the form

$$y_j = h_j^\epsilon(x, y'', t), \quad |x_j| < 2, \quad j = 1, \dots, m$$

for small ϵ and t . In particular, $Q_{\gamma^\epsilon(t)}$ is contained in M^ϵ .

We would like to show that $0 \in Q_{\gamma^\epsilon(t)}$. From (3.32), it follows that for small ϵ and generic t , $Q_{\gamma^\epsilon(t)}$ intersects a fixed $Q_{\gamma^\epsilon(t_0)}$ transversely along $S = Q_{\gamma^\epsilon(t)} \cap Q_{\gamma^\epsilon(t_0)}$. Fix a generic t . S contains points which can be arbitrarily close to the origin as ϵ tends to 0, and near the origin S is a closed complex submanifold of dimension $2n - 2m$. To show that $0 \in Q_{\gamma^\epsilon(t)}$ it suffices to verify that S contains the origin. Since $Q_{\gamma^\epsilon(t)}$ and $Q_{\gamma^\epsilon(t_0)}$ are complex submanifolds of codimension $m \leq n$, the span of their tangent spaces at each point in S is the whole space \mathbb{C}^{2n} . Therefore, the differentials dr_j vanish on S . On the other hand, we have

$$\partial_{x_j} r_j = \bar{y}_{\sigma(j)} + O(|z|^2), \quad \partial_{y_j} r_j = \bar{x}_{\sigma(j)} + O(|z|^2), \quad j = 1, \dots, m.$$

Near the origin the above equations define a (connected) smooth real submanifold S' of dimension $2n - 2m$. Obviously, S' contains the origin.

Since $S(\subset S')$ and S' have the same real dimension and both are closed sets, they must coincide near the origin. This shows that S , and hence $Q_{\gamma^\epsilon(t)}$, contains 0. Notice that the union of $Q_{\gamma^\epsilon(t)}$ contains an open subset U of $(M^\epsilon)^*$ with $\gamma^\epsilon \subset U$ and that $0 \in Q_w$ for $w \in U$. We conclude that Q_0 contains the open subset U of $(M^\epsilon)^*$; consequently, the complex variety Q_0 is the whole space \mathbb{C}^{2n} , since M^ϵ is generic. Therefore, $r_j(\xi, \eta, 0, 0) \equiv 0$ for $j = 1, \dots, m$.

Our next step is to find new coordinates for a fixed M^ϵ so that it has two Segre varieties forming part of coordinate subspaces. To this end, we fix a small ϵ so that M^ϵ contains two Segre varieties $Q_{\gamma^\epsilon(t')}$ and $Q_{\gamma^\epsilon(t'')}$ intersecting transversely at the origin. As shown above, $Q_{\gamma^\epsilon(t')}$, $Q_{\gamma^\epsilon(t'')}$ are given by $y' = g(x, y'')$ and $y' = h(x, y'')$, respectively. Moreover, $g(0) = h(0) = 0$. Since $Q_{\gamma(t')}$ is contained in M , then

$$r_j(x, g(x, y''), y'', \bar{x}, \bar{g}(\bar{x}, \bar{y}''), \bar{y}'') \equiv 0$$

for $j = 1, \dots, m$. Hence,

$$[g_j(x, y'')]_1 \bar{x}_{\sigma(j)} + \overline{[g_{\sigma(j)}(x, y'')]_1} x_j = 0.$$

This means that $[g_1]_1, \dots, [g_m]_1$ satisfy (3.23), so $[g_j]_1(x, y) = c_j x_j$. Similarly, $[h_j]_1(x, y) = d_j x_j$. Define the new coordinates

$$x_j^* = y_j - g_j(x, y''), \quad y_j^* = (y_j - h_j(x, y'')) / (c_j - d_j), \quad j = 1, \dots, m,$$

for which the linear terms are of the form (3.25). Now $M \equiv M^\epsilon$ contains two Segre varieties $x_1^* = \dots = x_m^* = 0$ and $y_1^* = \dots = y_m^* = 0$. For brevity, we replace z^* by z . In view of $r_j(z, 0) = r_j(0, \bar{z}) \equiv 0$ and the reality condition (3.21), one sees that M is given by (3.27) with

$$\begin{aligned} R_j(z, \bar{z}) &= \sum_{k,l=1}^m x_k \bar{y}_l a'_{jkl}(z, \bar{z}) + \sum_{k,l=1}^m \bar{x}_k y_l a''_{jkl}(\bar{z}, z) \\ &+ \sum_{k,l=1}^m x_k y_l b'_{jkl}(z, \bar{z}'') + \sum_{k,l=1}^m \bar{x}_k \bar{y}_l b''_{jkl}(\bar{z}, z''), \end{aligned}$$

where $a'_{jkl}(z, \bar{z})$ is independent of $x_1, \dots, x_{k-1}, \bar{y}_1, \dots, \bar{y}_{l-1}$, and $a''_{jkl}(\bar{z}, z)$ is independent of $\bar{x}_1, \dots, \bar{x}_{k-1}, y_1, \dots, y_{l-1}$. Obviously b'_{jkl}, b''_{jkl} , independent of $x_1, \dots, x_{k-1}, y_1, \dots, y_{l-1}$, are unique, and $b''_{\sigma(j)kl}(\bar{z}, z'') = \bar{b}'_{jkl}(\bar{z}, z'')$ by the reality condition $\bar{R}_j = R_{\sigma(j)}$. Put $a_{jkl} = (a'_{jkl} + \bar{a}''_{\sigma(j)kl}) / 2$. Then the reality condition $\bar{R}_j = R_{\sigma(j)}$ implies that R_j are given by (3.27)–(3.29). Note that $a_{jkl}(z, \bar{z})$ remains independent of $x_1, \dots, x_{k-1}, \bar{y}_1, \dots, \bar{y}_{l-1}$.

To achieve (3.30), we need to change coordinates one more time. We shall leave the Segre varieties $x_1 = \dots = x_m = 0$ and $y_1 = \dots = y_m = 0$

unchanged. Thus, we consider a holomorphic transformation

$$\psi: x_j \rightarrow x_j + u_j(z), \quad y_j \rightarrow y_j + v_j(z), \quad j = 1, \dots, m$$

with

$$(3.34) \quad u_j|_{x'=0} = 0, \quad v_j|_{y'=0} = 0, \quad u_j, v_j = O(2).$$

Put $\psi^{-1}(M)$ in the form of (3.27)–(3.29) with a_{jkl}^* in place of a_{jkl} . Thus, from $r_j^* = r_j \circ \psi$ one obtains

$$(3.35) \quad a_j^*(z, \bar{z}) + \bar{a}_{\sigma(j)}^*(\bar{z}, z) = \bar{y}_{\sigma(j)} u_j + x_j \bar{v}_{\sigma(j)} + y_j \bar{u}_{\sigma(j)} + \bar{x}_{\sigma(j)} v_j + u_j \bar{v}_{\sigma(j)} + \bar{u}_{\sigma(j)} v_j + a_j(z^*, \bar{z}^*) + \bar{a}_{\sigma(j)}(\bar{z}^*, z^*)$$

for $z^* = (x' + u, x'', y' + v, y'')$ and

$$a_j(z, \bar{z}) = \sum x_k \bar{y}_l a_{jkl}(z, \bar{z}), \quad a_j^*(z, \bar{z}) = \sum x_k \bar{y}_l a_{jkl}^*(z, \bar{z}).$$

Comparing the coefficients of $\bar{y}_{\sigma(j)} x^\alpha y^\beta$ and $x_j \bar{x}^\alpha \bar{y}^\beta$ in (3.35), respectively, one gets

$$\sum_{k=1}^m x_k a_{jk\sigma(j)}^*(z, 0) = u_j + \sum_{k=1}^m (x_k + u_k(z)) a_{jk\sigma(j)}(x + u, y + v, 0),$$

$$\sum_{l=1}^m \bar{y}_l a_{jjl}^*(0, \bar{z}) = \bar{v}_{\sigma(j)} + \sum_{l=1}^m (\bar{y}_l + \bar{v}_l(\bar{z})) a_{jjl}(0, \bar{x} + \bar{u}, \bar{y} + \bar{v}).$$

Now, the implicit function theorem implies that there exist unique $u, v = O(2)$ satisfying

$$(3.36) \quad \sum_{k=1}^m x_k a_{jk\sigma(j)}^*(z, 0) = \sum_{l=1}^m \bar{y}_l \bar{a}_{jjl}^*(0, \bar{z}) = 0, \quad j = 1, \dots, m.$$

Obviously, the unique solution (u, v) satisfies (3.34) also. Note that in the decompositions (3.28), $a_{jk\sigma(j)}^*(z, 0)$ is independent of x_1, \dots, x_{k-1} and $a_{jjl}^*(0, \bar{z})$ is independent of $\bar{y}_1, \dots, \bar{y}_{l-1}$. Now identities (3.36) yield (3.30). This completes the proof of the proposition. □

Proof of Theorem 3.6. — Following [6], we want to show that the Levi-flat set M , defined by (3.27)–(3.30), is the one given by (3.22).

Consider a real manifold in M parameterized by $x' = (1, \dots, 1)$, $x'' = y'' = 0$ and

$$y_j(t) = \begin{cases} it_j + \alpha_j(t), & \sigma(j) = j, \\ t_j + it_{\sigma(j)} + \alpha_j(t), & \sigma(j) > j, \\ -t_{\sigma(j)} + it_j + \alpha_j(t), & \sigma(j) < j \end{cases}$$

with $t \in \mathbb{R}^m$, $\alpha_j(0) = 0$ and $\alpha_{\sigma(j)}(t) = \bar{\alpha}_j(t)$. By the implicit function theorem, such α_j exist.

The Segre variety $Q_{\gamma(t)}$ is defined by

$$y_j + \bar{y}_{\sigma(j)}(t)x_j + R_j(z, \bar{\gamma}(t)) = 0, \quad j = 1, \dots, m.$$

By rescaling the coordinates, one may assume that R_j are small functions. Hence, $Q_{\gamma(t)}$, intersecting with a fixed neighborhood of the origin, is a smooth complex submanifold given by

$$y_j = h_j(x, y'', t), \quad h_j(x, y'', 0) = h_j(0, t) = 0$$

for $j = 1, \dots, m$, where the last identity comes from $0 \in Q_{\gamma(t)}$.

Since $Q_{\gamma(t)}$ is contained in M , we have

$$(3.37) \quad 0 \equiv x_j \bar{h}_{\sigma(j)}(\bar{x}, \bar{y}'', t) + \bar{x}_{\sigma(j)} h_j(x, y'', t) \\ + \sum \{x_k \bar{h}_l(\bar{x}, \bar{y}'', t) a_{jkl}(z, \bar{z}) + \bar{x}_k h_l(x, y'', t) \bar{a}_{\sigma(j)kl}(\bar{z}, z)\} \\ + \sum \{x_k h_l(\bar{x}, \bar{y}'', t) b_{jkl}(z, z'') + \bar{x}_k \bar{h}_l(\bar{x}, \bar{y}'', t) \bar{b}_{\sigma(j)kl}(\bar{z}, z'')\}$$

for $z = (x, h(x, y''), y'')$. Expanding the right-hand side of (3.37) as power series in $x, y'', \bar{x}, \bar{y}''$ and collecting quadratic terms yields

$$[h_j]_1(x, y'', t) = c_j(t)x_j, \quad c_{\sigma(j)}(t) + \bar{c}_j(t) = 0.$$

Now looking at (3.37), we see that the homogeneous terms $x^\alpha y''^\beta \bar{x}_{\sigma(j)}$ for $|\alpha| + |\beta| = 2$ give us

$$[h_j]_2(x, y'', t) = - \sum x_k \bar{c}_{\sigma(j)}(t) [a_{jk\sigma(j)}]_1(x, [h]_1(x, y'', t), y'', 0) \\ - \sum [h_l]_1(x, y'', t) [\bar{a}_{\sigma(j)\sigma(j)l}]_1(0, x, [h]_1(x, y'', t), y'').$$

From (3.30) it follows that $[h_j]_2 \equiv 0$. Assuming for the sake of induction that $[h_j]_2 = \dots = [h_j]_N = 0$ for $1 \leq j \leq m$, one gets from (3.30) and (3.37) that

$$[h_j]_{N+1}(x, y'', t) = \sum x_k \bar{c}_{\sigma(j)}(t) [a_{jk\sigma(j)}]_N(x, y, 0) \\ + \sum [h_l]_1(x, y'', t) [\bar{a}_{\sigma(j)\sigma(j)l}]_N(0, x, y) \equiv 0$$

for $y' = [h]_1(x, y'', t)$. This shows that $h_j(x, y'', t)$ are linear in x , and that $Q_{\gamma(t)}$ is given by

$$y_j = c_j(t)x_j, \quad c_{\sigma(j)}(t) + \bar{c}_j(t) = 0, \quad j = 1, \dots, m.$$

Therefore, M contains a portion of Q ; consequently, M and Q coincide as they are irreducible. The proof of the theorem is complete. \square

4. Invariants of Levi-flat sets and symplectic transformations.

Results in the preceding section allow one to choose much simpler defining functions of Levi-flat sets in suitable holomorphic coordinates. In this section we shall use (more restrictive) symplectic coordinates to study the defining functions for the Levi-flat sets that are invariant under a holomorphic symplectic mapping.

This section is divided into two parts: We shall first complete the proofs of Theorem 1.1 and Theorem 1.3 after we obtain the first-integrals from Levi-flat invariant sets. We then turn to a meromorphic eigenfunction problem, arising from Levi-flat sets, and formulate Theorem 4.9, from which Theorem 1.2 follows. The proof of Theorem 4.9 is given in Section 5.

From now on, Levi-flat sets in \mathbb{C}^{2n} are of codimension n . We shall also take into account that in section 3 we ignored the symplectic invariants μ_j and used a simpler permutation σ defined by (3.1). Therefore, we need to return to the 4 types of singularities, which involve μ_j (and a more general σ in case of (4.21) below).

Let σ be a permutation of $1, \dots, n$ satisfying

$$(4.1) \quad \sigma(j) = j, \quad 1 \leq j \leq K; \quad \sigma(j) \neq j = \sigma^2(j), \quad j > K.$$

We first consider $Q \subset \mathbb{C}^{2n}$ defined by

$$(4.2) \quad q_j(\xi, \eta) + \mu_j \bar{q}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, n$$

with

$$(4.3) \quad \begin{aligned} \mu_j &= 1, \quad 1 \leq j \leq K; & \mu_{\sigma(j)} &= \bar{\mu}_j, \quad |\mu_j| = 1, \quad j > K; \\ q_j(\xi, \eta) &= \begin{cases} \sum_{k=1}^n a_{jk} \xi_k \eta_k, & 1 \leq j \leq K, \\ \xi_j \eta_{\sigma(j)}, & K < j \leq K + 2L = n \end{cases} \end{aligned}$$

with $(a_{ij})_{1 \leq i, j \leq K}$ of rank K . Note again that we allow $K = 0$, or n as special cases.

With the above notations, consider a Levi-flat real analytic set of the form

$$(4.4) \quad q_j(\xi, \eta) + \mu_j \bar{q}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, n$$

with

$$(4.5) \quad R_{\sigma(j)}(\xi, \eta, \bar{\xi}, \bar{\eta}) = \mu_j \bar{R}_j(\bar{\xi}, \bar{\eta}, \xi, \eta).$$

Let $\xi' = \sqrt{\mu}\xi \equiv (\sqrt{\mu_1}\xi_1, \dots, \sqrt{\mu_n}\xi_n), \eta' = \eta$. Then equations (4.4)–(4.5), after multiplied by $\sqrt{\mu_j}$, are of the form

$$\tilde{q}_j(\xi', \eta') + \bar{q}_{\sigma(j)}(\bar{\xi}', \bar{\eta}') + \tilde{R}_j(\xi', \eta', \bar{\xi}', \bar{\eta}') = 0$$

with

$$\tilde{q}_j(\xi', \eta') = \sqrt{\mu_j}q_j(\sqrt{\mu}\xi', \eta'), \quad \tilde{R}_j(\xi', \eta', \bar{\xi}', \bar{\eta}') = \sqrt{\mu_j}R_j(\sqrt{\mu}\xi', \eta').$$

Now Theorem 3.1 says that the set is actually given by

$$\tilde{q}_j(\xi', \eta') + \bar{q}_{\sigma(j)}(\bar{\xi}', \bar{\eta}') + \tilde{H}_j(\xi', \eta') + \bar{H}_{\sigma(j)}(\bar{\xi}', \bar{\eta}') = 0.$$

Returning to the original coordinates and putting

$$H_j(\xi, \eta) = \sqrt{\mu_j}\tilde{H}_j(\sqrt{\mu}\xi, \eta),$$

one sees that the set is of the form

$$(4.6) \quad M: r_j = q_j(\xi, \eta) + \mu_j\bar{q}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) + H_j(\xi, \eta) + \mu_j\bar{H}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) = 0$$

for $j = 1, \dots, n$, where σ, q_j , and μ_j are given by (4.1) and (4.3), and $H_j(\xi, \eta) = O(3)$ are holomorphic.

THEOREM 4.1. — *Let M, H_j, q_j be as in (4.6). Let φ be a holomorphic symplectic mapping defined by (1.1)–(1.2) with $\varphi(M) = M$, and Φ a formal symplectic transformation such that $\Phi\varphi\Phi^{-1}$ is in the Birkhoff normal form (1.3). Then $|\lambda_{\sigma(j)}| = |\lambda_j|$ and*

$$(q_j + H_j) \circ \Phi^{-1}(\xi, \eta) = q_j(\xi, \eta) + \hat{H}_j(\xi\eta, \xi''\eta''), \quad j = 1, \dots, n$$

with $\hat{H}_j(\xi\eta, \xi''\eta'') = O(|(\xi, \eta)|^4)$ and $\xi''\eta'' = (\xi_{K+1}\eta_{\sigma(K+1)}, \dots, \xi_m\eta_{\sigma(m)})$.

Anticipating the proof of Theorem 4.1, we first prove Theorem 1.1 and Theorem 1.3.

Proof of Theorem 1.1. — Let M be the Levi-flat real analytic set defined by

$$\Re \left\{ \sum_{k=1}^n a_{jk}\xi_k\eta_k \right\} + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, n,$$

where the matrix (a_{jk}) is of rank n and $R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = O(3)$ are real-valued convergent power series. In Section 3 (Theorem 3.1) we have simplified the equation of M , (from now on) given by

$$\Re \{q_j(\xi, \eta) + H_j(\xi, \eta)\} = 0, \quad j = 1, \dots, n$$

with $H_j(\xi, \eta) = O(3)$ holomorphic, and $q_j = \sum_{k=1}^n a_{jk}\xi_k\eta_k$.

Theorem 4.1 establishes that if a symplectic mapping φ , given by (1.1)–(1.2) leaves the set M invariant, and if Φ is a formal symplectic transformation that puts φ into its Birkhoff normal form, then $(q_j + H_j) \circ \Phi^{-1}$ are power series which depend only on the products $\xi_j \eta_j$. (Indeed under our present hypotheses q_j depend only on $\xi \eta$ and σ is just the identity.) These power series are thus invariant under the action of the formally normalized mapping. Hence the functions $q_j + H_j$ are invariant, under the action of φ . Since $(a_{jk})_{1 \leq j, k \leq n}$ is nonsingular, a theorem of Vey [18] says that φ is normalizable by convergent transformations. The proof of Theorem 1.1 is complete. \square

Proof of Theorem 1.3. — Let r be the defining function of M , and q the quadratic form of r . Note that the order of r might be one, which can however be ruled out by applying the argument below to r^2 .

Since M is Levi-flat and $\mathcal{Q}: q = 0$ is a hypersurface in \mathbb{C}^2 , then \mathcal{Q} is also Levi-flat [6]. Next, we want to show that \mathcal{Q} is invariant under Λ . Consider first the case that M is the union of two distinct smooth real hypersurfaces. In this case, $r = r_1 r_2$ with $r_1(0) = r_2(0) = 0$. Since $r \circ \varphi$ vanishes on $r_1 = 0$ and on $r_2 = 0$, then r_1, r_2 divide $r \circ \varphi$. In particular, $r \circ \varphi = ur$. Thus, $q \circ \Lambda = u(0)q$, i.e., $\Lambda(\mathcal{Q}) = \mathcal{Q}$. Next, we assume that M is irreducible. Then either r is irreducible, or $r = ur_1^2$ with $r_1(0) = 0$. When r is irreducible, $r(\xi, \eta, \bar{\xi}, \bar{\eta})$ is also irreducible as a germ of holomorphic function in $\xi, \eta, \bar{\xi}, \bar{\eta}$, because M is of codimension 1 (e.g., see [6]). Since $r \circ \varphi$ vanishes on M^c , then r divides $r \circ \varphi$, which implies that $\Lambda(\mathcal{Q}) = \mathcal{Q}$. When $r = ur_1^2$, it is clear that r_1 divides $r_1 \circ \varphi$. Denote by $a\xi + b\eta$ the linear part of r_1 . Then $a\lambda\xi + b\bar{\lambda}\eta = u(0)(a\xi + b\eta)$. Since $u(0)$ is real, the above identity contradicts that not all a, b are zero. (Note that the contradiction also implies that r starts with quadratic terms, as claimed at the beginning of the proof.) Therefore, \mathcal{Q} is invariant under Λ .

Applying the above argument to \mathcal{Q} , one sees that all germs of real analytic functions vanishing on \mathcal{Q} are divisible by q , i.e., that $\mathcal{I} = \mathcal{I}_{\mathcal{Q}}$ is spanned by the real quadratic form q . Hence, \mathcal{I} is contained in one of E_{ij} and E_1 . Since the former contain no Λ -invariant \mathbb{C} -linear subspace of dimension one, then \mathcal{I} is contained in E_1 . Hence, we get $q(\xi, \eta, \bar{\xi}, \bar{\eta}) = \Re\{a\xi\eta\} + b|\xi|^2 + c|\eta|^2$. If $b = c = 0$, Theorem 1.1 says that φ is normalizable by holomorphic symplectic transformations.

If b or c is not zero, a result in [6] says that $a = 0$. Thus, we see that

M is given by

$$(4.7) \quad b|\xi|^2 + c|\eta|^2 + R(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad b \cdot c < 0, \quad b, c \in \mathbb{R}$$

with $R(\xi, \eta, \bar{\xi}, \bar{\eta}) = O(3)$ real-valued. By a change of symplectic coordinates, one may assume $b = -c$. Of course the proof of the theorem is complete, by applying Theorem 1.2. \square

Alternatively, one can prove the theorem without using Theorem 1.2 when M is given by (4.7), for which we shall prove the next result. \square

PROPOSITION 4.2. — *Let φ be a holomorphic symplectic mapping of \mathbb{C}^2 of the form*

$$\xi' = \lambda\xi + O(2), \quad \eta' = \lambda^{-1}\eta + O(2), \quad \lambda^2 \neq 1.$$

Let $M \subset \mathbb{C}^2$ be a Levi-flat real analytic hypersurface given by (4.7). Assume that as germs of real analytic sets, $\varphi(M) = M$. Then $|\lambda| = 1$, and there exists a single change of holomorphic symplectic coordinates that puts M into the complex cone $|\xi|^2 - |\eta|^2 = 0$ and linearizes φ simultaneously.

Proof. — Notice that for this proposition, we allow λ to be a root of unity. Also, it is obvious that $|\lambda| = 1$, since $r(\xi, \eta, \bar{\xi}, \bar{\eta}) = b|\xi|^2 + c|\eta|^2 + R(\xi, \eta, \bar{\xi}, \bar{\eta})$ is irreducible and $r \circ \varphi = dr$ for some analytic real function d .

Since M is Levi-flat, a theorem in [6] says that there is a holomorphic transformation ψ_1 such that $\psi_1(M)$ is the complex cone $|\xi|^2 - |\eta|^2 = 0$. (See also Theorem 3.6 in Section 3.) Next, we want to find a holomorphic mapping

$$\psi_2: \xi' = p(\xi, \eta)\xi, \quad \eta' = p(\xi, \eta)\eta$$

such that the Jacobian determinant of ψ_2 satisfies

$$(4.8) \quad \det(D\psi_2) = \{\det(D\psi_1)\} \circ \{\psi_1^{-1}\}.$$

Thus, $\psi = \psi_2 \circ \psi_1$ is symplectic, and $\psi(M)$ is still the cone $|\xi|^2 - |\eta|^2 = 0$. Write the right-hand side of (4.8) as $a_0^2(1 + u(\xi, \eta))$ with $a_0 \neq 0$ a constant, and $u(0) = 0$. We seek a solution of the form $p(\xi, \eta) = a_0(1 + v(\xi, \eta))$. Thus, (4.8) becomes

$$(4.9) \quad \xi v_\xi + \eta v_\eta + 2v = (\xi v_\xi + v)(\eta v_\eta + v) - \xi v_\xi \cdot \eta v_\eta + u.$$

Comparing coefficients, one sees that there exists a unique formal power series solution $v(\xi, \eta)$ with $v(0) = 0$. Let u_{ij}, v_{ij} be the coefficients of u, v , respectively. Put

$$u^* = \sum |u_{ij}| \xi^i \eta^j, \quad w = \sum (i + j + 2) |v_{ij}| \xi^i \eta^j.$$

Obviously, $\xi v_\xi, \eta v_\eta, \xi v_\xi + v, \eta v_\eta + v$ are majorized by w , for which we write $\xi v_\xi \prec w$, etc. Thus, (4.9) implies that

$$w \prec 2w^2 + u^*, \quad w(0) = u^*(0) = 0.$$

By the Cauchy majorant argument, w , and hence v , is convergent.

Now $\varphi^* = \psi\varphi\psi^{-1}$ is a holomorphic mapping sending the complex cone $|\xi|^2 - |\eta|^2 = 0$ into itself. In particular, φ^* sends a Segre variety of the cone into another Segre variety, i.e., φ^* sends a complex line (inside the cone) through the origin to another complex line. Hence, one first sees that φ^* leaves the eigenspaces of its linear part invariant. Thus, $\lambda \neq \bar{\lambda}$ implies that

$$\varphi^*: \xi \rightarrow f(\xi, \eta)\lambda\xi, \quad \eta \rightarrow g(\xi, \eta)\bar{\lambda}\eta$$

with $f(0) = g(0) = 1$. Furthermore, g/f is holomorphic near the origin, and remains constant on each complex line in the cone. Therefore, $f \equiv g$ on the cone, and hence near the origin in \mathbb{C}^2 . This shows that φ^* is of the form $(\xi, \eta) \rightarrow f(\xi, \eta)(\lambda\xi, \bar{\lambda}\eta)$ with $f(0) = 1$. Since $\det(\partial\varphi^*/\partial(\xi, \eta)) = 1$, we obtain $f(\xi, \eta) \equiv 1$. In particular, $\psi \circ \varphi \circ \psi^{-1} = \varphi^*$ is a linear transformation. □

We now turn to the proof Theorem 4.1. Let us first prove the following.

LEMMA 4.3. — *Let $\mathcal{Q} \subset \mathbb{C}^{2n}$ be defined by (4.2)–(4.3), and let h be a holomorphic homogeneous polynomial in ξ, η . Assume that h is real-valued on \mathcal{Q} . Then h is a polynomial in q_1, \dots, q_n .*

Proof. — By changing linear coordinates, we may assume that $\mu_j = -1$, and that $\sigma(j) = L + j$ for $j = K + 1, \dots, K + L$. Put $\xi = (\xi', \xi''), \eta = (\eta', \eta'')$ with

$$\begin{aligned} \xi'' &= (\xi_{K+1}, \dots, \xi_n) = (x_{K+1}, \dots, x_{K+L}, y_{K+1}, \dots, y_{K+L}), \\ \eta'' &= (\eta_{K+1}, \dots, \eta_n) = (z_{K+1}, \dots, z_{K+L}, w_{K+1}, \dots, w_{K+L}). \end{aligned}$$

Assume first that h depends only on ξ'', η'' . Expand $h(x, y, z, w) = \sum h_{abcd}x^a y^b z^c w^d$. On \mathcal{Q} , one has $w_j = \bar{y}_j \bar{z}_j x_j^{-1}$ for $j = K + 1, \dots, K + L$. Since h is real on \mathcal{Q} , then

$$\sum h_{abcd}x^{a-d}y^b z^c \bar{y}^d \bar{z}^d = \sum \bar{h}_{abcd}y^d z^d \bar{x}^{a-d} \bar{y}^b \bar{z}^c.$$

Fix a, b, c, d with $h_{abcd} \neq 0$. Comparing two sides of the above identity, one sees that there exist a', b', c', d' such that

$$(a - d, b, c, 0, d, d) = (0, d', d', a' - d', b', c').$$

Hence, $a = d$ and $b = c$. This shows that $h(\xi'', \eta'') = \sum h_{abba}(\xi''\eta''_\sigma)^{(a,b)}$.

For the general case, we shall prove by induction on l that there exist decompositions

$$(4.10) \quad h(\xi, \eta) = \sum_{|\beta| < l} c_{\alpha\beta}(\xi'', \eta'') \xi'^\alpha \eta'^\beta + \sum_{|\gamma| \geq l} c_\gamma(\xi''\eta''_\sigma) q_1^{\gamma_1} \cdots q_K^{\gamma_K}$$

for $l = 0, 1, \dots$. Fix ξ'', η'' satisfying equations (4.2) for $j > K$, and let ξ', η' satisfy

$$\sum_{k=1}^n a_{jk} \xi_k \eta_k = t_j, \quad j = 1, \dots, K$$

for $t_j \in \mathbb{R}$. Solving for η_1, \dots, η_K yields

$$(4.11) \quad \eta_j = \xi_j^{-1} b_j(t) + \xi_j^{-1} \sum_{k=K+1}^n c_{jk} \xi_k \eta_k, \quad j = 1, \dots, K,$$

where $t \rightarrow b(t) = (b_1(t), \dots, b_K(t))$ is a nonsingular linear transformation. The decompositions (4.10) are trivial for $l > \deg h$. Assuming that the decompositions (4.10) hold for $l > k$, we want to show that (4.10) for $l = k$ is valid. From (4.10)–(4.11) one sees that terms of order k in t are given by

$$\sum_{|\beta|=k} c_{\alpha\beta}(\xi'', \eta'') \xi'^{\alpha-\beta} b^\beta(t).$$

For fixed ξ'', η'' , the above summation is real-valued for $t \in \mathbb{R}^K$ and $\xi' \in \mathbb{C}^K$. Since the transformation $t \rightarrow b(t)$ is nonsingular, we know that on \mathcal{Q}

$$(4.12) \quad \begin{aligned} c_{\alpha\beta}(\xi'', \eta'') &= 0, \quad \alpha \neq \beta, |\beta| = k, \\ \Im \left\{ \sum_{|\beta|=k} B_\alpha^\beta c_{\beta\beta}(\xi'', \eta'') \right\} &= 0, \quad |\alpha| = k, \end{aligned}$$

where B_α^β are the coefficients of the linear transformation

$$(4.13) \quad t^\beta \rightarrow b^\beta(t) = \sum_{|\alpha|=k} B_\alpha^\beta t^\alpha, \quad |\beta| = k.$$

Now (4.12) implies that for α with $|\alpha| = k$, $\sum_{|\beta|=k} B_\alpha^\beta c_{\beta\beta}(\xi'', \eta'')$ are polynomials in $\xi''\eta''_\sigma$. Since the linear transformation (4.13) is nonsingular, then $c_{\beta\beta}(\xi'', \eta'')$ are power series in $\xi''\eta''_\sigma$ for $|\beta| = k$. Rewrite (4.10) as

$$h(\xi, \eta) = \sum_{|\beta| < k} \tilde{c}_{\alpha\beta}(\xi'', \eta'') \xi'^\alpha \eta'^\beta + \sum_{|\gamma| \geq k} c_\gamma(\xi''\eta''_\sigma) q_1^{\gamma_1} \cdots q_K^{\gamma_K}.$$

By induction, the proof of the lemma is complete. □

To state the next result, we need some notation. For a power series $f(\xi, \eta, \bar{\xi}, \bar{\eta})$, denote by $P_0 f$ the sum of terms in the power series expansion of f , which are not of the form $(\xi\eta)^a(\bar{\xi}\bar{\eta})^b(\xi''\eta''_\sigma)^c(\bar{\xi}''\bar{\eta}''_\sigma)^d$.

LEMMA 4.4. — *Let r_1, \dots, r_n be the formal power series in (4.6) with q_j being (4.3). Let d_1, \dots, d_n be power series in $\xi, \eta, \bar{\xi}, \bar{\eta}$ of order at least k . Assume that*

$$[P_0 d_1]_k q_1 + \dots + [P_0 d_n]_k q_n = 0.$$

Then there exist formal power series $\tilde{d}_1, \dots, \tilde{d}_n$ such that

$$(4.14) \quad d_1 r_1 + \dots + d_n r_n = \tilde{d}_1 r_1 + \dots + \tilde{d}_n r_n$$

with $\text{ord } \tilde{d}_j \geq k$ and $[P_0 \tilde{d}_j]_k = 0$.

Proof. — Choose \mathbb{C} -linear combinations r_1^*, \dots, r_K^* of r_1, \dots, r_K so that

$$r_j^*(\xi, \eta, \bar{\xi}, \bar{\eta}) = \xi_j \eta_j + \sum_{k>K} b_{jk} \xi_k \eta_k + \sum_{k=1}^n c_{jk} \overline{\xi_k \eta_k} + O(3).$$

It suffices to prove the lemma for

$$r_1^*, \dots, r_K^*, r_{K+1}, \dots, r_n.$$

For brevity, we drop the superscript of r_j^* . For $j > 1$, decompose $[d_j]_k = u_j q_1 + [u'_j]_k$ so that $[u'_j]_k$ contains only terms $\xi^{\alpha_1} \eta^{\beta_1} \bar{\xi}^{\gamma_1} \bar{\eta}^{\delta_1}$ with $\alpha_1 = 0$, or $\beta_1 = 0$. Put

$$u'_j = d_j - u_j r_1, \quad j > 1.$$

Then

$$d_1 r_1 + \dots + d_n r_n = \tilde{d}_1 r_1 + u'_2 r_2 + \dots + u'_n r_n.$$

Comparing terms $\xi^{\alpha_1} \eta^{\beta_1} \bar{\xi}^{\gamma_1} \bar{\eta}^{\delta_1}$ with $\alpha_1, \beta_1 \geq 1$, one sees that $[P_0 \tilde{d}_1]_k = 0$, and hence

$$[P_0 u'_2]_k q_2 + \dots + [P_0 u'_n]_k q_n = 0.$$

Inductively, one determines $\tilde{d}_2, \dots, \tilde{d}_K$ so that

$$d_1 r_1 + \dots + d_n r_n = \tilde{d}_1 r_1 + \dots + \tilde{d}_K r_K + u''_{K+1} r_{K+1} + \dots + u''_n r_n$$

and $[P_0 \tilde{d}_1]_k = \dots = [P_0 \tilde{d}_K]_k = 0$. To find \tilde{d}_{K+1} , we decompose $u''_j = v_j r_{K+1} + v'_j$ for $j > K + 1$ so that $[v'_j]_k$ contains only terms of the form $\xi^{\alpha_{K+1}} \eta^{\beta_{K+1}} \bar{\xi}^{\gamma_{K+1}} \bar{\eta}^{\delta_{K+1}}$ with $\alpha_{K+1} = 0$ or $\beta_{\sigma(K+1)} = 0$. Thus,

$$u''_{K+1} r_{K+1} + \dots + u''_n r_n = \tilde{d}_{K+1} r_{K+1} + v'_{K+2} r_{K+2} + \dots + v'_n r_n.$$

Comparing the terms of the form $\xi^\alpha \eta^\beta \bar{\xi}^\gamma \bar{\eta}^\delta$ with $\alpha_{K+1}, \beta_{\sigma(K+1)} \geq 1$ yields $[P_0 \bar{d}_{k+1}]_k = 0$. Now the proof of the lemma is complete by induction. \square

Proof of Theorem 4.1. — By Proposition 3.5, $r_1 \circ \varphi, \dots, r_n \circ \varphi$ are in the ideal generated by r_1, \dots, r_n . In particular,

$$q_j(\lambda \xi, \lambda^{-1} \eta) + \mu_j \bar{q}_{\sigma(j)}(\bar{\lambda} \xi, \bar{\lambda}^{-1} \eta) = \sum_{k=1}^n c_{jk} (q_k(\xi, \eta) + \mu_k q_{\sigma(k)}(\bar{\xi}, \bar{\eta}))$$

for some constants c_{jk} , where

$$\lambda \xi \equiv (\lambda_1 \xi_1, \dots, \lambda_n \xi_n), \quad \lambda^{-1} \eta \equiv (\lambda_1^{-1} \eta_1, \dots, \lambda_n^{-1} \eta_n).$$

Since $(a_{jk})_{1 \leq j, k \leq K}$ is nonsingular, then q_1, \dots, q_n are linearly independent. Now, a simple computation shows that $c_{jj} = 1$ for $j = 1, \dots, K$,

$$\lambda_j \lambda_{\sigma(j)}^{-1} = c_{jj} = \bar{\lambda}_{\sigma(j)} \bar{\lambda}_j^{-1}$$

for $j = K + 1, \dots, n$, and $c_{jk} = 0$ for $j \neq k$. In particular, $|\lambda_{\sigma(j)}| = |\lambda_j|$. Now, we have decompositions

$$(4.15) \quad \Delta r_j \equiv r_j \circ \varphi - \lambda_j \lambda_{\sigma(j)}^{-1} r_j = \sum D_{jk} r_k, \quad 1 \leq j \leq m$$

with

$$(4.16) \quad \min_k \{\text{ord } D_{jk}\} = \min \{\text{ord } \Delta r_j\} - 2 > 0.$$

By abuse of notation, we replace $\Phi \varphi \Phi^{-1}$ by φ , and $\Phi(M)$ by M , etc. Then M is of the form (4.6), while H_1, \dots, H_n are formal power series.

We need to show that

$$(4.17) \quad [P_0 H_j]_{d+1} = 0, \quad j = 1, \dots, n.$$

For the purpose of induction we shall also prove that there exist decompositions

$$(4.18) \quad r_j \circ \varphi - \lambda_j \lambda_{\sigma(j)}^{-1} r_j = \sum \tilde{D}_{jk} r_k, \quad j = 1, \dots, n$$

with

$$(4.19) \quad [P_0 \tilde{D}_{ij}]_{d-1} = 0.$$

Obviously, (4.17)–(4.19) hold for $d = 1$ when formulae (4.18) are replaced by (4.15). Assume for induction that $[\tilde{D}_{ij}]_k$ for $k < d$ have been determined such that (4.18)–(4.19) hold. We would like to show that (4.18)–(4.19) hold if d is replaced by $d + 1$. For brevity we shall temporarily replace $[H_j]_{d+2}$ by H_j , and $[D_{jk}]_d$ by D_{jk} , etc. Then by (4.17)–(4.19), terms of order $d + 2$ in (4.18) yield

$$(4.20) \quad P_0 K_j(\xi, \eta) + \mu_j P_0 \bar{K}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) = \sum P_0 D_{jk} \cdot [r_k]_2$$

with

$$K_j(\xi, \eta) = H_j(\lambda\xi, \lambda^{-1}\eta) - \lambda_j \lambda_{\sigma(j)}^{-1} H_j(\xi, \eta).$$

Obviously, the real and imaginary parts of the left-hand side of (4.20), i.e., $\Re\{P_0 K_j + \bar{\mu}_j P_0 K_{\sigma(j)}\}$ and $\Im\{-i P_0 K_j + i \bar{\mu}_j K_{\sigma(j)}\}$, vanish on \mathcal{Q} . From Lemma 4.3, it follows that K_j are polynomials in q_1, \dots, q_n ; in particular, $P_0 K_j = 0$. Now the non-resonance condition (1.2) yields $P_0 H_j = 0$, which gives us (4.17) with $d + 1$ in place of d .

We now have

$$\sum P_0[\tilde{D}_{jk}]_d \cdot [r_k]_2 = 0.$$

In view of Lemma 4.4, one can modify $[\tilde{D}_{ij}]_d$ so that (4.18)–(4.19) hold when d is replaced by $d + 1$. This completes the proof of the theorem. \square

The rest of section is to study the Levi-flat real analytic sets $M \subset \mathbb{C}^{2n}$ of the form

$$(4.21) \quad q_j(\xi, \eta, \bar{\xi}, \bar{\eta}) + R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) = 0, \quad j = 1, \dots, n$$

with

$$q_j = \begin{cases} |\xi_j|^2 - \mu_j |\eta_{\sigma(j)}|^2, & 1 \leq j \leq K, \\ \xi_j \bar{\xi}_{\sigma(j)} - \mu_j \bar{\eta}_j \eta_{\sigma(j)}, & K < j \leq n, \end{cases}$$

where μ_j and σ (a permutation of $1, \dots, n$) satisfy

$$(4.22) \quad \begin{aligned} 1 \leq \sigma(j) \leq K, \quad \mu_{\sigma(j)} = 1/\mu_j > 0, \quad 1 \leq j \leq K; \\ \sigma(j) = \sigma^{-1}(j) \neq j, \quad |\mu_j| = 1, \quad \mu_{\sigma(j)} = \bar{\mu}_j, \quad K < j \leq n \end{aligned}$$

for some $0 \leq K \leq n + 1$. We further assume that R_j satisfy the reality conditions

$$\begin{aligned} R_j(\xi, \eta, \bar{\xi}, \bar{\eta}) &= \bar{R}_j(\bar{\xi}, \bar{\eta}, \xi, \eta), \quad 1 \leq j \leq K; \\ R_{\sigma(j)}(\xi, \eta, \bar{\xi}, \bar{\eta}) &= \bar{R}_j(\bar{\xi}, \bar{\eta}, \xi, \eta), \quad K < j \leq n. \end{aligned}$$

With the above assumptions and the change of coordinates $\xi_j \rightarrow \xi_j, \eta_j \rightarrow \sqrt{\mu_{\sigma^{-1}(j)}} \eta_j$, we can then apply Theorem 3.6. Thus, M is given by

$$(4.23) \quad 0 = r_j = \begin{cases} F_j(\xi, \eta) \bar{F}_j(\bar{\xi}, \bar{\eta}) - \mu_j G_{\sigma(j)}(\xi, \eta) \bar{G}_{\sigma(j)}(\bar{\xi}, \bar{\eta}), & j \leq K, \\ F_j(\xi, \eta) \bar{F}_{\sigma(j)}(\bar{\xi}, \bar{\eta}) - \mu_j G_{\sigma(j)}(\xi, \eta) \bar{G}_j(\bar{\xi}, \bar{\eta}), & j > K, \end{cases}$$

where σ, μ_j are given by (4.22), and

$$(4.24) \quad \begin{aligned} F_j(\xi, \eta) &= \xi_j + f_j(\xi, \eta), & G_j(\xi, \eta) &= \eta_j + g_j(\xi, \eta), \\ f_j(\xi, \eta) &= O(2), & g_j(\xi, \eta) &= O(2). \end{aligned}$$

Introduce meromorphic functions

$$(4.25) \quad m_j(\xi, \eta) = \frac{\xi_j + f_j(\xi, \eta)}{\eta_{\sigma(j)} + g_{\sigma(j)}(\xi, \eta)}, \quad j = 1, \dots, n.$$

PROPOSITION 4.5. — *Let M be a real analytic set given by (4.22)–(4.24). Then m_1, \dots, m_n depend only on M . If M is invariant under φ defined by (1.1)–(1.2), then $|\lambda_{\sigma(j)}| = 1/|\lambda_j|$ and*

$$(4.26) \quad m_j \circ \varphi = \lambda_j \lambda_{\sigma(j)} m_j, \quad j = 1, \dots, n.$$

Proof. — We first want to show that μ_j, m_j are uniquely determined by M . Assume that M is also given by (4.23)–(4.24) with $r_j^*, \mu_j^*, f_j^*, g_j^*$ in place of r_j, μ_j, f_j, g_j , respectively. Define m_j^* as in (4.25). By the Weierstrass preparation theorem, we may assume that the coefficients of g_j, g_j^* satisfy the normalizing condition

$$(4.27) \quad g_{j\alpha\beta} = 0, \quad \beta_j \neq 0.$$

We need to show that $\mu_j^* = \mu_j, f_j^* = f_j$, and $g_j^* = g_j$. Since $r_j^* - r_j$ vanishes on M , it follows from Proposition 3.7 that

$$(4.28) \quad r_j^* - r_j = \sum D_{jk} r_k$$

with

$$(4.29) \quad \min\{\text{ord } D_{jk}\} = \text{ord}(r_j^* - r_j) - 2.$$

Comparing the quadratic terms in (4.28) gives $\mu_j^* = \mu_j$ and $D_{jk}(0) = 0$.

Assume for induction that

$$\Delta f_j \equiv f_j^* - f_j = O(d), \quad \Delta g_j \equiv g_j^* - g_j = O(d), \quad D_{jk} = O(d - 1).$$

For brevity, we replace $[f_j]_d$ by f_j , and $[D_{jk}]_{d-1}$ by D_{jk} , etc. In (4.28) terms of the form $\xi^\alpha \eta^\beta \bar{\xi}^\gamma \bar{\eta}^\delta$ with $|\alpha| + |\beta| = d$ and $|\gamma| + |\delta| = 1$ give us

$$(4.30) \quad \bar{\xi}_j \Delta f_j - \mu_j \bar{\eta}_{\sigma(j)} \Delta g_{\sigma(j)} = \sum D_{jk}(\xi, \eta, 0, 0)[r_k]_2, \quad j \leq K,$$

$$(4.31) \quad \bar{\xi}_{\sigma(j)} \Delta f_j - \mu_j \bar{\eta}_j \Delta g_{\sigma(j)} = \sum D_{jk}(\xi, \eta, 0, 0)[r_k]_2, \quad j > K.$$

On \mathcal{Q} , the right-hand sides of (4.30)–(4.31) vanish and one can put

$$(4.32) \quad \begin{aligned} \eta_{\sigma(j)} &= \rho_j \bar{\xi}_j / \sqrt{\mu_j}, & |\rho_j| &= 1, & j &\leq K; \\ \eta_j &= \sqrt{\mu_j} \rho_j \bar{\xi}_j, & \rho_j \bar{\rho}_{\sigma(j)} &= 1, & j &> K. \end{aligned}$$

Note that $\Delta g_{\sigma(j)}$ is independent of $\eta_{\sigma(j)}$. Inserting (4.32) into the right-hand side of (4.30) and discarding terms containing $\bar{\xi}_j$ yields $\Delta g_{\sigma(j)}(\xi, \eta) = 0$, where η is given by (4.32). This shows that the holomorphic function $\Delta g_{\sigma(j)}$ vanishes on \mathcal{Q} . Since \mathcal{Q} contains a totally real submanifold in \mathbb{C}^{2n} of maximal dimension, then $\Delta g_j = 0$, i.e., $[g_j^* - g_j]_d = 0$ for $j \leq K$. Similarly, inserting (4.32) into the right hand side of (4.31) and discarding terms containing $\bar{\xi}_{\sigma(j)}$ yields $[g_j^* - g_j]_d = 0$ for $K < j \leq n$. Now (4.30)–(4.31)

read $\Delta f_j(\xi, \eta) = 0$, in which η is given by (4.32). This shows that $f_j^* = f_j$. Now formula (4.29) yields $[D_{jk}]_{d-1} = 0$. Thus, under the normalizing condition (4.27) we obtain by induction that $[f_j]_d = [f_j^*]_d$ and $[g_j^*]_d = [g_j]_d$. Therefore, μ_j, m_j are uniquely determined by M .

Now, we can show that $|\lambda_{\sigma(j)}| = 1/|\lambda_j|$ and m_j satisfy (4.26). Note that $M = \varphi(M)$ is also given by (4.23) with μ_j being replaced by $|\lambda_j \lambda_{\sigma(j)}|^2 \mu_j$ and F_j, G_j being replaced by

$$F_j^*(\xi, \eta) = \lambda_j^{-1} F_j \circ \varphi(\xi, \eta) = \xi_j + O(2),$$

$$G_j^*(\xi, \eta) = \lambda_j G_j \circ \varphi(\xi, \eta) = \eta_j + O(2),$$

respectively. By the above uniqueness result, we have $|\lambda_{\sigma(j)} \lambda_j| = 1$ and $F_j^*/G_{\sigma(j)}^* = m_j$, which gives us (4.26). The proof of the proposition is complete. □

The above proposition leads us to the question: If a holomorphic symplectic mapping (1.1)–(1.2) admits n meromorphic eigenfunctions (4.25), can it be put into its Birkhoff normal form by a convergent transformation?

The rest of this section and Section 5 are devoted to this eigenfunction problem.

Before we turn to the eigenfunction problem, we first remark that a holomorphic symplectic mapping of the form (1.1)–(1.2) satisfies $|\lambda_{\sigma(j)}| \cdot |\lambda_j| = 1$, when it admits the Levi-flat real analytic set (4.23). Although the existence of eigenfunctions (4.25) puts no extra restriction on the eigenvalues of the mapping, it restricts the higher order terms in the Birkhoff normal, as shown in the following.

THEOREM 4.6. — *Let φ be a holomorphic symplectic mapping defined by (1.1)–(1.2). Assume that φ admits n meromorphic eigenfunctions m_j given by (4.25). Let Φ be a formal symplectic transformation such that*

$$\Phi\varphi\Phi^{-1}: (\xi, \eta) \rightarrow (\lambda_j \xi_j e^{H_{\zeta_j}(\zeta)}, \lambda_j^{-1} \eta_j e^{-H_{\zeta_j}(\zeta)}) + O(k).$$

Then

$$m_j \circ \Phi^{-1}(\xi, \eta) = \frac{\xi_j \kappa_j(\xi\eta) + \widehat{f}_j(\xi, \eta)}{\eta_{\sigma(j)} + \widehat{g}_{\sigma(j)}(\xi, \eta)}, \quad \min\{\text{ord } \widehat{f}_j, \text{ord } \widehat{g}_{\sigma(j)}\} \geq k,$$

where κ_j are power series in $\xi_1 \eta_1, \dots, \xi_n \eta_n$ with $\kappa_j(0) = 1$. Moreover, the formal power series $\omega_1, \dots, \omega_n$ in the normal form (1.3) of φ satisfy

$$(4.33) \quad \omega_j + \omega_{\sigma(j)} = 0.$$

Proof. — For brevity, we replace $m_j \circ \Phi^{-1}$ by m_j , and φ by $\Phi \circ \varphi \circ \Phi^{-1}$. Put m_j in the form (4.25), where f_j, g_j are now formal power series with g_j satisfying the normalizing condition (4.27).

From $m_j \circ \varphi = \lambda_j \lambda_{\sigma(j)} m_j$, we get

$$(4.34) \quad \begin{aligned} & (\eta_{\sigma(j)} + g_{\sigma(j)})(\xi_j e^{\omega_j} + \lambda_j^{-1} f_j \circ \varphi) \\ &= (\xi_j + f_j)(\eta_{\sigma(j)} e^{-\omega_{\sigma(j)}} + \lambda_{\sigma(j)} g_{\sigma(j)} \circ \varphi) + O(k + 1). \end{aligned}$$

Assume for induction that

$$(4.35) \quad \begin{aligned} f_j(\xi, \eta) &= \xi_j \kappa_j(\xi \eta) + O(d), \quad g_j(\xi, \eta) = O(d), \\ \omega_j(\xi \eta) + \omega_{\sigma(j)}(\xi \eta) &= O(|(\xi, \eta)|^{d-1}) \end{aligned}$$

hold for some $d < k - 1$. We would like to show that the above identities hold if d is replaced by $d + 1$. Setting $\eta_{\sigma(j)} = 0$ in (4.34), one gets from (4.35) and the normalizing condition (4.27) that

$$g_{\sigma(j)}(\xi, \eta) = \lambda_{\sigma(j)} g_{\sigma(j)}(\lambda \xi, \lambda^{-1} \eta) + O(d + 1).$$

Hence, $g_j(\xi, \eta) = O(d + 1)$. Now, terms of order $d + 1$ in (4.34) give us

$$(4.36) \quad \xi_j [\omega_j + \omega_{\sigma(j)}]_{(d-1)/2}(\xi \eta) + \lambda_j^{-1} [f_j]_d(\lambda \xi, \lambda^{-1} \eta) = [f_j]_d,$$

where $[\omega_j + \omega_{\sigma(j)}]_{(d-1)/2}$ is treated as zero if $(d - 1)/2$ is not an integer. Set

$$e_j = (0, \dots, 1, \dots, 0) \in \mathbb{Z}^n,$$

where 1 is the j -th entry. Collecting terms $\xi^\alpha \eta^\beta$ for $\alpha - \beta = e_j$ yields

$$[\omega_j + \omega_{\sigma(j)}]_{(d-1)/2}(\xi \eta) = 0.$$

We now have $[f_j]_d(\lambda \xi, \lambda^{-1} \eta) = \lambda_j [f_j]_d(\xi, \eta)$. Therefore, $[f_j]_d(\xi, \eta)$ contains only terms $\xi^\alpha \eta^\beta$ with $\alpha - \beta = e_j$, and (4.35) hold if d is replaced by $d + 1$. By induction, (4.15) holds for $d = k$. This shows that $\min\{\text{ord } \hat{f}_j, \text{ord } \hat{g}_j\} \geq k$, as stated in the theorem.

The identity (4.33) follows from the above result (with $k = \infty$) and the existence of Φ that normalizes φ . The proof of the theorem is complete. □

As a consequence of Proposition 4.5 and Proposition 4.6, we know that if φ has an invariant Levi-flat real analytic set defined by (4.21) with $\sigma^d = \text{Id}$ for some odd integer, then (4.33) implies that all $\omega_j = 0$, that is that φ is formally linearizable. By a theorem of Rüssmann [14], we obtain the following.

COROLLARY 4.7. — *Let φ be a holomorphic symplectic mapping defined by (1.1)–(1.2). Assume that φ admits an invariant Levi-flat real*

analytic set given by (4.21) with $\sigma^k = \text{Id}$ for some odd integer k . If the eigenvalues λ_j of φ satisfy the Diophantine condition

$$|\lambda^\alpha - \lambda_j| \geq c/|\alpha|^\tau, \quad |\alpha| > 1, j = 1, \dots, n$$

for some positive constant c, τ , then φ is linearizable by holomorphic symplectic mappings.

For $n = 1$, see Proposition 4.2 with weaker hypotheses.

To state the next theorem, we need some notation. Let σ be a permutation of $1, \dots, n$, and let m_j , given by (4.25), be meromorphic eigenfunctions of φ . Theorem 4.6 implies that if Φ normalizes φ , then

$$m_j \circ \Phi^{-1}(\xi, \eta) = \xi_j \kappa_j(\xi\eta)/\eta_{\sigma(j)}.$$

For $1 \leq j \leq n$, recall that τ_j is the smallest positive integer with $\sigma^{\tau_j}(j) = j$. Define

$$(4.37) \quad \tau_j \kappa = 1, \quad \text{for } \tau_j = 1; \quad \delta_j \kappa = \prod_{k=0}^{\tau_j-1} \kappa_{\sigma^k(j)}^{(-1)^k}, \quad \text{for } \tau_j > 1.$$

One readily sees that

$$(4.38) \quad \delta_j \kappa \cdot \delta_{\sigma(j)} \kappa = \begin{cases} 1, & \text{if } \tau_j \text{ is one or even,} \\ \kappa_j^2, & \text{otherwise.} \end{cases}$$

The dependence of $\delta_j \kappa$ on the formal normalizing transformation Φ is described as the following.

PROPOSITION 4.8. — Let $\delta_1 \kappa, \dots, \delta_n \kappa$ be as above. Fix j with $1 \leq j \leq n$. The formal power series $\delta_j \kappa$ is independent of the choice of Φ if τ_j is even, but $\delta_j \kappa \equiv 1$ for some formal normalizing transformation Φ if τ_j is odd.

Proof. — Let $\tilde{\Phi}$ be another formal symplectic mapping which transforms φ into (1.3). Put $m_j \circ \tilde{\Phi}^{-1} = \xi_j \tilde{\kappa}_j / \eta_{\sigma(j)}$. For $(\xi', \eta') = \tilde{\Phi} \circ \Phi^{-1}(\xi, \eta)$, one has

$$(4.39) \quad \xi'_j \tilde{\kappa}_j(\xi'\eta') / \eta'_{\sigma(j)} = \xi_j \kappa_j / \eta_{\sigma(j)}.$$

Since $\tilde{\Phi} \circ \Phi^{-1}$ preserves the normal form of φ , the formal theory of Birkhoff's normal form says that

$$\tilde{\Phi} \circ \Phi^{-1}: \xi'_j = \xi_j e^{G_{c_j}(\xi\eta)}, \quad \eta'_j = \eta_j e^{-G_{c_j}(\xi\eta)}, \quad j = 1, \dots, n,$$

where G is a formal power series in $\zeta = \xi\eta$. Now (4.39) yields

$$e^{G_{c_j}(\zeta) + G_{c_{\sigma(j)}}(\zeta)} \tilde{\kappa}_j(\zeta) = \kappa_j(\zeta),$$

which gives us $\delta_j \tilde{\kappa}(\zeta) = \delta_j \kappa(\zeta)$ if τ_j is even. When τ_j is odd, we get

$$e^{2G_{c_j(\zeta)}} \delta_j \tilde{\kappa}(\zeta) = \delta_j \kappa(\zeta).$$

Solving the above equation for a formal power series G , one obtains a formal normalizing transformation $\tilde{\Phi}$ such that $\delta_j \tilde{\kappa} \equiv 1$. The proof of the proposition is complete. □

We now state the following theorem.

THEOREM 4.9. — *Let φ, m_1, \dots, m_n be as in Theorem 4.6, and let $\delta_j \kappa$ be the associated formal power series. Assume that $\sigma^2 = \text{Id}$, and that there exists a formal transformation Φ normalizing φ such that all $\delta_j \kappa \circ \Phi$ are convergent. Then φ is normalizable by holomorphic symplectic mappings.*

In conclusion of this section, note that Theorem 1.2 follows from Proposition 4.5 and Theorems 4.6 and 4.9.

5. Normalization of holomorphic symplectic mappings with meromorphic eigenfunctions.

Recall that Proposition 4.5 provides us the relationship between the Levi-flat invariant sets of the form (4.23) of a holomorphic symplectic mapping and meromorphic eigenfunctions of the mapping. The relationship between meromorphic eigenfunctions and Birkhoff normal forms of holomorphic symplectic mappings might be interesting in its own right.

In Section 4, we introduced the following eigenfunctions:

$$(5.1) \quad m_j = \frac{\xi_j + f(\xi, \eta)}{\eta_{\sigma(j)} + g_{\sigma(j)}(\xi, \eta)}, \quad j = 1, \dots, n,$$

where σ is a permutation of $1, \dots, n$, and $f_j(\xi, \eta) = O(3), g_j(\xi, \eta) = O(3)$ are holomorphic. We should mention that we have no example of holomorphic symplectic mappings having eigenfunctions m_j , for which the Birkhoff normalization diverges.

The purpose of this section is to prove Theorem 4.9. However, we need a further reduction for the theorem, which is a necessary step for us to be able to apply the KAM method. Recall from Section 4 (Theorem 4.6) that if a formal holomorphic symplectic mapping Φ normalizes the symplectic mapping φ , then $m_j \circ \Phi^{-1}$ are of the form

$$(5.2) \quad \xi_j \kappa_j(\xi \eta) / \eta_{\sigma(j)}, \quad j = 1, \dots, n.$$

Assuming that $\sigma^2 = \text{Id}$ and all $\delta_j \circ \Phi$ are convergent, we now modify m_j such that

$$(5.3) \quad \delta_j \kappa \equiv 1, \quad j = 1, \dots, n.$$

If $\sigma(j) = j$, we simply put $\tilde{m}_j = m_j$. Otherwise, say $\sigma(1) = 2$, we put

$$\tilde{m}_1 = m_1/(\delta_1 \kappa \circ \Phi), \quad \tilde{m}_2 = m_2.$$

Then $\delta_1 \tilde{m} = \tilde{m}_1/\tilde{m}_2 = 1 = \delta_2 \tilde{m}$. Let us still denote by m_j the modified meromorphic functions \tilde{m}_j . In such a way we achieve (5.3).

Since $\sigma^2 = \text{Id}$, Proposition 4.8 implies that (5.3) remains true under all formal symplectic transformations that normalize φ . Also, (5.3) is equivalent to $\kappa_j = \kappa_{\sigma(j)}$, since $\delta_j \kappa = \kappa_j/\kappa_{\sigma(j)}$ when $\sigma^2 = \text{Id}$.

Thus, Theorem 4.9 is reduced to the following special case.

THEOREM 5.1. — *Let φ be a holomorphic symplectic mapping given by (1.1)–(1.2). Let m_1, \dots, m_n be n meromorphic eigenfunctions of φ of the form (5.1) with $\sigma^2 = \text{Id}$. Let Φ be a formal transformation that normalizes φ and transforms κ_j into (5.2). Assume that $\kappa_{\sigma(j)} = \kappa_j$ for $j = 1, \dots, n$. Then φ can be transformed into its Birkhoff normal form by some holomorphic symplectic transformation.*

We now proceed to prove Theorem 5.1, by applying the KAM method. Starting with the holomorphic symplectic mapping φ defined by (1.1)–(1.2), we construct a sequence of holomorphic symplectic mappings ϕ_m as follows. Put $\varphi_1 = \varphi$. Let Φ_1 be the unique normalized formal transformation transforming φ_1 into its normal form. Let $\sum \lambda_j \xi_j \eta'_j + H_1(\xi, \eta')$ be the generating function of φ_1 , that is,

$$\varphi_1: \xi'_j = \lambda_j \xi_j + \partial_{\eta'_j} H_1(\xi, \eta'), \quad \eta'_j = \lambda_j^{-1} \eta_j - \lambda_j^{-1} \partial_{\xi_j} H_1(\xi, \eta').$$

Let $\sum \xi_j \eta'_j + S_1(\xi, \eta')$ be the generating function of Φ_1 . From the formal theory of Birkhoff's normal form, one knows that $\text{ord } S_1 = d_0 \equiv \text{ord } PH_1$ for

$$PH_1(\xi, \eta') = \sum_{\alpha \neq \beta} H_{1,\alpha\beta} \xi^\alpha \eta'^\beta.$$

Let ϕ_1 be the holomorphic symplectic mapping generated by the truncated power series

$$\tilde{S}_1(\xi, \eta') = \sum_j \xi_j \eta'_j + \sum_{k=d_0}^{2d_0-3} [S_1]_k(\xi, \eta')$$

with

$$[S_1]_k(\xi, \eta') \equiv \sum_{\alpha+\beta=k} S_{1,\alpha\beta} \xi^\alpha \eta'^\beta.$$

Define $\varphi_2 = \phi_1 \circ \varphi_1 \circ \phi_1^{-1}$. Repeating the above construction, we find holomorphic symplectic transformations φ_k, ϕ_k with $\varphi_{k+1} = \phi_k \circ \varphi_k \circ \phi_k^{-1}$. Let $\sum \lambda_j \xi_j \eta'_j + H_k(\xi, \eta')$ be the generating function of φ_k and put $d_k = \text{ord } PH_k$.

If one of d_k is infinite, proof of Theorem 5.1 is trivial; so we will assume that all d_k are finite. Then

$$d_{k+1} \geq 2d_k - 2, \quad d_0 \geq 3.$$

In particular, $d_k - 2 \geq 2^k, k = 0, 1, \dots$. It is clear that as $k \rightarrow \infty$ the limit of $\phi_k \circ \phi_2 \circ \dots \circ \phi_1$ is a formal symplectic mapping ϕ such that $\phi \circ \varphi \circ \phi^{-1}$ is in the Birkhoff normal form. We shall complete the proof of Theorem 5.1 by showing the convergence of ϕ .

To avoid functional equations involving small divisors, we shall determine ϕ_1 through eigenfunctions m_j . *Changing notations*, replace d_0, \tilde{S}_1 and ϕ_1 by d, S, ϕ , respectively. We also rewrite

$$(5.4) \quad m_j(\xi, \eta) = \frac{\xi_j \kappa_j(\xi \eta) + f_j(\xi, \eta)}{\eta_{\sigma(j)} + g_{\sigma(j)}(\xi, \eta)},$$

where f_j, g_j satisfy

$$(5.5) \quad f_{j\alpha\beta} = 0, \quad \alpha - \beta = e_j; \quad g_{j\alpha\beta} = 0, \quad \beta_j \neq 0,$$

in which $e_j = (0, \dots, 1, \dots, 0)$. Theorem 4.6 implies that $\text{ord } f_j, \text{ord } g_j \geq d - 1$.

One knows that

$$(5.6) \quad \phi: \xi'_j = \xi_j + u_j(\xi, \eta), \quad \eta'_j = \eta_j + v_j(\xi, \eta), \quad j = 1, \dots, n$$

is determined by

$$(5.7) \quad \xi'_j = \xi_j - S_{\eta_j}(\xi, \eta'), \quad \eta'_j = \eta_j + S_{\xi_j}(\xi, \eta'),$$

where S , according to the normalizing condition on Φ , satisfies

$$(5.8) \quad S_{\alpha\alpha} = 0, \quad \text{ord } S = d.$$

One also has

$$u_j(\xi, \eta) = -S_{\eta_j}(\xi, \eta) + O(2d - 3), \quad v_j(\xi, \eta) = S_{\xi_j}(\xi, \eta) + O(2d - 3).$$

Set

$$(5.9) \quad m_j(\xi, \eta) = \hat{m}_j \circ \phi = \frac{(\xi_j + u_j) \hat{\kappa}_j \circ \phi(\xi, \eta) + \hat{f}_j \circ \phi(\xi, \eta)}{\eta_{\sigma(j)} + v_{\sigma(j)} + \hat{g}_{\sigma(j)} \circ \phi(\xi, \eta)}$$

with \hat{f}_j, \hat{g}_j satisfying (5.5). By Theorem 4.6, we have

$$\text{ord } \hat{f}_j, \text{ord } \hat{g}_j \geq 2d - 3; \quad \text{ord}(\hat{\kappa}_j - \kappa_j) \geq (d - 2)/2.$$

The linearized equations of (5.9) are

$$(5.10) \quad \eta_{\sigma(j)} S_{\eta_j}(\xi, \eta) + \xi_j S_{\xi_{\sigma(j)}}(\xi, \eta) = p_j(\xi, \eta), \quad j = 1, \dots, n$$

with

$$(5.11) \quad p_j = \xi_j \eta_{\sigma(j)} \nabla_{\zeta} \log \kappa_j \cdot (\xi S_{\xi} - \eta S_{\eta}) \\ + \xi_j g_{\sigma(j)} - \eta_{\sigma(j)} f_j / \kappa_j + \xi_j \eta_{\sigma(j)} (\hat{\kappa}_j / \kappa_j - 1) + O(2d - 2).$$

In particular,

$$(5.12) \quad \xi_j S_{\xi_j} + \eta_j S_{\eta_j} = p_j(\xi, \eta), \quad \text{for } \sigma(j) = j.$$

For $\sigma(j) \neq j$, rewrite (5.10) as

$$\xi_j^{-1} S_{\eta_j} + \eta_{\sigma(j)}^{-1} S_{\xi_{\sigma(j)}} = \nabla_{\zeta} \log \kappa_j \cdot (\xi S_{\xi} - \eta S_{\eta}) + \tilde{p}_j$$

with

$$(5.13) \quad \tilde{p}_j(\xi, \eta) = g_{\sigma(j)} / \eta_{\sigma(j)} - f_j / (\xi_j \kappa_j) + \hat{\kappa}_j / \kappa_j - 1 + \dots,$$

in which the omitted terms are of the form $\xi^{\alpha} \eta^{\beta} / \xi_j \eta_{\sigma(j)}$ with $|\alpha| + |\beta| \geq 2d - 2$. Using (5.3), we obtain

$$\sum_{k=0}^{\tau_j-1} (-1)^k \left(\frac{S_{\eta_{\sigma^k(j)}}}{\xi_{\sigma^k(j)}} + \frac{S_{\xi_{\sigma^{k+1}(j)}}}{\eta_{\sigma^{k+1}(j)}} \right) = \sum_{k=0}^{\tau_j-1} (-1)^k \tilde{p}_{\sigma^k(j)} \equiv \Delta_j \tilde{p};$$

in other words,

$$(5.14) \quad \frac{S_{\eta_j}}{\xi_j} - (-1)^{\tau_j} \frac{S_{\xi_j}}{\eta_j} + \sum_{k=1}^{\tau_j-1} (-1)^k \left(\frac{S_{\eta_{\sigma^k(j)}}}{\xi_{\sigma^k(j)}} - \frac{S_{\xi_{\sigma^k(j)}}}{\eta_{\sigma^k(j)}} \right) = \Delta_j \tilde{p}$$

for $\sigma(j) \neq j$.

LEMMA 5.2. — *Let $\sum \xi_j \eta_j + S(\xi, \eta)$, satisfying (5.8), be the generating function of the unique formal mapping Φ that normalizes φ . Then $S \equiv [S]_d + \dots + [S]_{2d-3}$ is uniquely determined by (5.12) and (5.14).*

Proof. — Let S be the solution to (5.12) and (5.14), which satisfies the normalization condition (5.8). Fix α, β with $\alpha - \beta \neq 0$ and choose j so that

$$(5.15) \quad |\alpha_j - \beta_j| = \max\{|\alpha_k - \beta_k|\}.$$

If $\sigma(j) = j$, then (5.12) implies that

$$(5.16) \quad S_{\alpha\beta} = p_{j\alpha\beta} / (\alpha_j + \beta_j).$$

Assume now that $\sigma(j) \neq j$. To determine $S_{\alpha\beta}$, we need to consider all coefficients of $\xi^{\alpha'} \eta^{\beta'}$ in (5.14), for which

$$\alpha' - \beta' = \alpha - \beta; \quad \alpha'_k \geq -1, \quad \beta'_k \geq -1, \quad k = 1, \dots, n.$$

Put $S_{\alpha\beta} = 0$ for $\alpha_k \leq -1$ or $\beta_k \leq -1$. From (5.14) we obtain

$$(\beta'_j - (-1)^{\tau_j} \alpha'_j) S_{\alpha', \beta'} + \sum_{k=1}^{\tau_j-1} (-1)^k (\beta_{\sigma^k(j)} - \alpha_{\sigma^k(j)}) S_{\alpha' + e_{\sigma^k(j)} - e_j, \beta' + e_{\sigma^k(j)} - e_j} = (\Delta_j \tilde{p})_{\alpha' - e_j, \beta' - e_j}.$$

If $\alpha'_j + \beta'_j = \min\{\tilde{\alpha}_j + \tilde{\beta}_j | \tilde{\alpha} - \tilde{\beta} = \alpha - \beta\}$, then

$$(5.17) \quad S_{\alpha' \beta'} = \frac{1}{\beta'_j - (-1)^{\tau_j} \alpha'_j} p_{j, \alpha' - e_j, \beta' - e_j}.$$

Assume that $S_{\alpha' \beta'}$ have been determined for all α', β' with $\alpha' - \beta' = \alpha - \beta$ and $\alpha'_j + \beta'_j < l$. Then for α', β' with $\alpha'_j + \beta'_j = l$, we have

$$(5.18) \quad S_{\alpha' \beta'} = \sum_{k=1}^{\tau_j-1} (-1)^{k-1} \frac{\beta_{\sigma^k(j)} - \alpha_{\sigma^k(j)}}{\beta'_j - (-1)^{\tau_j} \alpha'_j} S_{\alpha' - e_j + e_{\sigma^k(j)}, \beta' - e_j + e_{\sigma^k(j)}} + \frac{1}{\beta'_j - (-1)^{\tau_j} \alpha'_j} (\Delta_j \tilde{p})_{\alpha' - e_j, \beta' - e_j}.$$

Thus, (5.16)–(5.18) determine all $S_{\alpha' \beta'}$ for $\alpha' - \beta' = \alpha - \beta$. This completes the proof of the lemma. \square

We should point out that the proof of Lemma 5.2 does not depend on any assumption on $\delta_j \kappa$, and that the convergence of all $\delta_j \kappa$, as assumed in Theorem 4.9, would not give us an immediate good control of S due to the recursive formulae (5.18) when some $\tau_j \geq 2$. To get estimates for S , it is crucial that all $\nabla_\zeta \log \delta_j \kappa$ vanish, which is obtained via (5.3) for the modified eigenfunctions. Even if $\delta_j \kappa \equiv 1$, the recursive formulae (5.18) lead to another problem: The radius of convergence would shrink by a constant factor of $\tau_j - 1$ if $\tau_j > 1$. Therefore, only for $\tau_j = 1, 2$ shall we have the control of the radius of convergence that is useful in iteration.

We need some notation. Let $f = (f_1, \dots, f_m)$ be an m -tuple of holomorphic functions defined on $\Delta_r \subset \mathbb{C}^n$. Expand $f_j(z) = \sum f_{j\alpha} z^\alpha$ and let

$$f^*(z) = \sum f_{|\alpha|}^* z^\alpha, \quad f_k^* = \max\{|f_{j\alpha}|; |\alpha| = k, j = 1, \dots, m\};$$

$$\|f\|_r = \sup\{|f_j(z)|; z \in \Delta_r, j = 1, \dots, m\}.$$

In particular, one has $\xi^* = \xi_j^* = \sum(\xi_j + \eta_j)$, where (ξ, η) are coordinates of \mathbb{C}^{2n} . We also put

$$(\nabla f)^* = (\partial_{z_1} f, \dots, \partial_{z_n} f)^*, \quad \|\nabla f\|_r = \|(\partial_{z_1} f, \dots, \partial_{z_n} f)\|_r.$$

Given holomorphic functions f and g , one has

$$(f + g)^* \prec f^* + g^*; \quad f^* \prec g^*, \quad \text{if } f \prec g.$$

Also

$$(5.19) \quad f^* \prec z^*(\nabla f)^*, \quad \text{if } f(0) = 0.$$

LEMMA 5.3. — *Let f be a holomorphic function defined on $\Delta_r \subset \mathbb{C}^n$. Then $\|f^*\|_{(1-\theta)r} \leq \|f\|_r/\theta^{n+1}$.*

Proof. — Since $|f_\alpha| \leq \|f\|_r/r^{|\alpha|}$, then $\|f^*\|_{(1-\theta)r}$ is bounded by

$$\|f\|_r \sum_{\alpha} (1-\theta)^{|\alpha|} = \|f\|_r \sum_{k \geq 0} \binom{n+k}{n} (1-\theta)^k = \|f\|_r/\theta^{n+1}.$$

This completes the proof of the lemma. □

LEMMA 5.4. — *Let ϕ, S be given by (5.6)–(5.8). Assume that*

$$(5.20) \quad \|S\|_r \leq \theta^2 r^2/(4n), \quad 0 < \theta < 1.$$

Then

$$(5.21) \quad \begin{aligned} &\phi, \phi^{-1}: \Delta_{(1-k\theta)r} \rightarrow \Delta_{(1-(k-1)\theta)r}, \\ &\|\phi^{-1} - \text{Id}\|_{(1-k\theta)r} \leq \|S\|_{(1-(k-2)\theta)r}/(r\theta), \quad k \geq 2. \end{aligned}$$

Proof. — Fix $(\xi, \eta) \in \Delta_{(1-k\theta)r}$. Consider the mapping

$$T: \xi'_j \rightarrow \xi_j - S_{\eta'_j}(\xi, \eta'), \quad \eta'_j \rightarrow \eta_j - S_{\xi_j}(\xi, \eta').$$

From (5.20) it follows that T maps $\Delta_{(1-(k-1)\theta)r}$ into itself and

$$\|\nabla T\|_{(1-\theta)r} \leq 1/(4n).$$

Hence, with the norm $\|(\xi, \eta)\| = \max\{|\xi_j|, |\eta_j|\}$, T is a contraction map. By the fixed-point theorem, equations (5.7) determine a mapping $\phi: \Delta_{(1-k\theta)r} \rightarrow \Delta_{(1-(k-1)\theta)r}$. Applying the above argument to the mapping $\xi_j \rightarrow \xi'_j + S_{\eta'_j}(\xi, \eta'), \eta_j \rightarrow \eta'_j - S_{\xi_j}(\xi, \eta')$, one gets $\phi^{-1}: \Delta_{(1-k\theta)r} \rightarrow \Delta_{(1-(k-1)\theta)r}$. Consequently, we have

$$\|\phi^{-1} - \text{Id}\|_{(1-k\theta)r} \leq \|\nabla S\|_{(1-(k-1)\theta)r} \leq \|S\|_{(1-(k-2)\theta)r}/(r\theta).$$

The proof of the lemma is complete. □

Let $\kappa_j(\zeta_1, \dots, \zeta_n)$ be as in (5.4). Put

$$\kappa = (\kappa_1, \dots, \kappa_n), \quad \kappa^{-1} = (\kappa_1^{-1}, \dots, \kappa_n^{-1}).$$

LEMMA 5.5. — *Let φ and S be as in Lemma 5.2, and let ϕ be given by (5.7). Assume that*

$$(5.22) \quad \|(\kappa^{-1})^*\|_{r^2} < 2, \quad r^2\|(\nabla_\zeta \kappa)^*\|_{r^2} \leq 1/(16n^2).$$

There exist constants $c_1, c_2 > 1$ such that if

$$(5.23) \quad \|f\|_r + \|g\|_r \leq r\theta^{2n+4}/c_1,$$

then

$$(5.24) \quad \begin{aligned} \phi, \phi^{-1}: \Delta_{(1-k\theta)r} &\rightarrow \Delta_{(1-(k-2)\theta)r}, \quad k \geq 2, \\ \|\phi^{-1} - \text{Id}\|_{(1-2\theta)r} &\leq c_2(\|f\|_r + \|g\|_r)/\theta^{2n+3}. \end{aligned}$$

Proof. — We first introduce the following notation:

$$\widehat{S}(\xi, \eta) = \sum |S_{\alpha\beta}| \xi^\alpha \eta^\beta$$

which will be used only in the proof of this lemma. Fix α, β with $\alpha \neq \beta$ and let j be defined by (5.15). Note that (5.15) implies that

$$|\{\xi_k S_{\xi_k} - \eta_k S_{\eta_k}\}_{\alpha'\beta'}| \leq |\alpha_j - \beta_j| \widehat{S}_{\alpha'\beta'}, \quad k = 1, \dots, n$$

for $\alpha' - \beta' = \alpha - \beta$. Hence, for $\sigma(j) = j$ one gets from (5.16) that

$$\begin{aligned} |S_{\alpha\beta}| &\leq \{\xi_j \eta_j (\nabla_\zeta \kappa)^*(\kappa^{-1})^* \widehat{S} + \xi_j g_j^* + \eta_j (\kappa_j^{-1})^*(f_j)^*\}_{\alpha\beta} \\ &\leq \{\xi^* \eta^* (\nabla_\zeta \kappa)^*(\kappa^{-1})^* \widehat{S} + \xi^*(\kappa^{-1})^*(f^* + g^*)\}_{\alpha\beta}. \end{aligned}$$

If $\sigma(j) \neq j$, (5.17)–(5.18) yield

$$\begin{aligned} |S_{\alpha\beta}| &\leq (1 + \min\{\alpha_j, \beta_j\}) \{\xi^* g^* + \eta^*(\kappa^{-1})^* f^*\}_{\alpha\beta} \\ &\leq (|\alpha| + |\beta|) \{\xi^* g^* + \eta^*(\kappa^{-1})^* f^*\}_{\alpha\beta} \\ &= \{(\xi \cdot \partial_\xi + \eta \cdot \partial_\eta)(\xi^*(\kappa^{-1})^* f^* + \xi^* g^*)\}_{\alpha\beta}. \end{aligned}$$

Notice that we have used the essential assumption that $\tau_j = 2$ to obtain the first inequality above. Thus,

$$(5.25) \quad \widehat{S} \prec \frac{1}{1 - \xi^* \eta^* (\nabla_\zeta \kappa)^*(\kappa^{-1})^*} (\text{Id} + \xi \partial_\xi + \eta \partial_\eta)(\xi^*(\kappa^{-1})^*(f^* + g^*)).$$

From (5.19) we get $\|\xi^* \eta^* (\nabla_\zeta \kappa)^*(\kappa^{-1})^*\|_r < 1/2$ and

$$\begin{aligned} \|(\xi \cdot \partial_\xi + \eta \cdot \partial_\eta)(\xi^*(\kappa^{-1})^*(f^* + g^*))\|_{(1-\theta/2)r/\theta} &\leq 8n \|\xi^*(f^* + g^*)\|_{(1-\theta/2)r/\theta} \\ &\leq \tilde{c}_1 r (\|f\|_r + \|g\|_r) / \theta^{2n+2} \end{aligned}$$

for some constant \tilde{c}_1 . From (5.25) it follows that

$$\|\widehat{S}\|_{(1-\theta)r} \leq c_1 r (\|f\|_r + \|g\|_r) / (4n\theta^{2n+2}).$$

Now, Lemma 5.4 implies that the mapping ϕ , determined by (5.6)–(5.7), satisfies (5.24), provided (5.22)–(5.23) hold. \square

The following is a special case of the Weierstrass preparation theorem.

LEMMA 5.6. — *Let f be a holomorphic function on $\Delta_r \subset \mathbb{C}^{2n}$. Assume that $0 < \theta < 1/2$ and*

$$(5.26) \quad \|f\|_r \leq r\theta/12, \quad f(\xi, \eta) = O(2).$$

Fix $1 \leq j \leq n$. There is a unique decomposition

$$(5.27) \quad \eta_j + f(\xi, \eta) = (1 + u(\xi, \eta))(\eta_j + g(\xi, \eta)), \quad u(0) = 0,$$

where u, g are holomorphic on $\Delta_{(1-\theta)r}$ with g satisfying the normalizing condition (4.27). Moreover, for some constant $\tilde{c}_2 > 1$ one has

$$(5.28) \quad \|g^*\|_{(1-\theta)r} \leq \tilde{c}_2 \|f\|_r / \theta^{2n+1}, \quad \|u^*\|_{(1-\theta)r} \leq \tilde{c}_2 \|f\|_r / (r\theta^{2n+2}).$$

Proof. — From (5.26) it follows that for $\xi_j \in \Delta_r$ there exists a unique solution $\eta_j = -g(\xi, \eta)$ to $\eta_j + f(\xi, \eta) = 0$, given by

$$\begin{aligned} g(\xi, \eta) &= -\frac{1}{2\pi i} \int_{|\eta_j|=r} \eta_j \frac{1 + f_{\eta_j}(\xi, \eta)}{\eta_j + f(\xi, \eta)} d\eta_j \\ &= -\frac{1}{2\pi i} \int_{|\eta_j|=r} \frac{\eta_j f_{\eta_j}(\xi, \eta) - f(\xi, \eta)}{\eta_j + f(\xi, \eta)} d\eta_j, \quad (\xi, \eta) \in \Delta_r \subset \Delta^{2n-1}. \end{aligned}$$

Thus, Cauchy inequalities imply that

$$(5.29) \quad \|g\|_{(1-\theta/2)r} \leq 2(2\|f\|_r/\theta + \|f\|_r) \leq 6\|f\|_r/\theta.$$

Now the desired estimate for g^* in (5.28) follows from Lemma 5.3.

To estimate u^* , note that $u = (f - g)/(\eta_j + g(\xi, \eta))$. From (5.26) and (5.29) it follows that $\|u\|_{(1-\theta/2)r} \leq 4(\|f\|_r + \|g\|_{(1-\theta/2)r})/r \leq 28\|f\|_r/(r\theta)$. In view of Lemma 5.3, we obtain (5.27). This completes the proof of the lemma. □

PROPOSITION 5.7. — *Let ϕ be as in Lemma 5.5. Let $\hat{\kappa}_j, \hat{f}_j, \hat{g}_j$ be as in (5.9). There exist constants $c_3, c_4 > 1$ such that*

$$(5.30) \quad \|\hat{f}\|_{(1-6\theta)r} + \|\hat{g}\|_{(1-6\theta)r} \leq c_3(1 - \theta)^d (\|f\|_r + \|g\|_r) / \theta^{6n+6},$$

$$(5.31) \quad \begin{aligned} \|(\nabla_\zeta \hat{\kappa})^*\|_{((1-6\theta)r)^2} &\leq \|(\nabla_\zeta \kappa)^*\|_{r^2} + c_3(\|f\|_r + \|g\|_r) / (r^3 \theta^{7n+8}), \\ \|(\hat{\kappa}^{-1})^*\|_{((1-6\theta)r)^2} &\leq \|(\kappa^{-1})^*\|_{r^2} + c_3(\|f\|_r + \|g\|_r) / (r\theta^{7n+7}) \end{aligned}$$

provided (5.22) holds, and

$$(5.32) \quad \|f\|_r + \|g\|_r \leq r\theta^{6n+6}/c_4.$$

Proof. — *Changing the notation in (5.6), we write*

$$\phi^{-1}: \xi'_j = \xi_j + u_j(\xi, \eta), \quad \eta'_j = \eta_j + v_j(\xi, \eta).$$

We first want to express \hat{f}_j, \hat{g}_j in terms of f_j, g_j and S . Start with the decomposition

$$\eta_j + v_j(\xi, \eta) + g_j \circ \phi^{-1}(\xi, \eta) = (1 + a_j)(\eta_j + \hat{g}_j(\xi, \eta)).$$

From (5.24), one sees that

$$\|v_j + g_j \circ \phi^{-1}\|_{(1-2\theta)r} \leq 2c_2(\|f\|_r + \|g\|_r)/\theta^{2n+3}.$$

In view of (5.26) and Lemma 5.6, we get from (5.28) that

$$(5.33) \quad \begin{aligned} \|a_j\|_{(1-3\theta)r} &\leq 2c_2\tilde{c}_2(\|f\|_r + \|g\|_r)/((1-2\theta)r\theta^{4n+5}), \\ \|\hat{g}_j\|_{(1-3\theta)r} &\leq 2c_2\tilde{c}_2(\|f\|_r + \|g\|_r)/\theta^{4n+4} \end{aligned}$$

for $c_4 > 24c_2$. Since $\text{ord } \hat{g}_j \geq 2d - 3 \geq d$, the Schwarz lemma yields

$$(5.34) \quad \|\hat{g}_j\|_{(1-3\theta)r} \leq 2c_2\tilde{c}_2 \left(\frac{1-4\theta}{1-3\theta} \right)^d (\|f\|_r + \|g\|_r)/\theta^{4n+4}.$$

For $0 < \theta < 1/6$ and $c_4 > c_2\tilde{c}_2$, (5.32)–(5.33) imply that

$$(5.35) \quad \|a_j\|_{(1-3\theta)r} < 1/2.$$

We now decompose

$$((\xi_j + u_j)\kappa_j \circ \phi^{-1} + f_j \circ \phi^{-1})/(1 + a_{\sigma(j)}) = \xi_j \hat{\kappa}_j + \hat{f}_j.$$

Then

$$(5.36) \quad \begin{aligned} \xi_j(\hat{\kappa}_j - \kappa_j) + \hat{f}_j &= \{\xi_j(\kappa_j \circ \phi^{-1} - \kappa_j) + u_j \kappa_j \circ \phi^{-1} \\ &\quad + f_j \circ \phi^{-1} - \xi_j \kappa_j a_{\sigma(j)}\}/(1 + a_{\sigma(j)}). \end{aligned}$$

We also have

$$(5.37) \quad \begin{aligned} \|\kappa_j \circ \phi^{-1} - \kappa_j\|_{(1-2\theta)r} &\leq 2nr \|\nabla_\zeta \kappa_j\|_{((1-\theta)r)^2} \cdot \|\phi^{-1} - \text{Id}\|_{(1-2\theta)r} \\ &\leq c_2(\|f\|_r + \|g\|_r)/(r\theta^{2n+3}), \end{aligned}$$

where the last inequality comes from (5.22) and (5.24). Using (5.33) and (5.37), one gets from (5.36) that

$$\|\xi_j(\hat{\kappa}_j - \kappa_j) + \hat{f}_j\|_{(1-3\theta)r} \leq c_5(\|f\|_r + \|g\|_r)/\theta^{4n+5}$$

for some constant $c_5 > 1$. Note that \hat{f}_j^* and $\xi_j(\hat{\kappa}_j - \kappa_j)^*$ are majorized by $(\xi_j(\hat{\kappa}_j - \kappa_j)(\xi\eta) + \hat{f}_j(\xi, \eta))^*$. Thus,

$$\|\hat{f}_j^*\|_{(1-4\theta)r} \leq c_5(\|f\|_r + \|g\|_r)/\theta^{6n+6},$$

$$(5.38) \quad \|(\hat{\kappa}_j - \kappa_j)^*\|_{((1-4\theta)r)^2} \leq c_5(\|f\|_r + \|g\|_r)/(r(1-4\theta)\theta^{6n+6}).$$

Since $\text{ord } \hat{f}_j, \text{ord } \hat{g}_j \geq d$, the Schwarz lemma gives

$$(5.39) \quad \|\hat{f}\|_{(1-5\theta)r} \leq c_5 \left(\frac{1-5\theta}{1-4\theta} \right)^d (\|f\|_r + \|g\|_r) / \theta^{6n+6}.$$

Obviously, (5.34) and (5.39) yield (5.30) for $c_3 > 2c_2\tilde{c}_2 + c_5$. Note that $(1-\theta/2)(1-4\theta)^2 > (1-5\theta)^2$. Using (5.38) and Cauchy inequalities, we get

$$\|\nabla_\zeta(\hat{\kappa} - \kappa)\|_{((1-5\theta)r)^2} \leq \tilde{c}_5(\|f\|_r + \|g\|_r) / (r^3\theta^{6n+7}).$$

In view of (5.19), we obtain

$$\|(\nabla_\zeta(\hat{\kappa} - \kappa))^*\|_{((1-6\theta)r)^2} \leq \tilde{c}_5(\|f\|_r + \|g\|_r) / (r^3\theta^{7n+8})$$

for some larger \tilde{c}_5 . The above inequality yields the first half of (5.31) for $c_3 > \tilde{c}_5$. For the second half of (5.31), note that

$$\hat{\kappa}_j^{-1} - \kappa_j^{-1} = \kappa_j \sum_{k \geq 1} \left(\frac{\kappa_j - \hat{\kappa}_j}{\kappa_j} \right)^k.$$

Thus, (5.38) and (5.32) yields

$$\begin{aligned} \|(\hat{\kappa}^{-1} - \kappa^{-1})^*\|_{((1-6\theta)r)^2} &\leq \frac{2}{\theta^{n+1}} \sum_{k \geq 1} \left(\frac{2c_5(\|f\|_r + \|g\|_r)}{r(1-4\theta)\theta^{6n+6}} \right)^k \\ &\leq 8c_5(\|f\|_r + \|g\|_r) / (r\theta^{7n+7}). \end{aligned}$$

This completes the proof of the proposition. □

Put

$$(5.40) \quad r_k = \frac{1}{2} \left(1 + \frac{1}{k+1} \right) r, \quad k = 0, 1, \dots$$

Rewrite the above as

$$r_{k+1} = (1 - 6\theta_k)r_k, \quad \theta_k = \frac{1}{6(k+2)^2}, \quad k = 0, 1, \dots$$

Let us first prove a numerical result.

LEMMA 5.8. — *Let r_k, θ_k be given as above, and let $0 < r_0 = r < 1$. Let b_k, B_k, K_k be nonnegative numbers satisfying*

$$(5.41) \quad \begin{aligned} b_{k+1} &\leq c_3(1-\theta_k)^{2^k} b_k / \theta_k^{6n+6}, \\ B_{k+1} &\leq B_k + c_3 b_k / (r_k^3 \theta_k^{7n+8}), \quad K_{k+1} \leq K_k + c_3 b_k / (r_k \theta_k^{7n+7}) \end{aligned}$$

for $k \geq 0$. Let c_4 be the constant given in Proposition 5.7. There exists $c_6 > 1$, independent of r , such that if

$$(5.42) \quad b_0 \leq r_0 \theta_0^{6n+6} / c_6, \quad r_0^2 B_0 \leq 1 / (32n^2), \quad K_0 < 3/2,$$

then for all $k > 0$

$$(5.43) \quad b_k \leq r_k \theta_k^{6n+6} / c_4, \quad r_k^2 B_k \leq 1 / (16n^2), \quad K_k < 2.$$

Proof. — Put

$$\hat{b}_k = r_k \theta_k^{7n+9} / (32n^2 c_3 c_4), \quad k = 0, 1, \dots$$

It is easy to see that

$$\hat{b}_{k+1} / \hat{b}_k = (1 - 6\theta_k) \theta_{k+1}^{7n+9} / \theta_k^{7n+9} \geq (1 - 6\theta_0) \theta_1^{7n+9} / \theta_0^{7n+9} \equiv c_7.$$

On the other hand, one has

$$b_{k+1} / b_k \leq c_3 (1 - \theta_k)^{2k} / \theta_k^{6n+6} \rightarrow 0, \quad k \rightarrow \infty.$$

Hence, there exists N , independent of r , such that

$$(5.44) \quad b_{k+1} / b_k \leq \hat{b}_{k+1} / \hat{b}_k, \quad k \geq N.$$

Set

$$c_8 = 1 + \max_{k \leq N} c_3 (1 - \theta_k)^{2k} / \theta_k^{6n+6}.$$

Then (5.42) implies that

$$b_k \leq b_0 c_8^N \leq r_0 \theta_0^{6n+6} c_8^N / c_6, \quad k \leq N.$$

This shows that for sufficiently large c_6 , one has $b_k \leq \hat{b}_k$, $k = 0, \dots, N$. Now (5.44) implies that $b_k \leq \hat{b}_k$ for all k . In particular, the estimates (5.43) hold for b_k . Also,

$$r_{k+1}^2 B_{k+1} \leq r_0^2 B_0 + \sum_{j=1}^k \frac{c_3 b_j}{r_j \theta_j^{7n+8}} \leq 1 / (32n^2) + \sum_{j=1}^{\infty} \theta_j / (32n^2) < 1 / (16n^2)$$

and $K_{k+1} \leq K_0 + \sum_{j=1}^k c_3 b_j / r_j \theta_j^{7n+7} < 2$. This completes the proof of the lemma. □

We now complete the proof of Theorem 4.9 as follows. With the notation introduced at the beginning of this section, we would like to show the convergence of a subsequence of $\phi_1^{-1} \circ \phi_2^{-1} \circ \dots \circ \phi_k^{-1}$. To this end, it suffices to show that the sequence is bounded, or that

$$(5.45) \quad \phi_k^{-1}: \Delta_{r_k} \rightarrow \Delta_{r_{k-1}}, \quad k = 1, 2, \dots$$

are well-defined for suitable $r_0 = r$.

Start with

$$m_j^{(1)}(\xi, \eta) = m_j(\xi, \eta) = \frac{\xi_j \kappa_j^{(1)}(\xi, \eta) + f_j^{(1)}(\xi, \eta)}{\eta_{\sigma(j)} + g_{\sigma(j)}^{(1)}(\xi, \eta)}.$$

Set

$$b_0 = \|f^{(1)}\|_{r_0} + \|g^{(1)}\|_{r_0}, B_0 = \|(\nabla_\zeta \kappa^{(1)})^*\|_{r_0^2}, K_0 = \|((\kappa^{(1)})^{-1})^*\|_{r_0^2}.$$

Choose $r_0 = r$ so small that initial condition (5.42) holds. Applying Lemma 5.5 to φ_1 and ϕ_1 gives us (5.45) for ϕ_1 . Put $m_j^{(2)} = m_j^{(1)} \circ \phi_1^{-1}$ and

$$b_1 = \|f^{(2)}\|_{r_1} + \|g^{(2)}\|_{r_1}, B_1 = \|(\nabla_\zeta \kappa^{(2)})^*\|_{r_1^2}, K_1 = \|((\kappa^{(2)})^{-1})^*\|_{r_1^2}.$$

Since $d_k > 2^k$, Proposition 5.7 says that inequality (5.41) holds for $k = 1$. Consequently, Lemma 5.8 implies that the initial condition (5.43) holds for $k = 1$. Recursively, one sees that the mappings (5.45) are well-defined for all k . This completes the proof of Theorem 4.9.

6. Hamiltonian vector fields.

This section is to apply results for holomorphic symplectic mappings obtained in previous sections to holomorphic Hamiltonian systems.

Consider a holomorphic Hamiltonian function

$$(6.1) \quad H(\xi, \eta) = \sum \lambda_j \xi_j \eta_j + h(\xi, \eta), \quad h(\xi, \eta) = O(3).$$

The corresponding Hamiltonian system of H is

$$(6.2) \quad d\xi_j/dt = \lambda_j \xi_j + H_{\eta_j}, \quad d\eta_j/dt = -\lambda_j \eta_j - H_{\xi_j}, \quad j = 1, \dots, n.$$

The formal Birkhoff normal form of H is then given by

$$(6.3) \quad \widehat{H}(\xi, \eta) = \sum \lambda_j \xi_j \eta_j + \widehat{h}(\xi, \eta),$$

where \widehat{h} is a formal power series in products $\xi_1 \eta_1, \dots, \xi_n \eta_n$, provided $\lambda = (\lambda_1, \dots, \lambda_n)$ satisfies the non-resonance condition

$$(6.4) \quad \alpha \cdot \lambda = \alpha_1 \lambda_1 + \dots + \alpha_n \lambda_n \neq \lambda_j, \quad \alpha \in \mathbb{Z}^n$$

for all $\alpha \neq e_j = (0, \dots, 1, \dots, 0)$. The formal flow of the Hamiltonian (6.3) is given by

$$\widehat{\varphi}_t: \xi'_j = \xi_j e^{t\widehat{H}_{\xi_j}}, \quad \eta'_j = \eta_j e^{-t\widehat{H}_{\eta_j}}, \quad j = 1, \dots, n$$

for $t \in \mathbb{C}$. We say that a real or complex time t is *non-exceptional* with respect to $\lambda_1, \dots, \lambda_n$, if the eigenvalues $e^{t\lambda_1}, \dots, e^{t\lambda_n}$ satisfy the non-resonance condition (1.2). Note that, given $\lambda_1, \dots, \lambda_n$ satisfying (6.4), the exceptional t values are countable.

We need the following.

PROPOSITION 6.1. — *Let H be the Hamiltonian function (6.1) with λ_j satisfying (6.4), and let φ_t be its Hamiltonian flow. Assume that t is non-exceptional. If φ_t is in its Birkhoff normal form, so is H .*

Proof. — Replacing H by tH , one may assume that $t = 1$. We need to prove that H is a function of $\xi_1\eta_1, \dots, \xi_n\eta_n$. Put

$$(6.5) \quad H(x, y) = \tilde{H}(\xi_1\eta_1, \dots, \xi_n\eta_n) + H_k(\xi, \eta),$$

where $H_k(\xi, \eta) = O(k+1)$, and all terms in the power series expansion of \tilde{H} have order at most k in ξ, η . Note that for two holomorphic vector fields $v(\xi, \eta), \tilde{v}(\xi, \eta)$ with $v(0) = \tilde{v}(0) = 0$ and $\tilde{v}(\xi, \eta) - v(\xi, \eta) = O(k)$, the corresponding flows φ_t and $\tilde{\varphi}_t$ satisfy

$$\tilde{\varphi}_t(\xi, \eta) - \varphi_t(\xi, \eta) = O(|(\xi, \eta)|^k)$$

uniformly for $|t| < T < \infty$. The flow φ_t of (6.5) is of the form

$$\begin{aligned} \xi_j(t) &= \xi_j e^{t\tilde{H}_{\zeta_j}(\xi\eta)} + \sum a_{j\alpha\beta}(t)\xi^\alpha\eta^\beta, \\ \eta_j(t) &= \eta_j e^{-t\tilde{H}_{\zeta_j}(\xi\eta)} + \sum b_{j\alpha\beta}(t)\xi^\alpha\eta^\beta \end{aligned}$$

with

$$a_{j\alpha\beta}(t) = b_{j\alpha\beta}(t) \equiv 0, \quad |\alpha| + |\beta| < k.$$

For $\alpha + \beta = k$, one gets

$$da_{j\alpha\beta}/dt - \lambda_j a_{j\alpha\beta} = (\beta_j + 1)H_{\alpha, \beta + e_j} e^{(\alpha - \beta) \cdot \lambda t}, \quad \alpha - \beta \neq e_j$$

with $a_{j\alpha\beta}(0) = 0$. Hence

$$a_{j\alpha\beta}(t) = (\beta_j + 1) \frac{e^{(\alpha - \beta) \cdot \lambda t} - e^{\lambda_j t}}{(\alpha - \beta) \cdot \lambda - \lambda_j} H_{\alpha, \beta + e_j}$$

for $|\alpha| + |\beta| = k, \alpha - \beta \neq e_j$. Since φ_1 is in the normal form, then $a_{j\alpha\beta}(1)$ vanish for $\alpha - \beta \neq e_j$ and $|\alpha| + |\beta| = k$. Hence, $H_{\alpha\beta} = 0$ for $\alpha \neq \beta, |\alpha| + |\beta| = k + 1$, and $\beta \neq 0$. Computing the coefficients of $b_{j(\alpha - e_j)0}$ yields $H_{\alpha 0} = 0$ for $|\alpha| = k + 1$. Thus, $[H]_{k+1}$ is a power series in $\xi_1\eta_1, \dots, \xi_n\eta_n$. By induction, H is a power series in $\xi_1\eta_1, \dots, \xi_n\eta_n$. The proof of the proposition is complete. \square

With the above observation, we now introduce a notion. Let M be the germ of a set at $0 \in \mathbb{C}^{2n}$. We say that the germ M is *weakly invariant* under the flow φ_t , if the germ $\varphi_t(M)$ is contained in M for some non-exceptional $t \in \mathbb{C}$. From Theorems 1.1 and 4.1 and Proposition 6.1, we obtain the following.

COROLLARY 6.2. — *Let φ_t be a holomorphic Hamiltonian flow defined by (6.2) and (6.4), and let M be a Levi-flat real analytic set defined by (1.4). Assume that M is weakly invariant under the flow φ_t . Then the Hamiltonian system (6.2) is normalizable by convergent holomorphic symplectic mappings. Moreover, M is invariant under φ_t for all $t \in \mathbb{C}$.*

From Theorem 4.6, we have the following.

COROLLARY 6.3. — *Let M be a Levi-flat analytic set defined by (1.5), and let φ_t be the holomorphic Hamiltonian flow of (6.2) and (6.4). If M is invariant under φ_{t_0} for some non-exceptional t_0 , then M is invariant under φ_{st_0} for all $s \in \mathbb{R}$.*

Proof. — By Theorem 3.6, M is actually given by $r_1 = \dots = r_n = 0$ with r_j being given by (4.23). Let $m_j = F_j(\xi, \eta)/G_{\sigma(j)}(\xi, \eta)$ be the meromorphic functions defined by (4.25). Take a formal symplectic transformation $\Phi = \text{Id} + O(2)$ which transforms the Hamiltonian (6.1) into the Birkhoff normal form. Fix a non-exceptional t_0 such that $\varphi_{t_0}(M) = M$. By Theorem 4.6, we know that $m_j \circ \Phi^{-1}$ is an eigenfunction of the flow $\Phi \circ \varphi_t \circ \Phi^{-1}$ with eigenvalue $e^{t(\lambda_j + \lambda_{\sigma(j)})}$. Hence, m_j is also an eigenfunction of φ_t with the same eigenvalue. Thus, $F_j \circ \varphi_t = u_j e^{t\lambda_j} F_j$, $G_{\sigma(j)} \circ \varphi_t = u_j e^{-t\lambda_{\sigma(j)}} G_{\sigma(j)}$, where $u_j = 1 + O(1)$ are holomorphic functions dependent of t . Note that $|e^{t\lambda_j}| = |e^{-t\lambda_{\sigma(j)}}|$ hold for $t = t_0$, and hence holds for all $t = st_0$ when s is real. Now, it is straightforward that for each real s , $r_j \circ \varphi_{st_0} = v_j r_j$ for some $v_j \neq 0$. In particular, M is invariant under φ_{st_0} . This completes the proof of the corollary. \square

Finally, Theorem 1.2 gives us the following.

COROLLARY 6.4. — *Let M be as in Theorem 1.2, and let φ_t be as in Corollary 6.3. Assume that M is weakly invariant under the flow φ_t . The Hamiltonian system (6.2) is normalizable by holomorphic symplectic mappings.*

Appendix: Birkhoff normal forms in terms of time-one mappings of formal Hamiltonian systems and formal generating functions.

There are two ways to formulate the Birkhoff normal form for holomorphic symplectic mappings, of which one is of the form (1.3). The

other normal form can be described in terms of generating functions. For the sake of the reader, we derive one normal form from another in this appendix.

Let $(\xi', \eta') = \varphi(\xi, \eta)$ be a holomorphic symplectic mapping (1.1). Regarding the mapping as a graph over ξ, η' , on which the 1-form $\sum(\eta'_j d\xi_j + \xi'_j d\eta'_j)$ is closed, one has

$$(7.1) \quad \xi'_j = \lambda_j \xi_j + S_{\eta'_j}(\xi, \eta'), \quad \eta_j = \lambda_j \eta'_j + S_{\xi_j}(\xi, \eta')$$

for some convergent power series $S(\xi, \eta') = O(3)$. Assume that λ_j satisfy the non-resonance condition (1.2). By comparing the coefficients as in the Hamiltonian case (see [4], p. 85), there is a formal power series $T(\xi, \eta') = O(3)$ such that for the formal symplectic mapping Φ determined by

$$\xi'_j = \xi_j + T_{\eta'_j}(\xi, \eta'), \quad \eta_j = \eta'_j + T_{\xi_j}(\xi, \eta'),$$

$\hat{\varphi} = \Phi \circ \varphi \circ \Phi^{-1}$ is of the form (7.1) with

$$(7.2) \quad S = \hat{S}(\xi\eta')$$

being a formal power series in the products $\zeta_j = \xi_j \eta'_j$. Thus, the formal mapping given by (7.1)–(7.2) can also be referred as the Birkhoff normal form of φ .

Next, we want to put the formal mapping (7.1)–(7.2) into the form (1.3). From (7.1)–(7.2), one has

$$\xi'_j = \lambda_j \xi_j (1 + \lambda_j^{-1} \hat{S}_{\zeta_j}(\xi\eta')), \quad \eta'_j = \lambda_j^{-1} \eta_j (1 + \lambda_j^{-1} \hat{S}_{\zeta_j}(\xi\eta'))^{-1}$$

for $j = 1, \dots, n$. Hence, $\xi_j \eta_j$ are invariant by $\hat{\varphi}$. Solving for η'_j from the last n equations above, one sees that

$$(7.3) \quad \xi'_j = \lambda_j \xi_j e^{G_j(\xi\eta)}, \quad \eta'_j = \lambda_j^{-1} \eta_j e^{-G_j(\xi\eta)}$$

for some formal power series $G_j(\xi\eta)$ with $G_j(0) = 0$. Note that G_j are determined by

$$(7.4) \quad e^{G_j(\zeta)} = 1 + \lambda_j^{-1} \hat{S}_{\zeta_j}(\lambda^{-1} \zeta e^{G(\zeta)}), \quad j = 1, \dots, n,$$

where $\lambda^{-1} \zeta e^{G(\zeta)} \equiv (\lambda_1^{-1} \zeta_1 e^{G_1(\zeta)}, \dots, \lambda_n^{-1} \zeta_n e^{G_n(\zeta)})$. On the other hand, (7.3) yields

$$\sum d\xi'_j \wedge d\eta'_j - \sum d\xi_j \wedge d\eta_j = d \sum G_j(\xi_j \eta_j) d(\xi_j \eta_j).$$

The left-hand side vanishes. Hence, $G_j = H_{\zeta_j}$ for some formal power series H in ζ .

Conversely, if $(\xi', \eta') = \hat{\varphi}(\xi, \eta)$ is given by (1.3), then $\hat{\varphi}$ preserves the products $\xi_j \eta_j$ and is of the form

$$\xi'_j = \lambda_j \xi_j (1 + K_j(\xi \eta')), \quad \eta'_j = \lambda_j \eta'_j (1 + K_j(\xi \eta'))$$

with

$$(7.5) \quad 1 + K_j(\zeta) = e^{\omega_j(\lambda \zeta K(\zeta))}, \quad j = 1, \dots, n.$$

Now, one has

$$\sum \xi'_j d\eta'_j + \eta'_j d\xi'_j = d\left(\sum \lambda_j \xi_j \eta'_j\right) + \sum \lambda_j K_j(\xi \eta') d(\xi_j \eta'_j).$$

Hence, the last summand is closed, from which it follows that $K_j = \lambda_j^{-1} \hat{S}_{\zeta_j}(\zeta)$ for some formal power series \hat{S} . This shows that $\hat{\varphi}$ is of the form (7.1)–(7.2). From (7.4)–(7.5), one can also see that the convergence of \hat{S} implies the convergence of H , and vice versa.

Therefore, two normal forms (1.3) and (7.1)–(7.2) are equivalent, i.e., if one of the normal forms is realized by a holomorphic symplectic transformation, so is the other.

BIBLIOGRAPHY

- [1] M. ARTIN, On the solutions of analytic equations, *Invent. Math.*, 5 (1968), 277–291.
- [2] E. BEDFORD, Holomorphic continuation of smooth functions over Levi-flat hypersurfaces, *Trans. Amer. Math. Soc.*, 232 (1977), 323–341.
- [3] G.D. BIRKHOFF, Surface transformations and their dynamical applications, *Acta Math.*, 43 (1920), 1–119.
- [4] G.D. BIRKHOFF, *Dynamical Systems*, AMS Coll. Publ., vol. 9, 1927, reprinted 1966.
- [5] F. BRUHAT and H. CARTAN, Sur la structure des sous-ensembles analytiques réels, *C. R. Acad. Sci. Paris*, 244 (1957), 988–991.
- [6] D. BURNS and X. GONG, Singular Levi-flat real analytic hypersurfaces, *Amer. J. Math.*, 121, n° 1 (1999), 23–53.
- [7] H. CARTAN, Variétés analytiques réelles et variétés analytiques complexes, *Bull. Soc. Math. France*, 85 (1957), 77–99.
- [8] K. DIEDERICH and J.E. FORNAESS, Pseudoconvex domains with real-analytic boundary, *Ann. Math. (2)*, 107, n° 2 (1978), 371–384.
- [9] L.H. ELIASSON, Normal forms for Hamiltonian systems with Poisson commuting integrals—elliptic case, *Comment. Math. Helv.*, 65, n° 1 (1990), 4–35.
- [10] H. ITO, Convergence of Birkhoff normal forms for integrable systems, *Comment. Math. Helv.*, 64 (1989), 412–461.
- [11] J.K. MOSER, The analytic invariants of an area-preserving mapping near a hyperbolic fixed point, *Comm. Pure Appl. Math.*, 9 (1956), 673–692.

- [12] H. RÜSSMANN, Über die Existenz einer Normalform inhaltstreuer elliptischer Transformationen, *Math. Ann.*, 137 (1959), 64–77.
- [13] H. RÜSSMANN, Über das Verhalten analytischer Hamiltonscher Differentialgleichungen in der Nähe einer Gleichgewichtslösung, *Math. Ann.*, 154 (1964), 285–300.
- [14] H. RÜSSMANN, Über die Normalform analytischer Hamiltonscher Differentialgleichungen in der Nähe einer Gleichgewichtslösung, *Math. Ann.*, 169 (1967), 55–72.
- [15] B. SEGRE, Intorno al problem di Poincaré della rappresentazione pseudo-conform, *Rend. Acc. Lincei*, 13 (1931), 676–683.
- [16] C.L. SIEGEL, On integrals of canonical systems, *Ann. Math.*, 42 (1941), 806–822.
- [17] C.L. SIEGEL, Über die Existenz einer Normalform analytischer Hamiltonscher Differentialgleichungen in der Nähe einer Gleichgewichtslösung, *Math. Ann.*, 128 (1954), 144–170.
- [18] J. VEY, Sur certains systèmes dynamiques séparables, *Amer. J. Math.*, 100 (1978), 591–614.
- [19] S.M. WEBSTER, On the mapping problem for algebraic real hypersurfaces, *Invent. Math.*, 43, n° 1 (1977), 53–68.

Manuscrit reçu le 11 avril 2000,
accepté le 26 juillet 2000.

Xianghong GONG,
Oklahoma State University
Department of Mathematics
Stillwater, OK 74078 (USA).

Current address:
University of Wisconsin - Madison
Department of Mathematics
Madison, WI 53706 (USA).
gong@math.wisc.edu