



ANNALES

DE

L'INSTITUT FOURIER

Indranil BISWAS & Norbert HOFFMANN

A Torelli theorem for moduli spaces of principal bundles over a curve

Tome 62, n° 1 (2012), p. 87-106.

http://aif.cedram.org/item?id=AIF_2012__62_1_87_0

© Association des Annales de l'institut Fourier, 2012, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

A TORELLI THEOREM FOR MODULI SPACES OF PRINCIPAL BUNDLES OVER A CURVE

by Indranil BISWAS & Norbert HOFFMANN (*)

ABSTRACT. — Let X and X' be compact Riemann surfaces of genus ≥ 3 , and let G and G' be nonabelian reductive complex groups. If one component $\mathcal{M}_G^d(X)$ of the coarse moduli space for semistable principal G -bundles over X is isomorphic to another component $\mathcal{M}_{G'}^{d'}(X')$, then X is isomorphic to X' .

RÉSUMÉ. — Soient X et X' des surfaces de Riemann compactes de genre au moins 3, et G et G' des groupes complexes réductifs non abéliens. Si une composante $\mathcal{M}_G^d(X)$ de l'espace des modules de G -fibrés principaux semi-stables sur X est isomorphe à une composante $\mathcal{M}_{G'}^{d'}(X')$, alors X est isomorphe à X' .

Introduction

Let X be a compact connected Riemann surface of genus g_X . The classical Torelli theorem says that the isomorphism class of X is uniquely determined by the isomorphism class of the Jacobian $\text{Pic}^0(X)$, together with its canonical principal polarization Θ .

Several authors have studied nonabelian analogues, replacing Jacobians by moduli spaces of vector bundles. Suppose $g_X \geq 3$. Given a line bundle L on X , let $\mathcal{M}_{n,L}(X)$ denote the moduli space of semistable vector bundles E over X of rank n with fixed determinant $\bigwedge^n E \cong L$. Then the isomorphism

Keywords: Principal bundle, moduli space, Torelli theorem.

Math. classification: 14D20, 14C34.

(*) We are very grateful to the referee for pointing out a gap in an earlier version of this paper, and also for some valuable suggestions. We thank D. Prasad for a useful conversation. The first author wishes to thank the Freie Universität Berlin and the Institute for Mathematical Sciences at Chennai for hospitality. The second author was supported by the SFB 647: Raum - Zeit - Materie. He thanks TIFR (Mumbai) and IMSc (Chennai) for hospitality while some part of this work was done.

class of X is uniquely determined by the isomorphism class of the projective variety $\mathcal{M}_{n,L}(X)$.

This was first proved for $n = 2$ and $\deg(L)$ odd, by Mumford-Newstead [10, p. 1201, Corollary] and Tyurin [19, Theorem 1].

It was then proved more generally for n coprime to $\deg(L)$, by Tyurin [20, Theorem 1] and Narasimhan-Ramanan [12, Theorem 3].

Finally, this statement was proved in full generality by Kouvidakis-Pantev [6, Theorem E], and then later for $g_X \geq 4$ also by Hwang-Ramanan [5, Theorem 5.1] and by Sun [18, Corollary 1.3].

These proofs used either an intermediate Jacobian of the moduli space [10, 12], or Higgs bundles and the Hitchin map [6, 5], or rational curves on the moduli spaces [19, 20, 18].

Our aim here is to address a similar question for the moduli spaces of principal bundles. Given a connected reductive complex linear algebraic group G , let $\mathcal{M}_G^d(X)$ denote the moduli space of semistable principal G -bundles over X of topological type $d \in \pi_1(G)$.

THEOREM 0.1. — *Let G and G' be nonabelian connected reductive complex groups. Let $d \in \pi_1(G)$ and $d' \in \pi_1(G')$ be given. Let X and X' be compact Riemann surfaces of genus ≥ 3 . If $\mathcal{M}_G^d(X)$ is isomorphic to $\mathcal{M}_{G'}^{d'}(X')$, then X is isomorphic to X' .*

For the proof, our strategy is fundamentally different from the earlier ones. Whereas [6] and [5] start from the stable locus, our starting point is the strictly semistable locus. We show that it lies in the singular locus and is characterized by its type of singularities. Using powers of its anticanonical line bundle, we map it to a projective space. The fibers of this map allow us to reconstruct the Jacobian of X and its principal polarization. Then the usual Torelli theorem applies.

This method does not work if the strictly semistable locus is empty. But this case is rare, and can in fact be reduced to the earlier results on moduli spaces of vector bundles in the coprime case.

The structure of this paper is as follows. Section 1 deals with quotient singularities arising from a group action. We give a criterion to distinguish the case of a finite group from the torus \mathbb{C}^* .

In Section 2, we collect some basic facts about the moduli spaces $\mathcal{M}_G^d(X)$. We use several tools valid only in characteristic 0, like representations of $\pi_1(X)$, and the stability of induced bundles.

Section 3 describes the strictly semistable locus in $\mathcal{M}_G^d(X)$. We relate it to moduli spaces of principal H -bundles for Levi subgroups H of maximal

parabolic subgroups in G . Furthermore, we characterize it by the singularities it contains. As a byproduct, we show that the smooth locus of $\mathcal{M}_G^d(X)$ coincides with the regularly stable locus.

In the final Section 4, we prove Theorem 0.1, by reconstructing the compact Riemann surface X from the projective variety $\mathcal{M}_G^d(X)$. In fact we prove a slightly stronger result, namely that X is determined up to isomorphy by the smooth locus of $\mathcal{M}_G^d(X)$; see Theorem 4.1.

1. Some quotient singularities

Let S and S' be schemes of finite type over \mathbb{C} . We say that S is near a closed point $s_0 \in S$ *analytically isomorphic* to S' near a closed point $s'_0 \in S_0$ if for some open neighborhoods $s_0 \in U \subseteq S$ and $s'_0 \in U' \subseteq S'$ in the Euclidean topology, there exists a biholomorphic isomorphism $U \rightarrow U'$ that maps s_0 to s'_0 .

We always denote by $S^{\text{sing}} \subseteq S$ the singular locus of S . Let G be a reductive linear algebraic group over \mathbb{C} . We assume that G acts linearly on a finite-dimensional complex vector space V . Let

$$p : V \longrightarrow S := V//G = \text{Spec}(\text{Sym}(V^*)^G)$$

be the GIT-quotient, and put $s_0 := p(0) \in S$.

LEMMA 1.1. — *Let G be finite and nontrivial. Suppose that the fixed locus $V^g \subseteq V$ has codimension ≥ 2 for all $g \in G$ with $g \neq 1$.*

- i) *The quotient $S = V//G$ is singular in $s_0 = p(0)$.*
- ii) *If $s_0 \in U \subseteq S$ is an open neighborhood in the Euclidean topology such that $U \setminus S^{\text{sing}}$ is connected, then $\pi_1(U \setminus S^{\text{sing}})$ is nontrivial.*

Proof. — Using the assumption on $\text{codim } V^g$, [11, Lemma 4.4] implies

$$p^{-1}(S^{\text{sing}}) = \bigcup_{g \neq 1} V^g.$$

This proves (i). In the situation of (ii), $p^{-1}(U \setminus S^{\text{sing}})$ is connected because s_0 has only one inverse image. Hence $p^{-1}(U \setminus S^{\text{sing}})$ is a nontrivial covering of $U \setminus S^{\text{sing}}$, which is therefore not simply connected. \square

LEMMA 1.2. — *Let $G = \mathbb{C}^*$ act linearly on the finite-dimensional complex vector space V , with associated weight space decomposition*

$$V = \bigoplus_{m \in \mathbb{Z}} V_m.$$

Suppose that V_{-1} and V_1 both have dimension ≥ 2 .

- i) The quotient $S = V//\mathbb{C}^*$ is singular in $s_0 = p(0)$.
- ii) Every neighborhood $s_0 \in U \subseteq S$ in the Euclidean topology contains an open neighborhood $s_0 \in U' \subseteq U$ such that $U' \setminus S^{\text{sing}}$ is connected and $\pi_1(U' \setminus S^{\text{sing}})$ is trivial.

Proof. — It is easy to verify $S = (V_+ \oplus V_-)//\mathbb{C}^* \times V_0$ with

$$V_+ := \bigoplus_{m>0} V_m \quad \text{and} \quad V_- := \bigoplus_{m<0} V_m.$$

Replacing V by $V_+ \oplus V_-$ if necessary, we may thus assume $V_0 = 0$ without loss of generality. Then the GIT-stable locus $V^{\text{stab}} \subseteq V$ is

$$V^{\text{stab}} = V \setminus (V_+ \cup V_-),$$

and the image $p(V_+ \cup V_-)$ of $V \setminus V^{\text{stab}}$ is just the point $s_0 \in S$. Let

$$V^{\mu_n} = \bigoplus_{m \in n\mathbb{Z}} V_m \subseteq V$$

be the fixed locus of the finite subgroup $\mu_n \subseteq \mathbb{C}^*$. We claim

$$(1.1) \quad p^{-1}(S^{\text{sing}}) = V_+ \cup V_- \cup \bigcup_{n \geq 2} V^{\mu_n}.$$

To check this, we first consider a stable point $v \in V^{\text{stab}}$. Its image $p(v) \in S$ is smooth if and only if its stabilizer in \mathbb{C}^* is trivial, according to Luna's étale slice theorem [8] and Lemma 1.1.

Suppose next that $V^{\mu_n} \cap V^{\text{stab}}$ is non-empty for some $n \geq 2$. Then it is dense in V^{μ_n} , but also contained in the closed subset $p^{-1}(S^{\text{sing}})$. Hence $0 \in V^{\mu_n} \subseteq p^{-1}(S^{\text{sing}})$, which implies $s_0 \in S^{\text{sing}}$. It follows that $p^{-1}(S^{\text{sing}})$ contains the unstable locus $V_+ \cup V_-$ in this case.

Now suppose $V^{\mu_n} \subseteq V_+ \cup V_-$ for all $n \geq 2$. Then the canonical map

$$\tilde{S} := (V \setminus \bigcup_{n \geq 2} V^{\mu_n})/\mathbb{C}^* \longrightarrow S$$

is surjective. Here \mathbb{C}^* acts freely, so the set \tilde{S} is a (not necessarily separated) complex manifold. Applying [11, Lemma 4.4] to this map, we get $s_0 \in S^{\text{sing}}$. Hence $p^{-1}(S^{\text{sing}})$ contains $V_+ \cup V_-$ also in this case. This proves the claim (1.1), and in particular part (i) of the lemma.

Let a Euclidean neighborhood $s_0 \in U \subseteq S$ be given. For each $m \in \mathbb{Z}$, we choose a basis $(x_{m,i})_{1 \leq i \leq \dim V_m}$ of the vector space $(V_m)^*$. The \mathbb{C} -algebra $\text{Sym}(V^*)^{\mathbb{C}^*}$ is generated by finitely many nonconstant monomials f_1, \dots, f_N in the variables $x_{m,i}$. They provide a closed embedding $S \hookrightarrow \mathbb{C}^N$ with $s_0 \mapsto 0$. Hence U contains the open neighborhood

$$s_0 \in U' := \{s \in S : |f_n(s)| < \varepsilon \text{ for } n = 1, \dots, N\}$$

for some $\varepsilon > 0$. Because the f_n are monomials, the open subset

$$p^{-1}(U') = \{v \in V : |f_n(v)| < \varepsilon \text{ for } n = 1, \dots, N\} \subseteq V$$

is a star-shaped neighborhood of 0, in the sense that $\lambda v \in p^{-1}(U')$ holds for all $v \in p^{-1}(U')$ and $\lambda \in [0, 1]$. In particular, $p^{-1}(U')$ is contractible. Since we are only removing finitely many linear subspaces of complex codimension ≥ 2 , it follows that $p^{-1}(U' \setminus S^{\text{sing}})$ is connected and simply connected. Consequently, $U' \setminus S^{\text{sing}}$ is also connected and simply connected, because

$$p : p^{-1}(U' \setminus S^{\text{sing}}) \longrightarrow U' \setminus S^{\text{sing}}$$

is a fibration with connected fiber \mathbb{C}^* . □

2. Moduli spaces of principal bundles

Let X be a compact connected Riemann surface of genus $g_X \geq 3$. Let G from now on be a reductive connected linear algebraic group over \mathbb{C} . We denote by

$$\mathcal{M}_G = \mathcal{M}_G(X)$$

the coarse moduli space of semistable algebraic principal G -bundles E over X . Its connected components

$$\mathcal{M}_G^d = \mathcal{M}_G^d(X) \subseteq \mathcal{M}_G(X)$$

are irreducible normal projective varieties of dimension

$$\dim \mathcal{M}_G^d = (g_X - 1) \dim_{\mathbb{C}} G + \dim_{\mathbb{C}} Z_G$$

where Z_G denotes the center of G . These connected components are indexed by elements $d \in \pi_1(G)$, which correspond to topological types of algebraic principal G -bundles E over X . In the case $G = \text{GL}_n$ of vector bundles, the topological type $d \in \pi_1(\text{GL}_n) = \mathbb{Z}$ is their degree.

The construction of such moduli spaces is carried out in [15]; see also [3] and [16] for generalizations to higher dimensions. We recall from [15] that one has a dense open subscheme

$$\mathcal{M}_G^{\text{stab}} \subseteq \mathcal{M}_G$$

whose closed points correspond bijectively to isomorphism classes of stable principal G -bundles over X . Closed points in \mathcal{M}_G correspond bijectively to isomorphism classes of polystable principal G -bundles over X , and also to S -equivalence classes of semistable principal G -bundles over X , in the sense of [15, Definition 3.6].

Let \mathfrak{g} be the Lie algebra of G . Given a principal G -bundle E over X , we denote the associated adjoint vector bundle over X by

$$\mathrm{ad}(E) := E \times^G \mathfrak{g} := (E \times \mathfrak{g})/G.$$

If E is semistable, then $\mathrm{ad}(E)$ is also semistable [15, Corollary 3.18]. The vector space

$$H^1(X, \mathrm{ad}(E))$$

parameterizes infinitesimal deformations of E according to standard deformation theory. On it, we have an adjoint action of the automorphism group $\mathrm{Aut}(E)$. If E is polystable, then the GIT-quotient

$$H^1(X, \mathrm{ad}(E)) // \mathrm{Aut}(E)$$

is near 0 analytically isomorphic to \mathcal{M}_G near the point $[E]$. This is a standard consequence of Luna's étale slice theorem [8].

The center Z_G of G is a normal subgroup of $\mathrm{Aut}(E)$ for every principal G -bundle E over X . E is stable or polystable or semistable if and only if the induced principal G/Z_G -bundle E/Z_G over X is so, since both bundles have the same set of reductions to parabolic subgroups; cf. [14, Proposition 7.1]. If E is stable, then $\mathrm{Aut}(E)/Z_G$ is finite by [14, Proposition 3.2]. If moreover $\mathrm{Aut}(E) = Z_G$, then E is called *regularly stable*; in this case, \mathcal{M}_G is smooth at $[E]$.

The canonical exact sequence of reductive groups

$$1 \longrightarrow [G, G] \longrightarrow G \longrightarrow G/[G, G] \longrightarrow 1$$

shows that $\pi_1([G, G])$ is the torsion subgroup of $\pi_1(G)$. We put

$$\mathcal{M}'_G := \bigcup_{d \in \pi_1([G, G])} \mathcal{M}_G^d \subseteq \mathcal{M}_G.$$

If G is semisimple, then $\mathcal{M}'_G = \mathcal{M}_G$.

Choose a maximal compact subgroup $K_G \subseteq G$, and a base-point $x_0 \in X$. Let $\tilde{X} \rightarrow X$ be the universal covering of X . Given a group homomorphism $\rho : \pi_1(X, x_0) \rightarrow K_G$, the principal G -bundle

$$E_\rho := \tilde{X} \times^\rho G := (\tilde{X} \times G)/\pi_1(X, x_0)$$

is polystable, and its moduli point $[E_\rho] \in \mathcal{M}_G$ is contained in \mathcal{M}'_G . In this way, closed points in \mathcal{M}'_G correspond bijectively to homomorphism $\rho : \pi_1(X, x_0) \rightarrow K_G$ up to conjugacy [15, Corollary 3.15.1].

LEMMA 2.1. — *The set of points $[E_\rho] \in \mathcal{M}'_G$ coming from homomorphisms $\rho : \pi_1(X, x_0) \rightarrow K_G$ with dense image is Zariski-dense in \mathcal{M}'_G .*

Proof. — There are only countably many conjugacy classes of closed subgroups $K \subsetneq K_G$, according to [1, Proposition 10.12]. For any given closed subgroup $K \subsetneq K_G$, the conjugacy classes of homomorphisms $\rho : \pi_1(X, x_0) \rightarrow K_G$ factoring through K form a closed real analytic subset of \mathcal{M}'_G of a smaller dimension. Due to the Baire category theorem, the union of these subsets in \mathcal{M}'_G has empty interior (with respect to the Euclidean topology). This shows that the homomorphisms $\rho : \pi_1(X, x_0) \rightarrow K_G$ with dense image are dense in \mathcal{M}'_G for the Euclidean topology, and a fortiori for the Zariski topology. \square

LEMMA 2.2. — *Consider the polystable principal G -bundle E_ρ over X given by a homomorphism $\rho : \pi_1(X, x_0) \rightarrow K_G$. Its automorphism group $\text{Aut}(E_\rho)$ is canonically isomorphic to*

$$C_G(\rho) := \{g \in G : g \cdot \rho(\omega) = \rho(\omega) \cdot g \text{ for all } \omega \in \pi_1(X, x_0)\}.$$

Proof. — Every element $g \in C_G(\rho)$ defines a $\pi_1(X, x_0)$ -equivariant automorphism of the trivial bundle $(\tilde{X} \times G)$ over \tilde{X} , which descends to an automorphism of E_ρ . This embeds $C_G(\rho)$ into $\text{Aut}(E_\rho)$.

Now let $\varphi \in \text{Aut}(E_\rho)$ be given. Choose an embedding $G \hookrightarrow \text{GL}_r$ that maps K_G to the unitary group U_r . Then φ defines a nonzero holomorphic section of the unitary vector bundle

$$\tilde{X} \times^\rho \text{End}(\mathbb{C}^r)$$

over X . This vector bundle is polystable of degree 0, so the section lies in a trivial direct summand \mathcal{O}_X . Such a trivial direct summand corresponds, under the Narasimhan-Seshadri correspondence, to a line in $\text{End}(\mathbb{C}^r)$ on which $\pi_1(X, x_0)$ acts trivially.

Hence the section in question comes from a fixed point in $\text{End}(\mathbb{C}^r)$. But this fixed point has to be in the image of $G \hookrightarrow \text{GL}_r \subseteq \text{End}(\mathbb{C}^r)$. This shows that φ indeed comes from an element of $C_G(\rho)$. \square

PROPOSITION 2.3. — *Let E be a principal G -bundle over X . Suppose that E is stable, but not regularly stable.*

- i) *The point $[E] \in \mathcal{M}_G$ is singular.*
- ii) *There is a neighborhood $[E] \in U \subseteq \mathcal{M}_G$ in the Euclidean topology such that for every open neighborhood $[E] \in U' \subseteq U$ with $U' \setminus \mathcal{M}_G^{\text{sing}}$ connected, $\pi_1(U' \setminus \mathcal{M}_G^{\text{sing}})$ is nontrivial.*

Proof. — \mathcal{M}_G is near $[E]$ analytically isomorphic to the quotient of

$$V := H^1(X, \text{ad}(E))$$

modulo the finite group $\text{Aut}(E)/Z_G$ near the image of $0 \in V$. Let an element $\varphi \in \text{Aut}(E) \setminus Z_G$ be given. Using Lemma 1.1, it suffices to verify that the fixed locus $V^\varphi \subseteq V$ has codimension ≥ 2 .

The image $K_{G/Z_G} \subseteq G/Z_G$ of $K_G \subseteq G$ is again a maximal compact subgroup. Since E is stable, the induced principal G/Z_G -bundle E/Z_G is also stable. Hence we may assume $E/Z_G = E_\rho$ for some group homomorphism $\rho : \pi_1(X, x_0) \rightarrow K_{G/Z_G}$. The canonical homomorphism $\text{Aut}(E) \rightarrow \text{Aut}(E/Z_G)$ has kernel Z_G , so we get an embedding

$$\text{Aut}(E)/Z_G \hookrightarrow \text{Aut}(E/Z_G) = C_{G/Z_G}(\rho) \subseteq G/Z_G$$

using also Lemma 2.2. Let $Z_\varphi \subseteq G/Z_G$ denote the nontrivial finite subgroup generated by the image of $\varphi \in \text{Aut}(E)$ under this map.

We choose a maximal torus $T \subseteq G$ such that T/Z_G contains Z_φ , and decompose \mathfrak{g} into Z_φ -eigenspaces \mathfrak{g}_χ with respect to characters $\chi : Z_\varphi \rightarrow \mathbb{C}^*$. The fundamental group $\pi_1(X, x_0)$ also acts on \mathfrak{g} , via the composition

$$\pi_1(X, x_0) \longrightarrow K_{G/Z_G} \subseteq G/Z_G \longrightarrow \text{Aut}(\mathfrak{g})$$

of ρ with the adjoint action of G/Z_G on \mathfrak{g} , and we have

$$\text{ad}(E) = \tilde{X} \times^\rho \mathfrak{g}.$$

Because this action of $\pi_1(X, x_0)$ on \mathfrak{g} commutes with the adjoint action of $Z_\varphi \subseteq G/Z_G$ on \mathfrak{g} , we get a vector bundle decomposition

$$\text{ad}(E) = \bigoplus_{\chi: Z_\varphi \rightarrow \mathbb{C}^*} \text{ad}(E)_\chi \quad \text{with} \quad \text{ad}(E)_\chi = \tilde{X} \times^\rho \mathfrak{g}_\chi.$$

Since Z_φ is nontrivial, there is a root $\alpha : T/Z_G \rightarrow \mathbb{C}^*$ of G which is nontrivial on Z_φ . Then Z_φ acts on the root spaces \mathfrak{g}_α and $\mathfrak{g}_{-\alpha}$ by nontrivial characters. Hence the eigenspace \mathfrak{g}_{χ_0} for the trivial character $\chi_0 : Z_\varphi \rightarrow \mathbb{C}^*$ has codimension ≥ 2 in \mathfrak{g} . Thus the subbundle

$$\text{ad}(E)_{\chi_0} \subseteq \text{ad}(E)$$

is a direct summand of corank ≥ 2 . Because E is stable, $\text{ad}(E)$ is semistable of degree 0. It follows that $\text{ad}(E)_{\chi_0}$ also has degree 0. Thus Riemann-Roch implies that the linear subspace

$$V^\varphi = H^1(X, \text{ad}(E)_{\chi_0}) \subseteq V = H^1(X, \text{ad}(E))$$

has codimension $\geq 2(g_X - 1) \geq 4$. So Lemma 1.1 applies. \square

3. The strictly semistable locus

Let G still be a reductive connected linear algebraic group over \mathbb{C} . Let H be a Levi subgroup of a maximal parabolic subgroup $P \subsetneq G$. Let $K_H \subseteq H$ be a maximal compact subgroup. We choose a Borel subgroup $B \subseteq P$ of G , and a maximal torus $T \subseteq B \cap H$ of G . Let

$$\alpha_1, \dots, \alpha_r : T \longrightarrow \mathbb{C}^* \quad \text{and} \quad \alpha_1^\vee, \dots, \alpha_r^\vee : \mathbb{C}^* \longrightarrow T$$

be the simple roots and coroots of G . The quotient of $\text{Hom}(\mathbb{C}^*, T)$ modulo its subgroup $\mathbb{Z} \cdot \alpha_1^\vee + \dots + \mathbb{Z} \cdot \alpha_r^\vee$ is isomorphic to $\pi_1(G)$.

The Dynkin diagram of H is given by removing one simple root α_i from the Dynkin diagram of G . The centers of G and H are

$$Z_G = \bigcap_j \ker(\alpha_j) \quad \text{and} \quad Z_H = \bigcap_{j \neq i} \ker(\alpha_j);$$

cf. [4, (13.4)]. It follows that α_i induces an isomorphism

$$Z_H/Z_G \xrightarrow{\sim} \mathbb{C}^*.$$

The simple coroots of H are the α_j^\vee with $j \neq i$, so the sequence

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\cdot \alpha_i^\vee} \pi_1(H) \longrightarrow \pi_1(G) \longrightarrow 0$$

is exact. Hence the induced map $\pi_1([H, H]) \rightarrow \pi_1([G, G])$ is injective.

Given a principal H -bundle E over X , we denote by

$$E_G := E \times^H G = (E \times G)/H \longrightarrow X$$

the principal G -bundle obtained by extending the structure group of E using the inclusion map $H \hookrightarrow G$. If E is semistable and its topological type $e \in \pi_1(H)$ is torsion, then E_G is semistable [15, Lemma 3.5.11]. Sending $[E]$ to $[E_G]$ thus defines a morphism of projective schemes

$$i_H : \mathcal{M}'_H \longrightarrow \mathcal{M}'_G.$$

The automorphism group of H acts on \mathcal{M}_H , by extension of the structure group. An inner automorphism of H acts trivially on \mathcal{M}_H , so the group $\text{Out}(H)$ of outer automorphisms also acts on \mathcal{M}_H .

We denote the normalizer of H in G by $N_G(H)$. It acts on H by conjugation in G . Let Γ_H denote the image of $N_G(H)$ in $\text{Out}(H)$. The induced action of Γ_H on $Z_H/Z_G \cong \mathbb{C}^*$ is effective, since the centralizer of Z_H/Z_G in G/Z_G is known to be H/Z_G . This yields an embedding

$$\Gamma_H \hookrightarrow \text{Aut}(\mathbb{C}^*) = \{\pm 1\}.$$

The morphism i_H is Γ_H -invariant, so it descends to a morphism

$$\bar{i}_H : \mathcal{M}'_H/\Gamma_H \longrightarrow \mathcal{M}'_G.$$

Note that Γ_H acts trivially on $\pi_1([H, H])$, because $N_G(H)$ acts trivially on $\pi_1([G, G])$. Hence Γ_H acts on each component $\mathcal{M}_H^e \subseteq \mathcal{M}'_H$.

PROPOSITION 3.1. — *Assume that G is semisimple. Then the normalization of the strictly semistable locus*

$$\mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}} \subset \mathcal{M}_G$$

is given by the disjoint union

$$\bar{i} := \coprod \bar{i}_H : \mathcal{M}'_H/\Gamma_H \longrightarrow \mathcal{M}'_G = \mathcal{M}_G$$

over a set of representatives H for conjugacy classes of Levi subgroups of maximal parabolic subgroups in G .

Proof. — As \mathcal{M}'_H is normal, its quotient \mathcal{M}'_H/Γ_H is also normal, due to [9, p. 5]. The image of \bar{i} is contained in $\mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$ by construction. Conversely, every closed point in $\mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$ has the form

$$[E_G] \in \mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$$

for some semistable principal H -bundle E over X such that E_G is also semistable. Let $e \in \pi_1(H)$ be the topological type of E . Let

$$(3.1) \quad \chi_H : H \longrightarrow \mathbb{C}^*$$

denote the generator of $\text{Hom}(H, \mathbb{C}^*) \cong \mathbb{Z}$ which is dominant for P . If $\langle \chi_H, e \rangle > 0$, then the reduction $E_P := E \times^H P$ of E_G to P would violate semistability by [14, Lemma 2.1]; if $\langle \chi_H, e \rangle < 0$, then the reduction E_Q of E_G to the opposite parabolic subgroup $H \subseteq Q \subseteq G$ would do so. These contradictions show $\langle \chi_H, e \rangle = 0$, which means $e \in \pi_1([H, H])$. This proves that the image of \bar{i} is equal to $\mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$.

The projective morphism i_H is also affine due to Ramanathan's lemma [15, Lemma 4.8.1, Remark 4.8.2 and Lemma 5.1]. Hence the morphisms i_H , \bar{i}_H and \bar{i} are all finite. It remains to prove that \bar{i} is generically injective. For that, we use Lemma 2.1.

Let H' be a Levi subgroup of another maximal parabolic subgroup $P' \subseteq G$. Let $K_{H'} \subseteq H'$ again be a maximal compact subgroup. Suppose that two points

$$[E_\rho] \in \mathcal{M}'_H \quad \text{and} \quad [E_{\rho'}] \in \mathcal{M}'_{H'}$$

have the same image in \mathcal{M}_G for some homomorphisms

$$\rho : \pi_1(X, x_0) \longrightarrow K_H \quad \text{and} \quad \rho' : \pi_1(X, x_0) \longrightarrow K_{H'}.$$

Then there is an element $g \in G$ such that

$$g\rho(\omega)g^{-1} = \rho'(\omega)$$

holds for all $\omega \in \pi_1(X, x_0)$. Now assume moreover that the image of ρ is dense in K_H . Then we can conclude

$$gK_Hg^{-1} \subseteq K_{H'},$$

and hence $gHg^{-1} \subseteq H'$, so

$$gHg^{-1} = H'$$

by maximality. Thus H and H' are conjugate in G , so we may assume $H = H'$, and then we get $g \in N_G(H)$. It follows that the two points $[E_\rho]$ and $[E_{\rho'}]$ are indeed in the same Γ_H -orbit. \square

We return to the general case where G is reductive. Let

$$\Sigma_G \subseteq \mathcal{M}_G^{\text{sing}}$$

denote the set of all singular closed points $[E] \in \mathcal{M}_G$ such that every Euclidean neighborhood $[E] \in U \subseteq \mathcal{M}_G$ contains an open neighborhood $[E] \in U' \subseteq U$ for which $U' \setminus \mathcal{M}_G^{\text{sing}}$ is connected and simply connected. Proposition 2.3 shows $\Sigma_G \subseteq \mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$.

PROPOSITION 3.2. — *Let E be a principal H -bundle over X such that E_G is polystable with $\text{Aut}(E_G) = Z_H$. Then the point $[E_G] \in \mathcal{M}_G$ is in Σ_G .*

Proof. — Luna’s étale slice theorem [8] implies that \mathcal{M}_G is near the point $[E_G]$ analytically isomorphic to the GIT-quotient

$$H^1(X, \text{ad}(E_G)) // \text{Aut}(E_G)$$

near 0. Here $\text{Aut}(E_G) = Z_H$ acts on the vector space

$$V := H^1(X, \text{ad}(E_G)) = H^1(X, E \times^H \mathfrak{g})$$

via the adjoint action of Z_H on \mathfrak{g} . Since $Z_G \subseteq Z_H$ acts trivially, the isomorphism $\alpha_i : Z_H/Z_G \rightarrow \mathbb{C}^*$ yields an action of \mathbb{C}^* on \mathfrak{g} . Let $\mathfrak{g}_m \subseteq \mathfrak{g}$ denote the \mathbb{C}^* -eigenspace of weight $m \in \mathbb{Z}$. Since the adjoint action of H on \mathfrak{g} commutes with \mathbb{C}^* , we get a vector bundle decomposition

$$\text{ad}(E_G) = \bigoplus_{m \in \mathbb{Z}} \text{ad}(E_G)_m \quad \text{with} \quad \text{ad}(E_G)_m := E \times^H \mathfrak{g}_m.$$

As E_G is semistable, $\text{ad}(E_G)$ is semistable of degree 0. Consequently, $\text{ad}(E_G)_m$ also has degree 0. Due to Riemann-Roch, the \mathbb{C}^* -eigenspace

$$V_m = H^1(X, \text{ad}(E_G)_m) \subseteq V$$

has dimension $\geq (g_X - 1) \dim \mathfrak{g}_m$. Using $\mathfrak{g}_{\pm\alpha_i} \subseteq \mathfrak{g}_{\pm 1}$, we conclude

$$\dim V_{\pm 1} \geq g_X - 1 \geq 2.$$

Then Lemma 1.2 completes the proof. \square

COROLLARY 3.3. — *The strictly semistable locus $\mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$ is the Zariski closure of the subset $\Sigma_G \subseteq \mathcal{M}_G$.*

Proof. — The image $K_{H/Z_G} \subseteq H/Z_G$ of $K_H \subseteq H$ is again a maximal compact subgroup. Let E be a principal H -bundle over X such that the principal H/Z_G -bundle E/Z_G comes from a homomorphism

$$\rho : \pi_1(X, x_0) \longrightarrow K_{H/Z_G}.$$

Then the principal G/Z_G -bundle E_G/Z_G comes from the composition

$$\pi_1(X, x_0) \xrightarrow{\rho} K_{H/Z_G} \hookrightarrow K_{G/Z_G}$$

where $K_{G/Z_G} \subseteq G/Z_G$ is an arbitrary maximal compact subgroup containing K_{H/Z_G} . Hence E_G/Z_G is polystable, so E_G is also polystable.

If moreover the image of ρ is dense in K_{H/Z_G} , then Lemma 2.2 yields

$$\text{Aut}(E_G/Z_G) = C_{G/Z_G}(\rho) = Z_H/Z_G$$

since the centralizer of H/Z_G in G/Z_G is known to be Z_H/Z_G . This implies that the two canonical embeddings

$$Z_H \hookrightarrow \text{Aut}(E_G) \quad \text{and} \quad \text{Aut}(E_G)/Z_G \hookrightarrow \text{Aut}(E_G/Z_G)$$

are both isomorphisms. Using Proposition 3.2, it follows that the point $[E_G] \in \mathcal{M}_G$ is in Σ_G . This shows that $\Sigma_G \subseteq \mathcal{M}_G$ contains the inverse image of every point in \mathcal{M}_{G/Z_G} that comes from a homomorphism $\rho : \pi_1(X, x_0) \rightarrow K_{H/Z_G}$ with dense image. Such points are dense in

$$\mathcal{M}_{G/Z_G} \setminus \mathcal{M}_{G/Z_G}^{\text{stab}}$$

by Lemma 2.1 and Proposition 3.1, so Σ_G is dense in $\mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$. \square

COROLLARY 3.4. — *The smooth locus of \mathcal{M}_G consists precisely of the moduli points $[E] \in \mathcal{M}_G$ of regularly stable principal G -bundles E over X .*

Proof. — The singular locus $\mathcal{M}_G^{\text{sing}} \subseteq \mathcal{M}_G$ is closed and contains Σ_G , so it contains the strictly semistable locus $\mathcal{M}_G \setminus \mathcal{M}_G^{\text{stab}}$ due to the previous Corollary 3.3. The rest follows from Proposition 2.3.i. \square

4. Reconstructing the Riemann surface

In this section, we prove the following main result of this paper.

THEOREM 4.1. — *Let X, X' be compact connected Riemann surfaces of genus $g_X, g_{X'} \geq 3$. Let G, G' be nonabelian connected reductive linear algebraic groups over \mathbb{C} . Let $d \in \pi_1(G)$ and $d' \in \pi_1(G')$ be given.*

If the smooth locus of $\mathcal{M}_G^d(X)$ is isomorphic to the smooth locus of $\mathcal{M}_{G'}^{d'}(X')$, then X is isomorphic to X' .

This theorem is proved here by reconstructing X from the smooth locus of the variety $\mathcal{M}_G^d(X)$. Starting from a moduli space $\mathcal{M}_{G'}^{d'}(X')$ with isomorphic smooth locus, this reconstruction will automatically yield an isomorphic Riemann surface, thereby proving $X' \cong X$.

Thus it suffices to consider only one Riemann surface X . We reconstruct X from the smooth locus of $\mathcal{M}_G^d = \mathcal{M}_G^d(X)$, in several steps.

Step 1.

Let $Z_G^0 \subseteq Z_G$ be the identity component. We put $\bar{G} := G/Z_G^0$. The projection $G \rightarrow \bar{G}$ induces a morphism

$$(4.1) \quad \text{pr} : \mathcal{M}_G^d \longrightarrow \mathcal{M}_{\bar{G}}^{\bar{d}}$$

where $\bar{d} \in \pi_1(\bar{G})$ denotes the image of $d \in \pi_1(G)$.

LEMMA 4.2. — *We have $\text{pr}_*(\mathcal{O}_{\mathcal{M}_G^d}) = \mathcal{O}_{\mathcal{M}_{\bar{G}}^{\bar{d}}}$.*

Proof. — The canonical embedding

$$\mathcal{O}_{\mathcal{M}_{\bar{G}}^{\bar{d}}} \hookrightarrow \text{pr}_*(\mathcal{O}_{\mathcal{M}_G^d})$$

turns the latter into a coherent sheaf of algebras over the former. Let

$$U \subseteq \mathcal{M}_{\bar{G}}^{\bar{d}}$$

be the regularly stable locus. This locus U is known to be open and non-empty; one way to see this is to use Corollary 3.4. Because $\mathcal{M}_{\bar{G}}^{\bar{d}}$ is normal, it suffices to prove the claim over U .

Since Z_G^0 is central in G , the multiplication map

$$Z_G^0 \times G \longrightarrow G$$

is a group homomorphism. It induces a morphism of projective varieties

$$\mathcal{M}_{Z_G^0}^0 \times \mathcal{M}_G^d \longrightarrow \mathcal{M}_G^d$$

by extension of the structure group. Thus the abelian variety

$$A := \mathcal{M}_{Z_G^0}^0$$

acts on \mathcal{M}_G^d . The map pr in (4.1) is A -invariant, and its restriction

$$\text{pr} : \text{pr}^{-1}(U) \longrightarrow U$$

is a principal A -bundle. As A is integral, the claim follows over U . \square

Let ω_G denote the dualizing sheaf of \mathcal{M}_G^d . By [13, Proposition 2.2] or [7, Theorem 2.8], ω_G is a line bundle on \mathcal{M}_G^d . Its restriction to the smooth locus $\mathcal{M}_G^d \setminus \mathcal{M}_G^{\text{sing}}$ is the canonical line bundle $\det(\Omega^1)$.

COROLLARY 4.3. — *Choose $n \gg 0$. Then the line bundle $\omega_G^{\otimes(-n)}$ on \mathcal{M}_G^d is globally generated, and the corresponding morphism*

$$\varphi_{\omega_G^{\otimes(-n)}} : \mathcal{M}_G^d \longrightarrow \mathbb{P}^N$$

factors into the morphism pr in (4.1), followed by a closed immersion.

Proof. — We have $\omega_G \cong \text{pr}^*(\omega_{\bar{G}})$ according to [13, p. 525–527]. Hence the projection formula gives an isomorphism

$$\text{pr}_*(\omega_G^{\otimes(-n)}) \cong \omega_{\bar{G}}^{\otimes(-n)} \otimes \text{pr}_*(\mathcal{O}_{\mathcal{M}_G^d}) = \omega_{\bar{G}}^{\otimes(-n)}$$

for any n . Taking global sections on both sides, we get

$$H^0(\mathcal{M}_G^d, \omega_G^{\otimes(-n)}) = H^0(\mathcal{M}_{\bar{G}}^{\bar{d}}, \omega_{\bar{G}}^{\otimes(-n)}).$$

Now use that $\omega_{\bar{G}}^{\otimes(-1)}$ is ample, according to [13, Corollary 2.1]. \square

This shows how to reconstruct the morphism pr in (4.1), and in particular its target $\mathcal{M}_{\bar{G}}^{\bar{d}}$, from the variety \mathcal{M}_G^d .

One can in fact reconstruct $\mathcal{M}_{\bar{G}}^{\bar{d}}$ from just the smooth locus of \mathcal{M}_G^d , by taking the closure of its image under the map

$$\varphi_{\omega_G^{\otimes(-n)}} : \mathcal{M}_G^d \setminus \mathcal{M}_G^{\text{sing}} \longrightarrow \mathbb{P}^N$$

given by the canonical line bundle ω_G on $\mathcal{M}_G^d \setminus \mathcal{M}_G^{\text{sing}}$.

Remark 4.4. — Every holomorphic section of ω_G over $\mathcal{M}_G^d \setminus \mathcal{M}_G^{\text{sing}}$ extends to \mathcal{M}_G^d by normality, and hence is algebraic. Thus it suffices to assume in Theorem 4.1 that the two smooth loci are biholomorphic.

For the remaining steps, we can thus assume that G is semisimple, $G \neq \{1\}$, and that we are given the projective variety \mathcal{M}_G^d .

Step 2.

Corollary 3.3 characterizes the locus $\mathcal{M}_G^d \setminus \mathcal{M}_G^{\text{stab}}$ in \mathcal{M}_G^d . We assume that it is non-empty, and consider an irreducible component of it. Its normalization is, due to Proposition 3.1, isomorphic to

$$\mathcal{M}_H^e/\Gamma_H$$

for some Levi subgroup H of a maximal parabolic subgroup $P \subsetneq G$, and some inverse image $e \in \pi_1([H, H])$ of d .

The group Γ_H acts on the line bundle ω_H over \mathcal{M}_H^e . Its square $\omega_H^{\otimes 2}$ descends to a line bundle $\omega_H^{\otimes 2}/\Gamma_H$ over \mathcal{M}_H^e/Γ_H according to Kempf's lemma [2, Théorème 2.3]. This line bundle is determined by the normal variety \mathcal{M}_H^e/Γ_H alone, due to the following lemma.

LEMMA 4.5. — *The line bundle $\omega_H^{\otimes 2}/\Gamma_H$ is over the smooth locus of \mathcal{M}_H^e/Γ_H isomorphic to the square of the canonical line bundle $\det(\Omega^1)$.*

Proof. — This is clear if Γ_H is trivial. So we may assume $\Gamma_H \cong \{\pm 1\}$. The character $\chi_H : H \rightarrow \mathbb{C}^*$ in (3.1) induces a Γ_H -equivariant morphism

$$(\chi_H)_* : \mathcal{M}_H^e \longrightarrow J = J(X) := \text{Pic}^0(X).$$

Since Γ_H acts effectively on Z_H/Z_G , it also acts effectively on the isogenous torus $H/[H, H]$. Hence $-1 \in \Gamma_H$ acts on J by the inverse map from the group structure on J . In particular, the fixed locus

$$(\mathcal{M}_H^e)^{\Gamma_H} \subseteq \mathcal{M}_H^e$$

is contained in the inverse image of the 2-division points in J , so its codimension is at least $g_X \geq 3$. Thus it suffices to check the lemma over U/Γ_H for the smooth open subscheme

$$U := \mathcal{M}_H^e \setminus ((\mathcal{M}_H^e)^{\text{sing}} \cup (\mathcal{M}_H^e)^{\Gamma_H})$$

where Γ_H acts freely. But here the claim follows simply from the fact that the canonical projection $U \rightarrow U/\Gamma_H$ is étale. □

Step 3.

Replacing G by H in (4.1), we get a canonical Γ_H -equivariant morphism $\text{pr} : \mathcal{M}_H^e \rightarrow \mathcal{M}_{\overline{H}}^{\overline{e}}$ with $\overline{H} := H/Z_H^0$. It induces a morphism

$$(4.2) \quad \overline{\text{pr}} : \mathcal{M}_H^e/\Gamma_H \longrightarrow \mathcal{M}_{\overline{H}}^{\overline{e}}/\Gamma_H.$$

LEMMA 4.6. — *There is a dense open subscheme $\overline{U} \subseteq \mathcal{M}_{\overline{H}}^{\overline{e}}/\Gamma_H$ such that*

$$\overline{\text{pr}} : (\overline{\text{pr}})^{-1}(\overline{U}) \longrightarrow \overline{U}$$

is an étale-locally trivial fibration, with fiber either the Jacobian J or its quotient $J/\{\pm 1\}$ modulo the inverse map from its group structure.

Proof. — Let $U \subseteq \mathcal{M}_H^{\overline{e}}$ be the regularly stable locus. The restriction

$$(4.3) \quad \text{pr} : \text{pr}^{-1}(U) \longrightarrow U$$

is a principal J -bundle, as in the proof of Lemma 4.2.

Suppose for the moment that the action of Γ_H on $\mathcal{M}_H^{\overline{e}}$ is effective. Then Γ_H acts freely on some dense open subscheme $U' \subseteq U$. The principal J -bundle pr in (4.3) descends to a principal J -bundle

$$\text{pr}^{-1}(U')/\Gamma_H \longrightarrow U'/\Gamma_H.$$

Hence the lemma holds for $\overline{U} := U'/\Gamma_H \subseteq \mathcal{M}_{\overline{H}}^{\overline{e}}/\Gamma_H$ in this case.

It remains to treat the case where the action of Γ_H on $\mathcal{M}_H^{\overline{e}}$ is not effective. This means $\Gamma_H \cong \{\pm 1\}$, and that Γ_H acts trivially on $\mathcal{M}_H^{\overline{e}}$.

We claim that in this case $\overline{U} := U \subseteq \mathcal{M}_H^{\overline{e}} = \mathcal{M}_{\overline{H}}^{\overline{e}}/\Gamma_H$ satisfies the conditions in the lemma. Replacing U by an étale covering and pulling back, we can choose a section σ of the principal J -bundle pr in (4.3). Let $\gamma \in \Gamma_H$ be the nontrivial element. Then

$$\gamma \cdot \sigma = \xi + \sigma$$

for some morphism $\xi : U \rightarrow J$. Refining the covering of U if necessary, and using the divisibility of J , we may assume $\xi = 2\xi'$. Then

$$\gamma \cdot (\xi' + \sigma) = \gamma \cdot \xi' + \gamma \cdot \sigma = -\xi' + (\xi + \sigma) = \xi' + \sigma,$$

since Γ_H acts by the inverse map on J . So $\xi' + \sigma$ is Γ_H -invariant.

This shows that the principal J -bundle pr in (4.3) admits étale-locally Γ_H -equivariant trivializations. Hence it descends to an étale-locally trivial fibration with fiber $J/\Gamma_H = J/\{\pm 1\}$ over U . □

Remark 4.7. — The second case, where the general fiber of $\overline{\text{pr}}$ is $J/\{\pm 1\}$, actually occurs. An example is $G = \text{Sp}_{2n+2}$ and $H \cong \mathbb{C}^* \times \text{Sp}_{2n}$. Here $\Gamma_H \cong \{\pm 1\}$ is nontrivial, but its action on $\overline{H} = \text{Sp}_{2n}$ is trivial.

COROLLARY 4.8. — *Choose $n \gg 0$. Then the line bundle $(\omega_H^{\otimes 2}/\Gamma_H)^{\otimes (-n)}$ on \mathcal{M}_H^e/Γ_H is globally generated, and the corresponding morphism*

$$\varphi_{(\omega_H^{\otimes 2}/\Gamma_H)^{\otimes (-n)}} : \mathcal{M}_H^e/\Gamma_H \longrightarrow \mathbb{P}^N$$

factors into the morphism $\overline{\text{pr}}$ in (4.2), followed by a closed immersion.

Proof. — Using Kempf’s lemma, the anti-ample line bundle $\omega_{\bar{H}}^{\otimes 2}$ on $\mathcal{M}_{\bar{H}}^{\bar{e}}$ descends to an anti-ample line bundle $\omega_{\bar{H}}^{\otimes 2}/\Gamma_H$ on $\mathcal{M}_{\bar{H}}^{\bar{e}}/\Gamma_H$ with

$$\bar{\mathrm{pr}}^*(\omega_{\bar{H}}^{\otimes 2}/\Gamma_H) \cong \omega_H^{\otimes 2}/\Gamma_H.$$

The corollary can now be proved exactly as Corollary 4.3, since Lemma 4.6 and the normality of $\mathcal{M}_{\bar{H}}^{\bar{e}}/\Gamma_H$ together imply

$$\bar{\mathrm{pr}}_*(\mathcal{O}_{\mathcal{M}_{\bar{H}}^{\bar{e}}/\Gamma_H}) = \mathcal{O}_{\mathcal{M}_{\bar{H}}^{\bar{e}}/\Gamma_H}. \quad \square$$

This shows how to reconstruct the morphism $\bar{\mathrm{pr}}$ in (4.2), and in particular its general fiber $J(X)$ or $J(X)/\{\pm 1\}$, from the variety $\mathcal{M}_{\bar{H}}^{\bar{e}}/\Gamma_H$.

It is known that $J(X)$ can be reconstructed from the Kummer variety $J(X)/\{\pm 1\}$, for example as its integral closure in the field of meromorphic functions on the two-sheeted cover of its smooth locus given by the unique maximal torsionfree subgroup in its fundamental group.

Step 4.

Starting from the variety \mathcal{M}_G^d , we have now reconstructed the Jacobian $J = J(X)$, together with a morphism $j_E : J \rightarrow \mathcal{M}_G^d$ given by a general point $[E] \in \mathcal{M}_H^e$. This morphism is the composition

$$j_E : J \cong \mathcal{M}_{Z_H^0} \xrightarrow{\cdot E} \mathcal{M}_H^e \xrightarrow{i_H} \mathcal{M}_G^d.$$

LEMMA 4.9. — *The first Chern class $c_1(j_E^* \omega_G) \in H^2(J, \mathbb{Z})$ is a negative multiple of the canonical principal polarization Θ on J .*

Proof. — According to [13, Proposition 2.2], ω_G is the pullback along

$$\mathrm{ad} : \mathcal{M}_G^d \longrightarrow \mathcal{M}_{\mathrm{SL}(\mathfrak{g})}$$

of the determinant of cohomology line bundle $\mathcal{L}_{\mathrm{det}}$ on $\mathcal{M}_{\mathrm{SL}(\mathfrak{g})}$. Points in $\mathcal{M}_{\mathrm{SL}(\mathfrak{g})}$ correspond to polystable vector bundles V over X with trivial determinant; we recall that the fiber of $\mathcal{L}_{\mathrm{det}}$ over such a point is the determinant of the cohomology of V , i. e. the vector space

$$\Lambda^{\mathrm{top}} H^0(X, V) \otimes \Lambda^{\mathrm{top}} H^1(X, V)^*$$

of dimension one. Let $\mathfrak{g}_m \subseteq \mathfrak{g}$ denote the eigenspace where $\mathbb{C}^* \cong Z_H^0$ acts with weight $m \in \mathbb{Z}$. The line bundle $j_E^* \omega_G$ is the pullback of $\mathcal{L}_{\mathrm{det}}$ along the morphism

$$J \longrightarrow \mathcal{M}_{\mathrm{SL}(\mathfrak{g})}$$

that sends a line bundle L of degree 0 over X to the vector bundle

$$\bigoplus_{m \in \mathbb{Z}} L^{\otimes m} \otimes (E \times^H \mathfrak{g}_m)$$

with trivial determinant over X . Computing the determinant of its cohomology, we conclude that

$$c_1(\mathcal{L}) = \sum_{m \in \mathbb{Z}} -m^2 \dim \mathfrak{g}_m \cdot \Theta \in H^2(J, \mathbb{Z}).$$

This is a strictly negative multiple of Θ , since $\mathfrak{g}_m \neq 0$ for some $m \neq 0$, because the action of $\mathbb{C}^* \cong Z_H^0$ on \mathfrak{g} is not trivial. \square

Hence we can also reconstruct the principal polarization Θ on $J(X)$; then the usual Torelli theorem gives back the Riemann surface X .

Step 5.

It remains to treat the case $\mathcal{M}_G^d \setminus \mathcal{M}_G^{\text{stab}} = \emptyset$. According to [14, Proposition 7.8], this can only happen if

$$G/Z_G \cong \text{PGL}_{n_1} \times \cdots \times \text{PGL}_{n_r}.$$

In particular, the universal covering of G is a homomorphism

$$(4.4) \quad \text{SL}_{n_1} \times \cdots \times \text{SL}_{n_r} \longrightarrow G.$$

Its kernel is a finite subgroup μ in the product of the centers $\mu_{n_i} \subseteq \text{SL}_{n_i}$.

For each i , we lift the image of $d \in \pi_1(G)$ in $\pi_1(\text{PGL}_{n_i}) \cong \mathbb{Z}/n_i\mathbb{Z}$ to an integer $d_i \in \mathbb{Z}$. Then the emptiness of the strictly semistable locus also implies that d_i is coprime to n_i for all i . Let L_i be a line bundle of degree d_i on X , and let \mathcal{M}_{n_i, L_i} denote the coarse moduli space of semistable vector bundles E over X with rank n_i and determinant L_i . The homomorphism (4.4) induces a finite surjective morphism

$$(4.5) \quad \text{pr} : \mathcal{M}_{n_1, L_1} \times \cdots \times \mathcal{M}_{n_r, L_r} \longrightarrow \mathcal{M}_G^d.$$

The smooth locus in \mathcal{M}_G^d coincides with the regularly stable locus by Corollary 3.3; we denote it by $U \subseteq \mathcal{M}_G^d$. The abelian group of principal μ -bundles over X is isomorphic to μ^{2g_X} . This group acts on the morphism (4.5). This turns the restriction

$$\text{pr} : \text{pr}^{-1}(U) \longrightarrow U$$

into an unramified Galois covering with group μ^{2g_X} . Since \mathcal{M}_{n_i, L_i} is a unirational smooth projective variety, it is in particular simply connected [17]. Thus $\text{pr}^{-1}(U)$ is also simply connected, as we are removing a closed subset

of complex codimension ≥ 2 . This shows that $\text{pr}^{-1}(U)$ is the universal covering of U . Taking the integral closure of \mathcal{M}_G^d in the field of meromorphic functions on $\text{pr}^{-1}(U)$, we can thus recover the morphism (4.5) from the variety \mathcal{M}_G^d in this case.

Because the Picard group of \mathcal{M}_{n_i, L_i} is isomorphic to \mathbb{Z} , the extremal rays in the nef cone of the variety

$$\mathcal{M}_{n_1, L_1} \times \cdots \times \mathcal{M}_{n_r, L_r}$$

correspond to the factors \mathcal{M}_{n_i, L_i} . Thus we can even reconstruct these factors \mathcal{M}_{n_i, L_i} from the variety \mathcal{M}_G^d .

The projective variety \mathcal{M}_{n_i, L_i} is known to determine the Riemann surface X up to isomorphism; cf. [20, Theorem 1] or [12, Theorem 3]. Thus we can reconstruct X from $\mathcal{M}_G^d(X)$ in this case as well.

This completes the proof of Theorem 4.1. \square

BIBLIOGRAPHY

- [1] S. ADAMS, “Reduction of cocycles with hyperbolic targets”, *Ergodic Theory Dyn. Syst.* **16** (1996), p. 1111-1145.
- [2] J.-M. DREZET & M. NARASIMHAN, “Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques”, *Invent. Math.* **97** (1989), p. 53-94.
- [3] T. GÓMEZ & I. SOLS, “Moduli space of principal sheaves over projective varieties”, *Ann. of Math.* **161** (2005), p. 1037-1092.
- [4] J. HUMPHREYS, *Linear algebraic groups*, Graduate Texts in Mathematics, vol. 21, Springer-Verlag, New York - Heidelberg - Berlin, 1975.
- [5] J.-M. HWANG & S. RAMANAN, “Hecke curves and Hitchin discriminant”, *Ann. Sci. École Norm. Sup.* **37** (2004), p. 801-817.
- [6] A. KOUVIDAKIS & T. PANTEV, “The automorphism group of the moduli space of semi stable vector bundles”, *Math. Ann.* **302** (1995), p. 225-268.
- [7] S. KUMAR & M. NARASIMHAN, “Picard group of the moduli spaces of G -bundles”, *Math. Ann.* **308** (1997), p. 155-173.
- [8] D. LUNA, “Slices étalés”, *Mém. Soc. Math. Fr.* **33** (1973), p. 81-105.
- [9] D. MUMFORD, *Geometric invariant theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 34, Springer-Verlag, Berlin, 1965.
- [10] D. MUMFORD & P. NEWSTEAD, “Periods of a moduli space of bundles on curves”, *Amer. Jour. Math.* **90** (1968), p. 1200-1208.
- [11] M. NARASIMHAN & S. RAMANAN, “Moduli of vector bundles on a compact Riemann surface”, *Ann. of Math.* **89** (1969), p. 14-51.
- [12] ———, “Deformations of the moduli space of vector bundles over an algebraic curve”, *Ann. of Math.* **101** (1975), p. 391-417.
- [13] T. PANTEV, “Comparison of generalized theta functions”, *Duke Math. J.* **76** (1994), p. 509-539.
- [14] A. RAMANATHAN, “Stable principal bundles on a compact Riemann surface”, *Math. Ann.* **213** (1975), p. 129-152.
- [15] ———, “Moduli for principal bundles over algebraic curves”, *Proc. Indian Acad. Sci. (Math. Sci.)* **106** (1996), p. 301-328 and 421-449.

- [16] A. SCHMITT, “Singular principal bundles over higher-dimensional manifolds and their moduli spaces”, *Int. Math. Res. Not.* (2002), p. 1183-1209.
- [17] J.-P. SERRE, “On the fundamental group of a unirational variety”, *Jour. London Math. Soc.* **34** (1959), p. 481-484.
- [18] X. SUN, “Minimal rational curves on moduli spaces of stable bundles”, *Math. Ann.* **331** (2005), p. 925-937.
- [19] A. TYURIN, “An analogue of the Torelli theorem for two-dimensional bundles over an algebraic curve of arbitrary genus (Russian)”, *Izv. Akad. Nauk SSSR Ser. Mat.* **33** (1969), p. 1149-1170, in English: *Math. USSR, Izv.* **3** (1971).
- [20] ———, “Analogues of Torelli’s theorem for multidimensional vector bundles over an arbitrary algebraic curve (Russian)”, *Izv. Akad. Nauk SSSR Ser. Mat.* **34** (1970), p. 338-365, in English: *Math. USSR, Izv.* **4** (1971).

Manuscrit reçu le 28 juillet 2009,
révisé le 3 août 2010,
accepté le 8 février 2011.

Indranil BISWAS
School of Mathematics,
Tata Institute of Fundamental Research,
Homi Bhabha Road, Bombay 400005,
India

indranil@math.tifr.res.in

Norbert HOFFMANN
Freie Universität Berlin,
Institut für Mathematik,
Arnimallee 3, 14195 Berlin,
Germany

norbert.hoffmann@fu-berlin.de