

ANNALES DE L'INSTITUT FOURIER

GUNNAR FORST

On a theorem of M. Itô

Annales de l'institut Fourier, tome 28, n° 2 (1978), p. 157-159

http://www.numdam.org/item?id=AIF_1978__28_2_157_0

© Annales de l'institut Fourier, 1978, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON A THEOREM OF M. ITO (*)

by Gunnar FORST

Let G be a locally compact abelian group and let N_0 be a convolution kernel on G (i.e. a positive measure on G) satisfying the principle of unicity of mass. A convolution kernel N on G is said to be a *divisor* of N_0 if there exists a convolution kernel N' on G such that $N_0 = N * N'$. If such an N' exists and is uniquely determined, it is denoted N_0/N ; this is the case if N' exists and is a Hunt kernel on G . The set of divisors of N_0 is denoted $D(N_0)$.

The following result ⁽¹⁾ was recently proved by M. Itô, cf. Théorème 1 in [1].

THEOREM. — *The set*

$$C(N_0) = \{N \in D(N_0) \mid N_0/N \text{ is a Hunt kernel on } G\}$$

is a convex cone.

The proof in [1] of this result is somewhat complicated, and it is the purpose of this note to give a simpler proof, mainly using the ideas from [1].

LEMMA 1 ([1], p. 292-293). — *Let N be a non-zero convolution kernel. In order that there exist a Hunt kernel N' and an N' -invariant convolution kernel η such that $N_0 = N * N' + \eta$, it is necessary and sufficient that $N + pN_0 \in D(N_0)$ for all $p > 0$. In the affirmative case N' and η are uniquely deter-*

(*) This work was done while visiting McGill University, Montréal, in October 1976, supported by NRC-grant A-3108.

(1) The statement of Théorème 1 in [1] is not true; the argument uses Corollaire 2 of [1] which is false. What the proof actually shows is what is formulated in the Theorem.

mined as the vague limits

$$N' = \lim_{p>0} (N_0 / (N + pN_0)) \quad \text{and} \quad \eta = \lim_{t>\infty} N_0 * \alpha_t',$$

where $(\alpha_i)_{i \geq 0}$ is the vaguely continuous convolution semigroup (vccs) on G associated with N' .

LEMMA 2 (cf. [1], p. 295-299). — Let N_i for $i = 1, 2$ be a Hunt kernel with associated vccs $(\alpha_i^i)_{i \geq 0}$. Suppose that the convolution $N_1 * N_2$ exists. Then $(\alpha_1^1 * \alpha_2^2)_{i \geq 0}$ defines a vccs and

$$N = \int_0^\infty \alpha_t^1 * \alpha_t^2 dt,$$

defines a Hunt kernel satisfying

$$N_1 * N_2 = N * (N_1 + N_2). \quad (*)$$

Proof. — For $s, t \geq 0$ we have $\alpha_s^1 * N_1 \leq N_1$ and $\alpha_t^2 * N_2 \leq N_2$ which implies that the convolution $\alpha_s^1 * \alpha_t^2$ exists. By the convergence Lemma in [1] p. 295, it is rather easy to see that the mapping $(s, t) \mapsto \alpha_s^1 * \alpha_t^2$ is vaguely continuous on $[0, \infty[\times [0, \infty[$, and in particular $(\alpha_t^1 * \alpha_t^2)_{t \geq 0}$ is a vccs. For $n > 0$ we have (vaguely)

$$\begin{aligned} \left(\int_0^n \alpha_t^1 * \alpha_t^2 dt \right) * (N_1 + N_2) &= \left(\int_0^n \alpha_t^1 * \alpha_t^2 dt \right) * \left(\int_0^\infty (\alpha_s^1 + \alpha_s^2) ds \right) \\ &= \int_0^\infty \left(\int_0^n \alpha_s^1 * \alpha_t^1 * \alpha_t^2 dt \right) ds + \int_0^\infty \left(\int_0^n \alpha_s^2 * \alpha_t^1 * \alpha_t^2 dt \right) ds \\ &\leq \int_0^\infty \left(\int_t^\infty \alpha_u^1 * \alpha_t^2 du \right) dt + \int_0^\infty \left(\int_t^\infty \alpha_t^1 * \alpha_u^2 du \right) dt \\ &= \int_0^\infty \int_0^\infty \alpha_s^1 * \alpha_t^2 ds dt = N_1 * N_2. \end{aligned}$$

This shows that $\int_0^\infty \alpha_t^1 * \alpha_t^2 dt$ converges vaguely, and equation (*) follows by a monotone convergence argument.]

Proof of the theorem. — We shall see that $C(N_0)$ is convex. Let $N_1, N_2 \in C(N_0)$ and let $(\alpha_i^i)_{i \geq 0}$, $i = 1, 2$, be the vccs associated with the Hunt kernel $N_i' = N_0 / N_i$, $i = 1, 2$. Let $(N_{i,p}')_{p>0}$, $i = 1, 2$, be the resolvent for N_i' ,

$$N_{i,p}' = \int_0^\infty e^{-pt} \alpha_t^i dt \quad \text{for } p > 0 \text{ and } i = 1, 2.$$

For $p > 0$ and $i = 1, 2$ we have $N_{i,p}' * (N_i + pN_0) = N_0$ which shows that for $p > 0$

$$N_{1,p}' * N_{2,p}' * N_0 \leq N_{1,p}' * \frac{1}{p} N_0 \leq \left(\frac{1}{p} \right)^2 N_0.$$

It follows that the convolution $N'_{1,p} * N'_{2,p}$ exists, and by Lemma 2 the convolution kernel

$$\tilde{N}_p = \int_0^\infty e^{-2pt} \alpha_t^1 * \alpha_t^2 dt$$

is well-defined and satisfies

$$N'_{1,p} * N'_{2,p} = \tilde{N}_p * (N'_{1,p} + N'_{2,p}).$$

Moreover we have

$$(N_1 + N_2 + 2pN_0) * \tilde{N}_p = N_0. \tag{**}$$

In fact,

$$\begin{aligned} &(N_1 + N_2 + 2pN_0) * \tilde{N}_p * N'_{1,p} * N'_{2,p} \\ &= (N_1 + pN_0) * N'_{1,p} * N'_{2,p} * \tilde{N}_p + (N_2 + pN_0) * N'_{2,p} * N'_{1,p} * \tilde{N}_p \\ &= N_0 * N'_{2,p} * \tilde{N}_p + N_0 * N'_{1,p} * \tilde{N}_p = N_0 * N'_{1,p} * N'_{2,p}, \end{aligned}$$

where all the convolutions exist, and this implies (**) because $N'_{1,p} * N'_{2,p}$ satisfies the principle of unicity of mass.

By Lemma 1 the Hunt kernel

$$\tilde{N} = \lim_{p \rightarrow 0} \tilde{N}_p = \int_0^\infty \alpha_t^1 * \alpha_t^2 dt$$

satisfies $\tilde{N} * (N_1 + N_2) + \eta = N_0$ for some \tilde{N} -invariant convolution kernel η , and since $N_0 = N_1 * N'_1 = N_2 * N'_2$ we have (as in [1], p. 302)

$$\eta = \lim_{t \rightarrow \infty} N_0 * \alpha_t^1 * \alpha_t^2 \leq \lim_{t \rightarrow \infty} N_0 * \alpha_t^2 = 0,$$

which shows that $\tilde{N} * (N_1 + N_2) = N_0$, i.e.

$$N_1 + N_2 \in C(N_0). | .$$

BIBLIOGRAPHY

[1] M. Itô, Sur le cône convexe maximum formé par des diviseurs d'un noyau de convolution et son application. *Ann. Inst. Fourier*, 25, 3-4 (1975), 289-308.

Manuscrit reçu le 17 février 1977
 Proposé par G. Choquet.

Gunnar FORST,
 Matematisk Institut
 Universitetsparken 5
 DK-2100 København Ø
 Denmark.