

JOSÉ MANUEL CARBALLÉS

**Characteristic homomorphism for  $(F_1, F_2)$ -foliated bundles over subfoliated manifolds**

*Annales de l'institut Fourier*, tome 34, n° 3 (1984), p. 219-245

[http://www.numdam.org/item?id=AIF\\_1984\\_\\_34\\_3\\_219\\_0](http://www.numdam.org/item?id=AIF_1984__34_3_219_0)

© Annales de l'institut Fourier, 1984, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

# CHARACTERISTIC HOMOMORPHISM FOR $(F_1, F_2)$ -FOLIATED BUNDLES OVER SUBFOLIATED MANIFOLDS

by José Manuel CARBALLÉS

## 1. Introduction.

Let  $(F_1, F_2)$  be a couple of foliations on a differentiable manifold  $M$  such that the leaves of  $F_1$  contain those of  $F_2$ ; we shall say such couple  $(F_1, F_2)$  a subfoliation on  $M$ . While Moussu [9], Feigin [5], Cordero-Gadea [3] and Cordero-Masa [4] have study the (exotic) characteristic homomorphism of a subfoliation  $(F_1, F_2)$  using the techniques of Bernstein-Rozenfeld, Bott-Haefliger and Lehmann, our aim in this paper is to present the construction of the characteristic homomorphism of  $(F_1, F_2)$  using the techniques and language of Kamber-Tondeur for foliated bundles.

Our study is based on the notion of  $(F_1, F_2)$ -foliated principal bundle. This is a principal bundle of the form  $P = P_1 + P_2 \rightarrow M$  of structure group  $G_1 \times G_2$  endowed with a foliated structure given by a connection of the form  $\omega = \omega_1 + \omega_2$  (called adapted connection sum) and where, for each  $i = 1, 2$ ,  $P_i \rightarrow M$  is an  $F_i$ -foliated principal bundle of structure group  $G_i$ , and  $\omega_i$  is an adapted connection in  $P_i$ . The most meaningful example of  $(F_1, F_2)$ -foliated bundle over  $M$  is a reduction of the bundle of linear frames of the so called normal bundle of  $(F_1, F_2)$  defined by  $\nu(F_1, F_2) = (F_1/F_2) \oplus \nu F_1$ . This vector bundle  $\nu(F_1, F_2)$  has been used in [4] in order to define the characteristic homomorphism of  $(F_1, F_2)$  adapting the Bott [2] well-known construction of the characteristic

homomorphism of a foliation; our construction of the characteristic homomorphism of an  $(F_1, F_2)$ -foliated principal bundle generalizes that of Cordero-Masa in the same way as Kamber-Tondeur theory of characteristic classes of foliated bundles generalizes Bott theory. This approach allows, moreover, to initiate the study of the holonomy homomorphism of a "leaf" of a subfoliation, in the line of Goldman's paper [6] for the leaf of a foliation.

The paper is structured as follows. In § 2, we introduce the basic definitions and deduce the filtration preserving properties of the Weil homomorphism  $k(\omega)$  of an adapted connection sum in an  $(F_1, F_2)$ -foliated bundle. As a particular consequence, the vanishing theorem for the normal bundle of a subfoliation [4], [5] is reobtained. These properties of  $k(\omega)$  are used in order to prove the vanishing of  $k(\omega)$  on a differential ideal  $I$  of the product Weil algebra  $W(\mathfrak{g}_1 \oplus \mathfrak{g}_2)$  (firstly considered by Feigin [5]) and thus, following Kamber-Tondeur's theory, we introduce the generalized characteristic homomorphism of an  $(F_1, F_2)$ -foliated principal bundle  $P$ :

$$\Delta_* = \Delta_{(F_1, F_2)}(P) : H(W(\mathfrak{g}, H)_I) \longrightarrow H_{DR}(M)$$

where  $H \subset G$  is a closed Lie subgroup such that  $P$  admits an  $H$ -reduction. We show that  $\Delta_*$  does not depend on the connection sum  $\omega$  and that it satisfies the usual functorial properties (i.e. naturality under pull-backs and  $\rho$ -extensions). We also deal with the case where  $\omega_1$  and  $\omega_2$  both are basic connections.

In § 3, we relate the generalized characteristic homomorphism  $\Delta_*(P)$  with the generalized characteristic homomorphism (as defined in [7]) of each  $P_i$ ,  $i = 1, 2$ . Taking into account that any adapted connection sum in  $P$  is  $F_2$ -adapted, we deduce some properties of the characteristic homomorphism as  $F_2$ -foliated bundle of an  $(F_1, F_2)$ -foliated bundle as well as of any  $F_2$ -extension of it. This section ends with the construction of the generalized characteristic homomorphism  $\Delta_*(P)$  when considering a foliation  $F$  as a subfoliation in the three possible forms.

In § 4 we apply the general results of Kamber-Tondeur on

the cohomology of  $g$ -DG-algebras in order to calculate the cohomology  $H(W(g, H)_1)$ . In particular, this allows to refine the characteristic homomorphism of  $(F_1, F_2)$  as defined in [4]. The algebra of secondary characteristic invariants is constructed and a geometric interpretation of the generalized characteristic homomorphism is also given for the general situation.

Finally, in § 5, we restrict the  $(F_1, F_2)$ -foliated bundle  $P$  to the leaves of each foliation  $F_i, i = 1, 2$ ; this leads us, on the one hand to a slightly generalization of Goldman's study, and, on the other, to define the holonomy homomorphism of a "leaf" of a subfoliation and to discuss an example of Reinhart [10].

Through all this paper, the manifolds, maps, etc, will be assumed differentiable of class  $C^\infty$ . Also, we shall adopt the notation of [7].

This paper is a part of the doctoral dissertation of the author who would like to acknowledge here his gratitude to L.A. Cordero for his guidance and encouragement.

## 2. Characteristic homomorphism of an $(F_1, F_2)$ -foliated bundle.

Let  $M$  be an  $n$ -dimensional differentiable manifold,  $TM$  its tangent bundle. Through all this paper, we always assume  $M$  endowed with a  $(q_1, q_2)$ -codimensional subfoliation  $(F_1, F_2)$ , that is, of a couple of integrable subbundles  $F_i$  of  $TM$  of dimension  $n - q_i, i = 1, 2$ , and  $F_2$  being a subbundle of  $F_1$ . Therefore, for each  $i, F_i$  defines a  $q_i$ -codimensional foliation on  $M, d = q_2 - q_1 \geq 0$  and the leaves of  $F_1$  contain those of  $F_2$ .

Let  $Q_i = TM/F_i$  be the normal bundle of  $F_i, i = 1, 2$ , and  $Q_0$  the quotient bundle  $F_1/F_2$ ; then, there is a short exact sequence of vector bundles, canonically associated to

$(F_1, F_2), 0 \longrightarrow Q_0 \xrightarrow{i} Q_2 \xrightarrow{\pi} Q_1 \longrightarrow 0$  and the vector bundle  $\nu(F_1, F_2) = Q_0 \oplus Q_1$  is called the normal bundle of  $(F_1, F_2)$ .

Let  $P_i(M, G_i)$  be an  $F_i$ -foliated principal bundle,  $i = 1, 2$ , and let  $\omega_i$  be an adapted connection. Let

$$P(M, G_1 \times G_2) = P_1(M, G_1) + P_2(M, G_2)$$

be the principal bundle sum of  $P_1$  and  $P_2$ ; then  $\omega = \omega_1 + \omega_2$  defines two partial connections in  $P$  and  $\omega$  is adapted to both; endowed with these two partial connections,  $P$  will be said  $(F_1, F_2)$ -foliated and  $\omega = \omega_1 + \omega_2$  an adapted connection sum. Let us remark that, in particular,  $P$  is  $F_2$ -foliated and if both  $\omega_1$  and  $\omega_2$  are basic, then  $\omega = \omega_1 + \omega_2$  is also basic with respect to  $F_2$ .

Let  $L(Q_i)$  be the frame bundle of  $Q_i$ ,  $i = 0, 1$ , and  $L(Q_1) + L(Q_0)$  the bundle sum. As it can be easily shown using the results in [4],  $L(Q_1) + L(Q_0)$  is  $(F_1, F_2)$ -foliated and it will be called the bundle of transverse frames of  $(F_1, F_2)$ . Other examples can be obtained as follows; let  $P_i \rightarrow M$  be a  $G_i$ -principal bundle,  $i = 1, 2$ , endowed with an  $F_i$ -foliated structure,  $F_i$  being the orbit foliation defined on  $M$  by a left almost free action of a Lie subgroup  $K_i \subset G_i$  (see 2.4 in [7]); then, if  $K_2 \subset K_1$ ,  $P = P_1 + P_2$  is an  $(F_1, F_2)$ -foliated bundle. In particular, if  $P \rightarrow M$  is a  $G$ -principal bundle which is  $F_1$ -foliated by the orbits of the action of a Lie subgroup  $K_1 \subset G$  on  $M$ , as above, then for each Lie subgroup  $K_2 \subset K_1$ , the bundle  $P + P$  is  $(F_1, F_2)$ -foliated.

Let  $P = P_1 + P_2$  be an  $(F_1, F_2)$ -foliated bundle over  $M$ ,  $\omega = \omega_1 + \omega_2$  an adapted connection sum. If we denote  $G = G_1 \times G_2$ , its Lie algebra by  $\mathfrak{g} = \mathfrak{g}_1 \oplus \mathfrak{g}_2$  and  $k(\omega), k(\omega_1), k(\omega_2)$  the respective Weil homomorphisms, the following commutative diagram allows to write  $k(\omega) = k(\omega_1) \otimes k(\omega_2)$ :

$$\begin{array}{ccc}
 W(\mathfrak{g}) & \xrightarrow{L} & W(\mathfrak{g}_1) \otimes W(\mathfrak{g}_2) \\
 \downarrow k(\omega) & & \downarrow k(\omega_1) \otimes k(\omega_2) \\
 & & \Omega(P_1) \otimes \Omega(P_2) \\
 & & \downarrow \pi \\
 \Omega(P) & \xleftarrow{\bar{\Delta}^*} & \Omega(P_1 \times P_2)
 \end{array}$$

where  $L$  denotes the canonical isomorphism,  $\pi$  is defined by  $\pi(\alpha \otimes \beta) = p_1^* \alpha \wedge p_2^* \beta$ ,  $p_i: P_1 \times P_2 \rightarrow P_i$  the canonical projection, and  $\bar{\Delta}^*$  being induced by the canonical homomorphism  $\bar{\Delta}: P = P_1 + P_2 \rightarrow P_1 \times P_2$ .

Using  $L: W(g) \cong W(g_1) \otimes W(g_2)$ , the canonical even decreasing filtration of  $W(g)$  by G-DG-ideals can be written as

$$F^{2p}W(g) = \bigoplus_{j > p} \Lambda^* g^* \otimes S^j g^* \\ = \bigoplus_{j_1 + j_2 > p} \Lambda^* g^* \otimes S^{j_1} g_1^* \otimes S^{j_2} g_2^*, \quad p \geq 0$$

and we can define a new even decreasing filtration of  $W(g)$ , also by G-DG-ideals, by

$$'F^{2p}W(g) = \bigoplus_{j > p} \Lambda^* g^* \otimes S^j g_1^* \otimes S^* g_2^*, \quad p \geq 0.$$

Also,  $\Omega^*(P)$  has two decreasing filtrations by G-DG-ideals defined by the sheaves  $\underline{Q}_i^*$ ,  $i = 1, 2$ , of local 1-forms annihilating the foliation  $F_i$  on the base space  $M$ ; they are given by

$$F^p \Omega(P) = \Gamma(P, \pi^* \Lambda^p \underline{Q}_2^* \cdot \Omega_p), \\ 'F^p \Omega(P) = \Gamma(P, \pi^* \Lambda^p \underline{Q}_1^* \cdot \Omega_p), \quad p \geq 0.$$

Then, the Weil homomorphism  $k(\omega)$  of an adapted connection sum  $\omega = \omega_1 + \omega_2$  is filtration-preserving, that is

$$k(\omega) (F^{2p}W(g)) \subset F^p \Omega(P), \quad p \geq 0,$$

and if  $\omega_1$  and  $\omega_2$  are basic, then

$$k(\omega) (F^{2p}W(g)) \subset F^{2p} \Omega(P), \quad p \geq 0.$$

Moreover, one easily proves

**PROPOSITION 2.1.** — *Let  $\omega = \omega_1 + \omega_2$  be an adapted connection sum in  $P = P_1 + P_2$ . Then  $k(\omega) ('F^{2p}W(g)) \subset 'F^p \Omega(P)$ ,  $p \geq 0$ . If  $\omega_1$  and  $\omega_2$  are basic, then  $k(\omega) ('F^{2p}W(g)) \subset 'F^{2p} \Omega(P)$ ,  $p \geq 0$ .*

**COROLLARY 2.2.** — *For an adapted connection sum  $\omega = \omega_1 + \omega_2$ ,*

$$k(\omega) F^{2(q_2+1)}W(g) = 0, \quad k(\omega) 'F^{2(q_1+1)}W(g) = 0.$$

If  $\omega_1$  and  $\omega_2$  are basic,

$$k(\omega) F^{2((q_2/2)+1)} W(g) = 0, \quad k(\omega) {}'F^{2((q_1/2)+1)} W(g) = 0.$$

If we now consider the algebras of G-basic elements, we obtain similar properties for the Chern-Weil homomorphism  $h(\omega) : I(G) = I(G_1 \times G_2) \rightarrow \Omega(M)$  with respect to the following filtrations of  $I(G)$  and  $\Omega(M)$ :

$$F^{2p} I(G) = \bigoplus_{j \geq p} I^{2j}(G), \quad {}'F^{2p} I(G) = \bigoplus_{j \geq p} I^{2j}(G_1) \otimes I^{2j}(G_2), \quad p \geq 0$$

$$F^p \Omega(M) = \Gamma(M, \Lambda^p \underline{Q}_2^* \cdot \Omega_M),$$

$${}'F^p \Omega(M) = \Gamma(M, \Lambda^p \underline{Q}_1^* \cdot \Omega_M), \quad p \geq 0.$$

That is, since  $F^{q_2+1} \Omega(M) = 0$  and  ${}'F^{q_1+1} \Omega(M) = 0$ , we have

COROLLARY 2.3. — *Let  $\omega = \omega_1 + \omega_2$  be an adapted connection sum in an  $(F_1, F_2)$ -foliated bundle  $P = P_1 + P_2$ , and let  $h(\omega)$  denote the Chern-Weil homomorphism of  $P$ . Then*

$$h(\omega) F^{2(q_2+1)} I(G) = 0, \quad h(\omega) {}'F^{2(q_1+1)} I(G) = 0.$$

If, moreover,  $\omega_1$  and  $\omega_2$  are basic, then

$$h(\omega) F^{2((q_2/2)+1)} I(G) = 0, \quad h(\omega) {}'F^{2((q_1/2)+1)} I(G) = 0.$$

In particular, if  $P$  is the bundle of transverse frames of  $(F_1, F_2)$ , then Corollary 2.3 is the Vanishing Theorem for subfoliations stated in [4].

Next, let  $I \subset W(g)$  be the G-DG-ideal given by

$$I = F^{2(q_2+1)} W(g) + {}'F^{2(q_1+1)} W(g). \tag{2.1}$$

Then, by virtue of Corollary 2.2,  $I \subset \text{Ker}(k(\omega))$  and there is an induced G-DG-homomorphism  $k(\omega) : W(g)_I = W(g)/I \rightarrow \Omega(P)$ .

For any subgroup  $H \subset G$ , there is the relative ideal  $I_H$  of  $W(g, H) = W(g)_H$ , and thus if we construct

$$W(g, H)_I = W(g, H)/I_H = (W(g)_I)_H,$$

we can consider the induced DG-homomorphism

$$k(\omega)_H : W(g, H)_I \rightarrow \Omega(P)_H.$$

Now, if we assume  $H$  to be closed and  $P$  having an  $H$ -reduction given by a section  $s : M \rightarrow P/H$  of the induced map  $\hat{\pi} : P/H \rightarrow M$ ,

we can construct a DG-homomorphism as the composition

$$\Delta(\omega) = s^* \circ k(\omega)_H : W(g, H)_I \longrightarrow \Omega(P)_H \cong \Omega(P/H) \longrightarrow \Omega(M).$$

DEFINITION 2.4. — We shall call generalized characteristic homomorphism of the  $(F_1, F_2)$ -foliated bundle  $P$  the homomorphism  $\Delta_* = \Delta_{(F_1, F_2)}(P) : H(W(g, H)_I) \longrightarrow H_{DR}(M)$  induced by  $\Delta(\omega)$  in cohomology.

Remark. — If both  $\omega_1$  and  $\omega_2$  are basic connections, then  $k(\omega)$  vanishes on the ideal

$$I' = F^{2(lq_2/2)+1} W(g) + {}'F^{2(lq_1/2)+1} W(g)$$

and the generalized characteristic homomorphism of  $P$  will be  $\Delta_* : H(W(g, H)_I) \longrightarrow H_{DR}(M)$  because, under these conditions,  $\Delta(\omega)$  factorizes through  $p : W(g, H)_I \longrightarrow W(g, H)_{I'}$ , the canonical projection induced by the injection  $I \subset I'$ .

$\Delta_* = \Delta_{(F_1, F_2)}(P)$  is independent of the choice of  $\omega = \omega_1 + \omega_2$  in the following sense. Let  $\omega^0 = \omega_1^0 + \omega_2^0, \omega^1 = \omega_1^1 + \omega_2^1$  be two adapted connections sum in  $P$ . Let an  $H$ -reduction of  $P$  be given by a section  $s : M \longrightarrow P/H$ , and

$$\Delta_*^i = \Delta(\omega^i)_* : H(W(g, H)_I) \longrightarrow H_{DR}(M)$$

the homomorphism constructed using the connection  $\omega^i, i = 0, 1$ . Then,

PROPOSITION 2.5. —  $\Delta_*^0 = \Delta_*^1$ .

Proof. — Let  $f : M \times [0, 1] \longrightarrow M$  be the canonical projection, and let  $f^{-1}(F_k), k = 1, 2$ , the foliation inverse image of  $F_k$  via  $f$ . If  $P = P_1 + P_2$  is an  $(F_1, F_2)$ -foliated bundle over  $M$  then the inverse image  $P' = f^*(P) = f^*(P_1) + f^*(P_2)$  of  $P$  via  $f$  is  $f^{-1}(F_1, F_2) = (f^{-1}(F_1), f^{-1}(F_2))$ -foliated. Moreover, the connection  $\bar{\omega}$  given by

$$\bar{\omega}(X) = t(f^* \omega^1)(X) + (1 - t)(f^* \omega^0)(X), X \in T_{(u, t)}(P')$$

is obviously an adapted connection sum in  $P'$ .

On the other hand, if  $j_t : M \longrightarrow M \times [0, 1]$  is the canonical injection  $j_t(x) = (x, t)$ , for each  $t \in [0, 1]$ , then  $j_t^*(P') = P$  for any  $t \in [0, 1]$ ,  $\bar{j}_0^* \bar{\omega} = \omega^0, \bar{j}_1^* \bar{\omega} = \omega^1$  where  $\bar{j}_t : P \longrightarrow P'$



denotes the canonical lift of  $j_t$ . Thus, using  $\bar{\omega}$  to construct the generalized characteristic homomorphism of  $P'$ :  $\bar{\Delta}_* = \Delta_*(\bar{\omega})$ , we have  $\Delta_*^i = (j_i^*)_{DR} \circ \bar{\Delta}_*$ ,  $i = 0, 1$ . But, since  $(j_0^*)_{DR} = (j_1^*)_{DR}$ , then  $\Delta_*^0 = \Delta_*^1$ .

It is clear from the construction that  $\Delta_*$  depends a priori upon the H-reduction of  $P$  given by  $s$ . However, this construction is visibly independent of  $s$  if the closed subgroup  $H \subset G$  contains a maximal compact subgroup of  $G$ .

$\Delta_*$  has also the following properties of functoriality.

(A)  $\Delta_*$  is functorial under pullbacks.

This means more precisely the following. Let  $(F'_1, F'_2)$  and  $(F_1, F_2)$  be  $(q_1, q_2)$ -codimensional subfoliations on  $M'$  and  $M$  respectively, and let  $f: M' \rightarrow M$  be a differentiable map such that  $f_*(F'_i) \subset F_i$ ,  $i = 1, 2$ . Let  $P = P_1 + P_2$  be an  $(F_1, F_2)$ -foliated bundle over  $M$ , and let

$$P' = f^*P = f^*P_1 + f^*P_2$$

be the inverse image of  $P$  via  $f$ . Since each  $f^*P_i$  is  $F'_i$ -foliated ([1], Prop. 1.7), then  $P'$  is, in fact, an  $(F'_1, F'_2)$ -foliated bundle over  $M'$ . Then, if  $H \subset G$  is a closed subgroup and  $s: M \rightarrow P/H$  the section given an H-reduction of  $P$ ,  $s' = f^*s: M' \rightarrow P'/H$  gives an H-reduction of  $P'$  and we can easily prove

PROPOSITION 2.6. —  $\Delta_*(P') = f_{DR}^* \circ \Delta_*(P)$ .

It is clear that this result is applied in the particular case of  $f$  being transversal to the subfoliation  $(F_1, F_2)$  on  $M$  [4].

(B)  $\Delta_*$  is functorial under  $\rho$ -extensions.

This means more precisely the following. Let

$$\rho = (\rho_1, \rho_2): G = G_1 \times G_2 \rightarrow G' = G'_1 \times G'_2$$

a homomorphism of product Lie groups, that is, each  $\rho_i: G_i \rightarrow G'_i$  a Lie group homomorphism,  $i = 1, 2$ . If  $P$  is an  $(F_1, F_2)$ -foliated principal bundle over  $M$  and  $\omega$  an adapted connection sum in  $P$ , then  $P'$ , the extension of  $P$  by  $\rho$ , is  $(F_1, F_2)$ -foliated and  $\omega'$ , extension of  $\omega$  by  $\rho$ , is an adapted connection sum in  $P'$ .

Let  $H, H'$  be closed subgroups of  $G$  and  $G'$ , respectively,

such that  $\rho(H) \subset H'$ ; let  $I'$  and  $I$  be the ideals of  $W(g')$  and  $W(g)$  given by (2.1). Since  $W(d\rho)$  is graduation-preserving, then  $W(d\rho)(I') \subset I$  and diagram (4.72) in [7] can be used to state


PROPOSITION 2.7. —  $\Delta_*(P') = \Delta_*(P) \circ W(d\rho)^*$ .

3. Relation between  $\Delta_*(P)$  and  $\Delta_*(P_i)$ ,  $i = 1, 2$ .

Between the generalized characteristic homomorphism  $\Delta_*(P)$  of an  $(F_1, F_2)$ -foliated principal bundle  $P = P_1 + P_2$  and the generalized characteristic homomorphism  $\Delta_*(P_i)$  ([7]) of the  $F_i$ -foliated principal bundle  $P_i$ ,  $i = 1, 2$ , there exists a canonical relation given as follows.

Let  $\rho_i: G = G_1 \times G_2 \rightarrow G_i$  be the canonical projection,  $H_i \subset G_i$  a closed subgroup,  $i = 1, 2$ , and  $H = H_1 \times H_2 \subset G$ . Let  $s: M \rightarrow P/H$  be a section defining an  $H$ -reduction of  $P$  and let  $s_i: M \rightarrow P_i/H_i$  be the induced section defining an induced  $H_i$ -reduction of  $P_i$ . Then.

PROPOSITION 3.1. — *The diagram*



is commutative for each  $i = 1, 2$ . In fact, this diagram is also commutative at the cochain level.

*Proof.* — Since  $P_i$  is isomorphic (as  $F_i$ -foliated bundle) to the  $\rho_i$ -extension of  $P$ , and because  $\omega_i = (\rho_i)^* \omega$  is an adapted connection in  $P_i$ ,  $\omega = \omega_1 + \omega_2$  being an adapted connection sum in  $P$ , the following diagram commutes for each  $i = 1, 2$ :

$$\begin{array}{ccc}
 W(g_i, H_i) & \xrightarrow{W(d\rho_i)} & W(g, H) \\
 \downarrow k(\omega_i)_{H_i} & & \downarrow k(\omega)_H \\
 \Omega(P_i/H_i) & & \Omega(P/H) \\
 \searrow S_i^* & & \swarrow S^* \\
 & \Omega(M) &
 \end{array} \tag{3.1}$$

and we are reduced to show that  $W(d\rho_i)(F^{2(q_i+1)}W(g_i)) \subset I$  for each  $i = 1, 2$ .

For  $i = 2$ , this follows easily because  $W(d\rho_i)$  preserves the bigraduation and then

$$W(d\rho_2)W^{p,2q}(g_2) \subset W^{p,2q}(g).$$

For  $i = 1$ , the result follows from the fact that

$$W(d\rho_1)(\Lambda^u g_1^* \otimes S^v g_1^*) \subset \Lambda^u g^* \otimes S^v g_1^* \otimes S^0 g_2^*, \quad u, v \geq 0$$

since  $(d\rho_1)^* : Sg_1^* \rightarrow Sg^* = Sg_1^* \otimes Sg_2^*$  is given by

$$(d\rho_1)^*(\alpha) = \alpha \otimes 1.$$

*Remarks.* - 1) Since both  $\omega = \omega_1 + \omega_2$  and  $\omega_i$  are  $F_2$ -adapted connections, we can truncate the Weil algebras in diagram (3.1) at the degree  $q_2$  and thus, going into cohomology, obtain a commutative diagram relating the generalized characteristic homomorphisms of  $P$  and  $P_i$  as  $F_2$ -foliated principal bundles.

2) We can use  $\omega = \omega_1 + \omega_2$  to construct the generalized characteristic homomorphism of the  $F_2$ -foliated bundle  $P$  :

$$\Delta_{F_2}(P) : H(W(g, H)_{q_2}) \rightarrow H_{DR}(M).$$

Then, taking into account that the inclusion  $F^{2(q_2+1)}W(g) \subset I$  induces a projection  $p : W(g, H)_{q_2} \rightarrow W(g, H)_1$ , we obtain a commutative diagram

$$\begin{array}{ccc}
 H(W(g, H)_{q_2}) & \xrightarrow{p^*} & H(W(g, H)_1) \\
 \searrow \Delta_{F_2}(P) & & \swarrow \Delta_{(F_1, F_2)}(P) \\
 & H_{DR}(M) &
 \end{array} \quad (3.2)$$

and, therefore,  $\text{Im } \Delta_{F_2}(P) \subset \text{Im } \Delta_{(F_1, F_2)}(P)$ .

3) Let  $\rho : G = G_1 \times G_2 \longrightarrow G'$  be a homomorphism of Lie groups and consider the structure of  $F_2$ -foliated bundle on the  $\rho$ -extension  $P' = \rho_* P$  induced by the structure of  $F_2$ -foliated bundle underlying the  $(F_1, F_2)$ -foliated structure of  $P = P_1 + P_2$ .

Then, for suitable closed subgroups  $H \subset G$ ,  $H' \subset G'$ , the functoriality under  $\rho$ -extensions of the generalized characteristic homomorphism of foliated bundles ([7]) implies that the following diagram is commutative

$$\begin{array}{ccc}
 H(W(g', H')_{q_2}) & \xrightarrow{W(d\rho)^*} & H(W(g, H)_{q_2}) \\
 \searrow \Delta_{F_2}(P') & & \swarrow \Delta_{F_2}(P) \\
 & H_{DR}(M) &
 \end{array}$$

which combined with (3.2) leads to the following

**PROPOSITION 3.2.** — *Let  $P' \longrightarrow M$  be an  $F_2$ -foliated principal bundle with structure group  $G'$  and let  $P = P_1 + P_2$  be an  $(F_1, F_2)$ -foliated  $G$ -reduction of  $P$ . Assume  $i : P \longrightarrow P'$  be  $F_2$ -foliated compatibly with the homomorphism*

$$\rho : G = G_1 \times G_2 \longrightarrow G',$$

*and let  $H, H'$  be closed subgroups of  $G, G'$  respectively, verifying the suitable hypothesis. Then, the generalized characteristic homomorphism  $\Delta_{F_2}(P')$  of  $P'$  as  $F_2$ -foliated bundle factorizes through the generalized characteristic homomorphism  $\Delta_{(F_1, F_2)}(P)$*

of  $P$  as  $(F_1, F_2)$ -foliated bundle, that is, the following diagram is commutative:

$$\begin{array}{ccc}
 H(W(g', H')_{q_2}) & \xrightarrow{p^* \circ W(d\rho)^*} & H(W(g, H)_1) \\
 \Delta_{F_2}(P') \searrow & & \swarrow \Delta_{(F_1, F_2)}(P) \\
 & H_{DR}(M) &
 \end{array}$$

*Example.* — Let  $P' \cong L(Q_2) \cong L(\nu(F_1, F_2))$  be the canonically  $F_2$ -foliated bundle of transverse frames of  $F_2$ , and  $P$  the  $(F_1, F_2)$ -foliated bundle of transverse frames of  $(F_1, F_2)$ , which is a (not  $F_2$ -foliated) reduction of  $P'$  compatible with the canonical homomorphism  $\rho: Gl(q_1, \mathbb{R}) \times Gl(d, \mathbb{R}) \rightarrow Gl(q_2, \mathbb{R})$ .

If we consider in  $P'$  the  $\rho$ -extension of the  $F_2$ -foliated structure of  $P$ , this is not the canonical  $F_2$ -foliated structure of  $P'$ ; but, as it can be easily shown using the Lemma 5.3 in [4] (see [8]), both are integrably homotopic. Then, for suitable  $H, H'$  the proposition 3.2 provides the corresponding commutative diagram. If, moreover,  $H = O(q_1) \times O(d)$  and  $H' = O(q_2)$ , then  $\Delta_{F_2}(P')$  is just the characteristic homomorphism of the foliation  $F_2$ , whereas  $\Delta_{(F_1, F_2)}(P)$  is the characteristic homomorphism of the subfoliation  $(F_1, F_2)$  [4], as it will be established later.

Now, let us remark that a  $q$ -codimensional foliation  $F$  on  $M$  can be considered as a subfoliation on  $M$  in three different ways;  $(C_1): F_1 = F_2 = F, q_1 = q_2 = q$ ;  $(C_2): F_1 = TM, F_2 = F, q_1 = 0, q_2 = q$ ;  $(C_3): F_1 = F, F_2 = 0, q_1 = q, q_2 = n$ . Then, all the previous results particularize to these cases as follows:

*Case  $(C_1)$ .* — Here an  $(F_1, F_2)$ -foliated bundle  $P = P_1 + P_2$  is, in fact, an  $F$ -foliated bundle, the ideal  $I$  coincides with  $F^{2(q+1)}W(g)$ ,  $p^*$  in diagram (3.2) is an isomorphism and  $\Delta_{(F, F)}(P) = \Delta_F(P)$ .

Case  $(C_2)$ . — Here,  $P = P_1 + P_2$  is the sum of a flat bundle  $P_1$  and an  $F$ -foliated bundle  $P_2$ ; since

$$I = F^{2(q+1)} W(g) + 'F^2 W(g),$$

making calculations we obtain


$$W(g)_1 \cong \bigoplus_{j=0}^q \Lambda^* g^* \otimes S^0 g_1^* \otimes S^j g_2^*$$

$$W(g)_q \cong W(g)_1 \oplus \left( \bigoplus_{j=0}^{q-1} \bigoplus_{t=0}^{q-j} \Lambda^* g^* \otimes S_1^t g^* \otimes S_2^j g^* \right)$$

and then  $p^*$  in diagram (3.2) is surjective. Hence

$$\text{Im}(\Delta_F(P)) = \text{Im}(\Delta_{(TM, F)}(P)).$$

Case  $(C_3)$ . — In this case,  $P = P_1 + P_2$  is simply an ordinary bundle (that is, 0-foliated) which is not necessarily  $F$ -foliated. Thus, if we take  $H = G$  in diagram (3.2) and denote  $\tau : I(G) \rightarrow I(G)_n$  the canonical projection, we have a commutative diagram



where  $h^*$  denotes the Chern-Weil homomorphism of  $P$ . Thus, we can assert the following: if  $P = P_1 + P_2$  where  $P_1$  is a foliated bundle, then the Chern-Weil homomorphism of  $P$  vanishes on  $\text{Ker}(p^* \circ \tau)$ . Again, since any connection in  $P_2$  is basic with respect to the foliation by points on  $M$ , if  $P_1$  admits a basic connection then  $h^*$  vanishes on the kernel of the composition

$$I(G) \xrightarrow{\tau'} I(G)_{[n/2]} \xrightarrow{p'^*} I(G)_{1'}.$$

**4. Difference construction for  $\Delta_{(F_1, F_2)}(P)$ . Secondary invariants.**

The computation of  $H(W(g, H)_I)$  can be done from the general results in [7], Chapter 5, from where we shall take the notation.

We assume throughout that  $G$  is either connected or  $I(G) \cong I(G_0) \equiv I(g)$  for the connected component  $G_0$  of  $G$ ; the closed subgroup  $H \subset G$  is assumed to have finitely many connected components.

Then, let us consider in the  $G$ -DG-algebra  $W(g)_I$  the canonical connection given by the projection  $k : W(g) \rightarrow W(g)_I$ .

If the pair  $(g, h)$  is reductive ( $h = \text{Lie algebra of } H$ ), in accordance with Theorem 5.82 in [7] there exists a homomorphism  $\zeta(W(g)_I, H) : A(W(g)_I, H) \rightarrow (W(g)_I)_H = W(g, H)_I$  which induces an isomorphism in cohomology. In this way, the generalized characteristic homomorphism  $\Delta_{(F_1, F_2)}(P)$  of  $P$  will have the same image as the composition

$$H(A(W(g)_I, H)) \xrightarrow[\cong]{\zeta(W(g)_I, H)_*} H(W(g, H)_I) \xrightarrow{\Delta_{(F_1, F_2)}(P)} H_{DR}(M)$$

induced by the cochain map  $\tilde{\Delta}(\omega) = \Delta(\omega) \circ \zeta(W(g)_I, H)$ . In fact, the evaluation of  $\tilde{\Delta}(\omega)$  on the complex

$$A(W(g)_I, H) = \Lambda P_g \otimes (W(g)_I)_g \otimes I(H) = \Lambda P_g \otimes I(G)_I \otimes I(H)$$

is equal to that of Theorem 5.95 in [7] for the case of a foliated bundle.

If we now assume the pair  $(g, h)$  to be special Cartan (CS), then, by Theorem 5.107 in [7], there is an isomorphism

$$\bar{\beta} : H(\hat{A}(W(g)_I)) \otimes_{I(g)} I(H) \xrightarrow{\cong} H(A(W(g)_I, H))$$

where  $\hat{A}(W(g)_I) = \Lambda \hat{P} \otimes (W(g)_I)_g$ . Thus  $\Delta_{(F_1, F_2)}(P)$  has the same image as the composition  $\Delta_{(F_1, F_2)}(P) \circ \zeta(W(g)_I, H) \circ \bar{\beta}$ . Then taking into account that  $\hat{A}(W(g)_I) \subset A(W(g)_I, H)$ , we consider the composition

$$\hat{\Delta}(\omega) : \hat{A}(W(g)_I) \rightarrow A(W(g)_I, H) \xrightarrow{\tilde{\Delta}(\omega)} \Omega(M)$$

and, thus, the characteristic homomorphism  $\Delta_{(F_1, F_2)}(P)$  will be realized by  $\hat{\Delta}_* \otimes h'_* : H(\hat{A}(W(g)_I)) \otimes_{I(G)} I(H) \longrightarrow H_{DR}(M), h'_*$  being the characteristic homomorphism of the H-reduction  $P'$  of  $P$ . See 5.112 in [7] for more details.

• In particular, let us assume that  $P = P_1 + P_2$  is the bundle of transverse frames of  $(F_1, F_2)$ , and take

$$H = O(q_1) \times O(d) \subset Gl(q_1, \mathbf{R}) \times Gl(d, \mathbf{R}) = G.$$

Since  $gl(q_1, \mathbf{R})$  and  $gl(d, \mathbf{R})$  are reductive Lie algebras and  $(gl(q_1, \mathbf{R}) \times gl(d, \mathbf{R}), o(q_1) \times o(d))$  is symmetric, this pair will be special Cartan and the previous construction can be used. Then,  $\Delta_{(F_1, F_2)}(P)$  can be considered as defined on  $H(\hat{A}_I) \otimes_{I(G)} I(H)$ , where  $\hat{A}_I = \hat{A}(W(g)_I) = \Lambda \hat{P} \otimes I(gl(q_1, \mathbf{R}) \times gl(d, \mathbf{R}))_I$ . But, as it happens in the case of the bundle of transverse frames of a foliation [7],  $H(\hat{A}_I) \otimes_{I(G)} I(H) \cong H(\hat{A}_I)$ , and then

$$\hat{\Delta}_{(F_1, F_2)}(P) = \hat{\Delta}_* : H(\hat{A}_I) \longrightarrow H_{DR}(M).$$

On the other hand,  $I(gl(q_1, \mathbf{R})) = \mathbf{R}[c_1, \dots, c_{q_1}]$ ,  $I(gl(d, \mathbf{R})) = \mathbf{R}[c'_1, \dots, c'_d]$  and  $\Lambda \hat{P} = \Lambda \hat{P}_1 \otimes \Lambda \hat{P}_2$ ,  $\hat{P}_i$  being the Samelson subspace of the pair  $(g_i, h_i)$ ,  $i = 1, 2$ ; since both pairs are special Cartan,

$$\Lambda \hat{P}_1 = \Lambda(y_1, y_2, \dots, y_{q'_1}), \quad \Lambda \hat{P}_2 = \Lambda(y'_1, y'_3, \dots, y'_{d'})$$

where  $y_i = \sigma c_i$ ,  $y'_i = \sigma' c'_i$  and  $q'_1 = 2[(q_1 + 1)/2] - 1$ ,  $d' = 2[(d + 1)/2] - 1$ ,  $\sigma$  and  $\sigma'$  being the suspension maps. Therefore

$$\hat{A}_I = \Lambda(y_1, y_3, \dots, y_{q'_1}) \otimes \Lambda(y'_1, y'_3, \dots, y'_{d'})$$

$$\otimes \frac{\mathbf{R}[c_1, \dots, c_{q_1}] \otimes \mathbf{R}[c'_1, \dots, c'_d]}{I_g}$$

where

$$I_g = I \cap I(g) = \langle \{\alpha \otimes \beta \in I^{j_1}(g_1) \otimes I^{j_2}(g_2) / j_1 > q_1 \text{ or } j_1 + j_2 > q_2\} \rangle.$$

That is,  $\hat{A}_I = WO_I$ , the graded differential algebra defined in [4]. Therefore, the generalized characteristic homomorphism of the bundle of transverse frames of the subfoliation  $(F_1, F_2)$  coincides with the characteristic homomorphism of  $(F_1, F_2)$  as defined in [4]

$$\lambda_{(F_1, F_2)}^* : H(WO_I) \longrightarrow H_{DR}(M).$$



From this point of view, the generalized characteristic homomorphism  $\Delta_{(F_1, F_2)}(P)$  of an  $(F_1, F_2)$ -foliated bundle  $P = P_1 + P_2$  generalizes the characteristic homomorphism of the subfoliation  $(F_1, F_2)$  in the same way as Kamber-Tondeur's characteristic homomorphism of a foliated bundle generalizes Bott's characteristic homomorphism of a foliation ([7], [2]).

In order to construct the algebra of secondary characteristic invariants, from now on, we shall consider an  $(F_1, F_2)$ -foliated bundle  $P = P_1 + P_2$ ,  $H \subset G$  a closed subgroup with finitely many connected components and such that the pair of Lie algebras  $(g, h)$  be reductive. Let us denote  $P'$  the  $H$ -reduction of  $P$  used to define the characteristic homomorphism  $\Delta_{(F_1, F_2)}(P)$  of  $P$  and, to simplify the notation, put  $A_1 = A(W(g)_1, H)$ .

Let  $p: A_1 \rightarrow I(G)_1 \otimes_{I(G)} I(H)$  the composition of the canonical projection along  $\Delta P_g$ ,  $\lambda: A_1 \rightarrow I(G)_1 \otimes I(H)$  with the canonical map.

**DEFINITION 4.1.** —  $H(K_1)$ , where  $K_1 = \text{Ker } p$ , is called the algebra of secondary characteristic invariants of  $P$ .

**PROPOSITION 4.2.** — There is a short exact sequence of algebras

$$0 \rightarrow H(K_1) \rightarrow H(W(g, H)_1) \rightarrow I(G)_1 \otimes_{I(G)} I(H) \rightarrow 0. \quad (4.1)$$

*Proof.* — Consider the short exact sequence of complexes

$$0 \rightarrow K_1 \rightarrow A_1 \rightarrow I(G)_1 \otimes_{I(G)} I(H) \rightarrow 0.$$

Then (4.1) appears by writing up the associated long exact sequence of homology whose connecting homomorphism is null, and because  $H(A_1) \cong H(W(g, H)_1)$ .

□

The non-triviality of  $\Delta_{(F_1, F_2)}(P)/_{H(K_1)}$  is a measure for the incompatibility of the  $(F_1, F_2)$ -foliated structure of  $P = P_1 + P_2$  with its  $H$ -reduction  $P'$ ; that is,

**PROPOSITION 4.3.** — Let  $P = P_1 + P_2$  be an  $(F_1, F_2)$ -foliated bundle,  $H = H_1 \times H_2 \subset G$  a closed subgroup and  $P'$  an  $H$ -reduction of  $P$  which is  $(F_1, F_2)$ -foliated and such that, if

$\iota: H \rightarrow G$  is the injection, then the  $(F_1, F_2)$ -foliated structure of  $P$  is, in fact, the  $\iota$ -extension of that of  $P'$ . Then

$$\Delta_{(F_1, F_2)}(P)/_{H(K_I)} = 0.$$

*Proof.* – Applying Proposition 2.7 to the homomorphism  $\iota: (H, H) \rightarrow (G, H)$  we obtain the commutative diagram

$$\begin{array}{ccc}
 H(W(g, H)_I) & \xrightarrow{\Delta_*(P)} & H_{DR}(M) \\
 \downarrow W(d\iota)^* & & \nearrow \Delta_*(P') \\
 H(W(h, H)_I) \cong I(H)_I & & 
 \end{array} \quad (4.2)$$

and hence  $\Delta_*(P)/_{\text{Ker}(W(d\iota)^*)} = 0$ .

Moreover, there is a commutative diagram

$$\begin{array}{ccc}
 W(g, H)_I & \xrightarrow{W(d\iota)} & I(H)_I \\
 \uparrow \xi(W(g)_I, H) & & \uparrow \psi \\
 A_I & \xrightarrow{\quad} & I(G)_I \otimes_{I(G)} I(H)
 \end{array}$$

where  $\psi$  is the canonical projection of

$$I(G)_I \otimes_{I(G)} I(H) \cong I(H)/I \cdot I(H)$$

onto  $I(H)_I = I(H)/I$ . Thus, going into cohomology, we obtain a factorization  $H(W(g, H)_I) \rightarrow I(G)_I \otimes_{I(G)} I(H) \rightarrow I(H)_I$  of the vertical homomorphism in (4.2). Then, because

$$H(K_I) = \text{Ker} \{H(W(g, H)_I) \rightarrow I(G) \otimes_{I(G)} I(H)\}$$

by virtue of Proposition 4.2, we have  $H(K_I) \subset \text{Ker}(W(d\iota)^*)$ .

□

Moreover, as in the usual case of foliated bundles [7], we have

PROPOSITION 4.4. — *There is a splitting homomorphism*

$$\kappa : I(G)_1 \otimes_{I(G)} I(H) \longrightarrow H(W(g, H)_1)$$

of the short exact sequence (4.1) and the composition  $\Delta_*(P) \circ \kappa$  is induced by the characteristic homomorphism of  $P'$  :

$$h_*(P') : I(H) \longrightarrow H_{DR}(M).$$

5. Restriction to the leaves.

In this section we shall discuss the restriction of an  $(F_1, F_2)$ -foliated bundle  $P = P_1 + P_2$  to the leaves of each foliation  $F_i$ ,  $i = 1, 2$ . In order to do that, let us previously discuss the restriction to the leaves of an  $F_2$ -foliated bundle.

So, let  $(F_1, F_2)$  be a  $(q_1, q_2)$ -codimensional subfoliation on  $M$ ,  $L$  a leaf of  $F_1$  and  $j : L \rightarrow M$  the canonical immersion. Since  $F_2 \subset F_1$ ,  $F_2$  induces on  $L$  a foliation which will be denoted by  $F_L$ ; note that  $\text{codim}(F_L) = d = q_2 - q_1$  while  $\text{codim}(F_2) = q_2$ . Obviously,  $j$  maps the leaves of  $F_L$  into leaves of  $F_2$ .

Now, let  $\pi : P \rightarrow M$  be a  $G$ -principal fibre bundle and denote  $P' = j^*P$  the inverse image of  $P$  via  $j$ . Then  $\pi' : P' \rightarrow L$ , the restriction of  $P$  to  $L$ , is a  $G$ -principal fibre bundle and we shall denote  $\bar{j} : P' \rightarrow P$  the canonical injection. The following result is known [1] :

PROPOSITION 5.1. — *If  $P$  is  $F_2$ -foliated then  $P'$  is  $F_L$ -foliated. Moreover, if  $\omega$  is an adapted connection in  $P$  then  $\bar{j}^*\omega$  is an adapted connection in  $P'$ .*

Precisely the latter condition allows to consider, using connections  $\omega$  and  $\bar{j}^*\omega$ , a commutative diagram

$$\begin{array}{ccc}
 H(W(g, H)_{q_2}) & \xrightarrow{\Delta_*(P)} & H_{DR}(M) \\
 p^* \downarrow & & \downarrow j^* \\
 H(W(g, H)_d) & \xrightarrow{\Delta_*(P')} & H_{DR}(L)
 \end{array} \tag{5.1}$$

where  $p: W(g, H)_{q_2} \rightarrow W(g, H)_d$  is the canonical projection ( $d \leq q$ ),  $H \subset G$  is a subgroup satisfying the usual hypothesis and  $\Delta_*(P)$ ,  $\Delta_*(P')$  are the generalized characteristic homomorphisms of  $P$  and  $P'$ .

For example,  $Q_0 = F_1/F_2$  (the normal bundle of  $F_2$  relative to  $F_1$ ) is an  $F_2$ -foliated vector bundle on account of the existence on it of the so-called Bott connection [4], [1]. Moreover,  $Q_L = TL/F_L$ , the normal bundle of  $F_L$ , is canonically isomorphic to  $j^*Q_0$  [1] in such way that the Bott connection in  $Q_0$  pulls back via  $j$  to the Bott connection in  $Q_L$ . Therefore, the frame bundle of  $Q_0$ ,  $P$ , is an  $F_2$ -foliated  $G1(d, \mathbb{R})$ -principal bundle, and  $P' = j^*P$  is precisely the bundle of transverse frames of  $F_L$ . Thus, through the corresponding isomorphisms, diagram (5.1) becomes:

$$\begin{array}{ccc}
 H(W(gl(d, \mathbb{R}), O(d))_{q_2}) & \xrightarrow{\Delta_*} & H_{DR}(M) \\
 p^* \downarrow & & \downarrow j^* \\
 H(W(gl(d, \mathbb{R}), O(d))_d) \cong H(WO_d) & \xrightarrow{\Delta'_*} & H_{DR}(L)
 \end{array}$$

where  $\Delta'_*$  is just the usual characteristic homomorphism of foliation  $F_L$  on  $L$ .

Next, we shall discuss the restriction to a leaf of  $F_2$ . Thus, provided that we do not need to use the foliation  $F_1$ , we shall assume only one foliation  $F$  on  $M$ ,  $L$  a leaf of  $F$  and  $j: L \rightarrow M$  the canonical immersion. Now if  $\pi: P \rightarrow M$  is a  $G$ -principal bundle and  $P' = j^*P$  is the inverse image of  $P$  via  $j$ , we have

**PROPOSITION 5.2.** — *Each  $F$ -foliated bundle structure on  $P$  determines a flat bundle structure on  $P'$  in such way that if  $\omega$  is an adapted connection in  $P$ , then  $\omega' = \bar{j}^*\omega$  is a flat connection in  $P'$ .*

Therefore, if we consider on  $M$  the subfoliation  $(F, F)$  then the foliation  $F_L$  induced on  $L$  is trivial, that is,  $F_L = TL$ ,

and taking into account that  $W(g, H)_0 \cong (\Lambda g^*)_H \cong \Lambda(g/h)^{*H}$ , diagram (5.1) becomes

$$\begin{array}{ccc}
 H(W(g, H)_q) & \xrightarrow{\Delta_*} & H_{DR}(M) \\
 p^* \downarrow & & \downarrow j^* \\
 H(g, H) & \xrightarrow{\Delta'_*} & H_{DR}(L)
 \end{array} \tag{5.2}$$

$\Delta'_*$  being the generalized characteristic homomorphism of  $P'$  as flat bundle [7].

*Example.* – Let  $P$  be the bundle of transverse frames of  $F$ . Then, if  $\nu F = TM/F$  is the normal bundle of  $F$ ,  $\nu L = \nu F/L$  is the normal bundle of the leaf  $L$  of  $F$  and  $P' = j^*P$  is just the bundle of frames of  $\nu L$ . Following Goldman [6], any connection in  $P$  adapted to its canonical structure of  $F$ -foliated bundle will be said a foliation connection, and a connection in  $P'$  obtained as inverse image of a foliation connection will be said a leaf connection. In fact, Goldman showed that there is an unique leaf connection which is flat, and one easily checks that  $\Delta'_*$  in diagram (5.2) is nothing but the so-called holonomy homomorphism of the leaf  $L$  [6].

Again, let  $(F_1, F_2)$  be a  $(q_1, q_2)$ -codimensional subfoliation on  $M$ ,  $L_1$  a leaf of  $F_1$ ,  $j_1: L_1 \rightarrow M$  the canonical immersion,  $F_{L_1}$  the foliation on  $L_1$  induced by  $F_2$ ,  $P = P_1 + P_2$  an  $(F_1, F_2)$ -foliated bundle on  $M$  and  $P' = j_1^*P$  its inverse image via  $j_1$ . Then, since  $P$  is also  $F_2$ -foliated we can apply to it all previous results; so, in particular, we can construct a diagram (5.1) for this  $P = P_1 + P_2$ . On the other hand,

$$P' = j_1^*P_1 + j_1^*P_2 ;$$

then, applying the previous results to each  $j_1^*P_i$ ,  $i = 1, 2$ , it follows that  $P'$  is  $(TL_1, F_{L_1})$ -foliated over  $L_1$ . Moreover, if  $\omega$  is an adapted connection sum in  $P$  then  $\omega' = j_1^*\omega$  is an adapted connection sum in  $P'$ . If  $I$  and  $I'$  are the ideals given by (2.1) for the pairs  $(q_1, q_2)$  and  $(0, d)$ , respectively, then

$I \subset I'$  and, for an appropriate subgroup  $H$ , we obtain a commutative diagram

$$\begin{array}{ccc}
 H(W(g, H)_I) & \xrightarrow{\Delta_{(F_1, F_2)}(P)} & H_{DR}(M) \\
 \downarrow p'^* & & \downarrow j_1^* \\
 H(W(g, H)_{I'}) & \xrightarrow{\Delta_{(TL_1, FL_1)}(P')} & H_{DR}(L_1)
 \end{array} \quad (5.3)$$

where  $p'^*$  is induced by the canonical projection. If we now combine (5.3) with (5.1) through (3.2), we obtain

$$\begin{array}{ccc}
 H(W(g, H)_{q_2}) & \xrightarrow{\Delta_{F_2}(P)} & H_{DR}(M) \\
 \downarrow p^* & \searrow q^* & \nearrow \Delta_{(F_1, F_2)}(P) \\
 & H(W(g, H)_I) & \downarrow j_1^* \\
 & \downarrow p'^* & \\
 H(W(g, H)_a) & \xrightarrow{\Delta_{FL_1}(P')} & H_{DR}(L_1) \\
 \searrow q'^* & \nearrow \Delta_{(TL_1, FL_1)}(P') & \\
 & H(W(g, H)_{I'}) & 
 \end{array} \quad (5.4)$$

Now, if  $L_2$  is a leaf of  $F_2$  and  $j_2: L_2 \rightarrow M$  is its canonical immersion, then  $(F_1, F_2)$  induces on  $L_2$  the trivial subfoliation  $(TL_2, TL_2)$ . Therefore, the restriction to  $L_2$  of an  $(F_1, F_2)$ -foliated bundle  $P = P_1 + P_2$  is a flat bundle, and hence we obtain a commutative diagram similar to (5.4):

$$\begin{array}{ccc}
 H(W(g, H)_{q_2}) & \xrightarrow{\Delta_{F_2}(P)} & H_{DR}(M) \\
 \downarrow p^* & \searrow q^* & \nearrow \Delta_{(F_1, F_2)}(P) \\
 & H(W(g, H)_I) & \downarrow j_2^* \\
 & \swarrow p'^* & \\
 H(g, H) & \xrightarrow{\Delta_*(P')} & H_{DR}(L_2)
 \end{array} \quad (5.5)$$

If we now assume that  $L_1$  contains  $L_2$ ,  $j_0: L_2 \rightarrow L_1$  being the canonical immersion with  $j_2 = j_1 \circ j_0$ , then, using Proposition 3.1 and taking the closed subgroups

$$H_1 \subset G_1, H_2 \subset G_2, H = H_1 \times H_2$$

and  $\rho_i: G = G_1 \times G_2 \rightarrow G_i$ ,  $i = 1, 2$ , the canonical projections, there is a commutative diagram

$$\begin{array}{ccccc}
 H(g_1, H_1) & \xrightarrow{\rho_1^*} & H(g, H) & \xleftarrow{\rho_2^*} & H(g_2, H_2) \\
 \downarrow \Delta_*(j_1^* P_1) & & \downarrow \Delta_*(P') & \swarrow \Delta_*(j_2^* P_2) & \\
 H_{DR}(L_1) & \xrightarrow{j_0^*} & H_{DR}(L_2) & & 
 \end{array} \quad (5.6)$$

All these results, when particularized in certain examples, provide a starting point for a study of the holonomy of the leaves of a subfoliation similar to that of Goldman [6] for the leaves of a foliation.

*Example.* – With the previous notations, let

$$P = L(Q_1) + L(Q_0)$$

be the bundle of transverse frames of  $(F_1, F_2)$  and let  $(L_1, L_2)$  be a “leaf” of  $(F_1, F_2)$  (that is,  $L_i$  leaf of  $F_i$  and  $L_2 \subset L_1$ ); then  $P' = j_2^* P = L(j_0^*(\nu L_1)) + L(j_2^*(Q_0))$  is a reduction of the bundle of frames of the “normal bundle of  $(L_1, L_2)$ ” defined as  $\nu(L_1, L_2) = j_0^*(\nu L_1) \oplus j_2^*(Q_0)$ ,  $\nu L_1$  being the normal bundle of the leaf  $L_1$  [6], that is,  $\nu L_1 = Q_1/L_1$ . With a terminology analogous to that of Goldman, we call leaf connection any connection in  $P'$  obtained by pull-back of any adapted connection sum in  $P$ . Then, the following proposition can be easily proved :

**PROPOSITION 5.3.** – *There exists a unique leaf connection in  $P'$ . Moreover, this connection is flat.*

Through this result, we can state easily vanishing and obstruction theorems for the leaves of a subfoliation similar

to those in [6] for the leaves of a foliation, because the real Pontrjagin ring of  $\nu(L_1, L_2)$  is trivial and this fact provides a necessary condition for a pair  $(N_1, N_2)$  of connected manifolds, with injective immersions  $j_i: N_i \rightarrow M, i = 1, 2, j_0: N_2 \rightarrow N_1$  and  $j_2 = j_1 \circ j_0$ , to be a leaf of a subfoliation on  $M$ .

Now, if  $P'' = L(Q_2)$  and

$$\rho: G1(q_1, \mathbf{R}) \times G1(d, \mathbf{R}) \rightarrow G1(q_2, \mathbf{R})$$

is the canonical homomorphism, then, using the example that follows Proposition 3.2 and taking a closed subgroup

$$H' \subset G1(q_2, \mathbf{R})$$

such that  $\rho(H) \subset H'$ , we first obtain a commutative diagram combining (5.2) with (5.5):


$$\begin{array}{ccc}
 H(W(gl(q_1, \mathbf{R}) \oplus gl(d, \mathbf{R}), H)_1) & \xrightarrow{\Delta_{(F_1, F_2)}(P)} & H_{DR}(M) \\
 \downarrow p'^* & \swarrow q^* \circ W(d\rho)^* & \nearrow \Delta_{F_2}(P'') \\
 & H(W(gl(q_2, \mathbf{R}), H')_{q_2}) & \\
 & \downarrow p^* & \downarrow j_2^* \\
 H(gl(q_1, \mathbf{R}) \oplus gl(d, \mathbf{R}), H) & \xrightarrow{\Delta_*(P')} & H_{DR}(L_2) \\
 \swarrow \rho^* & \nearrow \Delta_*(j_2^* P'') & \\
 & H(gl(q_2, \mathbf{R}), H') & 
 \end{array} \tag{5.7}$$

and then, taking into account diagram (5.6):

$$\begin{array}{ccc}
 H(gl(q_2, \mathbf{R}), H') & & \\
 \downarrow \rho^* & \searrow \Delta_*(j_2^* P'') & \\
 H(gl(q_1, \mathbf{R}) \oplus gl(d, \mathbf{R}), H) & \xrightarrow{\Delta_*(P')} & H_{DR}(L_2) \\
 \uparrow \rho_1^* & & \uparrow j_0^* \\
 H(gl(q_1, \mathbf{R}), H) & \xrightarrow{\Delta_*(j_1^* P_1)} & H_{DR}(L_1)
 \end{array} \tag{5.8}$$




Now, we assume  $H = O(q_1) \times O(d)$  and  $H' = O(q_2)$ . In this case Goldman shows that  $p^*$  in diagram (5.7) is the zero homomorphism and concludes that the secondary foliation classes of  $F_2$  vanish in the leaves  $L_2$ . Essentially with the same arguments, one can prove that the homomorphism  $p'^*$  in diagram (5.7) is also zero and assert that the restriction to  $L_2$  of every secondary subfoliation class of  $(F_1, F_2)$  vanishes. Moreover, the homomorphism  $\Delta_*(P')$  is similar to the holonomy homomorphism defined by Goldman, and hence it can be called the holonomy homomorphism of the leaf  $(L_1, L_2)$  and denoted by  $\phi_{F,L}^*$ . Then, diagram (5.8) relate the holonomy homomorphism of  $(L_1, L_2)$  with that of each  $L_i$ ,  $i = 1, 2$ . Through the canonical isomorphisms we obtain the following commutative diagram:



where  $\varrho_i = 2[(q_i + 1)/2] - 1$ ,  $i = 1, 2$ ;  $\varrho'_1 = 2[(d + 1)/2] - 1$ ;  $\varrho''_1 = \varrho_1$ .

Obviously, the case of a subfoliation with trivialized normal bundle can be also discussed; to do that, it suffices to take  $H'$  as the trivial subgroup, and the diagram (5.8) becomes




This result may be used in order to obtain topological obstructions to the existence of subfoliations. Reinhart [10] exhibits a first example of these obstructions which can be expressed in our language as follows.

Let  $(F_1, F_2)$  be a  $(1, 2)$ -codimensional subfoliation on a manifold  $M$  with trivialized normal bundle; suppose  $F_1$  defined by the global 1-form  $\alpha_2$  and  $F_2$  defined by the global 1-forms  $\alpha_1, \alpha_2$ . Hence there exist 1-forms  $\tau_{11}, \tau_{21}, \tau_{22}$  on  $M$  such that  $d\alpha_1 = \alpha_1 \wedge \tau_{11} + \alpha_2 \wedge \tau_{21}$ ,  $d\alpha_2 = \alpha_2 \wedge \tau_{22}$ .

If  $(L_1, L_2)$  is a leaf of  $(F_1, F_2)$ , let us consider the 1-forms on  $L_2$  given by

$$\tau_{11}^L = j_2^*(\tau_{11}), \quad \tau_{21}^L = j_2^*(\tau_{21}), \quad \tau_{22}^L = j_2^*(\tau_{22}).$$

In this case, the previous diagram writes, at the cochain level, as



and, from it, we obtain the following holonomy classes :

a) for  $L_1$  as leaf of  $F_1$  :

$$\phi_{F_1, L_1}^*(h_1) = [\tau_{22/L_1}] \in H_{DR}(L_1)$$

b) for  $L_2$  as leaf of  $F_2$  :

$$\phi_{F_2, L_2}^*(h_1) = [\tau_{11}^L + \tau_{22}^L] \in H_{DR}(L_2), \quad \phi_{F_2, L_2}^*(h_2) = 0$$

since  $h_2 \in \text{Ker } \rho^*$ . In fact, Reinhart shows the vanishing of  $\phi_{F_2, L_2}^*(h_2)$  through a direct computation.

c) for  $(L_1, L_2)$  as leaf of  $(F_1, F_2)$  :

$$\phi_{F, L}^*(h'_1) = [\tau_{11}^L] \in H_{DR}(L_2), \quad \phi_{F, L}^*(h''_1) = [\tau_{22}^L] \in H_{DR}(L_2)$$

$$\phi_{F, L}^*(h'_1 + h''_1) = [\tau_{11}^L + \tau_{22}^L] \in H_{DR}(L_2).$$

Now, by comparing with Reinhart results one can deduce :

1) the vanishing of certain holonomy classes of  $L_2$  follows from the fact that they are obtained from elements of  $\text{Ker } \rho^*$ .

2) the image of  $\Lambda(h'_1)$  by  $\phi_{F, L}^*$  gives holonomy classes which cannot be obtained if we consider each leaf separately.

## BIBLIOGRAPHY

- [1] G. ANDRZEJCZAK, Some characteristic invariants of foliated bundles, Institute of Mathematics, Polish Academy of Sciences, Preprint 182, Warszawa, 1979.
- [2] R. BOTT, Lectures on characteristic classes and foliations, *Lecture Notes in Math.*, Vol. 279, Springer, Berlin, 1972.
- [3] L.A. CORDERO and P.M. GADEA, Exotic characteristic classes and subfoliations, *Ann. Inst. Fourier*, Grenoble, 26-1 (1976), 225-237 ; errata, *ibid.* 27, fasc. 4 (1977).
- [4] L.A. CORDERO and X. MASA, Characteristic classes of subfoliations, *Ann. Inst. Fourier*, Grenoble, 31-2 (1981), 61-86.
- [5] B.L. FEIGIN, Characteristic classes of flags of foliations, *Funct. Anal. and its Appl.*, 9 (1975), 312-317.

- [6] R. GOLDMAN, The holonomy ring of the leaves of foliated manifolds, *J. Differential Geometry*, 11 (1976), 411-449.
- [7] F.W. KAMBER and Ph. TONDEUR, Foliated bundles and characteristic classes, *Lecture Notes in Math.*, Vol 493, Springer, Berlin, 1975.
- [8] X. MASA, Characteristic classes of subfoliations II, preprint.
- [9] R. MOUSSU, Sur les classes exotiques des feuilletages, *Lecture Notes in Math.*, Vol. 392, Springer, Berlin, 1974, 37-42.
- [10] B.L. REINHART, Holonomy invariants for framed foliations, *Lecture Notes in Math.*, Vol. 392, Springer, Berlin, 1974, 47-52.

Manuscrit reçu le 17 mai 1983  
révisé le 8 novembre 1983.

José Manuel CARBALLÉS  
Departamento de Geometría y Topología  
Facultad de Matemáticas  
Universidad de Santiago de Compostela  
Santiago de Compostela (Espagne).