

JOURNAL

de Théorie des Nombres
de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Solomon VISHKAUTSAN

Quadratic rational functions with a rational periodic critical point of period 3

Tome 31, n° 1 (2019), p. 49-79.

<http://jtnb.cedram.org/item?id=JTNB_2019__31_1_49_0>

© Société Arithmétique de Bordeaux, 2019, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

Quadratic rational functions with a rational periodic critical point of period 3

par SOLOMON VISHKAUTSAN

Avec un appendice par Michael Stoll

RÉSUMÉ. Nous établissons une classification complète des graphes des points rationnels prépériodiques des fonctions rationnelles de degré 2 ayant un point critique rationnel de période 3 sous les hypothèses suivantes : ces fonctions n’admettent pas de points de période supérieure à 5 et une certaine conjecture sur le nombre de points rationnels sur une courbe affine plane de genre 6 est vraie. Nous montrons qu’il y a exactement six graphes possibles et que les fonctions associées possèdent au plus onze points prépériodiques.

ABSTRACT. We provide a complete classification of possible graphs of rational preperiodic points of quadratic rational functions defined over the rationals with a rational periodic critical point of period 3, under two assumptions: that these functions have no periodic points of period at least 5 and the conjectured enumeration of rational points on a certain genus 6 affine plane curve. We show that there are exactly six such possible graphs, and that rational functions satisfying the conditions above have at most eleven rational preperiodic points.

1. Introduction

In this article we continue the classification of preperiodicity graphs of quadratic rational functions defined over \mathbb{Q} with a \mathbb{Q} -rational periodic critical point that was begun in [3]. Our aim in this article is to provide a complete classification of rational quadratic functions defined over \mathbb{Q} with a \mathbb{Q} -rational periodic critical point of period 3.

Let $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a rational function, defined over some base field K . A point $P \in \mathbb{P}^1$ is called *periodic* for ϕ if there exists a positive integer n such that $\phi^n(P) = P$. The minimal such n is called the *period* of P . A point $P \in \mathbb{P}^1$ is called *preperiodic* for ϕ if there exists a nonnegative integer m such that $\phi^m(P)$ is periodic for ϕ .

Manuscrit reçu le 12 janvier 2018, accepté le 6 juin 2018.

2010 *Mathematics Subject Classification.* 37P35, 37P05.

Mots-clefs. rational functions, preperiodic points, preperiodicity graphs, moduli curves.

Research for this article has been supported by the Minerva Foundation and by ERC Advanced Grant “TADMICAMT”.

One of the main motivations for this article is the following conjecture of Morton and Silverman.

Conjecture (Uniform Boundedness Conjecture [12]). *Let $\phi : \mathbb{P}^N \rightarrow \mathbb{P}^N$ be a morphism of degree $d \geq 2$ defined over a number field K . Then the number of K -rational preperiodic points of ϕ is bounded by a bound depending only on N and the degrees of K/\mathbb{Q} and ϕ .*

This powerful conjecture implies in particular Merel's theorem of uniform boundedness of torsion subgroups of the Mordell–Weil groups of rational points on elliptic curves (see [17, Remark 3.19]), as well as the conjectured uniform boundedness of torsion subgroups of abelian varieties (see [6]).

By Northcott's theorem [15], the set $\text{PrePer}(\phi, K)$ of K -rational preperiodic points of a morphism $\phi : \mathbb{P}^N \rightarrow \mathbb{P}^N$ defined over a number field K , is finite. It can therefore be given a finite directed graph structure, called the *preperiodicity graph* of ϕ , by drawing an arrow from P to $\phi(P)$ for each $P \in \text{PrePer}(\phi, K)$. The conjecture of Morton and Silverman implies in particular that the number of possible preperiodicity graphs for such ϕ is finite and depends only on the constants $\deg(\phi)$, $[K : \mathbb{Q}]$ and N . It is therefore natural to ask for the list of all possible preperiodicity graphs for a given family of endomorphisms of \mathbb{P}^N .

The main example of a classification of preperiodicity graphs for rational functions is the classification of preperiodicity graphs of quadratic polynomials over \mathbb{Q} by Poonen [16]. It relies on the following conjecture.

Conjecture (Flynn, Poonen and Schaefer [7]). *Let ϕ be a quadratic polynomial defined over \mathbb{Q} . Then ϕ has no \mathbb{Q} -rational periodic points of period greater than 3.*

Evidence for this conjecture can be found in several articles, including [7, 8, 14, 18].

Theorem (Poonen [16]). *Assuming the Flynn, Poonen and Schaefer conjecture, there exist exactly 12 possible preperiodicity graphs for quadratic polynomials defined over \mathbb{Q} , and a quadratic polynomial has at most nine \mathbb{Q} -rational preperiodic points.*

In the article [3], J.K. Cani and this author proposed a generalization of Poonen's classification of preperiodicity graphs of quadratic polynomials to the classification of preperiodicity graphs of quadratic rational functions with a rational periodic critical point. Recall that a point $P \in \mathbb{P}^1$ is called *critical* if $\phi'(P) = 0$ (if P or $\phi(P)$ are ∞ then we conjugate ϕ by an automorphism of \mathbb{P}^1 , in order to move P and $\phi(P)$ away from ∞). This is indeed a generalization of quadratic polynomials since the latter are exactly quadratic rational functions with a rational *fixed* critical point (up to conjugation by an automorphism of \mathbb{P}^1). To give a broader perspective, Milnor

defined in [11] families of algebraic curves $\text{Per}_n(\mu)$ whose points correspond to rational maps of degree 2 with a periodic point of period n , and multiplier μ . Thus, quadratic rational functions with a periodic critical point of period n correspond to the points of the curve $\text{Per}_n(0)$.

In [3], Canci and the author provided a full classification of preperiodicity graphs of quadratic rational functions defined over \mathbb{Q} with a \mathbb{Q} -rational periodic critical point of period 2, up to the following conjecture.

Conjecture 1 (Canci and Vishkautsan [3]). *Let ϕ be a quadratic rational function defined over \mathbb{Q} having a \mathbb{Q} -rational periodic critical point of period 2. Then ϕ has no \mathbb{Q} -rational periodic points of period greater than 2.*

Theorem (Canci and Vishkautsan [3]). *Assuming the conjecture above, there are exactly 13 possible preperiodicity graphs for quadratic rational functions defined over \mathbb{Q} with a \mathbb{Q} -rational periodic critical point of period 2. Moreover, the number of \mathbb{Q} -rational preperiodic points of such maps is at most 9.*

In order to obtain a similar classification for quadratic rational functions with a \mathbb{Q} -rational periodic critical point of period 3, we need to assume a similar conjecture to the conjecture of Flynn, Poonen and Schaefer and Conjecture 1. First, let us define a *critical n -cycle* to be the set of iterates of a periodic critical point of period n .

Conjecture 2. *Let ϕ be a quadratic rational function defined over \mathbb{Q} with a \mathbb{Q} -rational critical 3-cycle. Then ϕ has no \mathbb{Q} -rational periodic points of period greater than 2 lying outside of the critical cycle.*

As (slight) support for this conjecture, we show in Section 4.3 that a rational function defined over \mathbb{Q} with a \mathbb{Q} -rational critical 3-cycle cannot have a \mathbb{Q} -rational periodic point of period 3 or 4 lying outside of the critical 3-cycle. For the classification of quadratic rational functions with a critical 3-cycle, we also need to assume the following conjecture.

Conjecture 3. *The genus 6 affine plane curve defined by the affine equation*

$$(1.1) \quad u^5v^2 + 2u^4v^3 - u^4v^2 - u^4v + u^3v^4 - 4u^3v^2 + u^3 \\ + u^2v^4 - 4u^2v^3 + 3u^2v - 2uv^3 + 4uv^2 - u + v^2 - v = 0$$

has Jacobian of Mordell–Weil rank exactly 2, and the \mathbb{Q} -rational points on this curve are exactly $(-1, 0)$, $(1, 1)$, $(-1, 1)$, $(0, 0)$, $(0, 1)$, and $(1, 0)$.

In the appendix to this article M. Stoll proves Conjecture 3, conditional on standard conjectures including the BSD conjecture, by enumerating all rational points on the curve defined by (1.1) using the Chabauty–Coleman method.

Theorem 1. *Assuming Conjectures 2 and 3, there are exactly six possible preperiodicity graphs for quadratic rational functions defined over \mathbb{Q} with a \mathbb{Q} -rational critical 3-cycle. Moreover, the number of preperiodic points of such maps is at most eleven.*

The six possible preperiodicity graphs are listed below in Table 1.1, and for each graph we provide an example of a rational function which *realizes* this graph (i.e., whose preperiodicity graph is isomorphic to the given graph). We say that a graph is *realizable* if there exists a rational function that realizes it.

TABLE 1.1. Realizable quadratic rational functions with a \mathbb{Q} -rat. periodic critical point of period 3

ID	$\phi(z)$	Preperiodicity graph	genus
R3P0	$\frac{1}{(z-1)^2}$		N/A
R3P1	$\frac{2z^2 - z - 1}{2z^2}$		0
R3P2	$\frac{z^2 + 5z - 6}{z^2}$		0
R3P3	$\frac{5z^2 - 7z + 2}{5z^2}$		0
R3P4	$\frac{3z^2 - 5z + 2}{3z^2}$		0
R3P5	$\frac{5z^2 - 11z + 6}{5z^2}$		1

The method by which we prove Theorem 1 is by showing (Section 4.2) that the graphs in Tables 1.2, 1.3 and 1.4 below are *inadmissible*, i.e., that there exist no quadratic rational functions defined over \mathbb{Q} with preperiodicity graphs containing isomorphic subgraphs. We prove in Section 3 that this is sufficient to prove Theorem 1. The way we prove inadmissibility of a given graph is by constructing a *dynamical modular curve* whose

TABLE 1.2. Inadmissible graphs for quadratic rational functions with a critical 3-cycle with a modular curve of genus 1

ID	Preperiodicity graph	genus
N3E1		1
N3E2		1
N3E3		1

TABLE 1.3. Inadmissible graphs for quadratic rational functions with a critical 3-cycle with a modular curve of genus 2

ID	Preperiodicity graph	genus
N3M1		2
N3M2		2
N3M3		2

TABLE 1.4. Inadmissible graphs for quadratic rational functions with a critical 3-cycle with a modular curve of genus > 2

ID	Preperiodicity graph	genus
N3H1		3
N3H2		6
N3H3		5

\mathbb{Q} -rational points correspond to conjugacy classes (under conjugation by automorphisms of \mathbb{P}^1) of rational functions defined over \mathbb{Q} with a rational critical 3-cycle that *admit* the graph (i.e., whose preperiodicity graph contains an isomorphic copy of the graph). We then prove that the dynamical modular curve has no \mathbb{Q} -rational points.

It is worth mentioning that other generalizations of Poonen's theorem and classifications of preperiodicity graphs have appeared in the literature: the classification of preperiodicity graphs of quadratic polynomials over quadratic fields by Doyle, Faber and Krumm [5]; the classification (over \mathbb{Q}) of preperiodicity graphs of rational maps with an automorphism by Manes [10]; and the classification of all post-critically finite maps defined over \mathbb{Q} by Lukas, Manes and Yap [9]. The latter classification is used in fact in the proof of Theorem 1.

We split the nine inadmissible graphs needed for the proof of Theorem 1 into three groups, according to the genus of the corresponding dynamical modular curve.

Acknowledgments. The author deeply thanks Michael Stoll and Maarten Derickx, without whose help and advice this research would not have been completed. The author would also like to thank Benjamin Collas and Konstantin Jakob for many fruitful discussions during the preparation of this article. Finally the author would like to thank Fabrizio Catanese for his support and encouragement.

2. Preliminaries

2.1. Dynatomic polynomials. Let $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a rational function. We write

$$\phi = [F(X, Y), G(X, Y)]$$

using homogeneous polynomials F and G . Let

$$\phi^n(X, Y) = [F_n(X, Y), G_n(X, Y)]$$

be the n -th iterate of ϕ for $n \geq 1$, and define

$$\Phi_{\phi,n}(X, Y) = YF_n(X, Y) - XG_n(X, Y).$$

We then define the n -th *dynatomic polynomial* by

$$\Phi_{\phi,n}^*(X, Y) = \prod_{k|n} (\Phi_{k,\phi}(X, Y))^{\mu(n/k)},$$

where μ is the Moebius mu function (for the proof that $\Phi_{\phi,n}^*$ are actually polynomials, see [13, Theorem 2.1]).

It is easy to see that if P is a periodic point of period n then P is a root of $\Phi_{\phi,n}^*$. The converse is not true however, and a root of $\Phi_{\phi,n}^*$ can correspond to a periodic point of period strictly dividing n . For any $n \geq 1$, we call the roots of $\Phi_{\phi,n}^*$ periodic points of *formal period* n for ϕ .

In the article we identify (by abuse of notation) the dynatomic polynomials $\Phi_{\phi,n}^*(X, Y)$ with their dehomogenization $\Phi_{\phi,n}^*(z)$, where $z = \frac{X}{Y}$. By dehomogenizing, we ignore the possibility of ∞ being a root of the dynatomic polynomials. This will not be a problem in what follows, however.

2.2. Linear equivalence of rational functions. We say that two rational functions $\phi_1, \phi_2 : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ are *linearly equivalent* if there exists an $f \in \text{PGL}_2$ acting as a projective automorphism of \mathbb{P}^1 such that $\phi_2 = \phi_1^f = f^{-1}\phi_1f$.

Let ϕ_1, ϕ_2 be linearly equivalent. A point P is periodic of period n for ϕ_1 if and only if $f^{-1}(P)$ is periodic of period n for ϕ_2 (similarly for preperiodic points). When ϕ_1, ϕ_2 and f are all defined over the same base field K , then it is clear that the preperiodicity graphs of ϕ_1 and ϕ_2 are isomorphic.

Therefore when classifying realizable graphs of rational functions defined over \mathbb{Q} we are actually interested in the \mathbb{Q} -rational conjugacy classes of quadratic rational functions rather than in the individual maps.

2.3. Post-critically finite quadratic rational functions. A rational function $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is called *post-critically finite* (or PCF for short) if all of its critical points are preperiodic. A rational function has exactly $2d - 2$ critical points (counted with multiplicity; see [17], Section 1.2), and a quadratic rational function has exactly 2 *distinct* critical points.

Lukas, Manes and Yap [9] provided a complete classification of all quadratic post-critically rational functions over \mathbb{Q} and their $\overline{\mathbb{Q}}$ -conjugacy classes, as well as all possible preperiodicity graphs of these functions. In particular, they showed that a quadratic rational function defined over \mathbb{Q} with a \mathbb{Q} -rational critical 3-cycle is PCF if and only if both of its critical points lie in the same 3-cycle, and there is a unique conjugacy class of quadratic rational functions (over \mathbb{Q}) with this property, realizing the graph R3P0 in Table 1.1.

3. Sufficiency of the nine inadmissibility graphs

Proposition 1. *Assume Conjecture 2. Any quadratic rational function with a rational critical 3-cycle that does not realize one of the graphs in Table 1.1, must admit one of the graphs in Tables 1.2, 1.3 or 1.4.*

Proof. We can arrange the graphs from Tables 1.1 through 1.4 in a Hasse diagram with respect to the partial order of subgraph isomorphism:




FIGURE 3.1. Hasse diagram of inadmissible graphs

Due to the results of [9] mentioned in Section 2.3, we can restrict ourselves to non-PCF (see Section 2.3) quadratic rational functions with a rational critical 3-cycle. Lemma 3.2 in [3] implies that starting from the critical cycle graph R3P1, one can obtain any realizable graph of a non-PCF rational function defined with a rational critical 3-cycle, by recursively

adding vertices and arrows to the graph using either of the following two recursion steps:

- (1) *Adding a periodic cycle.* We add a new cycle C to the graph. For each vertex P in C we add a vertex Q not in the cycle and an arrow $Q \rightarrow P$.
- (2) *Adding preimages to a non-periodic point.* We add two vertices Q_1, Q_2 and arrows $Q_1 \rightarrow P, Q_2 \rightarrow P$, where P is a vertex not lying in any cycle in the graph.

Now it is easy to check that from any graph in the Hasse diagram, by using one of the recursive steps we either generate a graph that is already in the Hasse diagram, or contains (a subgraph isomorphic to) one of the nine graphs in Tables 1.2, 1.3 or 1.4. \square

4. Constructing dynamical modular curves from the graphs

4.1. Realizable graphs with a critical 3-cycle.

4.1.1. *R3P0.* Note that graph R3P0 implies that we have two points in the 3-cycle which are critical. Since any quadratic rational function has exactly two distinct critical points, this means any function admitting the graph is *post-critically finite* (or PCF for short), i.e., all its critical points are preperiodic.

As mentioned in Section 2.3, Lukas, Manes and Yap proved in [9] that there is only one possible preperiodicity graph for a PCF quadratic rational function with a critical three cycle, which is graph R3P0 in Table 1. Moreover, there is only one \mathbb{Q} -conjugacy class of quadratic rational functions that has this preperiodicity graph, with the following representative.


$$(4.1) \quad \phi_0(z) = \frac{1}{(z-1)^2}.$$

4.1.2. *R3P1.* We can parametrize the non-PCF \mathbb{Q} -conjugacy classes of quadratic rational functions with a rational critical 3-cycle using the following map.

$$(4.2) \quad a \mapsto \phi_a(z) = \frac{(a+1)z^2 - az - 1}{(a+1)z^2}, \quad a \neq 0, -1, -2.$$

For the special values $a = 0, -2$ we get representatives of the conjugacy class realizing the graph R3P0.

In fact, given any rational function ϕ defined over \mathbb{Q} with a \mathbb{Q} -rational critical 3-cycle, we can conjugate ϕ by an element of $\mathrm{PGL}_2(\mathbb{Q})$ to move the periodic critical point P to 0, its image to ∞ and $\phi^2(P)$ to 1. One can then check that we obtain a function ϕ_a for some $a \in \mathbb{Q}$.

FIGURE 4.1. Preperiodicity graph of ϕ_a

The preperiodicity graph of ϕ_a where $a \neq 0, -1, -2$ will contain the graph in Fig. 4.1; it is easy to check that the non-periodic preimages of 0 and 1 are $-\frac{1}{a+1}$ and $-\frac{1}{a}$, respectively.

4.1.3. R3P2. We take the parametrization ϕ_a of R3P1 and consider a root b of the first dynatomic polynomial of ϕ_a . Such a root must correspond to a fixed point of ϕ_a .

$$(4.3) \quad \Phi_{a,1}^*(b) := \Phi_{\phi_a,1}^*(b) = (1+a)b^3 + (-1-a)b^2 + ab + 1 = 0.$$

We solve this equation for a .

$$(4.4) \quad a = -\frac{b^3 - b^2 + 1}{b(b^2 - b + 1)}.$$

We then substitute this into the expression for ϕ_a and get the following parametrization of quadratic rational functions realizing R3P2.


$$(4.5) \quad b \mapsto \phi_b(z) = \frac{(b-1)z^2 + (b^3 - b^2 + 1)z - b^3 + b^2 - b}{(b-1)z^2},$$

where b cannot obtain the values 0, 1 in \mathbb{Q} . We remark that ϕ_b degenerates also when b is a root of $b^2 - b + 1 = 0$, but for this case $b \notin \mathbb{Q}$. Moreover, the solutions to the equations

$$b^3 - b^2 + 1 = 0 \text{ and } b^3 - b^2 + 2b - 1 = 0$$

determine $a = 0$ and $a = 2$, respectively, but these do not have \mathbb{Q} -rational solutions either.

Each map ϕ_b admits the graph in Fig. 4.2.

FIGURE 4.2. Preperiodicity graph of ϕ_b

It is easy to check that

$$-\frac{b^3 - b^2 + b}{b - 1}, \quad \frac{b^3 - b^2 + b}{b^3 - b^2 + 1} \text{ and } \frac{b^2 - b + 1}{(b - 1)^2}$$

are the non-periodic preimages of 0, 1 and b , respectively.

4.1.4. R3P3. We start again with ϕ_a parametrizing R3P1 (see Section 4.1.2). Recall that $-1/(a+1)$ is the non-periodic preimage of 0. Let c be a preimage of $-1/(a+1)$, i.e., $\phi(c) = -\frac{1}{a+1}$. This implies the following equation.

$$(4.6) \quad (a+2)c^2 - ac - 1 = 0.$$

We solve this equation for a .

$$(4.7) \quad a = -\frac{2c^2 - 1}{c^2 - c}.$$

By substituting this expression for a , we get the following parametrization.

$$(4.8) \quad c \mapsto \phi_c(z) = \frac{(c^2 + c - 1)z^2 - (2c^2 - 1)z + c^2 - c}{(c^2 + c - 1)z^2},$$

where c cannot obtain the values 0, 1 and $\frac{1}{2}$ (at the latter value we get $a = -2$). Moreover, c cannot be a solution to the equation

$$c^2 + c - 1 = 0,$$

as ϕ_c degenerates for these values of c ; this equation however has no rational solutions. Similarly, for $c^2 = -\frac{1}{2}$ we get $a = 0$, but again there are no rational c satisfying this condition.




FIGURE 4.3. Preperiodicity graph of ϕ_c

Each map ϕ_c admits the graph in Fig. 4.3. It is easy to check that $\frac{c-1}{2c-1}$ is the second preimage of $-\frac{1}{a+1} = \frac{c^2-c}{c^2+c-1}$, other than c .

4.1.5. R3P4. We start with ϕ_a realizing R3P1 and find the expression for the second dynatomic polynomial of ϕ_a . A root of this polynomial is a point of formal period 2 for ϕ_a .

$$(4.9) \quad \Phi_{a,2}^*(z) := \Phi_{\phi_a,2}^*(z) = (a+1)((a+1)z^2 + (1-a)z - 1)$$

A formal point of period 2 for ϕ_a is not of period 2 only if it is of period 1 and so a root of the first dynatomic polynomial $\Phi_{a,1}^*(z)$. One can find such

points by computing the roots of the resultant of the first two dynatomic polynomials.

$$\text{Res}(\Phi_{a,1}^*(z), \Phi_{a,2}^*(z)) = -(a+1)^4(a^2+2a+5)$$

We see that there are no \mathbb{Q} -rational values of a which produce points of formal period 2 but of actual period 1.

We see that any \mathbb{Q} -rational periodic point d is of period 2 for ϕ_a if and only if it is a solution of the following equation.

$$(4.10) \quad (a+1)d^2 + (1-a)d - 1 = 0.$$

We solve this equation for a .

$$(4.11) \quad a = -\frac{d^2 + d - 1}{d^2 - d}.$$

We get the following parametrization for quadratic rational functions admitting the graph R3P4.

$$(4.12) \quad d \mapsto \phi_d(z) = \frac{(2d-1)z^2 - (d^2+d-1)z + d^2-d}{(2d-1)z^2},$$

where d cannot obtain the values 0, 1 and $\frac{1}{2}$. When d is a solution to the equations

$$d^2 + d - 1 = 0 \quad \text{and} \quad d^2 - 3d + 1 = 0$$

we get $a = 0$ and $a = -2$ respectively; these equations have no rational solutions, however.

One can check that the non-periodic preimage of d is $-\frac{d}{d-1}$ and its periodic image is $\frac{d-1}{2d-1}$. The non-periodic preimage of $\frac{d-1}{2d-1}$ is $\frac{d-1}{d}$. The non-periodic preimage of 0 is $\frac{d^2-d}{2d-1}$ and the non-periodic preimage of 1 is $\frac{d^2-d}{d^2+d-1}$. Thus the rational function ϕ_d admits the graph in Fig. 4.4.




FIGURE 4.4. Preperiodicity graph of ϕ_c

4.1.6. R3P5. We start with ϕ_d realizing R3P4. Recall that d is a periodic point of period 2 and the value of its non-periodic preimage is $-\frac{d}{d-1}$.

Let w be a preimage of $-\frac{d}{d-1}$. Then $\phi(w) = -\frac{d}{d-1}$. This gives us the following equation.

$$(4.13) \quad (4d^2 - 4d + 1)w^2 - (d^3 - 2d + 1)w + d^3 - 2d^2 + d = 0.$$

Four of the solutions to this equation correspond to degenerate maps ϕ_d . These four points are

$$(d, w) = (0, 0), (0, 1), (1, 0), \left(\frac{1}{2}, 1\right).$$

The discriminant of (4.13) with respect to w is:

$$\begin{aligned} \Delta &= d^6 - 16d^5 + 44d^4 - 50d^3 + 28d^2 - 8d + 1 \\ &= (d-1)^2(d^4 - 14d^3 + 15d^2 - 6d + 1) \end{aligned}$$

Using the discriminant, we find that the affine plane curve defined by (4.13) is birational to the affine plane curve defined by

$$(4.14) \quad y^2 = x^4 - 14x^3 + 15x^2 - 6x + 1,$$

where

$$(4.15) \quad x = d, \quad y = \frac{2(4d^2 - 4d + 1)w - (d^3 - 2d + 1)}{d - 1}$$

or

$$(4.16) \quad w = \frac{(d-1)y + (d^3 - 2d + 1)}{2(4d^2 - 4d + 1)}.$$

We can simplify the curve defined by (4.14) further to get the elliptic curve

$$(4.17) \quad E : y^2 + xy + y = x^3 - x^2.$$

This elliptic curve is curve 53a1 in the Cremona database (see [4]), it has rank 1 with trivial torsion. We thus get infinitely many rational points on the curve, and thus infinitely many non-linearly equivalent rational functions defined over \mathbb{Q} admitting graph R3P5 (only finitely many \mathbb{Q} -rational points on E will be mapped to degenerate solutions of (4.13)).

4.2. Nonrealizable graphs with a critical 3-cycle.

4.2.1. N3E1. We start with the graph R3P1 by taking the parametrization ϕ_a we defined in Section 4.1.2. We recall that the general form of a quadratic rational function admitting R3P1 is

$$(4.18) \quad \phi_a(z) = \frac{(a+1)z^2 - az - 1}{(a+1)z^2}, \quad a \neq 0, -1, -2.$$

We will show that the first dynatomic polynomial of ϕ_a cannot split over \mathbb{Q} . This immediately implies that a quadratic rational function with a rational critical three-cycle can have at most one \mathbb{Q} -rational fixed point.

The first dynatomic polynomial of ϕ_a is

$$(4.19) \quad \Phi_{a,1}^*(z) := \Phi_{\phi_a,1}^*(z) = (-a-1)z^3 + (a+1)z^2 - az - 1.$$

In R3P2 we determined that a root b of this polynomial must satisfy

$$(4.20) \quad a = -\frac{b^3 - b^2 + 1}{b(b^2 - b + 1)}.$$

A necessary and sufficient condition for the cubic polynomial $\Phi_{a,1}^*$ to split over \mathbb{Q} is for it to have a \mathbb{Q} -rational root and for the discriminant to be a square in \mathbb{Q} . The latter condition is defined by the following equation.

$$(4.21) \quad u^2 = \Delta(\Phi_{a,1}^*) = -3a^4 - 16a^3 - 50a^2 - 60a - 23.$$

Substituting (4.20) into (4.21) we obtain

$$(4.22) \quad v^2 = (b-1)(b^3 + b^2 - b + 3),$$

where

$$v = \frac{b^2(b^2 - b + 1)^2 u}{2b^3 - 4b^2 + 2b - 1}.$$

The affine plane curve C defined by (4.22) is of genus 1. We denote by \overline{C} its projective closure in $\mathbb{P}_{[X:Y:Z]}^2$ where $b = X/Z$ and $v = Y/Z$. The curve \overline{C} contains (at least) the following two \mathbb{Q} -rational points:

$$[0 : 1 : 0], [1 : 0 : 1].$$

It is birational to the elliptic curve with Cremona reference 19a3 with minimal model

$$E : y^2 + y = x^3 + x^2 + x.$$

Remark 1. It is interesting to note that the elliptic curve with Cremona reference 19a3 appears several times in the classification of preperiodicity graphs of rational functions with a rational critical cycle. For example, in [3] it is proven that it is birational to two modular curves of quite distinct graphs with critical 2-cycles (denoted there by N2E2 and N2E3), and it appears again below in Section 4.3.2. Similarly, the elliptic curve with Cremona reference 17a4 appears both for N3E3 below (see Section 4.2.3

below) and for the proof of Proposition 4.3 in [3]. The elliptic curve with Cremona reference 11a3 appears for N3E2 below (see Section 4.2.2 below) and for the graph N2E1 in [3].

The birational map $\psi : \overline{C} \rightarrow E$ is given by

$$\begin{aligned} \psi([X : Y : Z]) = [2Z^2 - 2XZ, \\ X^2 - 2XZ + YZ + Z^2, \\ -2X^2 + 4XZ - 2Z^2]. \end{aligned}$$

The rational points in the locus of indeterminacy of ψ are exactly the two points we have already discovered on \overline{C} . The elliptic curve E has Mordell–Weil group of rank 0 and torsion subgroup isomorphic to $\mathbb{Z}/3\mathbb{Z}$. It therefore contains only three rational points, which are

$$[0 : 0 : 1], [0 : 1 : 0], [0 : -1 : 1].$$

The preimages of these points under the birational map ψ , together with ψ 's locus of indeterminacy give us the full set of points on \overline{C} , and it turns out that the two points we found on \overline{C} are the only rational points on it. Neither of these points corresponds to a non-degenerate quadratic rational function ϕ_b .

As an aside, we prove the following.

Proposition 2. *There exists a unique conjugacy class of quadratic rational functions with a critical 3-cycle whose first dynatomic polynomial has a square \mathbb{Q} -rational discriminant.*

Proof. The affine plane curve C_1 described by (4.21) has genus 1, and it is birational to the elliptic curve with Cremona reference 19a1 with the following minimal model.

$$(4.23) \quad E_1 : y^2 + y = x^3 + x^2 - 9x - 15.$$

We consider the projective closure \overline{C}_1 of C_1 in $\mathbb{P}_{[X:Y:Z]}^2$ where $a = \frac{X}{Z}$ and $u = \frac{Y}{Z}$. The birational map $\psi : \overline{C}_1 \rightarrow E_1$ is given by

$$\begin{aligned} \psi = [1491X^3 + 4176X^2Z - 133XYZ + 3821XZ^2 - 114YZ^2 + 1146Z^3, \\ 533X^3 + 2839X^2Z + 171XYZ + 3891XZ^2 + 95YZ^2 + 1565Z^3, \\ -686X^3 - 1764X^2Z - 1512XZ^2 - 432Z^3]. \end{aligned}$$

The elliptic curve E_1 has Mordell–Weil rank 0, and torsion subgroup isomorphic to $\mathbb{Z}/3\mathbb{Z}$. The three rational points on E are

$$(4.24) \quad [0 : 1 : 0], [5 : 9 : 1], [5 : -10 : 1].$$

The only rational preimages of these points under ψ , together with ψ 's points of indeterminacy are the four points

$$(4.25) \quad [-6/7 : -19/49 : 1], [-6/7 : 19/49 : 1], [0 : 1 : 0], [-1 : 0 : 1].$$

These are therefore the only rational points on the curve \overline{C}_1 . The third point is at infinity and is therefore a “*cusp*” (a point added to the dynamical modular curve to obtain a projective curve), while the fourth point corresponds to the case $a = -1$ for which ϕ_a is a degenerate map.

It remains to check the case $a = -\frac{6}{7}$. For this map we get the following first dynatomic polynomial

$$(4.26) \quad \Phi_{-\frac{6}{7},1}^*(z) = -\frac{1}{7}(z^3 - z^2 - 6z + 7)$$

whose discriminant $\frac{361}{2401} = (\frac{19}{49})^2$ is a square in \mathbb{Q} as expected. However, this polynomial is irreducible and therefore $\phi_{-\frac{6}{7}}$ has no \mathbb{Q} -rational fixed points. \square

4.2.2. N3E2. We start with the parametrization ϕ_a of R3P1. Recall that the non-periodic preimage of 1 is $-\frac{1}{a}$. Assume that $-\frac{1}{a}$ has a preimage t , i.e., $\phi(t) = -\frac{1}{a}$. This implies the following equation.

$$(4.27) \quad a^2t^2 - a^2t + 2at^2 + t^2 - a = 0.$$

The curve C_2 described by this equation is of genus 1. We take the projective closure \overline{C}_2 of C_2 in $\mathbb{P}_{[X:Y:Z]}^2$ where $a = \frac{X}{Z}$ and $t = \frac{Y}{Z}$. \overline{C}_2 contains (at least) the following 4 rational points:

$$(4.28) \quad [0 : 1 : 0], [0 : 0 : 1], [1 : 0 : 0], [-1 : 1 : 1].$$

The affine plane curve \overline{C}_2 is birational to the elliptic curve E with reference 11a3 in the Cremona database. Its minimal model is given by

$$(4.29) \quad E : y^2 + y = x^3 - x^2.$$

This elliptic curve has Mordell–Weil rank 0 and a torsion subgroup isomorphic to $\mathbb{Z}/5\mathbb{Z}$ and 5 rational points:

$$(4.30) \quad [0 : 1 : 0], [0 : 0 : 1], [0 : -1 : 1], [1 : 0 : 1], [1 : -1 : 1].$$

The birational map $\psi : \overline{C}_2 \rightarrow E$ is defined by

$$\begin{aligned} \psi = [& XY^3 - XY^2Z + Y^3Z - YZ^3, \\ & -XY^2Z + Y^3Z + XYZ^2 - 2Y^2Z^2 + Z^4, \\ & -Y^3Z]. \end{aligned}$$

The preimages of the rational points under ψ together with ψ 's points of indeterminacy ($[1 : 0 : 0], [0 : 1 : 0]$) are the four points of \overline{C}_2 we have already discovered. Therefore these are all the rational points on \overline{C}_2 .

None of these points corresponds to a non-degenerate map ϕ_a , therefore graph N3E2 is non-realizable.

4.2.3. N3E3. We start the quadratic rational function ϕ_c parametrizing R3P3 (note that we could also have started with R3P4).

$$(4.31) \quad \phi_c(z) = \frac{(c^2 + c - 1)z^2 - (2c^2 - 1)z + c^2 - c}{(c^2 + c - 1)z^2}, \quad c \neq 0, 1, \frac{1}{2}.$$

We consider the second dynatomic polynomial for ϕ_c .

$$\begin{aligned} \Phi_{c,2}^*(z) := \Phi_{\phi_c,2}^*(z) &= (c^4 + 2c^3 - c^2 - 2c + 1)z^2 \\ &\quad + (-3c^4 - 2c^3 + 5c^2 - 1)z + c^4 - 2c^2 + c \end{aligned}$$

This polynomial is divisible by $c^2 + c - 1$; since the map ϕ_c degenerates for the two values of c which are its roots, we can divide by it to get

$$(4.32) \quad c^2 z^2 - 3c^2 z + c^2 + cz^2 + cz - c - z^2 + z = 0.$$

This equation describes a genus 1 affine plane curve which we will denote by C_3 , and its projective closure in $\mathbb{P}_{[X:Y:Z]}^2$ by \overline{C}_3 , where $c = \frac{X}{Z}$ and $z = \frac{Y}{Z}$. The curve \overline{C}_3 contains (at least) the following 6 points:

$$(4.33) \quad [1 : 1 : 1], [0 : 1 : 0], [0 : 0 : 1], [0 : 1 : 1], [1 : 0 : 0], [1 : 0 : 1].$$

This curve is birational to the elliptic curve

$$(4.34) \quad E : y^2 + xy + y = x^3 - x^2 - x$$

with Cremona reference 17a4. This curve has Mordell–Weil rank 0 and torsion subgroup isomorphic to $\mathbb{Z}/4\mathbb{Z}$, and thus contains exactly four \mathbb{Q} -rational points which are

$$[0 : 0 : 1], [0 : 1 : 0], [0 : -1 : 1], [1 : -1 : 1].$$

The map ψ from \overline{C}_3 to E is defined by

$$\begin{aligned} \psi &= [XY^3 - 4XY^2Z + 4XYZ^2 + Y^2Z^2 - XZ^3 - 2YZ^3 + Z^4, \\ &\quad XY^2Z - 3XYZ^2 + XZ^3 + YZ^3 - Z^4, \\ &\quad -Y^3Z + 3Y^2Z^2 - 3YZ^3 + Z^4]. \end{aligned}$$

The points of indeterminacy of ψ are $[0 : 1 : 1], [0 : 1 : 0], [1 : 0 : 0]$.

The preimages of the four rational points of E under ψ together with ψ 's points of indeterminacy give us all the rational points of \overline{C}_3 , and these are exactly the six points that we have already discovered, none of which correspond to non-degenerate quadratic rational functions.

4.2.4. *N3M1.* We start with the parametrization ϕ_c of the graph R3P3. Recall that c is a preperiodic point whose image under ϕ_c is the non-periodic preimage of 0. Let w be a preimage of c under ϕ_c , i.e., $\phi(w) = c$. This implies

$$(4.35) \quad -c^3w^2 - 2c^2w + 2cw^2 + c^2 - w^2 - c + w = 0$$

This equation describes an affine plane curve that we denote by C_4 , and we denote its projective closure in $\mathbb{P}_{[X:Y:Z]}^2$ by \overline{C}_4 , where $c = \frac{X}{Z}$ and $w = \frac{Y}{Z}$. This curve is of genus 2 and contains (at least) the following five points.

$$(4.36) \quad [0 : 1 : 0], [0 : 0 : 1], [0 : 1 : 1], [1 : 0 : 0], [1 : 0 : 1].$$

The curve \overline{C}_4 is birational to the hyperelliptic curve given by

$$(4.37) \quad H : y^2 = x^6 - 2x^5 + 3x^4 - 4x^3 - x^2 + 2x + 1.$$

The birational map $\psi : \overline{C}_4 \rightarrow H$ is defined by

$$\psi = [-XZ + Z^2, 2X^3YZ^2 + 2X^2Z^4 - 4XYZ^4 + 2YZ^5 - Z^6, -XZ].$$

The latter curve H contains the following points (*in weighted homogeneous coordinates*):

$$(4.38) \quad [1 : -1 : 0], [1 : 1 : 0], [0 : -1 : 1], [0 : 1 : 1], [1 : 0 : 1].$$

The hyperelliptic curve has Jacobian with torsion subgroup isomorphic to $\mathbb{Z}/15\mathbb{Z}$, and a rank bound computation in MAGMA [1] tells us the Jacobian has Mordell–Weil rank 0.

The curve H has good reduction at 3, and therefore the Mordell–Weil group of the Jacobian $J(\mathbb{Q})$ of H injects into $J(\mathbb{F}_3)$. Therefore all the maps in the following commutative diagram are injective.

$$\begin{array}{ccc} H(\mathbb{Q}) & \longrightarrow & J(\mathbb{Q}) \\ \downarrow & & \downarrow \\ H(\mathbb{F}_3) & \longrightarrow & J(\mathbb{F}_3) \end{array}$$

However, one can easily check that the only rational points in $H(\mathbb{F}_3)$ are the reductions (mod 3) of the five points in $H(\mathbb{Q})$ we have already discovered. Therefore these five points are the only \mathbb{Q} -rational points on H .

Using ψ we pull back these five points to C_4 , and together with the indeterminacy points of ψ (these are $[0 : 1 : 0], [1 : 0 : 0]$) we find that the only rational points on C are the five points we found before. None of these points correspond to non-degenerate quadratic rational functions.

4.2.5. N3M2. We start with the parametrization ϕ_c of the graph R3P3, and consider a root w of its first dynatomic polynomial. Such a root must correspond to a fixed point of ϕ_c .

$$\begin{aligned} \Phi_{c,1}^*(w) := \Phi_{\phi_c,1}^*(w) &= (-c^2 - c + 1)w^3 + (c^2 + c - 1)w^2 \\ &\quad + (-2c^2 + 1)w + c^2 - c = 0. \end{aligned}$$

We denote by C_5 the affine plane curve defined by this equation, and by \overline{C}_5 its projective closure in $\mathbb{P}_{[X:Y:Z]}^2$, where $c = \frac{X}{Z}$ and $w = \frac{Y}{Z}$. This is a curve of genus 2 and contains (at least) the following four rational points

$$(4.39) \quad [0 : 1 : 0], [0 : 0 : 1], [1 : 0 : 0], [1 : 0 : 1].$$

This curve is birational to the following hyperelliptic curve

$$(4.40) \quad H_5 : y^2 = x^6 - 2x^5 + 5x^4 - 6x^3 + 10x^2 - 8x + 5.$$

The birational map $\psi : \overline{C}_5 \rightarrow H_5$ is defined by

$$\begin{aligned} \psi &= [-YZ + Z^2, \\ &\quad 2XY^3Z^2 - 2XY^2Z^3 + Y^3Z^3 + 4XYZ^4 - Y^2Z^4 - 2XZ^5 + Z^6, \\ &\quad -YZ]. \end{aligned}$$

One can use MAGMA to check that the Jacobian J of H_5 has Mordell–Weil torsion subgroup isomorphic to $\mathbb{Z}/5\mathbb{Z}$ and rank 0. The curve H_5 contains (at least) the following two rational points (in weighted homogeneous coordinates).

$$(4.41) \quad [1 : -1 : 0], [1 : 1 : 0].$$

The Jacobian J contains (at least) the following rational points in Mumford representation:

$$\begin{aligned} (1, 0, 0), (1, x^3 - x^2, 2), (1, -x^3 + x^2, 2), \\ (x^2 - x + 1, x - 1, 2), (x^2 - x + 1, -x + 1, 2). \end{aligned}$$

We denote by $P_+ = [1 : 1 : 0]$ and $P_- = [1 : -1 : 0]$, the two points at infinity. Using this notation we get the following divisor representations on H :

$$\begin{aligned} (1, 0, 0) &\equiv \text{identity}, \\ (1, x^3 - x^2, 2) &\equiv P_+ - P_-, \\ (1, -x^3 + x^2, 2) &\equiv P_- - P_+, \\ (x^2 - x + 1, x - 1, 2) &\equiv [\alpha : -\bar{\alpha} : 1] + [\bar{\alpha} : -\alpha : 1] - P_+ - P_-, \\ (x^2 - x + 1, -x + 1, 2) &\equiv [\alpha : \bar{\alpha} : 1] + [\bar{\alpha} : \alpha : 1] - P_+ - P_-, \end{aligned}$$

where $\alpha = \frac{1}{2} + \frac{\sqrt{-3}}{2}$. The only rational points appearing in these representations are the two we already found, and therefore they are the only rational points on H_5 .

Pulling back the two rational points of H_5 under ψ and adding its indeterminacy points, we find all rational points on C_5 , and these are exactly the four points we found before. None of these points correspond to non-degenerate quadratic rational functions.

4.2.6. *N3M3*. We start with R3P4 and consider a root w of first dynatomic polynomial of ϕ_d (note that we can also start with R3P2 and look at the second dynatomic polynomial of ϕ_b). Such a root corresponds to a fixed point of ϕ_d .

$$\begin{aligned} \Phi_{d,1}^*(w) := \Phi_{d,1}^*(w) &= (-2d + 1)w^3 + (2d - 1)w^2 \\ &\quad + (-d^2 - d + 1)w + d^2 - d = 0 \end{aligned}$$

The affine plane curve C_6 described by this equation is of genus 2. We denote its projective closure in $\mathbb{P}_{[X:Y:Z]}^2$ by \overline{C}_6 , where $d = \frac{X}{Z}$ and $w = \frac{Y}{Z}$. The curve \overline{C}_6 contains (at least) the following five rational points

$$(4.42) \quad [0 : 1 : 0], [0 : 0 : 1], [1 : 2 : 2], [1 : 0 : 0], [1 : 0 : 1],$$

and is birational to the hyperelliptic curve

$$(4.43) \quad H : y^2 = x^6 + 2x^5 + 5x^4 + 8x^3 + 12x^2 + 8x + 4$$

under the map $\psi : \overline{C}_6 \rightarrow H$ defined by

$$\psi = [Z, 2Y^3 + 2XYZ - 2Y^2Z - 2XZ^2 + YZ^2 + Z^3, -Y].$$

Using MAGMA one can compute that the hyperelliptic curve H has Jacobian with Mordell–Weil torsion subgroup isomorphic to $\mathbb{Z}/2\mathbb{Z}$ and rank bounded by 1. The curve H contains (at least) the following six rational points

$$(4.44) \quad [1 : -1 : 0], [1 : 1 : 0], [-1 : -2 : 1], [-1 : 2 : 1], [0 : -2 : 1], [0 : 2 : 1].$$

We denote by $P_+ = [1 : 1 : 0]$ and $P_- = [1 : -1 : 0]$, the two points at infinity. One can check that $P_+ - P_-$ is of infinite order, and therefore the Mordell–Weil rank of the Jacobian is exactly 1. Using the Chabauty–Coleman/ Mordell–Weil sieving algorithm (see Bruin and Stoll [2]) implementation in MAGMA, we determine that the six points we have already found are the only rational points on H . Pulling them back to \overline{C}_6 , we find (together with the indeterminacy point $[1 : 0 : 0]$ of ψ) that the only rational points on \overline{C}_6 are the five points we have already found. None of these correspond to non-degenerate quadratic rational functions.

4.2.7. N3H1. We start with R3P2. Recall that

$$(4.45) \quad \phi_b(z) = \frac{(b-1)z^2 + (b^3 - b^2 + 1)z - b^3 + b^2 - b}{(b-1)z^2}, \quad b \neq 0, 1$$

is a parametrization of R3P2, where b is the fixed point and $\frac{b^2-b+1}{(b-1)^2}$ is its non-periodic preimage. Assume w is a preimage of $\frac{b^2-b+1}{(b-1)^2}$, i.e., $\phi_b(w) = \frac{b^2-b+1}{(b-1)^2}$. This implies

$$(4.46) \quad bw^2 - (b^4 - 2b^3 + b^2 + b - 1)w + b^4 - 2b^3 + 2b^2 - b = 0.$$

We denote by C_7 the affine plane curve defined by this equation, and by \overline{C}_7 its projective closure in $\mathbb{P}_{[X:Y:Z]}^2$, where $b = \frac{X}{Z}$ and $w = \frac{Y}{Z}$. The curve \overline{C}_7 has genus 3, and contains (at least) the following rational points

$$(4.47) \quad [0 : 1 : 0], [0 : 0 : 1], [1 : 0 : 0], [1 : 0 : 1].$$

The curve \overline{C}_7 is birational to the following hyperelliptic curve

$$(4.48) \quad H : y^2 = 4x^7 - 11x^6 + 14x^5 - 7x^4 - 2x^3 + 6x^2 - 4x + 1.$$

The birational map $\psi : \overline{C}_7 \rightarrow H$ is defined by:

$$\psi = [-Z, X^4 - 2X^3Z + X^2Z^2 - 2XYZ^2 + XZ^3 - Z^4, X - Z].$$

The curve H contains (at least) the following five points:

$$(4.49) \quad [1 : 0 : 0], [0 : -1 : 1], [0 : 1 : 1], [1 : -1 : 1], [1 : 1 : 1].$$

Using MAGMA, one can compute that the curve H has Jacobian J with Mordell–Weil rank 0, and torsion subgroup of order bounded by 72, and a two-torsion subgroup isomorphic to $\mathbb{Z}/2\mathbb{Z}$.

Let $P = [0 : -1 : 1], Q = [1 : -1 : 1]$. The divisor class $[P-Q]$ has order 36 in the Jacobian. This means that if the torsion subgroup has order 72 then it is isomorphic to either $\mathbb{Z}/72\mathbb{Z}$ or $\mathbb{Z}/36\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. However, the second option is ruled out by the structure of the two-torsion subgroup. Therefore the only two options for the torsion subgroup are either $\mathbb{Z}/72\mathbb{Z}$ or $\mathbb{Z}/36\mathbb{Z}$.

The image of $[P-Q]$ under the map $J(\mathbb{Q}) \rightarrow J(\mathbb{F}_5) \cong \mathbb{Z}/72\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ is *not* divisible by 2. In fact, under the identification of $J(\mathbb{F}_5)$ with $\mathbb{Z}/72\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, $[P - Q]$ is mapped to the element $(38, 1)$, which is not divisible by 2.

Therefore the Mordell–Weil group is isomorphic to $\mathbb{Z}/36\mathbb{Z}$, and is generated by $[P - Q]$. The curve H has good reduction at 5 so that all the maps in the following commutative diagram are injective.

$$\begin{array}{ccc} H(\mathbb{Q}) & \longrightarrow & J(\mathbb{Q}) \\ \downarrow & & \downarrow \iota \\ H(\mathbb{F}_5) & \longrightarrow & J(\mathbb{F}_5) \end{array}$$

There are six points in $H(\mathbb{F}_5)$ (given in weighted homogeneous coordinates):

$$(4.50) \quad [1 : 0 : 0], [0 : 1 : 1], [0 : 4 : 1], [1 : 1 : 1], [1 : 4 : 1], [2 : 0 : 1].$$

Of these points, only the five points $[1 : 0 : 0], [0 : 1 : 1], [0 : 4 : 1], [1 : 1 : 1], [1 : 4 : 1]$ map to the image of the map ι in the diagram, and these exactly correspond to the five points we know on H . Since the map $H(\mathbb{Q}) \rightarrow H(\mathbb{F}_5)$ is injective, this proves that these are the only rational points on H .

Combining the preimages of ψ together with its locus of indeterminacy, we find all rational points on \overline{C}_7 , and these are exactly the four points we already discovered. None of these four points correspond to ϕ_b admitting the graph N3H1.

4.2.8. N3H2. For the proof that the graph N3H2 is non-admissible (up to standard conjectures) see the appendix by M. Stoll.

4.2.9. N3H3. We start with R3P5. Recall that graph R3P4 had the parametrization

$$(4.51) \quad \phi_d(z) = \frac{(2d-1)z^2 - (d^2 + d - 1)z + d^2 - d}{(2d-1)z^2}, \quad d \neq 0, 1, \frac{1}{2},$$

where d is a periodic point of (formal) period 2, and the graph R3P5 was represented by the following additional condition.

$$(4.52) \quad (4d^2 - 4d + 1)w^2 - (d^3 - 2d + 1)w + d^3 - 2d^2 + d = 0.$$

The point w was such that $\phi^2(w) = d$ and $\phi(w)$ is non-periodic. The second periodic point of period 2 is $\frac{d-1}{2d-1}$ and its non-periodic preimage is $\frac{d-1}{d}$. Let u be a rational preimage of $\frac{d-1}{d}$, i.e., $\phi(u) = \frac{d-1}{d}$. This implies

$$(4.53) \quad (2d-1)u^2 - (d^3 + d^2 - d)u + d^3 - d^2 = 0.$$

Together with (4.52), we get the following affine space curve parametrizing the graph N3H3:

$$(4.54) \quad C_1 : \begin{cases} (4d^2 - 4d + 1)w^2 - (d^3 - 2d + 1)w + d^3 - 2d^2 + d = 0 \\ (2d-1)u^2 - (d^3 + d^2 - d)u + d^3 - d^2 = 0. \end{cases}$$

Using the discriminant (with respect to u) we can bring the second equation to the form

$$(4.55) \quad v^2 = d^4 + 2d^3 - 9d^2 + 10d - 3$$

where

$$(4.56) \quad v = \frac{(4d-2)u - (d^3 + d^2 - d)}{d},$$

and we have already seen in Section 4.1.6 that (4.53) can be transformed in a similar way to

$$(4.57) \quad y^2 = d^4 - 14d^3 + 15d^2 - 6d + 1.$$

Thus the curve C_1 is birational to a curve C_2 in $\mathbb{A}_{(d,y,v)}^3$ defined by

$$(4.58) \quad C_2 : \begin{cases} y^2 = d^4 - 14d^3 + 15d^2 - 6d + 1 \\ v^2 = d^4 + 2d^3 - 9d^2 + 10d - 3. \end{cases}$$

The curve C_2 is of genus 5, and is a double-cover of the following curve:

$$(4.59) \quad H_1 : s^2 = (d^4 - 14d^3 + 15d^2 - 6d + 1)(d^4 + 2d^3 - 9d^2 + 10d - 3).$$

The curve H_1 has genus 3, and its equation can be simplified to the following model:

$$(4.60) \quad H_2 : y^2 = -3x^8 + 4x^7 - 2x^6 - 16x^5 + 11x^4 + 16x^3 - 2x^2 - 4x - 3.$$

This curve has automorphism group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. When quotienting the curve by the automorphism $[Z : -Y : -X]$, we get the following curve:

$$(4.61) \quad H_3 : y^2 = -3x^6 + 4x^5 - 26x^4 + 12x^3 - 55x^2 - 16x + 4.$$

This genus 2 hyperelliptic curve contains (at least) the following two points:

$$(4.62) \quad P_1 = [0 : -2 : 1], P_2 = [0 : 2 : 1].$$

Using MAGMA, one can compute that H_3 has Jacobian with Mordell–Weil rank bounded by 1, and torsion subgroup isomorphic to $\mathbb{Z}/2\mathbb{Z}$. The divisor class $[P_1 - P_2]$ has infinite order in the Jacobian, and therefore generates a finite index subgroup of the Mordell–Weil group. We can use the implementation of the Chabauty–Coleman/Mordell–Weil sieving algorithm (cf. Bruin and Stoll [2]) in MAGMA, and find that the only rational points on the curve are the two points we had already discovered.

We pull these two points all the way back C_1 to find the complete set of rational points on the latter curve (not including 4 points at infinity):

$$\left(\frac{1}{2}, 1, 1\right), (1, 0, 0), (0, 0, 0), (1, 1, 0), (0, 0, 1).$$

None of these points have a d value corresponding to a quadratic rational function ϕ_d realizing N3H3.

4.3. Support for Conjecture 1.

4.3.1. *Extra 3-cycle.* We compute the 3-rd dynatonic polynomial of ϕ_a .

$$\begin{aligned} \Phi_{a,3}^*(z) := \Phi_{\phi_a,3}^*(z) &= (a^5 + 5a^4 + 10a^3 + 10a^2 + 5a + 1)z^5 \\ &\quad + (-2a^5 - 7a^4 - 9a^3 - 4a^2 + a + 1)z^4 \\ &\quad + (a^5 - 7a^3 - 13a^2 - 10a - 3)z^3 \\ &\quad + (2a^4 + 5a^3 + 4a^2 + a)z^2 \\ &\quad + (a^3 + 3a^2 + 3a + 1)z \end{aligned}$$

We divide the dynatonic polynomial by the factor $(a+1)z(z-1)$ to get the irreducible polynomial

$$P_3 := a^3z^3 - a^3z^2 + 3a^2z^3 - 2a^2z + 3az^3 + 2az^2 - 2az - a + z^3 + 2z^2 - z - 1$$

The curve C_3 defined by the equation $P_3 = 0$ contains the four points

$$[0 : 1 : 0], [-1 : 0 : 1], [1 : 0 : 0], [-1 : 1 : 1].$$

Furthermore, C_3 is birational to the genus 2 hyperelliptic curve defined by

$$H_3 : y^2 = x^6 - 4x^5 + 6x^4 - 2x^3 + x^2 - 2x + 1$$

Using MAGMA, we can check that H_3 has Jacobian of Mordell–Weil rank 0. The curve H_3 has good reduction modulo 3, and its reduction has Jacobian isomorphic to the group $\mathbb{Z}/19\mathbb{Z}$. This means that Mordell–Weil group (of the Jacobian) of H_3 is either trivial or isomorphic to $\mathbb{Z}/19\mathbb{Z}$. However, one can check that the element on the Jacobian defined by the difference of the two points at infinity $[1 : -1 : 0] - [1 : 1 : 0]$ has order 19 and therefore the Jacobian of H_3 is isomorphic to $\mathbb{Z}/19\mathbb{Z}$.

We get the following commutative diagram where all the maps are injective.

$$\begin{array}{ccc} H_3(\mathbb{Q}) & \longrightarrow & J(\mathbb{Q}) \\ \downarrow & & \downarrow \\ H_3(\mathbb{F}_3) & \longrightarrow & J(\mathbb{F}_3) \end{array}$$

We check that

$$H_3(\mathbb{F}_3) = \{[1 : -1 : 0], [1 : 1 : 0], [0 : -1 : 1], [0 : 1 : 1], [1 : -1 : 1], [1 : 1 : 1]\}.$$

All of these points are reductions of points from $H_3(\mathbb{Q})$, and this implies that the \mathbb{Q} -rational points on H_3 are exactly the six points

$$[1 : -1 : 0], [1 : 1 : 0], [0 : -1 : 1], [0 : 1 : 1], [1 : -1 : 1], [1 : 1 : 1].$$

We pull these points back to C_3 , to find that the only rational points on C_3 are the four points we already discovered. None of which correspond to non-degenerate maps ϕ_a .

4.3.2. Extra 4-cycle. We compute the 4th dynatomic polynomial of ϕ_a , and divide by a factor of $(a+1)^4$, to get the following polynomial.

$$\begin{aligned}
 P_4(a, t) := & (a^6 + 6a^5 + 15a^4 + 20a^3 + 15a^2 + 6a + 1)z^{12} \\
 & + (-2a^7 - 19a^6 - 64a^5 - 107a^4 - 98a^3 - 49a^2 - 12a - 1)z^{11} \\
 & + (a^8 + 16a^7 + 77a^6 + 152a^5 + 128a^4 + 16a^3 - 46a^2 - 30a - 6)z^{10} \\
 & + (-3a^8 - 27a^7 - 58a^6 + 38a^5 + 248a^4 + 304a^3 + 165a^2 + 41a + 4)z^9 \\
 & + (3a^8 + 8a^7 - 61a^6 - 239a^5 - 240a^4 - 8a^3 + 111a^2 + 59a + 9)z^8 \\
 & + (-a^8 + 11a^7 + 64a^6 + 10a^5 - 251a^4 - 317a^3 - 126a^2 - 8a + 2)z^7 \\
 & + (-6a^7 + 12a^6 + 128a^5 + 133a^4 - 94a^3 - 153a^2 - 48a - 1)z^6 \\
 & + (-16a^6 - 5a^5 + 130a^4 + 156a^3 - a^2 - 42a - 8)z^5 \\
 & + (-25a^5 - 26a^4 + 78a^3 + 94a^2 + 12a - 7)z^4 \\
 & + (-25a^4 - 31a^3 + 26a^2 + 33a + 5)z^3 \\
 & + (-16a^3 - 20a^2 + 2a + 5)z^2 \\
 & + (-6a^2 - 7a - 1)z - a - 1.
 \end{aligned}$$

The equation $P_4 = 0$ describes an affine plane curve of genus 12 which we denote by C_4 . We follow the trace map method described in [14]: Let

$$tr_4(z) = z + \phi(z) + \phi^2(z) + \phi^3(z).$$

The image of the curve C_4 under the map $\Psi : (a, z) \mapsto (a, tr_4(z))$ is birational to the quotient curve $C_4 / \langle \sigma \rangle$, where σ is the automorphism of C_4 defined by $(a, z) \mapsto (a, \phi(z))$. When ϕ is a polynomial, the image of Ψ can be calculated by taking the resultant of $\Phi_{a,4}^*(z)$ and $t - tr_4(z)$ with respect to z . However, in our case $t - tr_4(z)$ is not a polynomial, since $tr_4(z)$ is a rational function in z . We fix this by clearing out the denominator of $tr_4(z)$; we denote the numerator of tr_4 by A and the denominator by B .

We compute the resultant $\text{Res}(P_4, Bt - A)$ and find the following irreducible factor.

$$\begin{aligned}
 \tilde{P}_4(a, t) = & a^7t - 2a^6t^2 + a^5t^3 - a^7 + 13a^6t - 17a^5t^2 + 5a^4t^3 \\
 & - 7a^6 + 53a^5t - 47a^4t^2 + 10a^3t^3 - 6a^5 + 80a^4t \\
 & - 60a^3t^2 + 10a^2t^3 + 43a^4 + 42a^3t - 38a^2t^2 + 5at^3 \\
 & + 95a^3 - 10a^2t - 11at^2 + t^3 + 89a^2 - 19at - t^2 + 42a - 6t + 9
 \end{aligned}$$

The degree 8 curve defined by $\tilde{P}_4 = 0$ is of genus 1. It is birational to the elliptic curve with Cremona reference 19a3 (this curve has already been encountered as being birational to the modular curve of N3E1, see the

remark in Section 4.2.1), and has the following minimal model

$$y^2 + y = x^3 + x^2 + x.$$

Its Mordell–Weil group is isomorphic to $\mathbb{Z}/3\mathbb{Z}$ and therefore there are exactly three rational points on this elliptic curve. These are

$$[0 : 0 : 1], [0 : 1 : 0], [0 : -1 : 1].$$

Pulling these points back to the curve $C_4/\langle\sigma\rangle$, we find all rational points on $C_4/\langle\sigma\rangle$ to be

$$[0 : 1 : 0], [1 : 0 : 0], [1 : 1 : 0].$$

All these points are “cusps” (the points added to the dynamical modular curve to make it projective), and therefore we can conclude that there are no quadratic rational functions ϕ_a with a 4-cycle.

References

- [1] W. BOSMA, J. CANNON & C. PLOUOST, “The Magma algebra system. I. The user language”, *J. Symb. Comput.* **24** (1997), no. 3-4, p. 235-265.
- [2] N. BRUIN & M. STOLL, “The Mordell–Weil sieve: proving non-existence of rational points on curves”, *LMS J. Comput. Math.* **13** (2010), p. 272-306.
- [3] J. K. CANCI & S. VISHKAUTSAN, “Quadratic maps with a periodic critical point of period 2”, *Int. J. Number Theory* **13** (2017), no. 6, p. 1393-1417.
- [4] J. CREMONA, “The elliptic curve database for conductors to 130000”, in *Algorithmic number theory*, Lecture Notes in Computer Science, vol. 4076, Springer, 2006, p. 11-29.
- [5] J. R. DOYLE, X. FABER & D. KRUMM, “Preperiodic points for quadratic polynomials over quadratic fields”, *New York J. Math.* **20** (2014), p. 507-605.
- [6] N. FAKHRUDDIN, “Questions on self maps of algebraic varieties”, *J. Ramanujan Math. Soc.* **18** (2003), no. 2, p. 109-122.
- [7] E. V. FLYNN, B. POONEN & E. F. SCHAEFER, “Cycles of quadratic polynomials and rational points on a genus-2 curve”, *Duke Math. J.* **90** (1997), no. 3, p. 435-463.
- [8] B. HUTZ & P. INGRAM, “On Poonen’s conjecture concerning rational preperiodic points of quadratic maps”, *Rocky Mt. J. Math.* **43** (2013), no. 1, p. 193-204.
- [9] D. LUKAS, M. MANES & D. YAP, “A census of quadratic post-critically finite rational functions defined over \mathbb{Q} ”, *LMS J. Comput. Math.* **17A** (2014), p. 314-329.
- [10] M. MANES, “ \mathbb{Q} -rational cycles for degree-2 rational maps having an automorphism”, *Proc. Lond. Math. Soc.* **96** (2008), no. 3, p. 669-696.
- [11] J. MILNOR, “Geometry and dynamics of quadratic rational maps”, *Exp. Math.* **2** (1993), no. 1, p. 37-83.
- [12] P. MORTON, “Rational periodic points of rational functions”, *Int. Math. Res. Not.* **1994** (1994), no. 2, p. 97-110.
- [13] ———, “Periodic points, multiplicities, and dynamical units”, *J. Reine Angew. Math.* **461** (1995), p. 81-122.
- [14] ———, “Arithmetic properties of periodic points of quadratic maps. II”, *Acta Arith.* **87** (1998), no. 2, p. 89-102.
- [15] D. G. NORTHCOTT, “Periodic points on an algebraic variety”, *Ann. Math.* **51** (1950), p. 167-177.
- [16] B. POONEN, “The classification of rational preperiodic points of quadratic polynomials over \mathbb{Q} : a refined conjecture”, *Math. Z.* **228** (1998), no. 1, p. 11-29.
- [17] J. H. SILVERMAN, *The arithmetic of dynamical systems*, Graduate Texts in Mathematics, vol. 241, European Mathematical Society, 2007.
- [18] M. STOLL, “Rational 6-cycles under iteration of quadratic polynomials”, *LMS J. Comput. Math.* **11** (2008), p. 367-380.

Appendix. Rational points on a curve of genus 6 par Michael Stoll

ABSTRACT. We determine the set of rational points on the curve of genus 6 that parameterizes quadratic maps $\phi: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ with an orbit of length 3 containing a critical point and a point P such that $\phi^{\circ 3}(P)$ has order 2 (but $\phi^{\circ 2}(P)$ is not periodic). The result is conditional on standard conjectures (including BSD) on the L -series of the Jacobian of the curve in question.

A.1. Introduction. We consider the curve C that classifies (up to conjugation by an automorphism of \mathbb{P}^1) quadratic maps $\phi: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ such that

- (1) ϕ has a cycle of length 3 containing a critical point, and
- (2) ϕ has a cycle of length 2, together with a marked third preimage of one of the two points in the cycle (whose second image is not in the cycle).

(See the introduction of the main article for the definitions of a critical point and n -cycle). These are exactly the type of maps described by the graph N3H2 in Table 4 of the main article. The goal is to show that no such (non-degenerate) ϕ are defined over \mathbb{Q} , which is equivalent to showing that all the rational points on C correspond to degenerate maps ϕ . To this end, we first derive an equation for C as a singular affine plane curve, then we construct its canonical model in \mathbb{P}^5 , which is a smooth projective curve D birational to C . Finally, we determine $D(\mathbb{Q})$ (to be able to do this, we have to assume some standard conjectures regarding the L -series of the Jacobian of D , including the BSD conjecture) and map the points back to C .

The arguments and computations are to a large extent parallel to those performed in [21] in a similar situation, so we give here a condensed description and refer the reader to [21] for more information.

A.2. The curve and its canonical model. We can fix the critical 3-cycle to be $0 \mapsto \infty \mapsto 1 \mapsto 0$ with 0 the critical point (see Section 4.1.2 in the main article for details). Then

$$\phi(z) = \frac{(a+1)z^2 - az - 1}{(a+1)z^2}$$

with a parameter a . The condition for the existence of a rational 2-cycle is then that $(a+1)^2 + 4$ is a square. The conic given by this condition can be parameterized, leading to

$$a = -\frac{4t}{t^2 - 1} - 1$$

and

$$\phi_t(z) = \frac{4tz^2 - (t^2 + 4t - 1)z + t^2 - 1}{4tz^2};$$

the 2-cycle contains the two points $\frac{t+1}{2}$ and $\frac{t-1}{2t}$, which are swapped by the automorphism (on the parameter space) $t \mapsto -1/t$. ϕ_t degenerates for $t = -1, 0, 1, \infty$.

The other preimage of $\frac{t+1}{2}$ is $-\frac{t+1}{t-1}$. We will require it to have a rational second preimage. The requirement of a rational (first) preimage results in a curve that is birational to the elliptic curve E with label 53a1 in the Cremona database, whose Mordell–Weil group is \mathbb{Z} (This is the curve parametrizing rational functions admitting the graph R3P5 in Table 1 of the main article; see Section 4.1.6 there). Requiring a second rational preimage results in a double cover of this curve, which (from setting $\phi_t^{\circ 2}(z) = -\frac{t+1}{t-1}$) can be given by the affine equation

$$\begin{aligned} 16t^2(t+1)z^4 + (t^2 + 4t - 1)(t^3 - 13t^2 - 5t + 1)z^3 \\ + (t^5 + 29t^4 + 34t^3 - 30t^2 - 3t + 1)z^2 \\ - 4(t-1)t(t+1)(t^2 + 4t - 1)z + 2(t-1)^2t(t+1)^2 = 0. \end{aligned}$$

A quick computation in Magma [19] shows that (the smooth projective model of) this curve has genus $g = 6$. Setting

$$t = \frac{u+1}{u-1} \quad \text{and} \quad z = \frac{1}{1-v}$$

results in the simpler equation

$$\begin{aligned} F(u, v) := u^5v^2 + 2u^4v^3 - u^4v^2 - u^4v + u^3v^4 - 4u^3v^2 + u^3 \\ + u^2v^4 - 4u^2v^3 + 3u^2v - 2uv^3 + 4uv^2 - u + v^2 - v = 0 \end{aligned}$$

for our curve C . In terms of u and v , the involution corresponding to the double cover $C \rightarrow E$ is given by $(u, v) \mapsto (u, u^{-1} - 1 - u - v)$.

Set

$$\omega_0 = \frac{du}{\frac{\partial F}{\partial v}(u, v)} = -\frac{dv}{\frac{\partial F}{\partial u}(u, v)}.$$

Then the following differentials are a basis of the space of regular differentials on C :

$$\begin{aligned} \omega_1 &= (u^3v + u^2v^2 - 2u^2v - uv^2 + v)\omega_0, \\ \omega_2 &= (u^2v + uv^2 - uv - v)\omega_0, \\ \omega_3 &= uv\omega_0, \\ \omega_4 &= (u^2 - u)\omega_0, \\ \omega_5 &= (u - 1)\omega_0, \\ \omega_6 &= \omega_0. \end{aligned}$$

The projective closure D of the image of the corresponding canonical map $C \rightarrow \mathbb{P}^5$ (with coordinates w_1, \dots, w_6) is then defined by the following six

quadratic equations:

$$\begin{aligned}
 -w_2w_5 + w_1w_6 &= 0, \\
 -w_2w_4 + w_1w_5 + w_2w_5 &= 0, \\
 -w_5^2 + w_4w_6 - w_5w_6 &= 0, \\
 -w_3^2 - w_3w_4 + w_1w_6 + w_2w_6 + w_3w_6 &= 0, \\
 w_1^2 + 2w_1w_2 - w_1w_4 - w_2w_4 + w_4^2 - w_1w_5 + w_4w_5 &= 0, \\
 w_1w_2 + 2w_2^2 - w_2w_4 - w_2w_5 + w_4w_5 + w_5^2 \\
 - w_1w_6 - w_2w_6 + w_4w_6 + w_5w_6 &= 0.
 \end{aligned}$$

The birational map $D \rightarrow C$ is given by

$$u = \frac{w_4 + w_5 + w_6}{w_5 + w_6}, \quad v = \frac{w_3}{w_5 + w_6}.$$

Using the `PointSearch` command of Magma, we find the following nine rational points on D :

$$\begin{aligned}
 P_1 &= (0 : 0 : 0 : 2 : -2 : 1), & P_6 &= (-2 : 1 : 0 : 0 : 0 : 0), \\
 P_2 &= (0 : 0 : 1 : 0 : 0 : 1), & P_7 &= (0 : 0 : 0 : 0 : 0 : 1), \\
 P_3 &= (0 : 0 : -1 : 2 : -2 : 1), & P_8 &= (0 : 0 : 1 : 0 : -1 : 1), \\
 P_4 &= (0 : 0 : 0 : 0 : -1 : 1), & P_9 &= (1 : -1 : 1 : 0 : -1 : 1). \\
 P_5 &= (1 : -1 : 0 : 0 : -1 : 1),
 \end{aligned}$$

We suspect that these are all the rational points. The proof of this claim will take up the remainder of this appendix.

We note that the images of these points on C are $(-1, 0)$, $(1, 1)$, $(-1, 1)$, $(0, 0)$, $(0, 1)$, a point at infinity, $(1, 0)$, and two times a point at infinity. Since $u = -1, 0, 1, \infty$ gives a degenerate ϕ and $v = \infty$ gives $z = 0$, which also makes ϕ degenerate, this will imply that there are no non-degenerate ϕ defined over \mathbb{Q} with the required properties.

A.3. The Jacobian. Working with the equations for $D \subset \mathbb{P}^5$, we find that D has bad reduction (at most) at the primes 3, 53 and 99 563. In each of these cases, the reduction is semistable, with a single component in the special fiber that has one split node for $p = 3$, two non-split nodes each defined over \mathbb{F}_{53} for $p = 53$, and one split node for $p = 99\,563$. (A node is *split*, if the two tangent directions are defined over the field of definition of the node.) So D is semistable, and its Jacobian J has conductor $N_J = 3 \cdot 53^2 \cdot 99\,563$. Since D maps non-trivially to the elliptic curve E of conductor 53, J is isogenous to a product $E \times A$, where A is an abelian variety over \mathbb{Q} of dimension 5 and with conductor $N_A = 3 \cdot 53 \cdot 99\,563 = 15\,830\,517$.

With a computation analogous to that leading to Lemma 4 in [21], we find that the torsion subgroup of $J(\mathbb{Q})$ has exponent dividing 2 (we did not try to find the torsion subgroup exactly) and that the differences of the nine known rational points on D generate a subgroup isomorphic to \mathbb{Z}^2 of $J(\mathbb{Q})$. So if we can show that $J(\mathbb{Q})$ has rank 2, then we know generators of a subgroup of finite index (and the rank is strictly less than the genus), so we can apply the Chabauty–Coleman method.

Since there is little hope to perform a successful Selmer group computation on J , which would give an upper bound for the rank, we follow the approach already used in [21] and assume that the L -series of J has an analytic continuation to all of \mathbb{C} , that the function

$$\Lambda(J, s) = N_J^{s/2} (2\pi)^{-6s} \Gamma(s)^6 L(J, s)$$

satisfies the functional equation $\Lambda(J, 2 - s) = w_J \Lambda(J, s)$ with the global root number $w_J = \pm 1$, and that the Birch and Swinnerton–Dyer conjecture holds for J . For the computations, it is better to work with the L -series of A , since its conductor N_A is smaller than N_J . According to Magma’s implementation of L -series, it requires the coefficients of $L(A, s)$ up to $n \approx 105\,000$ for a precision of 20 decimal digits. We find the coefficients of the Euler factors of the L -series of J up to the required bound by counting the \mathbb{F}_q -points on D for all prime powers q below the bound. This is most efficiently done on the affine model C , by keeping track of the points at infinity modulo p and of what happens at the singular points (and some care has to be taken at the bad primes). In this way and after dividing by the Euler factors of E , we obtain within a few hours the relevant coefficients for the computation. We then check (using Magma’s `CheckFunctionalEquation`) that the data we have computed is compatible with the expected functional equation for $L(A, s)$ with root number $w_A = -1$ (but not with $w_A = 1$). So we can safely assume that $L(J, s)$ satisfies the functional equation with root number $w_J = w_A w_E = (-1)(-1) = 1$. This is also in agreement with the expectation that the global root number should be equal to the product of the local root numbers, which in the semistable case is $(-1)^{g+s}$, where s is the total number of Frobenius orbits of split nodes. Here $g = 6$ and $s = 2$, so the root number should indeed be 1. We then evaluate the derivative of $L(A, s)$ at $s = 1$ numerically and find a clearly nonzero value of $\approx 0.026803015530623712948$. So according to the BSD conjecture, the rank of $A(\mathbb{Q})$ should be 1 and the rank of $J(\mathbb{Q})$ should therefore be 2.

A.4. Applying the Chabauty–Coleman method. By the results of the previous section (assuming BSD for J), we now know that the differences of the known rational points provide us with generators of a finite-index subgroup of $J(\mathbb{Q})$. To apply Chabauty–Coleman, we have to fix a prime p . We choose $p = 5$, because 5 is a prime of good reduction and the

set of known rational points maps bijectively to the points in $D(\mathbb{F}_5)$ under reduction. This latter observation implies that it will be enough to show that each residue class mod 5 in $D(\mathbb{Q}_5)$ contains at most one rational point. By the results of [20], this follows when we can show that for each point in $D(\mathbb{F}_5)$, there is a differential in the annihilator of $J(\mathbb{Q})$ whose reduction mod 5 does not vanish there.

We first have to find a basis of this annihilator in the space of regular differentials on D over \mathbb{Q}_5 . We first fix one of our known rational points P as a base-point; we have to choose it in such a way that its reduction \bar{P} mod 5 is non-special in the sense that the Riemann–Roch space of $6\bar{P}$ is one-dimensional. We check that P_4 satisfies this requirement. We then use the reduction map $J(\mathbb{Q}) \rightarrow J(\mathbb{F}_5)$ to find two independent points in its kernel, which we represent by divisors of the form $\mathcal{D}_j - 6P_4$, where \mathcal{D}_1 and \mathcal{D}_2 are effective of degree 6. Since \bar{P}_4 is non-special, it follows that \mathcal{D}_j reduces mod 5 to $6\bar{P}_4$ for $j = 1, 2$. We choose $t = 1 + w_5/w_6$ as a uniformizer at P_4 (that reduces to a uniformizer at \bar{P}_4), express the differentials $\omega_1, \dots, \omega_6$ as power series in t times dt , integrate formally, and use the method explained in [21] to compute the relevant integrals modulo 5^3 (modulo 5^2 would actually be sufficient). They are all multiples of 5, so we divide them by 5 and reduce the 2×6 -matrix obtained modulo 5. Its kernel gives the reduction of annihilator, which in our case is generated by the reductions of $\omega_1 + \omega_4$, $\omega_2 - \omega_4$, ω_5 and ω_6 . Since the locus of vanishing of a linear combination of the ω_j on D is given by the corresponding hyperplane section, all we have to do is to check that the line defined in $\mathbb{P}_{\mathbb{F}_5}^5$ by $w_1 + w_4 = w_2 - w_4 = w_5 = w_6 = 0$ does not meet $D_{\mathbb{F}_5}$, which is easily verified. This concludes the proof.

References for the Appendix

- [19] W. BOSMA, J. CANNON & C. PLAYOUST, “The Magma algebra system. I. The user language”, *J. Symb. Comput.* **24** (1997), no. 3-4, p. 235-265.
- [20] M. STOLL, “Independence of rational points on twists of a given curve”, *Compos. Math.* **142** (2006), no. 5, p. 1201-1214.
- [21] ———, “Rational 6-cycles under iteration of quadratic polynomials”, *LMS J. Comput. Math.* **11** (2008), p. 367-380.

Solomon VISHKAUTSAN
 Tel-Hai Academic College, Israel
E-mail: wishcow@gmail.com
URL: http://wishcow.com

Michael STOLL
 Mathematisches Institut, Universität Bayreuth, 95440 Bayreuth, Germany
E-mail: Michael.Stoll@uni-bayreuth.de
URL: http://www.computeralgebra.uni-bayreuth.de