

JOURNAL

de Théorie des Nombres
de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Shiang TANG

Motivic Galois representations valued in Spin groups

Tome 33, n° 1 (2021), p. 197-221.

<http://jtnb.centre-mersenne.org/item?id=JTNB_2021__33_1_197_0>

© Société Arithmétique de Bordeaux, 2021, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.centre-mersenne.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.centre-mersenne.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.centre-mersenne.org/>

Motivic Galois representations valued in Spin groups

par SHIANG TANG

*Dedicated to the memory of my mother,
Que Ling Ping, 1954/07/08 – 2019/10/28*

RÉSUMÉ. Soit m un entier tel que $m \geq 7$ et $m \equiv 0, 1, 7 \pmod{8}$. Nous construisons des systèmes strictement compatibles de représentations l -adiques $\Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{\mathbb{Q}}_l) \xrightarrow{\mathrm{spin}} \mathrm{GL}_N(\overline{\mathbb{Q}}_l)$ qui sont potentiellement automorphes et motiviques. Comme application, dans certains cas nous donnons une réponse positive au problème de Galois inverse pour les groupes spinoriels sur \mathbb{F}_p . Pour m impair, nous comparons nos exemples avec le travail de A. Kret et S. W. Shin ([18]), qui étudie les représentations galoisiennes automorphes à valeurs dans GSpin_m .

ABSTRACT. Let m be an integer such that $m \geq 7$ and $m \equiv 0, 1, 7 \pmod{8}$. We construct strictly compatible systems of representations of $\Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{\mathbb{Q}}_l) \xrightarrow{\mathrm{spin}} \mathrm{GL}_N(\overline{\mathbb{Q}}_l)$ that are potentially automorphic and motivic. As an application, we prove instances of the inverse Galois problem for the \mathbb{F}_p -points of the spin groups. For odd m , we compare our examples with the work of A. Kret and S. W. Shin ([18]), which studies automorphic Galois representations valued in GSpin_m .

1. Introduction

In [28, 8.4], Serre asks whether there are motives (over \mathbb{Q} , say), whose motivic Galois groups are equal to a given semisimple (or more generally, reductive) algebraic group G . This question and its variants have been studied by many people, including N. Katz, M. Dettweiler, S. Reiter, Z. Yun, S. Patrikis and others; see for example [32], [24] and [7]. Most of the results in the literature concern exceptional algebraic groups. In this paper, we study a weaker version of Serre’s question for spin groups. We find spin groups interesting because their faithful representations have large dimensions and they do not occur in the étale cohomology of smooth projective varieties in any obvious way. In [18], Kret and Shin prove the existence of

Manuscrit reçu le 12 juin 2020, accepté le 13 décembre 2020.

2010 *Mathematics Subject Classification*. 11F80.

Mots-cléfs. Galois representations, spin groups, inverse Galois problem, automorphic representations.

Galois representations into GSpin_{2n+1} corresponding to suitable cuspidal automorphic representations of GSp_{2n} over totally real fields. Their Galois representations are motivic in the sense that they occur in the cohomology of certain Shimura varieties. Studying Spin_{2n+1} -valued Galois representations then appears to be a simple matter of passing from a reductive group to its (semisimple) derived subgroup, but the distinction can indeed be subtle: for example, when $n = 1$, $\mathrm{GSpin}_3 = \mathrm{GL}_2$, it is well-known that holomorphic modular forms (or elliptic curves) give rise to p -adic Galois representations $\rho : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}}_p)$ that are odd, i.e. $\det \rho(c) = -1$; in particular, they do not land in SL_2 , nor can they be twisted into SL_2 in any obvious way. In fact, two-dimensional geometric Galois representations that are even (i.e. $\det \rho(c) = 1$) are expected to come from Maass forms by the Fontaine–Mazur–Langlands conjectures.

Our main theorem is the following:

Theorem 1.1. (*Proposition 4.1 and Theorem 4.2*) *Let m be an integer such that $m \geq 7$ and $m \equiv 0, 1, 7 \pmod{8}$. There exists a strictly compatible system*

$$R_{\lambda} : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{GL}_N(\overline{M}_{\lambda})$$

with distinct Hodge–Tate weights and with coefficients in a number field M such that for a density one set of rational primes l and for $\lambda|l$, $R_{\lambda} = \mathrm{spin} \circ r_{\lambda}$, where $r_{\lambda} : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{M}_{\lambda})$ is a homomorphism with Zariski-dense image and $\mathrm{spin} : \mathrm{Spin}_m \rightarrow \mathrm{GL}_N$ is the spin representation. Moreover, $\{R_{\lambda}\}$ is potentially automorphic and motivic in the following sense:

- *There exist a totally real extension F^+/\mathbb{Q} and a regular L -algebraic, cuspidal automorphic representation Π of $\mathrm{GL}_N(\mathbb{A}_{F^+})$ such that*

$$R_{\lambda}|_{\Gamma_{F^+}} \cong r_{\Pi, \iota_l},$$

where $\iota_l : \overline{\mathbb{Q}}_l \xrightarrow{\sim} \mathbb{C}$ is a fixed field isomorphism.

- *There is a smooth projective variety X/\mathbb{Q} and integers i and j such that R_{λ} is a $\Gamma_{\mathbb{Q}}$ -subrepresentation of $H^i(X_{\overline{\mathbb{Q}}}, \overline{\mathbb{Q}}_l(j))$.*

This is the spin-analog of the main result of [7]. As an application, we prove new instances of the inverse Galois problem:

Theorem 1.2. (*Corollary 4.3*) *For $m \geq 7$, $m \equiv 0, 1, 7 \pmod{8}$, $\mathrm{Spin}_m(\mathbb{F}_p)$ is the Galois group of a finite Galois extension of \mathbb{Q} for p belonging to a set of rational primes of positive density.*

Let us explain the strange-looking congruence condition on m appeared in Theorem 1.1. Let T be a maximal split torus of Spin_m and let $W = N(T)/T$ be the Weyl group. Then the integer m satisfies the above congruence condition if and only if the longest element $w^0 \in W$ acts as -1 on $X^{\bullet}(T)$ and w^0 has a representative in $N(T)$ of order 2 ([1, Section 3]; see

also Lemma 2.3). Having such an involution at hand is crucial in standard Galois deformation theoretic arguments: one typically constructs an appropriate mod p representation $\bar{r} : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{\mathbb{F}}_p)$ which is odd in the sense of [14, Definition 1.2] (which will hold if the complex conjugation maps into the conjugacy class of the above involution in Spin_m), then deform it to a geometric (in the sense of Fontaine–Mazur) characteristic zero representation using either Ramakrishna style techniques (originated in [26] and sublimed in [14]) or Khare–Wintenberger style arguments ([16, Section 3]). Oddness of \bar{r} is crucial in both methods.

Due to the automorphic nature of our construction, the Hodge–Tate weights of the compatible system in Theorem 1.1 are distinct. In contrast, suppose w^0 acts as -1 on $X^\bullet(T)$ but it does not lift to an involution in $N(T)$ (which happens if and only if $m \equiv 3, 4, 5 \pmod{8}$), then we do not expect Galois representations $r : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{\mathbb{Q}}_l)$ such that $\mathrm{spin} \circ r : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{GL}_N(\overline{\mathbb{Q}}_l)$ comes from a pure motive with *regular* Hodge structure. In fact, the spin representation $\mathrm{spin} : \mathrm{Spin}_m \rightarrow \mathrm{GL}_N$ is valued in Sp_N for $m \equiv 3, 4, 5 \pmod{8}$ by Lemma 1.3. If there were such a motive, let V be its real Hodge structure such that $V \otimes_{\mathbb{R}} \mathbb{C} = \bigoplus_{p+q=w} V_{p,q}$ with $\overline{V}_{p,q} = V_{q,p}$ and $w \in \mathbb{Z}$. By symmetry of the symplectic torus and regularity, we have $\dim V_{p,q} = \dim V_{-p,-q} = 1$ for all p, q appearing in the direct sum. In particular, $w = 0$ and the complex conjugation action corresponds to the longest element in the Weyl group of Sp_N , which lifts to an order 4 element in the normalizer of the torus of Sp_N , a contradiction.¹ Assuming the Fontaine–Mazur conjecture, we can express this in purely Galois theoretic terms: for $m \equiv 3, 4, 5 \pmod{8}$, there should not exist l -adic Galois representations

$$\Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{\mathbb{Q}}_l) \xrightarrow{\mathrm{spin}} \mathrm{GL}_N(\overline{\mathbb{Q}}_l)$$

that are unramified everywhere, potentially semistable at l with distinct Hodge–Tate weights. For example, let $m = 3$, then $\mathrm{Spin}_3 \cong \mathrm{SL}_2$ and the spin representation is the canonical injection $\mathrm{SL}_2 \rightarrow \mathrm{GL}_2$. In this case, our speculation follows from a theorem of Calegari ([9, Theorem 1.2]) under some mild hypotheses.

Methods and organization of this paper. The method we use in proving Theorem 1.1 is very similar to that of [7]. In Section 2, we begin by constructing a mod p representation $\bar{r} : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{\mathbb{F}}_p)$ (for m satisfying the congruence condition in Theorem 1.1) related to the action of the Weyl group of Spin_m on the weight space of the spin representation for which $\mathrm{spin} \circ \bar{r}$ satisfies the assumptions of the automorphic lifting theorems in [3]. Succeeding in constructing such a representation requires, in

¹This follows from Lemma 3.1 (with $\delta = 1$) and the proof of Lemma 4.12 of [1]. One can also verify this directly.

addition to the techniques used in [7, Section 2], a detailed calculation on the structure of Spin_m and the crucial observation that a certain part of the Weyl group acts transitively on the weight lattice of the spin representation (Lemma 2.1). We then deform \bar{r} to a geometric representation $r : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{\mathbb{Q}_p})$ that is Steinberg at a finite place using a version of the Khare–Wintenberger argument: similar arguments have appeared in [7, Section 3] and [25, Section 3], we present an axiomatized version in Section 3. Then [3, Theorem C] implies that $\mathrm{spin} \circ r$ is potentially automorphic and is part of a strictly compatible system $\{R_\lambda\}$. We wish to show that the Zariski closure of the image of R_λ equals Spin_m for *all* λ . Following [7, Section 4], we exploit the fact that the image of r contains a regular unipotent element of Spin_m and use ideas of Larsen and Pink ([22]), where the key is to show that R_λ is irreducible for all λ . This is proven for E_6 in [7] using elementary combinatorial properties of the formal character of E_6 , which does not carry over to Spin_m since the rank of the latter can get arbitrarily large. Instead, we get away with a weaker statement by invoking [3, Theorem D] (which relies on Larsen’s work [20]): this is where the density one condition in Theorem 1.1 came from.² This is done in Section 4. In Section 5, we show that $\{R_\lambda\}$ occurs in the cohomology of a smooth projective variety following [7, Section 5]. In Section 6, we compare our construction with the work of Kret and Shin ([18]): we explain how the main theorem of [18] yields a stronger version of Theorem 1.1 for $m \equiv 1, 7 \pmod{8}$.³ We also observe that for $m \equiv 3, 5 \pmod{8}$, [18] (in which the automorphic representations are regular L–algebraic) does not yield Spin_m –valued Galois representations (Lemma 6.3).

1.1. Notation. Let F be a field. Fix an algebraic closure \bar{F} of F and write Γ_F for the absolute Galois group $\mathrm{Gal}(\bar{F}/F)$ of F . If F is a number field, then for each place v of F , we fix an embedding $\bar{F} \rightarrow \bar{F}_v$ into an algebraic closure of F_v , which gives rise to an injective group homomorphism $\Gamma_{F_v} \rightarrow \Gamma_F$. For any finite place v , let k_v be the residue field of v and let $\mathrm{Fr}_v \in \Gamma_{k_v}$ be the arithmetic Frobenius. If H is a group (typically the points over a finite field or a p -adic field of a reductive algebraic group), and there is a continuous group homomorphism $r : \Gamma_F \rightarrow H$, we will sometimes write $r|_v$ for $r|_{\Gamma_{F_v}}$, the restriction of r to the decomposition group Γ_{F_v} . If H acts on a finite-dimensional vector space V , we write $r(V)$ for the Γ_F –module induced by precomposing this action with r . (Typically H will be a reductive algebraic group and V will be its Lie algebra equipped with the adjoint action of H .)

²One could conceivably impose additional local deformation conditions (for instance, those corresponding to supercuspidal representations) to force R_λ to be irreducible for all λ , but I do not know how to do this at present.

³We note that very recently Kret and Shin have extended their work to even orthogonal groups in [19]. The main theorem of [19] yields a stronger version of Theorem 1.1 for $m \equiv 0 \pmod{8}$ in a similar way.

Let $\kappa : \Gamma_F \rightarrow \mathbb{Z}_p^\times$ be the p -adic cyclotomic character and $\bar{\kappa}$ be its reduction modulo p . We will always assume $p \neq 2$, and our main theorems will make stronger hypotheses on p .

We recall here some deformation-theoretic terminology. Given a topologically finitely-generated profinite group Γ , a finite extension E/\mathbb{Q}_p with ring of integers \mathcal{O} and residue field k , a reductive algebraic group G defined over \mathcal{O} and a continuous homomorphism $\bar{r} : \Gamma \rightarrow G(k)$, let $R_{\mathcal{O}, \bar{r}}^\square$ be the universal lifting ring representing the functor sending a complete local noetherian \mathcal{O} -algebra R with residue field k to the set of lifts $r : \Gamma \rightarrow G(R)$ of \bar{r} . We will always leave the \mathcal{O} implicit, writing only $R_{\bar{r}}^\square$, and at various points in the argument we enlarge \mathcal{O} ; see [3, Lemma 1.2.1] for a justification of (the harmlessness of) this practice. We write $R_{\bar{r}}^\square \otimes \overline{\mathbb{Q}_p}$ for $R_{\mathcal{O}, \bar{r}}^\square \otimes_{\mathcal{O}} \overline{\mathbb{Q}_p}$ for any particular choice of \mathcal{O} , and again by [3, Lemma 1.2.1], $R_{\bar{r}}^\square \otimes \overline{\mathbb{Q}_p}$ is independent of the choice of E .

When K/\mathbb{Q}_p is a finite extension and Γ is Γ_K , we consider quotients of $R_{\bar{r}}^\square$ having fixed inertial type and p -adic Hodge type. The fundamental analysis here is due to Kisin ([17]), and the state of the art, and our point of reference, is [4], and we refer there for details. We will index p -adic Hodge types of deformations $r : \Gamma_K \rightarrow G(\mathcal{O})$ by collections $\underline{\mu}(r) = \{\mu(r, \tau)\}_{\tau : K \rightarrow \overline{\mathbb{Q}_p}}$ of (conjugacy classes of) Hodge–Tate co-characters, and write $R_{\bar{r}}^{\square, \underline{\mu}(r)}$ for the \mathbb{Z}_p -flat quotient of $R_{\bar{r}}^\square$ whose points in finite local E -algebras are precisely those of $R_{\bar{r}}^\square$ that are moreover potentially semi-stable with p -adic Hodge type $\underline{\mu}(r)$. We likewise consider the quotients with fixed inertial type σ , $R_{\bar{r}}^{\square, \underline{\mu}(r), \sigma}$, referring to [4, Section 3.2] for details.

We recall some basic facts on spin groups and spin representations, see [15, Lecture 20] for more details. Let $m \geq 3$ be an integer. Consider the symmetric form

$$(x, y) = x_1 y_m + x_2 y_{m-1} + \cdots + x_m y_1$$

on $V = \mathbb{Q}^m$ with associated quadratic form $Q(x) = (x, x)$. Let $C(Q)$ be the Clifford algebra associated to (V, Q) . It is equipped with an embedding $V \subset C(Q)$ which is universal for maps $f : V \rightarrow A$ into associative rings A satisfying $f(x)^2 = Q(x)$ for all $x \in V$. The algebra $C(Q)$ has a $\mathbb{Z}/2\mathbb{Z}$ -grading, $C(Q) = C(Q)^+ \oplus C(Q)^-$, induced from the grading on the tensor algebra. On the Clifford algebra $C(Q)$ we have an unique anti-involution $*$ that is determined by $(v_1 \cdots v_r)^* = (-1)^r v_r \cdots v_1$ for all $v_1, \dots, v_r \in V$. We define for all \mathbb{Q} -algebras R ,

$$\mathrm{GSpin}_m(R) = \{g \in (C(Q)^+ \otimes R)^\times : gVg^* \subset V\}.$$

Let $N : C(Q) \rightarrow \mathbb{Q}^\times$, $x \rightarrow xx^*$ be the Clifford norm. It induces a character $N : \mathrm{GSpin}_m \rightarrow \mathbb{G}_m$. We define Spin_m to be the kernel of N . The action of the group GSpin_m , resp. Spin_m stabilizes $V \subset C(Q)$, which induces a

surjection $\mathrm{GSpin}_m \twoheadrightarrow \mathrm{SO}_m$, resp. $\mathrm{Spin}_m \twoheadrightarrow \mathrm{SO}_m$. Denote by $\mu_2 = \{\pm 1\}$ the kernel of $\mathrm{Spin}_m \twoheadrightarrow \mathrm{SO}_m$.

Now we turn to spin representations. We keep the notation in the paragraph above with \mathbb{Q} replaced by \mathbb{C} . Let $\mathfrak{so}_m(\mathbb{C}) := \mathfrak{so}(Q) = \{X \in \mathrm{End}(V) : (Xv, w) + (v, Xw) = 0, \forall v, w \in V\}$. By [15, Lemma 20.7], there is a natural embedding of Lie algebras $\mathfrak{so}(Q) \rightarrow C(Q)^{\mathrm{even}}$. Let $n = \lfloor m/2 \rfloor$. The weight lattice of $\mathfrak{so}_m \mathbb{C}$ may be described as the submodule of $\sum_{1 \leq i \leq n} \mathbb{Q} \chi_i$ spanned by χ_1, \dots, χ_n and $\frac{1}{2}(\sum_i \chi_i)$. Suppose $m = 2n$ is even, write $V = W \oplus W'$, where W and W' are isotropic subspaces of (V, Q) . By [15, Lemma 20.9], there is an isomorphism of algebras $C(Q) \cong \mathrm{End}(\wedge^* W)$. We have $\wedge^* W = \wedge^{\mathrm{even}} W \oplus \wedge^{\mathrm{odd}} W$, it follows that $\mathfrak{so}(Q) \subset C(Q)^{\mathrm{even}} \cong \mathrm{End}(\wedge^{\mathrm{even}} W) \oplus \mathrm{End}(\wedge^{\mathrm{odd}} W)$. Hence we have two representations of $\mathfrak{so}_m \mathbb{C}$, $\wedge^{\mathrm{even}} W$ and $\wedge^{\mathrm{odd}} W$. We let $S = \wedge^{\mathrm{even}} W$ for even n , and let $S = \wedge^{\mathrm{odd}} W$ for odd n . By [15, Lemma 20.15], S is irreducible of dimension 2^{n-1} whose weights are $\frac{1}{2}(\sum_i \pm \chi_i)$ where the number of minus signs is even. We call S the spin representation of $\mathfrak{so}_m(\mathbb{C})$. Now suppose $m = 2n + 1$ is odd, write $V = W \oplus W' \oplus U$, where W and W' are isotropic subspaces of (V, Q) , and U is a one-dimensional subspace perpendicular to them. In this case, we have $\mathfrak{so}(Q) \subset C(Q)^{\mathrm{even}} \cong \mathrm{End}(\wedge^* W)$. We let $S = \wedge^* W$. By [15, Proposition 20.20], S is an irreducible representation of $\mathfrak{so}_m \mathbb{C}$ of dimension 2^n whose weights are $\frac{1}{2}(\sum_i \pm \chi_i)$. We again call S the spin representation of $\mathfrak{so}_m(\mathbb{C})$. The spin representation may be viewed as a group homomorphism $\mathrm{spin} : \mathrm{Spin}_m \rightarrow \mathrm{GL}_{2^n}$, resp. $\mathrm{spin} : \mathrm{Spin}_m \rightarrow \mathrm{GL}_{2^{n-1}}$ for odd m , resp. even m . The following fact is contained in [15, Exercise 20.38].

Lemma 1.3. *There is nondegenerate bilinear form β on S such that if $m = 2n+1$, β is symmetric when $n \equiv 0, 3 \pmod{4}$ and skew-symmetric otherwise; and if $m = 2n$ with n even, β is symmetric when $n \equiv 0 \pmod{4}$ and skew-symmetric otherwise. Therefore, if $m = 2n + 1$, the spin representation is valued in $\mathrm{SO}_{2^n} \mathbb{C}$ for $n \equiv 0, 3 \pmod{4}$ and valued in $\mathrm{Sp}_{2^n} \mathbb{C}$ for $n \equiv 1, 2 \pmod{4}$; and if $m = 2n$ with n even, the spin representation is valued in $\mathrm{SO}_{2^{n-1}} \mathbb{C}$ for $n \equiv 0 \pmod{4}$ and valued in $\mathrm{Sp}_{2^{n-1}} \mathbb{C}$ for $n \equiv 2 \pmod{4}$.*

2. Construction of residual Galois representations

Before we construct our mod p Galois representations into Spin_m , we make some preliminary calculations. We follow the notation of Section 1.1.

We have the following explicit description of the two-fold covering $\pi : \mathrm{Spin}_m \twoheadrightarrow \mathrm{SO}_m$ (see the proof of [15, Proposition 20.28]). Let $g \in \mathrm{SO}_m = \mathrm{SO}(Q)$, we may write g as a product of reflections $r_{w_1} \cdots r_{w_r}$ for $w_i \in V$ with $Q(w_i) = -1$ (r is necessarily even), then the two elements in $\pi^{-1}(g)$ are $\pm w_1 \cdots w_r$. We fix the following ordered basis for $V = \mathbb{Q}^m$: for $m = 2n$, $e_1, \dots, e_n, f_n, \dots, f_1$ satisfying $(e_i, f_i) = 1$ for $1 \leq i \leq n$, $(e_i, f_j) = 0$ for $1 \leq i \neq j \leq n$, and $(e_i, e_j) = (f_i, f_j) = 0$ for $1 \leq i, j \leq n$; for $m = 2n + 1$,

$e_1, \dots, e_n, u_0, f_n, \dots, f_1$ satisfying $(e_i, f_i) = 1$ for $1 \leq i \leq n$, $(e_i, f_j) = 0$ for $1 \leq i \neq j \leq n$, $(e_i, e_j) = (f_i, f_j) = 0$ for $1 \leq i, j \leq n$, $(u_0, u_0) = 1$, $(u_0, e_i) = (u_0, f_i) = 0$ for $1 \leq i \leq n$. With this basis, we have for any \mathbb{Q} -algebra R

$$\mathrm{SO}_m(R) = \{g \in \mathrm{SL}_m(R) : gJg^t = J\}$$

where $J = [a_{ij}]_{1 \leq i, j \leq m}$ whose nonzero entries are exactly $a_{i, m+1-i} = 1$ for $1 \leq i \leq m$. Then we have a standard maximal torus T_{SO} defined by $T_{\mathrm{SO}}(R) = \{\mathrm{diag}(t_1, \dots, t_n, 1, t_n^{-1}, \dots, t_1^{-1}) : t_i \in R^\times\}$ for any \mathbb{Q} -algebra R , where we omit the middle entry 1 when $m = 2n$ is even. We let $T = \pi^{-1}T_{\mathrm{SO}}$ be the corresponding maximal torus of Spin_m . The Weyl group W is then isomorphic to $N(T_{\mathrm{SO}})/T_{\mathrm{SO}}$, which is a semidirect product $D \rtimes S_n$, where D is described in what follows. We let D be $\{\pm 1\}^n$ when $m = 2n + 1$, and the subgroup of $\{\pm 1\}^n$ consisting of elements with an even number of minus signs when $m = 2n$. It acts on T_{SO} in the following way: For any element $\epsilon = (\epsilon_i) \in D$ and $t = \mathrm{diag}(t_1, \dots, t_n, 1, t_n^{-1}, \dots, t_1^{-1}) \in T_{\mathrm{SO}}$, define $t^\epsilon := \mathrm{diag}(t_1^{\epsilon_1}, \dots, t_n^{\epsilon_n}, 1, t_n^{-\epsilon_n}, \dots, t_1^{-\epsilon_1}) \in T_{\mathrm{SO}}$. The symmetric group S_n acts on T_{SO} by permuting the diagonal entries: $\forall \sigma \in S_n$, $t^\sigma := \mathrm{diag}(\sigma(t_1), \dots, \sigma(t_n), 1, \sigma(t_n)^{-1}, \dots, \sigma(t_1)^{-1})$.

Let w^0 be the longest element in W . When $m = 2n + 1$, w^0 acts as -1 on $X^\bullet(T_{\mathrm{SO}})$; when $m = 2n$, w^0 acts as -1 on $X^\bullet(T_{\mathrm{SO}})$ if and only if n is even. In both cases, $w^0 = (-1, \dots, -1) \in D$.

Lemma 2.1. *The group $D \leq W$ acts simply transitively on $\Lambda_{\mathrm{spin}} := \{\frac{1}{2}(\sum_{i=1}^n \pm \chi_i)\}$, the weights of the spin representation of Spin_m .*

Proof. This follows immediately from the explicit action of D on T described above. □

Lemma 2.2. *The group homomorphism $N(T_{\mathrm{SO}}) \rightarrow W$ admits a section $s : W \rightarrow N(T_{\mathrm{SO}})$. Indeed, for $\sigma \in S_n$, let M_σ be the corresponding standard $n \times n$ permutation matrix, then $s(\sigma) = \begin{pmatrix} M_\sigma & 0 \\ 0 & M_\sigma^{at} \end{pmatrix}$, where M_σ^{at} is the anti-diagonal transpose of M_σ ; for $\epsilon = (\epsilon_i) \in D$, $s(\epsilon) = d_\epsilon$, where d_ϵ is the $m \times m$ matrix obtained from the identity matrix I_m by swapping the i -th and the $(m + 1 - i)$ -th columns whenever $\epsilon_i = -1$ and leaving the rest of the columns unchanged when $m = 2n$ is even, and d_ϵ is the $m \times m$ matrix obtained from the identity matrix I_m by performing the column operations above and replacing the $(n + 1, n + 1)$ -entry with $(-1)^{|\{1 \leq i \leq n: \epsilon_i = -1\}|}$ when $m = 2n + 1$ is odd.*

By the lemma above, we may identify D with its isomorphic image in $N(T_{\mathrm{SO}})$. With this identification, let $\tilde{D} = \pi^{-1}D \subset \mathrm{Spin}_m$. We would like to calculate the structure of \tilde{D} . Suppose $m = 2n$ is even, for each $i \in [1, n]$, let $w_i = \frac{1}{\sqrt{2}}(e_i - f_i)$. A straightforward calculation shows that $Q(w_i) = -1$ and $r_{w_i} \in O_m$ equals the matrix obtained by swapping the i -th and the

$(m + 1 - i)$ -th columns of I_m . It follows that D is generated by $r_{w_i}r_{w_j}$ for $1 \leq i < j \leq n$, and hence \tilde{D} is generated by w_iw_j for $1 \leq i < j \leq n$. Note that $w_i^2 = -1$ and $w_iw_j = -w_jw_i$ for $i \neq j$, which imply that

$$(w_1 \cdots w_k)^2 = (-1)^{k(k+1)/2}$$

for $1 \leq k \leq n$. Now suppose $m = 2n + 1$ is odd, define w_i as above and let $w_0 := \sqrt{-1}u_0$, then $r_{w_0}r_{w_i} \in \mathrm{SO}_m$ equals the matrix obtained from I_m by swapping the i -th and the $(m + 1 - i)$ -th columns and replacing the $(n + 1, n + 1)$ -entry with -1 . It follows that $r_{w_0}r_{w_i}$, $1 \leq i \leq n$ generate D and hence $\omega_i := w_0w_i$, $1 \leq i \leq n$ generate \tilde{D} . Note that $\omega_i^2 = -1$ and $\omega_i\omega_j = -\omega_j\omega_i$ for $i \neq j$, which imply that

$$(\omega_1 \cdots \omega_k)^2 = (-1)^{k(k+1)/2}$$

for $1 \leq k \leq n$. We thus obtain

Lemma 2.3. *The sequence $1 \rightarrow \{\pm 1\} \rightarrow \tilde{D} \rightarrow D \rightarrow 1$ is nonsplit. Moreover, if $m = 2n$, w^0 acts as -1 on $X^\bullet(T)$ if and only if n is even, in which case it corresponds to the element $w_1 \cdots w_n \in \mathrm{Spin}_m$ under the map $N(T) \rightarrow W$. This element has order two if and only if $n \equiv 0 \pmod{4}$. If $m = 2n + 1$, w^0 always acts as -1 on $X^\bullet(T)$, and the corresponding element $\omega_1 \cdots \omega_n \in \mathrm{Spin}_m$ has order two if and only if $n \equiv 0, 3 \pmod{4}$.*

For the rest of this section, we make the following *parity assumption* on the integer m :

- If $m = 2n$ is even, then $n \equiv 0 \pmod{4}$;
- If $m = 2n + 1$ is odd, then $n \geq 3$ and $n \equiv 0, 3 \pmod{4}$;

equivalently, $m \geq 7$ and $m \equiv 0, 1, 7 \pmod{8}$.

By Lemma 2.3, $w^0 \in W$ then lifts to an order two element in $N(T)$; and it follows from Lemma 1.3 that the spin representation is

$$\mathrm{spin} : \mathrm{Spin}_m \rightarrow \mathrm{SO}_N$$

where $N = 2^n$ for odd m and $N = 2^{n-1}$ for even m .

We now begin to construct the mod p representation valued in Spin_m . We do this by first realizing \tilde{D} as a Galois group of some finite extension of \mathbb{Q} , and then modify the corresponding homomorphism $\Gamma_{\mathbb{Q}} \rightarrow N(T)_{\mathbb{F}_p} \subset \mathrm{Spin}_m(\mathbb{F}_p)$ so that after composing with spin , the resulting representation satisfies the assumptions of the potential automorphy theorems of [3].

Lemma 2.4. *There is a finite, totally real Galois extension L/\mathbb{Q} whose Galois group is isomorphic to D satisfying property (S_N) with $N = 2$ in the sense of [30, Definition 2.1.2], i.e. every prime p which is ramified in L/\mathbb{Q} satisfies*

- $p \equiv 1 \pmod{4}$.

- If v is a place of L above p , then the local extension L_v/\mathbb{Q}_p is totally ramified.

Proof. We have $D \cong (\mathbb{Z}/2\mathbb{Z})^n$ or $(\mathbb{Z}/2\mathbb{Z})^{n-1}$. We pick a prime $p_1 \equiv 1 \pmod{4}$. We will construct inductively p_1, \dots, p_n such that $p_i \equiv 1 \pmod{4}$ and $p_i \in (\mathbb{Q}_{p_j}^\times)^2$ for $i \neq j$. Suppose we have p_1, \dots, p_k ($k < n$) already and consider $\mathbb{Q}(\sqrt{-1}, \sqrt{p_1}, \dots, \sqrt{p_k})$. Chebotarev implies that there are infinitely many p that splits in this field. In particular, $-1, p_1, \dots, p_k \in (\mathbb{Q}_p^\times)^2$, which is equivalent to

$$\left(\frac{-1}{p}\right) = \left(\frac{p_1}{p}\right) = \dots = \left(\frac{p_k}{p}\right) = 1.$$

In particular, $p \equiv 1 \pmod{4}$ and quadratic reciprocity implies that

$$\left(\frac{p}{p_1}\right) = \dots = \left(\frac{p}{p_k}\right) = 1,$$

which is equivalent to $p \in (\mathbb{Q}_{p_j}^\times)^2$ for $1 \leq j \leq k$. Thus, if we take p_{k+1} to be such a prime, the induction can proceed. Now we let $L = \mathbb{Q}(\sqrt{p_1}, \dots, \sqrt{p_n})$. Then $\text{Gal}(L/\mathbb{Q}) = D$ and $L_{v_i} = \mathbb{Q}_{p_i}(\sqrt{p_i})$, which is a totally ramified extension of \mathbb{Q}_{p_i} . \square

Let $\underline{s} : \Gamma_{\mathbb{Q}} \rightarrow \text{Gal}(L/\mathbb{Q}) = D$ be the corresponding homomorphism produced by the above lemma. We need to lift \underline{s} to a surjective homomorphism $s : \Gamma_{\mathbb{Q}} \rightarrow \tilde{D}$ across the short exact sequence $1 \rightarrow \mu_2 \rightarrow \tilde{D} \rightarrow D \rightarrow 1$. The method we use is a minor modification of the argument in page 12 of [30]. Since the exact sequence of groups is nonsplit by Lemma 2.3, it suffices to find a lift $s : \Gamma_{\mathbb{Q}} \rightarrow \tilde{D}$ of \underline{s} , which will automatically be surjective. Suppose that \underline{s} has a lift to \tilde{D} everywhere locally (including ∞), then \underline{s} has a lift to \tilde{D} . In fact, \underline{s} induces a map $\underline{s}^* : H^2(D, \mu_2) \rightarrow H^2(\Gamma_{\mathbb{Q}}, \mu_2)$, and \underline{s} lifts to \tilde{D} if and only if $\underline{s}^*(\xi) = 0$, where ξ is the class of the extension $\tilde{D} \rightarrow D$. The natural map

$$H^2(\Gamma_{\mathbb{Q}}, \mu_2) \rightarrow \prod_{p \leq \infty} H^2(\Gamma_{\mathbb{Q}_p}, \mu_2)$$

is injective because $H^2(\Gamma_{\mathbb{Q}}, \mu_2) = \text{Br}(\mathbb{Q})[2]$ and $\text{Br}(\mathbb{Q}) \rightarrow \prod_{p < \infty} \text{Br}(\mathbb{Q}_p)$. It follows that if \underline{s} lifts everywhere locally, then it lifts globally. It thus remains to show that \underline{s} lifts everywhere locally. For $p < \infty$, $\underline{s}|_{\Gamma_{\mathbb{Q}_p}}$ lifts to \tilde{D} follows from property (S_2) by the exact same argument as in the last two paragraphs of page 12 of [30]; for $p = \infty$, $\underline{s}|_{\Gamma_{\mathbb{R}}}$ obviously lifts to \tilde{D} because $\underline{s}(c) = 1$ (c is the complex conjugation). Therefore, we obtain a surjection

$$s : \Gamma_{\mathbb{Q}} \twoheadrightarrow \tilde{D} \subset N(T)_{\mathbb{Z}}.$$

We would like to modify s so that $s(c)$ belongs to T and is “sufficiently odd” in the sense that $\dim(\mathfrak{so}_m)^{\text{Ad}s(c)} = |\Phi|/2$, where Φ is the root system

of \mathfrak{so}_m . When $s(c) \in \text{Spin}_m$ satisfies this equality, we call it a *split Cartan involution*, or a *Chevalley involution* in Spin_m .

Let us fix a Borel subgroup $B \supset T$ of Spin_m . It corresponds to a set of simple roots $\Delta \subset \Phi := \Phi(\text{Spin}_m, T)$. Let ρ^\vee be the half sum of positive coroots in Φ^\vee . By examining the Plates in [6], if Φ is of type B_n , ρ^\vee has coefficients in \mathbb{Z} if and only if $n \equiv 0, 3 \pmod 4$; and if Φ is of type D_n , ρ^\vee has coefficients in \mathbb{Z} if and only if $n \equiv 0, 1 \pmod 4$. In particular, ρ^\vee has coefficients in \mathbb{Z} for $m = 2n$ or $2n + 1$ satisfying the parity assumption we made before. It follows that for those m , $\rho^\vee \in X_\bullet(T)$, i.e. it is a well-defined cocharacter of T .

Let k be a field and G be an algebraic group defined over k , we denote by G_k or $G(k)$ the k -points of G . Observe that if $s : \Gamma_{\mathbb{Q}} \rightarrow N(T)_k$ is a group homomorphism and $\phi \in H^1(\Gamma_{\mathbb{Q}}, T_k)$, where $\Gamma_{\mathbb{Q}}$ acts on T_k via the composite $\text{Ad} \circ s$, then $\phi \cdot s : \Gamma_{\mathbb{Q}} \rightarrow N(T)_k$ is a group homomorphism.

Recall that we have an isomorphism $\mathbb{G}_m^n \xrightarrow{\sim} T_{SO}$,

$$(t_1, \dots, t_n) \mapsto \text{diag}(t_1, \dots, t_n, 1, t_n^{-1}, \dots, t_1^{-1})$$

(where we omit the middle 1 when m is even). We also have an isomorphism

$$T \xrightarrow{\sim} \{(z, t_1, \dots, t_n) \in \mathbb{G}_m^{n+1} : z^2 = t_1 \cdots t_n\}.$$

Let $T[2]$ be the group of \mathbb{Z} -points of T . It consists of elements of the form $(z, \epsilon_1, \dots, \epsilon_n)$, where $z = \pm 1$, $\epsilon_i = \pm 1$ and the number of -1 among ϵ_i is even. The group $\Gamma_{\mathbb{Q}}$ acts on $T[2]$ via $\text{Ad} \circ s$. This action factors through $\text{Gal}(L/\mathbb{Q}) \cong D$. Then $T[2]$ is a direct sum of two irreducible D -modules, $V_1 = \{(1, \epsilon_1, \dots, \epsilon_n) : \epsilon_i = \pm 1\}$ and $V_2 = \{(-1, \epsilon_1, \dots, \epsilon_n) : \epsilon_i = \pm 1\}$. Letting $S = \text{cs}(L/\mathbb{Q}) \cup \{\infty, 2\}$ ($\text{cs}(L/\mathbb{Q})$ denotes the set of places of \mathbb{Q} that split completely in L), [23, Theorem 9.2.3(v)] implies that for $i = 1, 2$, the natural maps

$$H^1(\Gamma_{\mathbb{Q}, S}, V_i) \rightarrow H^1(\Gamma_{\mathbb{R}}, V_i)$$

are surjective, and hence we have a surjection

$$H^1(\Gamma_{\mathbb{Q}, S}, T[2]) \twoheadrightarrow H^1(\Gamma_{\mathbb{R}}, T[2]) = \text{Hom}(\Gamma_{\mathbb{R}}, T[2])$$

(recall that $s(c) \in \mu_2$). Let $\phi \in H^1(\Gamma_{\mathbb{Q}, S}, T[2])$ be a class that maps to $c \mapsto s(c)\rho^\vee(-1)$ in $\text{Hom}(\Gamma_{\mathbb{R}}, T[2])$. We now replace s by $\phi \cdot s$. Then *having made this replacement*, $s(c) = \rho^\vee(-1)$, which is a split Cartan involution by [32, Lemma 2.3]. Let E be the fixed field of $\ker s$.

Now we are going to further modify s by an element in $H^1(\Gamma_{\mathbb{Q}}, T_{\mathbb{F}_p})$ (for a prime p) so that the resulting homomorphism $\Gamma_{\mathbb{Q}} \rightarrow N(T)_{\mathbb{F}_p} \subset \text{Spin}_m(\mathbb{F}_p)$ satisfies favorable conditions so that we can use the results of [3] to deform it to a geometric representation. The method we use here is extremely close to the proof of [7, Proposition 2.8]. We first establish some notation. For a fixed prime p , we write κ for the p -adic cyclotomic character and $\bar{\kappa}$ for its mod p reduction. For a prime l dividing $p - 1$ such that l^2 does not divide

$p - 1$, let $\text{pr}(l)$ be the canonical projection from \mathbb{F}_p^\times onto the l -torsion subgroup $\mathbb{F}_p^\times[l]$. Let $\bar{\kappa}[l] = \text{pr}(l) \circ \bar{\kappa} : \Gamma_{\mathbb{Q}} \rightarrow \mathbb{F}_p^\times[l]$.

Proposition 2.5. *Consider pairs of primes (l, p) such that p splits in E/\mathbb{Q} and $p - 1$ is divisible by l but not l^2 . Then there exist infinitely many primes l such that there exist infinitely many pairs (l, p) such that there exists a homomorphism*

$$\bar{r} : \Gamma_{\mathbb{Q}} \rightarrow T_{\mathbb{F}_p}[l] \cdot \tilde{D} \subset N(T)_{\mathbb{F}_p}$$

satisfying

- (1) $\bar{r}(c) = \rho^\vee(-1)$.
- (2) At p , fix any choice of integers n_α for $\alpha \in \Delta$, $\bar{r}|_p = \prod_{\alpha \in \Delta} \alpha^\vee \circ \bar{\kappa}[l]^{n_\alpha}$.
- (3) There is a prime q of order $l \pmod p$ such that $\bar{r}|_q$ is unramified with Frobenius mapping to $\rho^\vee(q) \in T_{\mathbb{F}_p}[l]$.

Moreover, in addition to the above, we can choose $p > 2(N + 1)$, $l \geq h_{\text{Spin}_m}$ (h_{Spin_m} is the Coxeter number of Spin_m), $\{n_\alpha\}$ and \bar{r} such that

- (4) $\text{spin} \circ \bar{r}|_p$ is a direct sum of distinct powers of $\bar{\kappa}[l]$.
- (5) For any Borel $B \supset T$, $H^0(\Gamma_{\mathbb{Q}_p}, \bar{r}(\mathfrak{g}/\mathfrak{b}))$ vanishes (where $\mathfrak{g} = \mathfrak{so}_m$ and \bar{r} acts on $\mathfrak{g}/\mathfrak{b}$ via the adjoint action).
- (6) $\text{spin} \circ \bar{r}|_{\Gamma_{\mathbb{Q}}(\mu_p)}$ is absolutely irreducible.

Proof. Let l be any odd prime such that E (the fixed field of $\ker s$) and $\mathbb{Q}(\mu_{l^2})$ are linearly disjoint over \mathbb{Q} (e.g. take l split in E/\mathbb{Q}). Later in the argument we will require l to be larger than some constant depending only on the group Spin_m . Now we take p to be any prime split in $E(\mu_l)$ but nonsplit in $E(\mu_{l^2})$ (such a prime exists by Chebotarev). In other words, p splits in E/\mathbb{Q} and $p - 1$ is divisible by l but not l^2 . In particular, E and $\mathbb{Q}(\mu_p)$ are linearly disjoint over \mathbb{Q} . By Chebotarev, there exists a prime q that splits in E and has order l in \mathbb{F}_p^\times . Let r be a prime that splits in $E(\mu_p)$. The Galois group $\Gamma_{\mathbb{Q}}$ acts on $T_{\mathbb{F}_p}[l]$ via $\text{Ad} \circ s$, which factors through $D \cong \text{Gal}(L/\mathbb{Q})$. Then $T_{\mathbb{F}_p}[l]$ decomposes into a direct sum of irreducible $\mathbb{F}_l[D]$ -modules: $T_{\mathbb{F}_p}[l] = \bigoplus W_i$. Let $\Sigma = \text{cs}(L/\mathbb{Q}) \cup \{\infty, l\}$ and $T = \{p, q, r\}$. Then [23, Theorem 9.2.3(v)] implies that the canonical homomorphisms

$$H^1(\Gamma_{\mathbb{Q}, \Sigma}, W_i) \rightarrow \bigoplus_{v \in T} H^1(\Gamma_{\mathbb{Q}_v}, W_i)$$

are surjective, from which it follows that the canonical homomorphism

$$H^1(\Gamma_{\mathbb{Q}, \Sigma}, T_{\mathbb{F}_p}[l]) \rightarrow \bigoplus_{v \in T} H^1(\Gamma_{\mathbb{Q}_v}, T_{\mathbb{F}_p}[l])$$

is surjective as well. We let $\phi \in H^1(\Gamma_{\mathbb{Q}, \Sigma}, T_{\mathbb{F}_p}[l])$ be a class such that $\phi|_p = \prod_{\alpha \in \Delta} \alpha^\vee \circ \bar{\kappa}[l]^{n_\alpha}$, $\phi|_q$ is unramified with $\phi(\text{Fr}_q) = \rho^\vee(q)$, and $\phi|_r$ is unramified with $\phi(\text{Fr}_r) = t$, where t is any element in $T_{\mathbb{F}_p}[l]$ such that $\lambda(t)$ for $\lambda \in \Lambda_{\text{spin}}$ are all distinct (such t exists provided that l is sufficiently

large). We set $\bar{r} = \phi \cdot s$. It follows immediately from the choice of ϕ that (2) and (3) hold. Since $s(c) = \rho^\vee(-1)$ and l is odd, (1) holds. It remains to address (4) through (6). We have

$$\text{spin} \circ \bar{r}|_p = \bigoplus_{\lambda \in \Lambda_{spin}} \bar{\kappa}[l] \sum_{\alpha \in \Delta} n_\alpha \langle \lambda, \alpha^\vee \rangle;$$

since Λ_{spin} is multiplicity-free, choosing $\{n_\alpha\}$ to be sufficiently general will make the exponents distinct. This proves (4). For (5), note that

$$\bar{r}|_p(\mathfrak{g}/\mathfrak{b}) = \bigoplus_{\beta \in \Phi^-} \bar{\kappa}[l] \sum_{\alpha \in \Delta} n_\alpha \langle \beta, \alpha^\vee \rangle.$$

By choosing l to be sufficiently large and $\{n_\alpha\}$ to be sufficiently general, we can ensure that the absolute values of the exponents are in $(0, l)$; in particular, $H^0(\Gamma_{\mathbb{Q}_p}, \bar{r}(\mathfrak{g}/\mathfrak{b}))$ vanishes. Finally, D acts transitively on Λ_{spin} by Lemma 2.1, so any nonzero submodule of $\text{spin} \circ \bar{r}$ has nonzero projection to each of the weight spaces; since E and $\mathbb{Q}(\mu_p)$ are linearly disjoint over \mathbb{Q} , the same is true for $\text{spin} \circ \bar{r}|_{\Gamma_{\mathbb{Q}(\mu_p)}}$. On the other hand, $\bar{r}(\text{Fr}_r) = t \in T_{\mathbb{F}_p}[l]$ acts via distinct characters on the different weight spaces of the spin representation by the choice of t . It follows that $\text{spin} \circ \bar{r}|_{\Gamma_{\mathbb{Q}(\mu_p)}}$ is absolutely irreducible, which shows (6). \square

3. Lifting Galois representations

In this section, we prove a lifting theorem using a version of the Khare–Wintenberger argument ([16]), similar to the arguments in [7, Section 3] and [25, Section 3]. We axiomatize their arguments and state our lifting theorem as general as we can. On the other hand, the main theorem of [14] allows one to deform a much larger class of G -valued mod p Galois representations to geometric representations in characteristic zero, yet it does not establish potential automorphy of the lifts.

Let G be a split connected semisimple group over \mathbb{Q} . Let T be a maximal split torus of G and let $\Phi = \Phi(G, T)$ be the root system of G . We assume that G satisfies the following:

- G contains a split Cartan involution, i.e. there is an element $\tau \in G$ of order two such that $\dim \mathfrak{g}^{\text{Ad}\tau} = |\Phi|/2$.
- G admits an irreducible faithful representation $R : G \rightarrow \text{GL}_N$ factoring through SO_N (the orthogonal group preserving the pairing $(x, y) = x_1y_1 + x_2y_2 + \cdots + x_Ny_N$ on \mathbb{Q}^N ; this is different from the pairing used in Section 1.1) such that the formal character of R is multiplicity-free.⁴

⁴This is a very restrictive assumption.

Let F^+ be a totally real field. Let $\bar{r} : \Gamma_{F^+} \rightarrow G(\overline{\mathbb{F}}_p)$ be a continuous representation. Let S be the union of all the archimedean places of F^+ , the places of F^+ dividing p and the places of F^+ at which \bar{r} is ramified. Suppose that \bar{r} satisfies the following:

- \bar{r} is odd in the sense that for $v|\infty$ with $c_v \in \Gamma_{F_v^+}$ the complex conjugation, $\bar{r}(c_v)$ is a split Cartan involution.
- For v not above p , $\bar{r}|_v$ has a characteristic zero lift $r_v : \Gamma_{F_v^+} \rightarrow G(\overline{\mathbb{Z}}_p)$.
- For $v|p$, $\bar{r}|_v$ has a characteristic zero lift $r_v : \Gamma_{F_v^+} \rightarrow G(\overline{\mathbb{Z}}_p)$ with some fixed Hodge type $\underline{\mu}(r_v)$ and inertial type σ_v such that $R \circ r_v$ is potentially diagonalizable in the sense of [3] with regular Hodge–Tate weights.⁵
- $R \circ \bar{r}|_{F^+(\mu_p)}$ is absolutely irreducible.
- p is sufficiently large relative to G .

We let \mathcal{O} be the ring of integers in a finite extension E of \mathbb{Q}_p , enlarged if necessary so that all of the above data are defined over \mathcal{O} .

We recall the definition of the Clozel–Harris–Taylor group scheme \mathcal{G}_n over \mathbb{Z} which is defined as the semidirect product $(\mathrm{GL}_n \times \mathrm{GL}_1) \rtimes \{1, j\}$ where $j(g, a)j = (a({}^t g)^{-1}, a)$, and the similitude character $\nu : \mathcal{G}_n \rightarrow \mathrm{GL}_1$ given by $\nu(g, a) = a$ and $\nu(j) = -1$. Suppose we have a homomorphism $r : \Gamma_{F^+} \rightarrow G(A)$ for some ring A . Let F/F^+ be a quadratic extension of F , and define $\rho(r) : \Gamma_{F^+} \rightarrow \mathcal{G}_N(R)$ as the composite

$$\Gamma_{F^+} \xrightarrow{R(r) \times \mathrm{res}_F} \mathrm{SO}_N(A) \times \mathrm{Gal}(F/F^+) \longrightarrow \mathcal{G}_N(A),$$

where the last map sends $g \in \mathrm{SO}_N(A)$ to its image in $\mathrm{GL}_N(A)$ and sends the nontrivial element of $\mathrm{Gal}(F/F^+)$ to j . By our choice of the pairing defining SO_N , $\rho(r)$ is a well-defined homomorphism.

We choose a quadratic CM extension F/F^+ such that F does not contain ζ_p , $R \circ \bar{r}|_{F(\mu_p)}$ remains irreducible, and all the finite places in S split in F/F^+ . We will define global deformation conditions for \bar{r} and $\rho(\bar{r})$ with respect to F/F^+ . Then we will show that there are natural finite maps between the corresponding deformation rings and use the \mathcal{O} –finiteness of the deformation ring for the group \mathcal{G}_N to conclude that the deformation ring for the group G is \mathcal{O} –finite; this, combined with a standard calculation of its Krull dimension (as in [4]), will imply that it has a $\overline{\mathbb{Q}}_p$ –point. For the \mathcal{G}_N –deformation problems, we will fix the multiplier: Let $\delta_{F/F^+} : \Gamma_{F^+} \rightarrow \{\pm 1\}$ be the quadratic character associated with F/F^+ ; we require that any local or global deformation of $\rho(\bar{r})$ composed with ν equals δ_{F/F^+} . In particular, since all the finite places $v \in S$ split in F/F^+ , $\rho(\bar{r})|_v$ and all its local

⁵For the regularity to hold, it is necessary that the formal character of R is multiplicity-free.

deformations are valued in the group GL_N . Thus we will omit δ_{F/F^+} from our notation when we discuss about local deformations.

Recall from the discussion in Section 1.1 that for each $v \in S$ we can consider the lifting rings $R_{\bar{r}|_v}^\square$ and $R_{\rho(\bar{r})|_v}^\square$. For $v \in S$ not above p , choose an irreducible component $\mathcal{C}(r_v)$, resp. $\mathcal{C}(\rho(r_v))$ of $R_{\bar{r}|_v}^\square \otimes \overline{\mathbb{Q}}_p$, resp. $R_{\rho(\bar{r})|_v}^\square \otimes \overline{\mathbb{Q}}_p$ containing r_v , resp. $\rho(r_v)$, such that under the natural map

$$\mathrm{Spec} R_{\bar{r}|_v}^\square \otimes \overline{\mathbb{Q}}_p \rightarrow \mathrm{Spec} R_{\rho(\bar{r})|_v}^\square \otimes \overline{\mathbb{Q}}_p,$$

$\mathcal{C}(r_v)$ maps to $\mathcal{C}(\rho(r_v))$.

Similarly, for $v|p$, the fixed inertial type and fixed p -adic Hodge type of r_v induces corresponding data for $\rho(r_v)$. Choose $\mathcal{C}(r_v)$ to be a potentially crystalline component mapping into a potentially crystalline component $\mathcal{C}(\rho(r_v))$ containing the potentially diagonalizable point $\rho(r_v)$ under the natural map

$$\mathrm{Spec} R_{\bar{r}|_v}^{\square, \mu(r_v), \sigma_v} \otimes \overline{\mathbb{Q}}_p \rightarrow \mathrm{Spec} R_{\rho(\bar{r})|_v}^{\square, \mu(\rho(r_v)), \rho(\sigma_v)} \otimes \overline{\mathbb{Q}}_p.$$

We now define the global deformation rings, for \bar{r} and $\rho(\bar{r})$, by considering lifts that locally lie on the irreducible components we have just specified. More precisely, following the formalism of [4, Section 4.2], we let R_G^{univ} be the quotient of the universal, unramified outside S deformation ring for \bar{r} corresponding to the fixed set of components $\{\mathcal{C}(r_v)\}_{v \in S}$. We similarly define $R_{\mathrm{GL}}^{\mathrm{univ}}$ corresponding to the local components $\{\mathcal{C}(\rho(r_v))\}$ (and fixed polarization δ_{F/F^+}). These rings all exist by absolute irreducibility of the respective residual representations, and by the discussion in [4, Section 4.2] ([4, Lemma 3.4.1] plays a key role here). By construction, there is a natural \mathcal{O} -algebra map

$$R_{\mathrm{GL}}^{\mathrm{univ}} \rightarrow R_G^{\mathrm{univ}}.$$

Lemma 3.1. *The map $R_{\mathrm{GL}}^{\mathrm{univ}} \rightarrow R_G^{\mathrm{univ}}$ is surjective.*

Proof. The tangent space of $R_{\mathrm{GL}}^{\mathrm{univ}}$ is a subspace of $H^1(\Gamma_{F^+, S}, \rho(\bar{r})(\mathfrak{gl}_N))$ and the tangent space of R_G^{univ} is a subspace of $H^1(\Gamma_{F^+, S}, \bar{r}(\mathfrak{g}))$. For p large enough, \mathfrak{gl}_N is by [29, Proposition 2] a semisimple G -module, so a fortiori $\bar{r}(\mathfrak{g})$ is a $\Gamma_{F^+, S}$ -direct summand of $\rho(\bar{r})(\mathfrak{gl}_N)$. It follows that the natural map $H^1(\Gamma_{F^+, S}, \bar{r}(\mathfrak{g})) \rightarrow H^1(\Gamma_{F^+, S}, \rho(\bar{r})(\mathfrak{gl}_N))$ is injective; the dual map is surjective, and we conclude by Nakayama's lemma that the map on universal deformation rings without local conditions is surjective. It then follows immediately that $R_{\mathrm{GL}}^{\mathrm{univ}} \rightarrow R_G^{\mathrm{univ}}$ is surjective as well. \square

By the proof of [3, Theorem 4.3.1], $R_{\mathrm{GL}}^{\mathrm{univ}}$ is \mathcal{O} -finite. Lemma 3.1 then implies that R_G^{univ} is \mathcal{O} -finite. We claim that R_G^{univ} has Krull dimension at least one. Indeed, this follows from [4, Theorem B]: the assumptions

there are satisfied since we have assumed that \bar{r} is odd, $R \circ \bar{r}|_{F+(\mu_p)}$ is irreducible (which, under our assumption on p , implies that $H^0(\Gamma_{F^+}, \bar{r}(\mathfrak{g})(1))$ vanishes), and the Hodge–Tate cocharacters are regular. Thus R_G^{univ} has a $\overline{\mathbb{Q}}_p$ -point, say $r : \Gamma_{F^+} \rightarrow G(\overline{\mathbb{Q}}_p)$. Then $R \circ r$ satisfies the hypotheses in [3, Theorem C]: (1) is clear; (2) holds since R is valued in SO_N by assumption; (3) holds because for $v|p$, $R \circ r|_v$ lies on the same (potentially crystalline) component as the potentially diagonalizable point $R \circ r_v = \rho(r_v)$; since potentially crystalline deformation rings are regular, there is a unique potentially crystalline component of $\text{Spec } R_{\rho(\bar{r})|_v}^{\square, \mu(\rho(r_v)), \rho(\sigma_v)} \otimes \overline{\mathbb{Q}}_p$ passing through $\rho(r_v)$. It follows that $R \circ r|_v$ is potentially diagonalizable. (4) holds by our assumption on \bar{r} . Therefore, Theorem C of loc. cit. implies that $R \circ r$ is potentially automorphic and belongs to a strictly compatible system of ℓ -adic representations. Let us denote this compatible system by $\{R_\lambda\}$, where $R_\lambda : \Gamma_{F^+} \rightarrow \text{GL}_N(\overline{M}_\lambda)$ is a continuous representation with M a number field and λ primes of M . Suppose that for $v|p$, the Hodge–Tate weights of $R \circ r_v$ (and hence $R \circ r|_v$) are extremely regular in the sense of [3, Section 2.1], then [3, Theorem D] (which ultimately relies on Larsen’s work [20]) and the standard Brauer induction argument (see [3, Theorem 5.5.1]) imply that for a density one set of rational primes l , R_λ is irreducible for $\lambda|l$.

We summarize the above discussion in the following:

Theorem 3.2. *Retain the assumptions on G and $\bar{r} : \Gamma_{F^+, S} \rightarrow G(\overline{\mathbb{F}}_p)$ imposed at the beginning of this section. Then \bar{r} has a lift $r : \Gamma_{F^+} \rightarrow G(\overline{\mathbb{Q}}_p)$ unramified outside S such that*

- (1) *For each $v \in S$ not lying above p , $r|_v := r|_{\Gamma_{F_v^+}}$ and r_v lie on the same irreducible component of $\text{Spec}(R_{\bar{r}|_v}^{\square} \otimes \overline{\mathbb{Q}}_p)$.*
- (2) *For each $v|p$, $r|_v$ and r_v lie on the same irreducible component of $\text{Spec } R_{\bar{r}|_v}^{\square, \mu(r_v), \sigma_v} \otimes \overline{\mathbb{Q}}_p$. Moreover, $R \circ r|_v$ is potentially diagonalizable in the sense of [3] with regular Hodge–Tate weights.*
- (3) *$R \circ r$ is potentially automorphic in the sense of [3].*
- (4) *$R \circ r$ is part of a strictly compatible system $R_\lambda : \Gamma_{F^+} \rightarrow \text{GL}_N(\overline{M}_\lambda)$ indexed by finite places λ of some number field M . If in addition, the Hodge–Tate weights of $R \circ r|_v$ are extremely regular in the sense of [3, Section 2.1], then for a density one set of rational primes l , R_λ is irreducible for $\lambda|l$.*

4. Compatible systems of Spin Galois representations

In this section, we use the lifting method of Section 3 to deform the representation $\bar{r} : \Gamma_{\mathbb{Q}} \rightarrow \text{Spin}_m(\mathbb{F}_p)$ in Proposition 2.5 to a characteristic zero representation satisfying favorable local conditions. Moreover, we will

show that the associated compatible system of Galois representations $\{R_\lambda\}$ has Spin_m monodromy for a density one set of rational primes l below λ .

For the rest of this paper, we assume that m satisfies the parity assumption in Section 2.

We need to specify the local deformations of \bar{r} , following the notation in Section 3. Let S be the union of ∞, p, q and the places of \mathbb{Q} at which \bar{r} is ramified. We choose a quadratic field F such that F does not contain ζ_p , $\mathrm{spin} \circ \bar{r}|_{F(\mu_p)}$ remains irreducible, and all the finite places in S split in F/\mathbb{Q} . Suppose that we have a homomorphism $r : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(A)$ for some ring A ; as in Section 3, we define $\rho(r) : \Gamma_{\mathbb{Q}} \rightarrow \mathcal{G}_N(A)$ as the composite (recall that the spin representation lands in SO_N for m satisfying the parity assumption)

$$\Gamma_{\mathbb{Q}} \xrightarrow{\mathrm{spin}(r) \times \mathrm{res}_F} \mathrm{SO}_N(A) \times \mathrm{Gal}(F/\mathbb{Q}) \longrightarrow \mathcal{G}_N(A),$$

where the last map sends $g \in \mathrm{SO}_N(A)$ to its image in $\mathrm{GL}_N(A)$ and sends the nontrivial element of $\mathrm{Gal}(F/\mathbb{Q})$ to j .

Deformation condition at p . Recall that $\bar{r}|_p = \prod_{\alpha \in \Delta} \alpha^\vee \circ \bar{\kappa}[l]^{n_\alpha}$. Fix

$$\chi_T = \prod_{\alpha \in \Delta} \alpha^\vee \circ (\kappa^{\tilde{n}_\alpha} \cdot [\bar{\kappa}]^{-n_\alpha} \cdot [\bar{\kappa}[l]^{n_\alpha})$$

lifting $\bar{r}|_p$, where $[\bar{\kappa}]$, resp. $[\bar{\kappa}[l]]$ denotes the Teichmüller lift of $\bar{\kappa}$, resp. $[\bar{\kappa}[l]]$, and $\tilde{n}_\alpha \equiv n_\alpha \pmod{l-1}$ are greater than one and sufficiently general ($\tilde{n}_\alpha > 1$ ensures that our characteristic zero lifts are potentially crystalline, cf. [24, Lemma 4.8]). We take r_p to be χ_T . Let $\mathcal{C}(r_p)$ be an ordinary potentially crystalline component of $\mathrm{Spec} R_{\bar{r}|_p}^{\square, \mu(r_p), \sigma(r_p)} \otimes \overline{\mathbb{Q}}_p$ containing r_p determined by the ordinary deformation condition of [24, Definition 4.1] with respect to the Borel $B \supset T$ associated to Δ (Proposition 2.5, (5) ensures that this deformation condition is well-defined). Note that the genericity of $\{\tilde{n}_\alpha\}$ for $\alpha \in \Delta$ and the fact that Λ_{spin} is multiplicity-free imply that $\mathrm{spin} \circ r_p$ has regular Hodge–Tate weights. We may and do assume that the integers \tilde{n}_α are chosen so that the Hodge–Tate weights of $\mathrm{spin} \circ r_p$ are extremely regular in the sense of [3, Section 2.1]. Let $\mathcal{C}(\rho(r_p))$ ($\rho(r_p) = \mathrm{spin} \circ r_p$) be the corresponding potentially crystalline component of $\mathrm{Spec} R_{\rho(r_p)}^{\square, \mu(\rho(r_p)), \rho(\sigma(r_p))} \otimes \overline{\mathbb{Q}}_p$.

Deformation condition at q . Define $\rho_q : \Gamma_{\mathbb{Q}_q} \rightarrow \mathrm{SL}_2(\mathbb{Z}_p)$ by $\rho_q(\sigma_q) = \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix}$ and $\rho_q(\tau_q) = \begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$, where τ_q is a topological generator of $\mathrm{Gal}(\mathbb{Q}_q^{\mathrm{tame}}/\mathbb{Q}_q^{\mathrm{unr}})$ and $\sigma_q \in \mathrm{Gal}(\mathbb{Q}_q^{\mathrm{tame}}/\mathbb{Q}_q)$ is a lift of the Frobenius element in $\mathrm{Gal}(\mathbb{Q}_q^{\mathrm{unr}}/\mathbb{Q}_q)$ satisfying $\sigma_q \tau_q \sigma_q^{-1} = \tau_q^q$. We set $r_q = \varphi \circ \rho_q$, where $\varphi : \mathrm{SL}_2 \rightarrow \mathrm{Spin}_m$ is the principal SL_2 homomorphism with respect to T and a Borel $B \supset T$ of Spin_m . Then r_q lifts $\bar{r}|_q$. Suppose that l is greater than h_{Spin_m} (see Proposition 2.5), then the Steinberg deformation condition

in [24, Section 4.3] (with respect to B) is well-defined, which corresponds to an irreducible component $\mathcal{C}(r_q)$ of $\text{Spec } R_{\bar{r}|_q}^\square \otimes \overline{\mathbb{Q}}_p$ passing through r_q . Let $\mathcal{C}(\rho(r_q))$ be the corresponding irreducible component of $\text{Spec } R_{\rho(\bar{r})|_q}^\square \otimes \overline{\mathbb{Q}}_p$.

Deformation conditions at other primes. Let v be a prime of \mathbb{Q} outside $\{p, q\}$ at which \bar{r} is ramified. We have $\bar{r}(I_v) \subset T_{\mathbb{F}_p} \cdot \tilde{D}$, which implies (since we are assuming p is sufficiently large) that $|\bar{r}(I_v)|$ is not divisible by p . We take the minimal prime to p deformation condition of [24, Section 4.4] at v , which determines irreducible components of the local deformation rings as before.

Proposition 4.1. *Let $\bar{r} : \Gamma_{\mathbb{Q}} \rightarrow \text{Spin}_m(\overline{\mathbb{F}}_p)$ be the mod p representation in Proposition 2.5 (recall that $m \geq 7$ and $m \equiv 0, 1, 7 \pmod{8}$). Then \bar{r} admits a characteristic zero lift $r : \Gamma_{\mathbb{Q}} \rightarrow \text{Spin}_m(\overline{\mathbb{Z}}_p)$ unramified outside a finite set of primes such that*

- (1) $r|_p$ is ordinary in the sense of [24, Section 4.1] with Hodge–Tate cocharacter $\prod_{\alpha \in \Delta} (\alpha^\vee)^{\tilde{n}_\alpha}$ and $R \circ r|_p$ is potentially diagonalizable in the sense of [3] with extremely regular Hodge–Tate weights (in the sense of [3, Section 2.1]).
- (2) $r|_q$ is Steinberg in the sense of [24, Section 4.3], which is moreover ramified.
- (3) The Zariski closure of the image of r equals Spin_m .
- (4) $R \circ r$ is potentially automorphic, i.e. there exist a totally real extension F^+/\mathbb{Q} and a regular L -algebraic, cuspidal automorphic representation Π of $\text{GL}_N(\mathbb{A}_{F^+})$ such that $\text{spin} \circ r|_{\Gamma_{F^+}} \cong r_{\Pi, \iota}$, where $\iota : \overline{\mathbb{Q}}_p \xrightarrow{\sim} \mathbb{C}$ is a fixed field isomorphism.

Proof. By Proposition 2.5, \bar{r} satisfies the hypotheses at the beginning of Section 3 with $F^+ = \mathbb{Q}$, $G = \text{Spin}_m$ and $R = \text{spin}$. (1) and (4) then follow immediately from Theorem 3.2. By the same theorem, $r|_q$ lies on $\mathcal{C}(r_q)$, and hence it is Steinberg. We will show that $r|_q$ is ramified. Since Π is cuspidal, Π_w is generic for all places w of F^+ , and therefore local–global compatibility ([10]) and [3, Lemma 1.3.2] imply that $\text{spin} \circ r|_{\Gamma_{F_w^+}}$ is a smooth point on its (generic fiber) local lifting ring. Let $w|q$ be any place of F^+ above q . Since $r|_{\Gamma_{F_w^+}}$ lies on the same irreducible component as the restriction $r_q|_{\Gamma_{F_w^+}}$ of the Steinberg-type lift constructed above, so do $\text{spin} \circ r|_{\Gamma_{F_w^+}}$ and $\text{spin} \circ r_q|_{\Gamma_{F_w^+}}$. The latter is visibly a smooth point, so by [3, Lemma 1.3.4(2)] (due to Choi) the inertial restrictions $\text{spin} \circ r|_{\Gamma_{F_w^+}}$ and $\text{spin} \circ r_q|_{\Gamma_{F_w^+}}$ are isomorphic. In particular, $r|_q$ is ramified, which proves (2). It remains to prove (3). Let G_r be the Zariski closure of r , which is reductive since r is irreducible. Moreover, (2) implies that G_r contains a regular

unipotent element of Spin_m . It then follows from a theorem of Dynkin ([27, Theorem A]) that the Lie algebra of G_r is one of \mathfrak{sl}_2 , \mathfrak{g}_2 and \mathfrak{so}_{2n+1} when $m = 2n + 1$ (where \mathfrak{g}_2 may occur only when $m = 7$); one of \mathfrak{sl}_2 , \mathfrak{g}_2 , \mathfrak{so}_{2n-1} and \mathfrak{so}_{2n} when $m = 2n$ (where \mathfrak{g}_2 may occur only when $m = 8$). The argument of [24, Lemma 7.8] then implies that $G_r = \mathrm{Spin}_m$ if the integers $\{\tilde{n}_\alpha\}_{\alpha \in \Delta}$ are taken to be sufficiently general. \square

Theorem 4.2. *Let r be as in Proposition 4.1. Then $\mathrm{spin} \circ r$ is part of a strictly compatible system $R_\lambda : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{GL}_N(\overline{M}_\lambda)$ indexed by finite places λ of some number field M with regular Hodge–Tate weights such that for a density one set of rational primes l , the Zariski closure of the image of R_λ equals Spin_m for $\lambda|l$.*

Proof. By Proposition 4.1(1) and Theorem 3.2(4), for a density one set Σ of rational primes l , R_λ is irreducible for $\lambda|l$. Then [21, Theorem 4] implies that for $l \in \Sigma$ and $\lambda|l$, R_λ factors through Spin_m . In fact, let G_λ be the Zariski closure of the image of R_λ . By Proposition 4.1(3), $G_\lambda = \mathrm{Spin}_m$ for some $\lambda|l$, so $(G_\lambda, \mathrm{std}_\lambda)$ (where std_λ denotes the canonical inclusion $G_\lambda \subset \mathrm{GL}_N$) is similar to $(\mathrm{Spin}_m, \mathrm{spin})$ in the sense of [21], i.e. they have the same formal characters (since the group of connected components and the formal character of the connected component of the monodromy group are independent of l ; see [22, Propositions 6.12 and 6.14]). If m is even, then [21, Theorem 4] implies that $G_\lambda = \mathrm{Spin}_m$; if $m = 2n + 1$ is odd, the only “basic similarity classes” in the sense of loc. cit. relevant to $(\mathrm{Spin}_{2n+1}, \mathrm{spin})$ are products $(\prod_{i=1}^k \mathrm{Spin}_{2n_i+1}, \boxtimes_i \mathrm{spin}_{2n_i+1})$ for some decomposition $n_1 + \cdots + n_k = n$, where spin_{2n_i+1} denotes the appropriate spin representation. This external product is simply the restriction of the 2^n -dimensional spin representation to $\prod \mathrm{Spin}_{2n_i+1} \subset \mathrm{Spin}_{2n+1}$, so in any case $G_\lambda \subset \mathrm{GL}_{2^n}$ factors (up to conjugacy) through $\mathrm{Spin}_{2n+1} \subset \mathrm{GL}_{2^n}$. It follows that R_λ factors as $\mathrm{spin}(r_\lambda)$ for some $r_\lambda : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{Spin}_m(\overline{M}_\lambda)$. It remains to show that when m is odd, $G_\lambda = \mathrm{Spin}_m$ for $\lambda|l$ with l belonging to a density one set of rational primes. By Proposition 4.1(2) and (4) and local–global compatibility ([10]), G_λ contains (up to conjugacy) the image under the spin representation of a regular unipotent element of Spin_m as long as λ is not above q . By [7, Lemma 3.5], r_λ (for λ above some $l \in \Sigma - \{q\}$) then has image containing a regular unipotent element. We deduce from Dynkin’s theorem ([27, Theorem A]) that the Zariski closure of the image of r_λ equals Spin_m . \square

Using some results of Larsen ([20]), we give an application of the above to the inverse Galois problem for the \mathbb{F}_p -points of spin groups.

Corollary 4.3. *For $m \geq 7$, $m \equiv 0, 1, 7 \pmod{8}$, $\mathrm{Spin}_m(\mathbb{F}_p)$ is the Galois group of a finite Galois extension of \mathbb{Q} for p belonging to a set of rational primes of positive density.*

Proof. Let $\{R_\lambda\}$ be as in Theorem 4.2. By [3, Lemma 5.3.1,(3)], after replacing M by a finite extension if necessary, we may assume that R_λ is valued in $\mathrm{GL}_N(M_\lambda)$. Let $\mathrm{cs}(M/\mathbb{Q})$ be the set of rational primes l that split completely in M/\mathbb{Q} . Then [20, Theorem 3.17] implies that there is a subset T of $\mathrm{cs}(M/\mathbb{Q})$ of relative density one, such that for $l \in T$ and $\lambda|l$, $R_\lambda(\Gamma_{\mathbb{Q}})$ is a hyperspecial maximal compact subgroup of $G_\lambda(\mathbb{Q}_l)$. By Theorem 4.2, $G_\lambda(\mathbb{Q}_l)$ equals $\mathrm{Spin}_m(\mathbb{Q}_l)$ for a density one set of rational primes l . It follows that $R_\lambda(\Gamma_{\mathbb{Q}}) = \mathrm{Spin}_m(\mathbb{Z}_l) \twoheadrightarrow \mathrm{Spin}_m(\mathbb{F}_l)$ for l belonging to a set of rational primes of positive density. \square

Remark 4.4. Using a completely analogous argument, one can show that for $n \geq 2$, $\mathrm{SO}_{2n+1}(\mathbb{F}_p)$, resp. $\mathrm{SO}_{4n}(\mathbb{F}_p)$ is the Galois group of a finite Galois extension of \mathbb{Q} for p belonging to a set of rational primes of positive density. We compare this with the main result in [33], which shows that the finite simple groups $\Omega_{2n+1}(p)$ and $\mathrm{P}\Omega_{4n}^+(p)$ both occur as the Galois group of a Galois extension of \mathbb{Q} for all integers $n \geq 2$ and all primes $p \geq 5$.

5. Realization of R_λ in the cohomology of algebraic varieties

Theorem 5.1. *The compatible system of Galois representations $\{R_\lambda\}$ in Theorem 4.2 is motivic in the following sense: there is a smooth projective variety X/\mathbb{Q} and integers i and j such that R_λ is a $\Gamma_{\mathbb{Q}}$ -subrepresentation of $H^i(X_{\overline{\mathbb{Q}}}, \overline{\mathbb{Q}}_l(j))$.*

The proof of this theorem is almost identical to the argument in [7, Section 5], except for a few minor modifications. However, for the reader's convenience, we reproduce here part of their argument. The main reference for this argument is [31]. In [7, Section 5], the dimension of $\{R_\lambda\}$ is odd, which falls into (Case ST) of [31]; in our case, the dimension of $\{R_\lambda\}$ (being a power of 2) is even, which falls into (Case END) of [31]. This is why we need to modify the argument of [7, Section 5] to prove Theorem 5.1. Fix an isomorphism $\iota_l : \overline{\mathbb{Q}}_l \xrightarrow{\sim} \mathbb{C}$; it is implicit in all of the constructions of [31]. The compatible system $\{R_\lambda\}$ is by construction potentially automorphic, i.e. there is a totally real field F^+/\mathbb{Q} such that $R_\lambda|_{\Gamma_{F^+}} \cong r_{l,\iota_l}(\Pi^0)$ for some cuspidal automorphic representation Π^0 of $\mathrm{GL}_N(\mathbb{A}_{F^+})$ (in the notation of [3, Theorem 2.1.1]). Let F/F^+ be a quadratic CM extension. Replacing Π^0 by its base change to F , we may assume that $R_\lambda|_{\Gamma_F} \cong r_{l,\iota_l}(\Pi^0)$ for a conjugate self-dual cuspidal automorphic representation Π^0 of $\mathrm{GL}_N(\mathbb{A}_F)$. Then $R_\lambda|_{\Gamma_F} \cong R_l(\Pi^{0,\vee})$ in the notation of [31, Theorem 7.5] (note that the normalizations for the local Langlands correspondence used in [3] and [31] differ by a dual: see [3, Theorem 2.1.1] and [31, Section 2.3]). Replacing Π^0 by its dual, we may write

$$R_\lambda|_{\Gamma_F} \cong R_l(\Pi^0).$$

We may further adjust the integers $\{\tilde{n}_\alpha\}$ in Proposition 4.1 so that the associated automorphic representation Π^0 is *slightly regular* (a.k.a. Shin regular) in the sense of [31] (this is necessary in Case END).

Let $n = N + 1$ (N is “ m ” in [31]).⁶ We recall the construction of $R_l(\Pi^0)$. Let E be an imaginary quadratic field not contained in F satisfying the four bulleted conditions in Step (II) of the proof of [31, Theorem 7.5]. Replace F by FE and Π^0 by $\mathrm{BC}_{FE/F}(\Pi^0)$. Then the triple (E, F, Π^0) satisfies the six bulleted conditions at the beginning of Step (I) of the proof of loc. cit. Let F' be an imaginary quadratic extension of F^+ satisfying the three bulleted conditions in Step (I) of loc. cit. Then replace F by FF' and Π^0 by $\mathrm{BC}_{FF'/F}(\Pi^0)$ so that [31, Proposition 7.4] applies to the triple (E, F, Π^0) . There is a Hecke character ψ of $\mathbb{A}_E^\times/E^\times$ such that, setting $\Pi = \psi \boxtimes \Pi^1$ (which is an automorphic representation of the group $\mathrm{GL}_1(\mathbb{A}_E) \times \mathrm{GL}_n(\mathbb{A}_F)$; Π^1 is constructed from Π^0 in [31, Section 7.1]), we have (in the notation of [31, Corollary 6.10]⁷)

$$R_l(\Pi^0) := R_l''(\Pi) := R_l'(\Pi) \otimes \mathrm{rec}_{l, l_i}((\varpi \circ N_{F/E}) \otimes |\cdot|^{1/2}),^8$$

where $R_l'(\Pi) := \tilde{R}_l'(\Pi) \otimes \mathrm{rec}_{l, l_i}(\psi^c)|_{\Gamma_F}$ and

$$C_G \cdot \tilde{R}_l'(\Pi) = \sum_{\pi^\infty \in \mathcal{R}_l(\Pi)} R_{\xi, l}^{n-1}(\pi^\infty)^{ss}$$

(see [31, (5.5), (6.23)]; G the unitary similitude group defined in [31, Section 5.1]). The rest of the argument is identical to that of [7, Section 5]: one shows that $\tilde{R}_l'(\Pi)$ is a subrepresentation of the cohomology of a smooth projective variety over F , which implies that $R_\lambda|_{\Gamma_F}$ (which is a twist of $\tilde{R}_l'(\Pi)$) is a subrepresentation of a smooth projective variety over F after possibly replacing F by a finite extension (using [13, IV. Proposition D.1]). It then follows from Frobenius reciprocity and the irreducibility of R_λ that R_λ is a subrepresentation of a smooth projective variety over \mathbb{Q} . This finishes the proof of Theorem 5.1.

6. Comparison with the work of Kret and Shin

In this section, we explain how the main theorem of [18] implies a stronger version of Theorem 1.1 in the case when $m = 2n + 1$ is odd. Moreover, the very recent preprint [19] implies a stronger version of Theorem 1.1 in the case when $m = 2n$ is even in a similar way.

⁶To keep our notation consistent with that of [31], we borrow the same letter n from the preceding sections which was used to denote the rank of the spin group. We hope this does not cause any confusion.

⁷In our case, $i = 1$, $m_i = n - 1$.

⁸ $\varpi : \mathbb{A}_E^\times/E^\times \rightarrow \mathbb{C}^\times$ is the Hecke character defined in [31, Section 3.1].

The maximal torus T of Spin_{2n+1} may be described as

$$T \xrightarrow{\sim} \{(z, t_1, \dots, t_n) \in \mathbb{G}_m^{n+1} : z^2 = t_1 \cdots t_n\}.$$

For $1 \leq i \leq n$, let $\chi_i \in X^\bullet(T)$ be the character defined by $\chi_i(z, t_1, \dots, t_n) = t_i$, and let $\lambda_i \in \frac{1}{2}X_\bullet(T)$ be the cocharacter defined by

$$(2\lambda_i)(t) = (t, 1, \dots, 1, t^2, 1, \dots, 1),$$

where t^2 is located at the $(i+1)$ -th entry. Then the character lattice $X^\bullet(T)$ is generated by χ_1, \dots, χ_n and $\frac{1}{2}(\sum_i \chi_i)$, where $\frac{1}{2}(\sum_i \chi_i)$ sends (z, t_1, \dots, t_n) to z ; and the cocharacter lattice $X_\bullet(T)$ equals $\{\sum_i m_i \lambda_i : m_i \in \mathbb{Z}, \sum m_i \in 2\mathbb{Z}\}$.

Note that $\mathrm{Sp}_{2n}^\vee = \mathrm{SO}_{2n+1}$, $\mathrm{PSp}_{2n}^\vee = \mathrm{Spin}_{2n+1}$ and $\mathrm{GSp}_{2n}^\vee = \mathrm{GSpin}_{2n+1}$.

Let π_∞ be a discrete series representation of $\mathrm{Sp}_{2n}(\mathbb{R})$ with a regular and tempered L-parameter $\phi_\infty : W_{\mathbb{R}} \rightarrow \mathrm{SO}_{2n+1}(\mathbb{C})$ defined by

$$\phi_\infty(z) = \mathrm{diag}(z^{m_1} \bar{z}^{-m_1}, \dots, z^{m_n} \bar{z}^{-m_n}, 1, z^{-m_n} \bar{z}^{m_n}, \dots, z^{-m_1} \bar{z}^{m_1})$$

for $z \in \mathbb{C}^\times$, $m_i \in \mathbb{Z}$ and $\phi_\infty(j) = J$, where $J = [a_{ij}]$ is the $(2n+1)$ by $(2n+1)$ matrix with $a_{n+1, n+1} = (-1)^n$, $a_{i, 2n-i+2} = 1$ for $i \neq n+1$, and $a_{ij} = 0$ otherwise. We want to lift ϕ_∞ to $\mathrm{Spin}_{2n+1}(\mathbb{C})$. Note that $\phi_\infty|_{\mathbb{C}^\times}$ has a unique lift

$$z \mapsto \left(\sum 2m_i \lambda_i \right) \left(\frac{z}{|z|} \right).$$

In particular, $-1 \mapsto (-1)^{\sum m_i}$ (we use -1 to also denote the nontrivial element in the center of Spin_{2n+1} by abuse of notation). The element j necessarily maps to a lift of $J \in \mathrm{SO}_{2n+1}(\mathbb{C})$ in $\mathrm{Spin}_{2n+1}(\mathbb{C})$, which is either $\omega_1 \cdots \omega_n$ or $-\omega_1 \cdots \omega_n$ (in the notation of Lemma 2.3 and the paragraph preceding it). We have $(\omega_1 \cdots \omega_n)^2 = (-1)^{n(n+1)/2}$. To have a well-defined homomorphism $\tilde{\phi}_\infty : W_{\mathbb{R}} \rightarrow \mathrm{Spin}_{2n+1}(\mathbb{C})$ lifting ϕ_∞ , we must ensure that $\tilde{\phi}_\infty(j)\tilde{\phi}_\infty(z)\tilde{\phi}_\infty(j)^{-1} = \tilde{\phi}_\infty(\bar{z})$ and $\tilde{\phi}_\infty(j)^2 = \tilde{\phi}_\infty(-1)$. The first equality is automatic. The second equality is by the above equivalent to $(-1)^{n(n+1)/2} = (-1)^{\sum m_i}$. Therefore, ϕ_∞ lifts to $\mathrm{Spin}_{2n+1}(\mathbb{C})$ if and only if $\sum m_i$ has the same parity as $n(n+1)/2$. On the automorphic side, this says π_∞ descends to a discrete series representation of $\mathrm{PSP}_{2n}(\mathbb{R})$ if and only if the integers m_i appearing in its L-parameter satisfies $\sum m_i \equiv \frac{n(n+1)}{2} \pmod{2}$.

Now we ask: Of the discrete series that descends to $\mathrm{PSP}_{2n}(\mathbb{R})$, are any of them L-algebraic in the sense of [8]? Consider their L-parameter $\tilde{\phi}_\infty|_{\mathbb{C}}$, $z \mapsto \left(\sum 2m_i \lambda_i \right) \left(\frac{z}{|z|} \right) = z \sum m_i \lambda_i \bar{z}^{-\sum m_i \lambda_i}$, we have $\sum m_i \lambda_i \in X_\bullet(T)$ if and only if $\sum m_i \in 2\mathbb{Z}$. Therefore we have

Lemma 6.1. *Let π_∞ and ϕ_∞ be as before. Then*

- (1) π_∞ descends to a discrete series representation of $\mathrm{PSP}_{2n}(\mathbb{R})$ if and only if $\sum m_i \equiv \frac{n(n+1)}{2} \pmod{2}$.

- (2) π_∞ descends to an L -algebraic discrete series representation of $\mathrm{PSP}_{2n}(\mathbb{R})$ if and only if $n \equiv 0, 3 \pmod{4}$ and $\sum m_i \in 2\mathbb{Z}$.

A trace formula argument shows that there are infinitely many cuspidal automorphic representations π of $\mathrm{PSP}_{2n}(\mathbb{A}_{\mathbb{Q}})$ such that its archimedean component π_∞ satisfies Lemma 6.1, (2) and π is Steinberg at a finite place of \mathbb{Q} (cf. [11]). Let $\tilde{\pi}$ be the cuspidal automorphic representation of $\mathrm{GSp}_{2n}(\mathbb{A}_{\mathbb{Q}})$ associated to π via the natural map $\mathrm{pr} : \mathrm{GSp}_{2n} \rightarrow \mathrm{PSP}_{2n}$. We want to use the main theorem of [18] to associate to $\tilde{\pi}$ GSpin_{2n+1} -valued Galois representations.

Lemma 6.2. *The infinity component $\tilde{\pi}_\infty$ of $\tilde{\pi}$ is L -cohomological in the sense of [18], i.e. $\tilde{\pi}_\infty \cdot |\lambda_0|^{\frac{n(n+1)}{4}}$ is cohomological in the sense of [18, Definition 1.12] (λ_0 corresponds to the similitude character of GSp_{2n}).*

Proof. Since π_∞ is a discrete series representation of $\mathrm{Sp}_{2n}(\mathbb{R})$ with infinitesimal character $\mu(\pi_\infty)$ (where $\mu(\pi_\infty)$ is determined by writing $\phi_\infty(z) = z^{\mu(\pi_\infty)} \bar{z}^{\nu(\pi_\infty)}$), it is $\xi_{\mu(\pi_\infty)-\rho}^\vee$ -cohomological ([5, Theorem V.3.3]), where $\xi_{\mu(\pi_\infty)-\rho}$ is the highest-weight $\mu(\pi_\infty) - \rho$ representation of $\mathrm{Sp}_{2n}(\mathbb{C})$, and ρ is the half-sum of positive roots for the choice of root basis used also to parametrize highest weights. We need an extension $\tilde{\xi}$ of $\xi_{\mu(\pi_\infty)-\rho}$ to $\mathrm{GSp}_{2n}(\mathbb{C})$ such that (letting K be a maximal compact subgroup of $\mathrm{Sp}_{2n}(\mathbb{R})$)

$$H^*(\mathfrak{gsp}_{2n}(\mathbb{C}), K \cdot Z_{\mathrm{GSp}_{2n}}(\mathbb{R}); \tilde{\pi}_\infty \cdot |\lambda_0|^{\frac{n(n+1)}{4}} \otimes \tilde{\xi}^\vee) \neq 0.$$

From the definition of this cohomology group, and the fact that $\pi_\infty \otimes \xi_{\mu(\pi_\infty)-\rho}^\vee$ has non-zero $(\mathfrak{sp}_{2n}(\mathbb{C}), K)$ -cohomology, this reduces to checking that we can find an extension $\tilde{\xi}^\vee$ of $\xi_{\mu(\pi_\infty)-\rho}^\vee$ such that $\tilde{\xi}$ and $\tilde{\pi}_\infty \cdot |\lambda_0|^{\frac{n(n+1)}{4}}$ have the same central character; note that we will only have to check this on $\mathbb{R}_{>0}^\times \subset Z_{\mathrm{GSp}_{2n}}(\mathbb{R})$ since $-1 \in K$. We choose $\tilde{\xi}$ to be the representation with highest weight $\mu(\tilde{\pi}_\infty) - \rho + \frac{n(n+1)}{4}\lambda_0 \in X^\bullet(T_{\mathrm{GSp}_{2n}})$, where $\mu(\tilde{\pi}_\infty) = \sum m_i \lambda_i$, which (since $\sum_i m_i$ is even) lies in $X_\bullet(T_{\mathrm{Spin}_{2n+1}}) = X^\bullet(T_{\mathrm{PSP}_{2n}}) \subset X^\bullet(T_{\mathrm{GSp}_{2n}})$, and both ρ and $\frac{n(n+1)}{4}\lambda_0$ are integral since $n \equiv 0, 3 \pmod{4}$. The lemma is now proven, since this highest weight representation has the same central character as $\tilde{\pi}_\infty \cdot |\lambda_0|^{\frac{n(n+1)}{4}}$. \square

Therefore, by [18, Theorem A], the cuspidal automorphic representation $\tilde{\pi}$ of $\mathrm{GSp}_{2n}(\mathbb{A}_{\mathbb{Q}})$ (for $n \equiv 0, 3 \pmod{4}$) has an associated weakly compatible system of Galois representations

$$\rho_{\tilde{\pi}, l} : \Gamma_{\mathbb{Q}} \rightarrow \mathrm{GSpin}_{2n+1}(\overline{\mathbb{Q}}_l)$$

satisfying the conclusions of loc. cit.⁹ In particular, since the central character of $\tilde{\pi}$ is trivial, [18, Theorem A (i)] implies that $\rho_{\tilde{\pi},l}$ is in fact valued in $\text{Spin}_{2n+1}(\overline{\mathbb{Q}}_l)$. Moreover, for sufficiently general $\{m_i\}_{1 \leq i \leq n}$, $\tilde{\pi}_\infty$ is spin-regular in the sense of [18], i.e. the L-parameter of $\tilde{\pi}_\infty$ maps to a regular parameter for GL_{2n} by the spin representation. In this case, the Zariski-closure of $\rho_{\tilde{\pi},l}$ equals Spin_{2n+1} , and thus we obtain a Spin_{2n+1} -compatible system of motivic Galois representations with full monodromy.

In contrast, the following lemma shows that for the remaining n , we do not expect the Galois representations associated to cuspidal automorphic representations of $\text{PSP}_{2n}(\mathbb{A}_{\mathbb{Q}})$ that are regular algebraic at infinity to land in Spin_{2n+1} by [8, Conjecture 5.3.4].

Lemma 6.3. *Suppose $n \equiv 1, 2 \pmod{4}$ (equivalently, $\frac{n(n+1)}{2}$ is odd) and π is a cuspidal automorphic representation of $\text{Sp}_{2n}(\mathbb{A}_{\mathbb{Q}})$ with trivial central character. Denote by π^b the associated representation of $\text{PSP}_{2n}(\mathbb{A}_{\mathbb{Q}})$. If π is regular algebraic,¹⁰ then π^b is C-algebraic but not L-algebraic.*

Proof. We can write the L-parameter ϕ_∞ of π_∞ as

$$\phi_\infty(z) = \text{diag}(z^{m_1} \bar{z}^{l_1}, \dots, z^{m_n} \bar{z}^{l_n}, 1, z^{-m_n} \bar{z}^{-l_n}, \dots, z^{-m_1} \bar{z}^{-l_1})$$

with all $m_i, l_i \in \mathbb{Z}$ and $z \in \mathbb{C}^\times$. Arthur’s classification of automorphic representations of classical groups ([2]) implies that π has a functorial transfer to a cuspidal automorphic representation of $\text{GL}_{2n+1}(\mathbb{A}_{\mathbb{Q}})$, so Clozel’s archimedean purity theorem ([12, Lemme 4.9]) implies that $m_i + l_i = 0$ for all i . Since the integers m_i are distinct and nonzero, it is then easy to see that ϕ_∞ is isomorphic to the L-parameter under the same name defined before Lemma 6.1. Since the L-parameter $\tilde{\phi}_\infty$ of π_∞^b lifts ϕ_∞ , the calculations preceding Lemma 6.1 imply that $\sum m_i \equiv \frac{n(n+1)}{2} \equiv 1 \pmod{2}$. A easy calculation shows that ρ^\vee (the half-sum of coroots of Spin_{2n+1}) equals $n\lambda_1 + (n-1)\lambda_2 + \dots + 1\lambda_n$ (which does not lie in $X_\bullet(T)$ since $\frac{n(n+1)}{2}$ is odd). It follows that $\sum m_i \lambda_i \in \rho^\vee + X_\bullet(T)$ and hence π_∞^b is C-algebraic but not L-algebraic ($\tilde{\phi}_\infty(z) = z \sum m_i \lambda_i \bar{z}^{-\sum m_i \lambda_i}$). \square

However, if π is not assumed to be regular at infinity, then π^b can be L-algebraic, in which case the expected Galois representations associated to it lands in Spin_{2n+1} ([8, Conjecture 3.2.1]).

Acknowledgments. I thank Stefan Patrikis for helpful conversations and for suggesting me to compare my construction with [18]. I thank Patrick Allen for carefully reading through a draft of this paper and making useful comments. Last but not least, I thank my mother, who was taken to

⁹Such a compatible system appears in the cohomology of a Shimura variety for a suitable inner form of GSp_{2n} by the construction of [18].

¹⁰Since the dual group SO_{2n+1} is adjoint, L-algebraicity is equivalent to C-algebraicity.

a better place on October 28, 2019, for her constant encouragements and unconditional support on my mathematical endeavors when I was an undergraduate student in China. Without those, I would not have had the opportunity to earn a doctorate in Mathematics in the United States, much less been equipped to write the present paper. This work is dedicated to the memory of her with enormous gratitude and deepest respect.

References

- [1] J. ADAMS & X. HE, “Lifting of elements of Weyl groups”, *J. Algebra* **485** (2017), p. 142-165.
- [2] J. ARTHUR, *The Endoscopic classification of representations orthogonal and symplectic groups*, Colloquium Publications, vol. 61, American Mathematical Society, 2013.
- [3] T. BARNET-LAMB, T. GEE, D. GERAGHTY & R. TAYLOR, “Potential automorphy and change of weight”, *Ann. Math.* **179** (2014), no. 2, p. 501-609.
- [4] R. BELLOVIN & T. GEE, “ G -valued local deformation rings and global lifts”, *Algebra Number Theory* **13** (2019), no. 2, p. 333-378.
- [5] A. BOREL & N. WALLACH, *Continuous Cohomology, Discrete Subgroups, and Representations of Reductive Groups*, Mathematical Surveys and Monographs, vol. 67, American Mathematical Society, 2000.
- [6] N. BOURBAKI, *Lie groups and Lie algebras, Chapters 4-6*, Springer, 2002.
- [7] G. BOXER, F. CALEGARI, M. EMERTON, B. LEVIN, K. MADAPUSI PERA & S. PATRIKIS, “Compatible systems of Galois representations associated to the exceptional group E_6 ”, *Forum Math. Sigma* **7** (2019), article no. e4 (29 pages).
- [8] K. BUZZARD & T. GEE, “The conjectural connections between automorphic representations and Galois representations”, in *Automorphic forms and Galois representations (Durham, 2011)*, London Mathematical Society Lecture Note Series, vol. 414, London Mathematical Society, 2011, p. 135-187.
- [9] F. CALEGARI, “Even Galois representations and the Fontaine–Mazur conjecture. II”, *J. Am. Math. Soc.* **25** (2012), no. 2, p. 533-554.
- [10] A. CARAIANI, “Local–global compatibility and the action of monodromy on nearby cycles”, *Duke Math. J.* **161** (2012), no. 12, p. 2311-2413.
- [11] L. CLOZEL, “On limit multiplicities of discrete series representations in spaces of automorphic forms”, *Invent. Math.* **83** (1986), no. 2, p. 265-284.
- [12] ———, “Motifs et formes automorphes: applications du principe de functorialité”, in *Automorphic forms, Shimura varieties, and L -functions. Vol. I (Ann Arbor, 1988)*, Perspectives in Mathematics, vol. 10, Academic Press Inc., 1990, p. 77-159.
- [13] P. DELIGNE, J. S. MILNE, A. OGUS & K.-Y. SHIH, *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics, vol. 900, Springer, 1982.
- [14] N. FAKHRUDDIN, C. KHARE & S. PATRIKIS, “Relative deformation theory and lifting irreducible Galois representations”, <https://arxiv.org/abs/1904.02374>, 2019.
- [15] W. FULTON & J. HARRIS, *Representation theory: a first course*, Graduate Texts in Mathematics, vol. 129, Springer, 2013.
- [16] C. KHARE & J.-P. WINTENBERGER, “Serre’s modularity conjecture (I)”, *Invent. Math.* **178** (2009), no. 3, p. 485-504.
- [17] M. KISIN, “Potentially semi-stable deformation rings”, *J. Am. Math. Soc.* **28** (2008), no. 2, p. 513-546.
- [18] A. KRET & S. W. SHIN, “Galois representations for general symplectic groups”, <https://arxiv.org/abs/1609.04223>, to appear in *J. Eur. Math. Soc.*, 2016.
- [19] ———, “Galois representations for even general special orthogonal groups”, <https://arxiv.org/abs/2010.08408>, 2020.
- [20] M. LARSEN, “Maximality of Galois actions for compatible systems”, *Duke Math. J.* **80** (1995), no. 3, p. 601-630.
- [21] M. LARSEN & R. PINK, “Determining representations from invariant dimensions”, *Invent. Math.* **102** (1990), no. 1, p. 377-398.

- [22] ———, “On ℓ -independence of algebraic monodromy groups in compatible systems of representations”, *Invent. Math.* **107** (1992), no. 3, p. 603-636.
- [23] J. NEUKIRCH, A. SCHMIDT & K. WINGBERG, *Cohomology of number fields*, 2nd corr. ed., Grundlehren der Mathematischen Wissenschaften, vol. 323, Springer, 2013.
- [24] S. PATRIKIS, “Deformations of Galois representations and exceptional monodromy”, *Invent. Math.* **205** (2016), no. 2, p. 269-336.
- [25] S. PATRIKIS & S. TANG, “Potential automorphy of GSpin_{2n+1} -valued Galois representations”, <https://arxiv.org/abs/1910.03164>, 2019.
- [26] R. RAMAKRISHNA, “Deforming Galois representations and the conjectures of Serre and Fontaine–Mazur”, *Ann. Math.* **1566** (2002), no. 1, p. 115-154.
- [27] J. SAXL & G. M. SEITZ, “Subgroups of algebraic groups containing regular unipotent elements”, *J. Lond. Math. Soc.* **55** (1997), no. 2, p. 370-386.
- [28] J.-P. SERRE, “Propriétés conjecturales des groupes de Galois motiviques et des représentations ℓ -adiques”, in *Motives (Seattle, 1991)*, Proceedings of Symposia in Pure Mathematics, vol. 55, American Mathematical Society, 1991, p. 377-400.
- [29] ———, “Sur la semi-simplicité des produits tensoriels de représentations de groupes”, *Invent. Math.* **116** (1994), no. 1, p. 513-530.
- [30] ———, *Topics in Galois theory*, CRC Press, 2016.
- [31] S. W. SHIN, “Galois representations arising from some compact Shimura varieties”, *Ann. Math.* **173** (2011), no. 3, p. 1645-1741.
- [32] Z. YUN, “Motives with exceptional Galois groups and the inverse Galois problem”, *Invent. Math.* **196** (2014), no. 2, p. 267-337.
- [33] D. ZYWINA, “The inverse Galois problem for orthogonal groups”, <https://arxiv.org/abs/1409.1151>, 2014.

Shiang TANG
1409 West Green Street
Urbana, IL 61801, United States
E-mail: shiangtang1989@gmail.com