

On the largest prime factor of $n! + 2^n - 1$

par FLORIAN LUCA et IGOR E. SHPARLINSKI

RÉSUMÉ. Pour un entier $n \geq 2$, notons $P(n)$ le plus grand facteur premier de n . Nous obtenons des majorations sur le nombre de solutions de congruences de la forme $n! + 2^n - 1 \equiv 0 \pmod{q}$ et nous utilisons ces bornes pour montrer que

$$\limsup_{n \rightarrow \infty} P(n! + 2^n - 1)/n \geq (2\pi^2 + 3)/18.$$

ABSTRACT. For an integer $n \geq 2$ we denote by $P(n)$ the largest prime factor of n . We obtain several upper bounds on the number of solutions of congruences of the form $n! + 2^n - 1 \equiv 0 \pmod{q}$ and use these bounds to show that

$$\limsup_{n \rightarrow \infty} P(n! + 2^n - 1)/n \geq (2\pi^2 + 3)/18.$$

1. Introduction

For any positive integer $k > 1$ we denote by $P(k)$ the largest prime factor of k and by $\omega(k)$ the number of distinct prime divisors of k . We also set $P(1) = 1$ and $\omega(1) = 0$.

It is trivial to see that $P(n! + 1) > n$. Erdős and Stewart [4] have shown that

$$\limsup_{n \rightarrow \infty} \frac{P(n! + 1)}{n} > 2.$$

This bound is improved in [7] where it is shown that the above upper limit is at least $5/2$, and that it also holds for $P(n! + f(n))$ with a nonzero polynomial $f(X) \in \mathbb{Z}[X]$.

Here we use the method of [7], which we supplement with some new arguments, to show that

$$\limsup_{n \rightarrow \infty} \frac{P(n! + 2^n - 1)}{n} > (2\pi^2 + 3)/18.$$

We also estimate the total number of distinct primes which divide at least one value of $n! + 2^n - 1$ with $1 \leq n \leq x$.

These results are based on several new elements, such as bounds for the number of solutions of congruences with $n! + 2^n - 1$, which could be of independent interest.

Certainly, there is nothing special in the sequence $2^n - 1$, and exactly the same results can be obtained for $n! + u(n)$ with any nonzero binary recurrent sequence $u(n)$.

Finally, we note that our approach can be used to estimate $P(n! + u(n))$ with an arbitrary linear recurrence sequence $u(n)$ (leading to similar, albeit weaker, results) and with many other sequences (whose growth and the number of zeros modulo q are controllable).

Throughout this paper, we use the Vinogradov symbols \gg , \ll and \asymp as well as the Landau symbols O and o with their regular meanings. For $z > 0$, $\log z$ denotes the natural logarithm of z .

Acknowledgments. During the preparation of this paper, F. L. was supported in part by grants SEP-CONACYT 37259-E and 37260-E, and I. S. was supported in part by ARC grant DP0211459.

2. Bounding the number of solutions of some equations and congruences

The following polynomial

$$(2.1) \quad F_{k,m}(X) = (2^k - 1) \prod_{i=1}^m (X + i) - (2^m - 1) \prod_{i=1}^k (X + i) + 2^m - 2^k$$

plays an important role in our arguments.

Lemma 2.1. *The equation*

$$F_{k,m}(n) = 0$$

has no integer solutions (n, k, m) with $n \geq 3$ and $m > k \geq 1$.

Proof. One simply notices that for any $n \geq 3$ and $m > k \geq 1$

$$\begin{aligned} (2^k - 1) \prod_{i=1}^m (n + i) &\geq 2^{k-1} (n + 1)^{m-k} \prod_{i=1}^k (n + i) \\ &\geq (n + 1) 2^{m-2} \prod_{i=1}^k (n + i) \geq 2^m \prod_{i=1}^k (n + i) \\ &> (2^m - 1) \prod_{i=1}^k (n + i). \end{aligned}$$

Hence, $F_{k,m}(n) > 0$ for $n \geq 3$. □

Let $\ell(q)$ denote the multiplicative order of 2 modulo an odd integer $q \geq 3$.

For integers $y \geq 0$, $x \geq y + 1$, and $q \geq 1$, we denote by $T(y, x, q)$ the set of solutions of the following congruence

$$T(y, x, q) = \{n \mid n! + 2^n - 1 \equiv 0 \pmod{q}, y + 1 \leq n \leq x\},$$

and put $T(y, x, q) = \#T(y, x, q)$. We also define

$$T(x, q) = T(0, x, q) \quad \text{and} \quad T(x, q) = T(0, x, q).$$

Lemma 2.2. *For any prime p and integers x and y with $p > x \geq y + 1 \geq 1$, we have*

$$T(y, x, p) \ll \max\{(x - y)^{3/4}, (x - y)/\ell(p)\}.$$

Proof. We assume that $p \geq 3$, otherwise there is nothing to prove. Let $\ell(p) > z \geq 1$ be a parameter to be chosen later.

Let $y + 1 \leq n_1 < \dots < n_t \leq x$ be the complete list of $t = T(y, x, p)$ solutions to the congruence $n! + 2^n - 1 \equiv 0 \pmod{p}$, $y + 1 \leq n \leq x$. Then

$$T(y, x, p) = \mathcal{U}_1 \cup \mathcal{U}_2,$$

where

$$\mathcal{U}_1 = \{n_i \in T(y, x, p) \mid |n_i - n_{i+2}| \geq z, i = 1, \dots, t - 2\},$$

and $\mathcal{U}_2 = T(y, x, p) \setminus \mathcal{U}_1$.

It is clear that $\#\mathcal{U}_1 \ll (x - y)/z$. Assume now that $n \in \mathcal{U}_2 \setminus \{n_{t-1}, n_t\}$. Then there exists a nonzero integers k and m with $0 < k < m \leq z$, and such that

$$n! + 2^n - 1 \equiv (n + k)! + 2^{n+k} - 1 \equiv (n + m)! + 2^{n+m} - 1 \equiv 0 \pmod{p}.$$

Eliminating 2^n from the first and the second congruence, and then from the first and the third congruence, we obtain

$$\begin{aligned} n! \left(\prod_{i=1}^k (n + i) - 2^k \right) + 2^k - 1 \\ \equiv n! \left(\prod_{i=1}^m (n + i) - 2^m \right) + 2^m - 1 \equiv 0 \pmod{p}. \end{aligned}$$

Now eliminating $n!$, we derive

$$(2^m - 1) \left(\prod_{i=1}^k (n + i) - 2^k \right) - (2^k - 1) \left(\prod_{i=1}^m (n + i) - 2^m \right) \equiv 0 \pmod{p},$$

or $F_{k,m}(n) \equiv 0 \pmod{p}$, where $F_{k,m}(X)$ is given by (2.1). Because $\ell(p) > z$, we see that for every $0 < k < m \leq z$ the polynomial $F_{k,m}(X)$ has a nonzero coefficient modulo p and $\deg F_{k,m} = m \leq z$, thus for every $0 < k < m < z$ there are at most z suitable values of n (since $p > x \geq y + 1 \geq 1$).

Summing over all admissible values of k and m , we derive $\#\mathcal{U}_2 \ll z^3 + 1$. Therefore

$$T(y, x, p) \leq \#\mathcal{U}_1 + \#\mathcal{U}_2 \ll (x - y)/z + z^3 + 1.$$

Taking $z = \min\{(x - y)^{1/4}, \ell(p) - 1\}$ we obtain the desired inequality. \square

Obviously, for any $n \geq p$ with $n! + 2^n - 1 \equiv 0 \pmod{p}$, we have $2^n \equiv 1 \pmod{p}$. Thus

$$(2.2) \quad T(p, x, p) \ll x/\ell(p).$$

Lemma 2.3. *For any integers $q \geq 2$ and $x \geq y + 1 \geq 1$, we have*

$$T(y, x, q) \leq \left(2 + O\left(\frac{1}{\log x}\right)\right) \frac{(x - y) \log x}{\log q} + O(1).$$

Proof. Assume that $T(y, x, q) \geq 6$, because otherwise there is nothing to prove. We can also assume that q is odd. Then, by the Dirichlet principle, there exist integers $n \geq 4$, $m > k \geq 1$, satisfying the inequalities

$$1 \leq k < m \leq 2 \frac{x - y}{T(y, x, q) - 4}, \quad y + 1 \leq n < n + k < n + m \leq x,$$

and such that

$$n! + 2^n - 1 \equiv (n + k)! + 2^{n+k} - 1 \equiv (n + m)! + 2^{n+m} - 1 \equiv 0 \pmod{q}.$$

Arguing as in the proof of Lemma 2.2, we derive $F_{m,k}(n) \equiv 0 \pmod{q}$. Because $F_{m,k}(n) \neq 0$ by Lemma 2.1, we obtain $|F_{m,k}(n)| \geq q$. Obviously, $|F_{m,k}(n)| = O(2^k x^m) = O((2x)^m)$. Therefore,

$$\log q \leq m(\log x + O(1)) \leq 2 \frac{(x - y)(\log x + O(1))}{T(y, x, p) - 4},$$

and the result follows. \square

Certainly, Lemma 2.2 is useful only if $\ell(p)$ is large enough.

Lemma 2.4. *For any x the inequality $\ell(p) \geq x^{1/2}/\log x$ holds for all except maybe $O(x/(\log x)^3)$ primes $p \leq x$.*

Proof. Put $L = \lfloor x^{1/2}/\log x \rfloor$. If $\ell(p) \leq L$ then $p|R$, where

$$R = \prod_{i=1}^L (2^i - 1) \leq 2^{L^2}.$$

The bound $\omega(R) \ll \log R/\log \log R \ll L^2/\log L$ concludes the proof. \square

We remark that stronger results are known, see [3, 6, 9], but they do not seem to be of help for our arguments.

3. Main Results

Theorem 3.1. *The following bound holds:*

$$\limsup_{n \rightarrow \infty} \frac{P(n! + 2^n - 1)}{n} \geq \frac{2\pi^2 + 3}{18} = 1.2632893 \dots$$

Proof. Assuming that the statement of the above theorem is false, we see that there exist two constants $\lambda < (2\pi^2 + 3)/18$ and μ such that the inequality $P(n! + 2^n - 1) < \lambda n + \mu$ holds for all integer positive n .

We let x be a large positive integer and consider the product

$$W = \prod_{1 \leq n \leq x} (n! + 2^n - 1).$$

Let $Q = P(W)$ so we have $Q \leq \lambda x + \mu$. Obviously,

$$(3.1) \quad \log W = \frac{1}{2}x^2 \log x + O(x^2).$$

For a prime p , we denote by s_p the largest power of p dividing at least one of the nonzero integers of the form $n! + 2^n - 1$ for $n \leq x$. We also denote by r_p the p -adic order of W . Hence,

$$(3.2) \quad r_p = \sum_{1 \leq s \leq s_p} T(x, p^s),$$

and therefore, by (3.1) and (3.2), we deduce

$$(3.3) \quad \sum_{\substack{p|W \\ p \leq Q}} \log p \sum_{1 \leq s \leq s_p} T(x, p^s) = \log W = \frac{1}{2}x^2 \log x + O(x^2).$$

We let \mathcal{M} be the set of all possible pairs (p, s) which occur on the left hand side of (3.3), that is,

$$\mathcal{M} = \{(p, s) \mid p|W, p \leq Q, 1 \leq s \leq s_p\},$$

and so (3.3) can be written as

$$(3.4) \quad \sum_{(p,s) \in \mathcal{M}} T(x, p^s) \log p = \frac{1}{2}x^2 \log x + O(x^2).$$

As usual, we use $\pi(y)$ to denote the number of primes $p \leq y$, and recall that by the Prime Number Theorem we have $\pi(y) = (1 + o(1))y/\log y$.

Now we introduce subsets $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3, \mathcal{E}_4 \in \mathcal{M}$, which possibly overlap, and whose contribution to the sums on the left hand side of (3.4) is $o(x^2 \log x)$. After this, we study the contribution of the remaining set \mathcal{L} .

- Let \mathcal{E}_1 be the set of pairs $(p, s) \in \mathcal{M}$ with $p \leq x/\log x$. By Lemma 2.3, we have

$$\begin{aligned} \sum_{(p,s) \in \mathcal{E}_1} T(x, p^s) \log p &\ll x \log x \sum_{(p,s) \in \mathcal{E}_1} \frac{1}{s} + \sum_{(p,s) \in \mathcal{E}_1} \log p \\ &\ll x \log x \sum_{p \leq x/\log x} \log(s_p + 1) \\ &\quad + \sum_{p \leq x/\log x} s_p \log p \ll x^2, \end{aligned}$$

because obviously $s_p \ll x \log x$.

- Let \mathcal{E}_2 be the set of pairs $(p, s) \in \mathcal{M}$ with $s \geq x/(\log x)^2$. Again by Lemma 2.3, and by the inequality

$$s_p \ll x \frac{\log x}{\log p},$$

we have

$$\begin{aligned} \sum_{(p,s) \in \mathcal{E}_2} T(x, p^s) \log p &\ll x \log x \sum_{(p,s) \in \mathcal{E}_2} \frac{1}{s} + \sum_{(p,s) \in \mathcal{E}_2} \log p \\ &\ll x \log x \sum_{p \leq Q} \sum_{x/(\log x)^2 \leq s \leq s_p} \frac{1}{s} + \sum_{p \leq Q} s_p \log p \\ &\ll x\pi(Q) \log x \log \log x \ll x^2 \log \log x, \end{aligned}$$

because $Q = O(x)$ by our assumption.

- Let \mathcal{E}_3 be the set of pairs $(p, s) \in \mathcal{M}$ with $\ell(p) \leq x^{1/2}/\log x$. Again by Lemmas 2.3 and 2.4, and by the inequality $s_p \ll x \log x$, we have

$$\begin{aligned} \sum_{(p,s) \in \mathcal{E}_3} T(x, p^s) \log p &\ll x \log x \sum_{(p,s) \in \mathcal{E}_2} \frac{1}{s} + \sum_{(p,s) \in \mathcal{E}_3} \log p \\ &\ll x \log x \sum_{\substack{p \leq Q \\ \ell(p) \leq x^{1/2}/\log x}} \sum_{1 \leq s \leq s_p} \frac{1}{s} \\ &\quad + \sum_{\substack{p \leq Q \\ \ell(p) \leq x^{1/2}/\log x}} s_p \log p \\ &\ll x(\log x)^2 \sum_{\substack{p \leq Q \\ \ell(p) \leq x^{1/2}/\log x}} 1 \ll x^2/\log x. \end{aligned}$$

- Let \mathcal{E}_4 be the set of pairs $(p, s) \in \mathcal{M} \setminus (\mathcal{E}_1 \cup \mathcal{E}_3)$ with $s < x^{1/4}$. By Lemma 2.2 and by (2.2), we have

$$\begin{aligned} \sum_{(p,s) \in \mathcal{E}_3} T(x, p^s) \log p &\ll x^{1/4} \sum_{p \leq Q} T(x, p) \log p \\ &\ll x^{1/4} \sum_{p \leq Q} \left(p^{3/4} + x/\ell(p) \right) \log p \\ &\ll x^{1/4} Q^{3/4} \sum_{p \leq Q} \log p \\ &\ll x^{1/4} Q^{7/4} \ll x^2. \end{aligned}$$

We now put $\mathcal{L} = \mathcal{M} \setminus (\mathcal{E}_1 \cup \mathcal{E}_2 \cup \mathcal{E}_3 \cup \mathcal{E}_4)$.

The above estimates, together with (3.4), show that

$$(3.5) \quad \sum_{(p,s) \in \mathcal{L}} T(x, p^s) \log p = \frac{1}{2} x^2 \log x + O(x^2 \log \log x).$$

The properties of the pairs $(p, s) \in \mathcal{L}$ can be summarized as

$$p > \frac{x}{\log x}, \quad \ell(p) \geq \frac{x^{1/2}}{\log x}, \quad \frac{x}{(\log x)^2} \geq s \geq x^{1/4}.$$

In what follows, we repeatedly use the above bounds.

We now remark that because by our assumption $P(n! + 2^n - 1) \leq \lambda n + \mu$ for $n \leq x$, we see that $T(x, p^s) = T(\lfloor (p - \mu)/\lambda \rfloor, x, p^s)$.

Thus, putting $x_p = \min\{x, p\}$, we obtain

$$T(x, p^s) = T(\lfloor (p - \mu)/\lambda \rfloor, x, p^s) = T(\lfloor (p - \mu)/\lambda \rfloor, x_p, p^s) + T(x_p, x, p^s).$$

Therefore,

$$(3.6) \quad \sum_{(p,s) \in \mathcal{L}} T(x, p^s) \log p = U + V,$$

where

$$U = \sum_{(p,s) \in \mathcal{L}} T(\lfloor (p - \mu)/\lambda \rfloor, x_p, p^s) \log p,$$

and

$$V = \sum_{(p,s) \in \mathcal{L}} T(x_p, x, p^s) \log p.$$

To estimate U , we observe that, by Lemma 2.3,

$$\begin{aligned} U &\leq (2 + o(1)) \log x \sum_{p \leq Q} \left(\binom{x_p - \frac{p - \mu}{\lambda}}{\quad} \sum_{x/\log x > s \geq x^{1/4}} \frac{1}{s} + O(1) \right) \\ &\leq (3/2 + o(1)) (\log x)^2 \sum_{p \leq Q} \binom{x_p - \frac{p - \mu}{\lambda}}{\quad} + O(x^2). \end{aligned}$$

Furthermore,

$$\begin{aligned} \sum_{p \leq Q} \binom{x_p - \frac{p - \mu}{\lambda}}{\quad} &= \sum_{p \leq x} \binom{p - \frac{p - \mu}{\lambda}}{\quad} + \sum_{x < p \leq Q} \binom{x - \frac{p - \mu}{\lambda}}{\quad} \\ &= \left(\frac{\lambda - 1}{2\lambda} + o(1) \right) \frac{x^2}{\log x} + \left(\frac{(\lambda - 1)^2}{2\lambda} + o(1) \right) \frac{x^2}{\log x} \\ &= \left(\frac{\lambda - 1}{2} + o(1) \right) \frac{x^2}{\log x}. \end{aligned}$$

Hence

$$(3.7) \quad U \leq \left(\frac{3(\lambda - 1)}{4} + o(1) \right) x^2 \log x.$$

We now estimate V . For an integer $\alpha \geq 1$ we let \mathcal{P}_α be the set of primes $p \leq Q$ with

$$\ell(p) = \dots = \ell(p^\alpha) \neq \ell(p^{\alpha+1}).$$

Thus, $\ell(p^{\alpha+1}) = \ell(p)p$.

Accordingly, let \mathcal{L}_α be the subset of pairs $(p, s) \in \mathcal{L}$ for which $p \in \mathcal{P}_\alpha$.

We see that if $(p, s) \in \mathcal{L}$ and $n \leq x$, then $p^2 > n$, and therefore the p -adic order of $n!$ is

$$\text{ord}_p n! = \left\lfloor \frac{n}{p} \right\rfloor.$$

For $p \in \mathcal{P}_\alpha$ we also have

$$\text{ord}_p(2^{\ell(p)} - 1) = \alpha.$$

Clearly, if $n \geq p$ then $\text{ord}_p(n! + 2^n - 1) > 0$ only for $n \equiv 0 \pmod{\ell(p)}$. Because $\ell(p^{\alpha+1}) = p\ell(p) \gg x^{3/2}/(\log x)^2 > x$, we see that, for $p \leq n \leq x$,

$$\text{ord}_p(2^n - 1) = \begin{cases} 0, & \text{if } n \not\equiv 0 \pmod{\ell(p)}, \\ \alpha, & \text{if } n \equiv 0 \pmod{\ell(p)}. \end{cases}$$

Therefore, for $n \leq \alpha p - 1$ and $n \equiv 0 \pmod{\ell(p)}$, we have

$$\text{ord}_p(n! + 2^n - 1) \leq \text{ord}_p n! < n/(p - 1) \ll \log x.$$

Thus, $T(x_p, \alpha p - 1, p^s) = 0$ for $(p, s) \in \mathcal{L}_\alpha$.

On the other hand, for $n \geq (\alpha + 1)p$, we have $\text{ord}_p(n!) > n/p - 1 \geq \alpha$. Hence, for $n \equiv 0 \pmod{\ell(p)}$, we derive

$$\text{ord}_p(n! + 2^n - 1) = \text{ord}_p(2^n - 1) = \alpha < n/p \ll \log x.$$

As we have mentioned $\text{ord}_p(n! + 2^n - 1) = 0$ for every $n \geq p$ with $n \equiv 0 \pmod{\ell(p)}$. Thus, $T((\alpha + 1)p, x, p^s) = 0$ for $(p, s) \in \mathcal{L}_\alpha$.

For $\alpha = 1, 2, \dots$, let us define

$$Y_{\alpha,p} = \min\{x, \alpha p - 1\} \quad \text{and} \quad X_{\alpha,p} = \min\{x, (\alpha + 1)p\}.$$

We then have

$$V = \sum_{\alpha=1}^{\infty} V_\alpha,$$

where

$$V_\alpha = \sum_{(p,s) \in \mathcal{L}_\alpha} T(x_p, x, p^s) \log p.$$

For every $\alpha \geq 1$, and $(p, s) \in \mathcal{L}_\alpha$, as we have seen,

$$T(x_p, x, p^s) = T(Y_{\alpha,p}, X_{\alpha,p}, p^s).$$

We now need the bound,

$$(3.8) \quad T(Y_{\alpha,p}, X_{\alpha,p}, p^s) \leq \frac{X_{\alpha,p} - Y_{\alpha,p}}{\ell(p)} + 1,$$

which is a modified version of (2.2). Indeed, if $Y_{\alpha,p} = x$ then $X_{\alpha,p} = x$ and we count solutions in an empty interval. If $Y_{\alpha,p} = \alpha p - 1$ (the other alternative), we then replace the congruence modulo p^s by the congruence modulo p and remark that because $n > Y_{\alpha,p} \geq p$ we have $n! + 2^n - 1 \equiv 2^n - 1 \pmod{p}$ and (3.8) is now immediate.

We use (3.8) for $x^{1/2}/(\log x)^2 \geq s \geq x^{1/4}$, and Lemma 2.3 for $x/(\log x)^2 > s \geq x^{1/2}/(\log x)^2$. Simple calculations lead to the bound

$$V_\alpha \leq (1 + o(1)) (\log x)^2 \sum_{p \in \mathcal{P}_\alpha} (X_{\alpha,p} - Y_{\alpha,p}) + O(x^2).$$

We now have

$$\sum_{p \in \mathcal{P}_\alpha} (X_{\alpha,p} - Y_{\alpha,p}) = \sum_{\substack{p \in \mathcal{P}_\alpha \\ p \leq x/(\alpha+1)}} (p + 1) + \sum_{\substack{p \in \mathcal{P}_\alpha \\ x/(\alpha+1) < p \leq (x+1)/\alpha}} (x - \alpha p + 1).$$

Thus, putting everything together, and taking into account that the sets \mathcal{P}_α , $\alpha = 1, 2, \dots$, are disjoint, we derive

$$\begin{aligned} V &\leq (1 + o(1)) (\log x)^2 \left(\sum_{p \leq x/2} p + \sum_{\alpha=1}^{\infty} \sum_{x/(\alpha+1) < p \leq (x+1)/\alpha} (x - \alpha p) \right) \\ &= (1 + o(1)) (\log x)^2 \\ &\quad \times \left(\frac{x^2}{8 \log x} + \frac{x^2}{\log x} \sum_{\alpha=1}^{\infty} \left(\frac{1}{\alpha(\alpha+1)} - \frac{2\alpha+1}{2\alpha(\alpha+1)^2} \right) \right) \\ &= (1 + o(1)) (\log x)^2 \left(\frac{x^2}{8 \log x} + \frac{x^2}{2 \log x} \sum_{\alpha=1}^{\infty} \frac{1}{\alpha(\alpha+1)^2} \right) \\ &= (1 + o(1)) (\log x)^2 \\ &\quad \times \left(\frac{x^2}{8 \log x} + \frac{x^2}{2 \log x} \sum_{\alpha=1}^{\infty} \left(\frac{1}{\alpha(\alpha+1)} - \frac{1}{(\alpha+1)^2} \right) \right) \\ &= (1 + o(1)) (\log x)^2 \left(\frac{x^2}{8 \log x} + \frac{x^2}{2 \log x} \left(2 - \frac{\pi^2}{6} \right) \right). \end{aligned}$$

Hence

$$(3.9) \quad V \leq \left(\frac{27 - 2\pi^2}{24} + o(1) \right) x^2 \log x.$$

Substituting (3.7) and (3.9) in (3.6), and using (3.5), we derive

$$\frac{3(\lambda - 1)}{4} + \frac{27 - 2\pi^2}{24} \geq \frac{1}{2},$$

which contradicts the assumption $\lambda < (2\pi^2 + 3)/18$, and thus finishes the proof. □

Theorem 3.2. *For any sufficiently large x , we have:*

$$\omega \left(\prod_{1 \leq n \leq x} (n! + 2^n - 1) \right) \gg \frac{x}{\log x}.$$

Proof. In the notation of the proof of Theorem 3.1, we derive from (3.2) and Lemma 2.3, that

$$r_p \ll \sum_{1 \leq s \leq s_p} \frac{x \log x}{s \log p} + 1 \ll \frac{x \log x \log(s_p + 1)}{\log p} + s_p.$$

Obviously $s_p \ll x \log x / \log p$, therefore $r_p \ll x(\log x)^2 / \log p$. Thus, for any prime number p ,

$$p^{r_p} = \exp \left(O \left(x(\log x)^2 \right) \right),$$

which together with (3.1) finishes the proof. \square

4. Remarks

We recall the result of Fouvry [5], which asserts that $P(p-1) \geq p^{0.668}$ holds for a set of primes p of positive relative density (see also [1, 2] for this and several more related results). By Lemma 2.4, this immediately implies that $\ell(p) \geq p^{0.668}$ for a set of primes p of positive relative density. Using this fact in our arguments, one can easily derive that actually

$$\limsup_{n \rightarrow \infty} \frac{P(n! + 2^n - 1)}{n} > \frac{2\pi^2 + 3}{18}.$$

However, the results of [5], or other similar results like the ones from [1, 2], do not give any effective bound on the relative density of the set of primes with $P(p-1) \geq p^{0.668}$, and thus cannot be used to get an explicit numerical improvement of Theorem 3.1.

We also remark that, as in [7], one can use lower bounds on linear forms in p -adic logarithms to obtain an “individual” lower bound on $P(n! + 2^n - 1)$. The *ABC-conjecture* can also be used in the same way as in [8] for $P(n! + 1)$.

References

- [1] R. C. BAKER, G. HARMAN, *The Brun-Titchmarsh theorem on average*. Analytic number theory, Vol. 1 (Allerton Park, IL, 1995), Progr. Math. **138**, Birkhäuser, Boston, MA, 1996, 39–103.
- [2] R. C. BAKER, G. HARMAN, *Shifted primes without large prime factors*. Acta Arith. **83** (1998), 331–361.
- [3] P. ERDŐS, R. MURTY, *On the order of $a \pmod{p}$* . Proc. 5th Canadian Number Theory Association Conf., Amer. Math. Soc., Providence, RI, 1999, 87–97.
- [4] P. ERDŐS, C. STEWART, *On the greatest and least prime factors of $n! + 1$* . J. London Math. Soc. **13** (1976), 513–519.
- [5] É. FOUVRY, *Théorème de Brun-Titchmarsh: Application au théorème de Fermat*. Invent. Math. **79** (1985), 383–407.
- [6] H.-K. INDLEKOFER, N. M. TIMOFEEV, *Divisors of shifted primes*. Publ. Math. Debrecen **60** (2002), 307–345.
- [7] F. LUCA, I. E. SHPARLINSKI, *Prime divisors of shifted factorials*. Bull. London Math. Soc. **37** (2005), 809–817.
- [8] M.R. MURTY, S. WONG, *The ABC conjecture and prime divisors of the Lucas and Lehmer sequences*. Number theory for the millennium, III (Urbana, IL, 2000), A K Peters, Natick, MA, 2002, 43–54.
- [9] F. PAPPALARDI, *On the order of finitely generated subgroups of $\mathbb{Q}^* \pmod{p}$ and divisors of $p-1$* . J. Number Theory **57** (1996), 207–222.
- [10] K. PRACHAR, *Primzahlverteilung*. Springer-Verlag, Berlin, 1957.

Florian LUCA
Instituto de Matemáticas
Universidad Nacional Autónoma de México
C.P. 58089, Morelia, Michoacán, México
E-mail : `fluca@matmor.unam.mx`

Igor E. SHPARLINSKI
Department of Computing
Macquarie University
Sydney, NSW 2109, Australia
E-mail : `igor@ics.mq.edu.au`