

# ANNALES DE L'I. H. P., SECTION A

W. M. OLIVA

## **On the chaotic behavior and the non-integrability of the four vortices problem**

*Annales de l'I. H. P., section A*, tome 55, n° 2 (1991), p. 707-718

[http://www.numdam.org/item?id=AIHPA\\_1991\\_\\_55\\_2\\_707\\_0](http://www.numdam.org/item?id=AIHPA_1991__55_2_707_0)

© Gauthier-Villars, 1991, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## **On the chaotic behavior and the non-integrability of the four vortices problem**

by

**W. M. OLIVA**

I.M.E., Universidade de São Paulo, Caixa Postal 20570,  
05508 São Paulo SP, Brazil

---

**ABSTRACT.** — The paper is based in a survey talk given by the author in 1990 about the vortex model. Some brief comments are made about the works of Ziglin and Khanin on the chaotic behavior, non-integrability and quasi-periodicity of the vortex motions in the case of four points vortices; some other ideas about a new approach to non-integrability are also presented.

**RÉSUMÉ.** — Cet article est basé sur un exposé de revue donné par l'auteur en 1990 sur le modèle de vortex. On commente brièvement les travaux de Ziglin et Khanin sur le comportement chaotique, la non-intégrabilité et la quasi-périodicité du modèle dans le cas de quatre vortices ponctuels. On présente aussi d'autres idées sur une nouvelle approche de la non intégrabilité.

---

### **1. THE VORTEX MODEL**

For detailed presentations about the vortex model in fluid mechanics we refer the reader to Chorin and Marsden (1979) and to Marchioro and Pulvirenti (1983).

Choose constants  $K_1, \dots, K_N$  and initial positions

$$\mathbf{x}_1 = (x_1, y_1), \dots, \mathbf{x}_N = (x_N, y_N)$$

in the plane. The velocity  $\mathbf{u}_j(\mathbf{x}, t)$  of  $\mathbf{x} = (x, y) \in \mathbf{R}^2$  due to the  $j$ th vortex is given by

$$\mathbf{u}_j(\mathbf{x}, t) = \left( -\frac{K_j}{2\pi} \frac{(y-y_j)}{|\mathbf{x}-\mathbf{x}_j|^2}, \frac{K_j}{2\pi} \frac{(x-x_j)}{|\mathbf{x}-\mathbf{x}_j|^2} \right), \quad (1)$$

if we ignore the other vortices. When all the vortices are moving, they produce the velocity field  $\mathbf{u}(\mathbf{x}, t) = \sum_{j=1}^N \mathbf{u}_j(\mathbf{x}, t)$  and since each vortex ought to move as it was carried by the net velocity field of the other vortices, then each  $\mathbf{x}_j$ ,  $j=1, \dots, N$ , moves according to the equations:

$$\left. \begin{aligned} \dot{x}_j &= -\frac{1}{2\pi} \sum_{i \neq j} \frac{K_i (y_j - y_i)}{r_{ij}^2} \\ \dot{y}_j &= \frac{1}{2\pi} \sum_{i \neq j} \frac{K_i (x_j - x_i)}{r_{ij}^2}, \quad r_{ij} = |\mathbf{x}_i - \mathbf{x}_j| \end{aligned} \right\} \quad (2)$$

or, equivalently, for  $i, j=1, \dots, N$ :

$$\left. \begin{aligned} K_j \dot{x}_j &= \frac{\partial H}{\partial y_j} \\ K_j \dot{y}_j &= -\frac{\partial H}{\partial x_j}, \quad H = -\frac{1}{4\pi} \sum_{i \neq j} K_i K_j \log |\mathbf{x}_i - \mathbf{x}_j|. \end{aligned} \right\} \quad (3)$$

The construction of the velocity field  $\mathbf{u}(\mathbf{x}, t)$  produces formal solutions of the Euler's equation in  $\mathbf{R}^2$  and have the property that the classical circulation theorems are satisfied (*see op. cit.* above).

Poincaré and Kirchoff observed that in a new system of coordinates of  $\mathbf{R}^{2N}$ ,  $(x'_\alpha, y'_\alpha)$ , given by

$$\left. \begin{aligned} x'_\alpha &= \sqrt{|K_\alpha|} x_\alpha \\ y'_\alpha &= \sqrt{|K_\alpha|} \text{sign}(K_\alpha) y_\alpha, \quad \alpha = 1, \dots, N, \end{aligned} \right\} \quad (4)$$

the system of ordinary differential equations (3) becomes a Hamiltonian system. The phase space is an open dense set of  $\mathbf{R}^{2N}$  since  $r_{ij} \neq 0$  (collisions of vortices are not allowed).

It is easy to see that, because of the symmetries of the Hamiltonian  $H$ , the system of equations (3) has the four first integrals:

$$I_1 = H, \quad I_2 = \sum_{\alpha=1}^N K_\alpha x_\alpha,$$

$$I_3 = \sum_{\alpha=1}^N K_\alpha y_\alpha \quad \text{and} \quad I_4 = \sum_{\alpha=1}^N K_\alpha (x_\alpha^2 + y_\alpha^2).$$

For the Poisson bracket corresponding to the canonical symplectic 2-form of  $\mathbf{R}^{2n}$ :  $\omega = \sum_{\alpha=1}^N dx'_\alpha \wedge dy'_\alpha$ , one has  $\{I_2^2 + I_3^2, I_4\} = 0$ . In fact,

$$\{I_2^2 + I_3^2, I_4\} = \sum_{i=1}^N \left[ \frac{\partial I_4}{\partial y'_i} \frac{\partial (I_2^2 + I_3^2)}{\partial x'_i} - \frac{\partial I_4}{\partial x'_i} \frac{\partial (I_2^2 + I_3^2)}{\partial y'_i} \right] = 0$$

[see Aref and Pomphrey (1982)]. The vortex systems, for  $N=2$  and  $N=3$ , are then Liouville integrable. The existence of four independent integrals of motion suggest, at first sight, that the integrability of system (3) is possible in the case of  $N=4$  vortices. But this is not, necessarily, the case because the integrals  $I_2, I_3$  and  $I_4$  do not commute, that is, their Poisson brackets do not vanish, since:

$$\{I_2, I_3\} = \sum_{\alpha} K_\alpha, \quad \{I_2, I_4\} = 2I_3 \quad \text{and} \quad \{I_3, I_4\} = -2I_2.$$

The case of positive *intensities* ( $K_\alpha > 0, \alpha=1, \dots, N$ ) implies that all solutions in  $\mathbf{R}^{2N}$  are bounded (since  $I_4 = \text{const.}$  is compact) and then defined for all time; in particular when  $N=2$  or 3 the phase space has regions foliated by invariant tori, then all the solutions define quasi-periodic motions. Using carefully KAM theory, Khanin (1982) showed that in the phase space of any system with an arbitrary number of vortices there exists a set of initial conditions of positive measure for which the motions of vortices are quasi-periodic (see Khanin 1982).

Questions related to chaotic behavior and non integrability of the four vortices problem will be considered in the next section.

In the paper the motions of vortices happen in an unbounded domain, that is,  $\Gamma = \mathbf{R}^2$ . The case of  $\Gamma$  bounded presents also many interesting questions; see for instance Dürr and Pulvirenti (1982), Marchioro and Pulvirenti (1983) and the survey paper Aref (1983).

## 2. THE PROBLEM OF FOUR VORTICES

Fix a system of coordinates in  $\mathbf{R}^2$  and consider four vortices  $P_\alpha$  of coordinates  $\mathbf{x}_\alpha = (x_\alpha, y_\alpha), \alpha=1, \dots, 4$ . Assume  $K_i = 1$  for  $i=1, 2, 3$  and

$K_4 = \varepsilon > 0$ . The equations of motion are given by

$$\left. \begin{aligned} \dot{x}_i &= \frac{\partial H}{\partial y_i} = \frac{\partial H_0}{\partial y_i} + \varepsilon \frac{\partial H_1}{\partial y_i}, \\ \dot{y}_i &= -\frac{\partial H}{\partial x_i} = -\frac{\partial H_0}{\partial x_i} - \varepsilon \frac{\partial H_1}{\partial x_i}, \quad i = 1, 2, 3, \\ \varepsilon \dot{x}_4 &= \frac{\partial H}{\partial y_4} = \varepsilon \frac{\partial H_1}{\partial y_4}, \quad \varepsilon \dot{y}_4 = -\frac{\partial H}{\partial x_4} = -\varepsilon \frac{\partial H_1}{\partial x_4}, \end{aligned} \right\} \quad (5)$$

where

$$H = H_0 + \varepsilon H_1, \quad H_0 = -\frac{1}{4\pi} (\log r_{12}^2 + \log r_{23}^2 + \log r_{13}^2),$$

$$H_1 = -\frac{1}{4\pi} (\log r_{14}^2 + \log r_{24}^2 + \log r_{34}^2)$$

and

$$r_{t,s} = |\mathbf{x}_t - \mathbf{x}_s|, \quad t, s = 1, \dots, 4.$$

Recall that  $x'_i = x_i$ ,  $y'_i = y_i$ ,  $x'_4 = \sqrt{\varepsilon} x_4$  and  $y'_4 = \sqrt{\varepsilon} y_4$  are canonical coordinates in  $\mathbf{R}^8$  and the system (5) becomes a Hamiltonian system in the new coordinates; the symplectic canonical form is

$$\omega = \sum_{i=1}^3 dx_i \wedge dy_i + dx'_4 \wedge dy'_4.$$

The motions for the 3-equal vortices are given by

$$\dot{x}_i = \frac{\partial H_0}{\partial y_i}, \quad \dot{y}_i = -\frac{\partial H_0}{\partial x_i}, \quad i = 1, 2, 3, \quad (6)$$

and the *restricted 4-vortices problem* corresponds to the motions of three vortices with equal unit intensities and the fourth vortex not influencing the three first ones. It is, precisely, given by system (5) as  $\varepsilon \rightarrow 0$ , that is, the decoupled system:

$$\left. \begin{aligned} \dot{x}_i &= \frac{\partial H_0}{\partial y_i}, \quad \dot{y}_i = -\frac{\partial H_0}{\partial x_i}, \quad i = 1, 2, 3, \\ \dot{x}_4 &= \frac{\partial H_1}{\partial y_4}, \quad \dot{y}_4 = -\frac{\partial H_1}{\partial x_4}. \end{aligned} \right\} \quad (7)$$

Ziglin, in two papers [Ziglin (1980) and Khanin (1982) – appendix], considered the question of nonintegrability of a problem of four point vortices. He started with system (7), that is, with the restricted 4-vortices problem. Let  $a_i$ ,  $i = 1, 2, 3$  be the sides of the triangle determined by the three unit vortices, and  $A_i$ ,  $i = 1, 2, 3$  be the angles opposite to the corresponding sides. Then the relative problem of the three vortices has

the following equations derived from (6) or (7)<sub>1</sub>:

$$\left. \begin{aligned} \dot{a}_1 &= \frac{1}{2\pi} \left( \frac{\sin A_3}{a_2} - \frac{\sin A_2}{a_3} \right) \\ \dot{a}_2 &= \frac{1}{2\pi} \left( \frac{\sin A_1}{a_3} - \frac{\sin A_3}{a_1} \right) \\ \dot{a}_3 &= \frac{1}{2\pi} \left( \frac{\sin A_2}{a_1} - \frac{\sin A_1}{a_2} \right) \end{aligned} \right\} \quad (8)$$

and admits the two independent first integrals:  $a_1 a_2 a_3 = c_1^3$  and  $a_1^2 + a_2^2 + a_3^2 = c_2^2$ .

Substituting  $a_3 = c_1^3/a_1 a_2$  into (8) he obtained the system  $\dot{a} = F(a, c_2)$ ,  $a = (a_1, a_2)$ , which has a center  $a_0 = (c_1, c_1)$ , an equilateral triangle; after this, he took the periodic solutions close to this elliptical fixed point which are given by a one-parameter family of periodic functions; choosing properly a small parameter for this family, he substituted these periodic functions in equations (7)<sub>2</sub> of the fourth vortex. This way he obtained a time dependent Hamiltonian system:

$$\frac{d\xi}{d\tau} = \frac{\partial F}{\partial \eta}, \quad \frac{d\eta}{d\tau} = -\frac{\partial F}{\partial \xi}, \quad (9)$$

where  $F = F(\xi, \eta, \tau, \mu) = F_0(\xi, \eta) + \mu F_1(\xi, \eta, \tau, \mu) + \dots$ . The unperturbed Hamiltonian system ( $\mu = 0$ ) is defined by  $F_0(\xi, \eta)$  and the corresponding phase portrait has a homoclinic fixed point. As usual [see Holmes (1980)], for  $\mu \neq 0$  it is necessary to examine system (9) in the extended phase space  $\{\xi, \eta, \tau \pmod{2\pi}\}$  and consider the Poincaré-map of the plane  $\{\tau \pmod{2\pi}\} = 0$  to itself, given by the cylindrical phase-flow. In general, for  $\mu \neq 0$ , the homoclinic orbit of the hyperbolic fixed point of this map will split in two transversal separatrices. In this case the perturbed Poincaré-map presents a chaotic behavior [see Moser (1973) and Smale (1967)] due to the occurrence of a horseshoe; in particular, no domain containing the closure of the trajectory of that transverse homoclinic point admits an analytic first integral for the Poincaré map. Thus, the proof of the non integrability reduces to checking that the separatrices intersect transversally. In Ziglin (1980) the author reduced the proof to the non vanishing of an integral (see Melnikov, 1963) and he succeeded in showing this, after making the evaluation by computer. In Khanin (1982 – appendix), Ziglin argued only using continuity and stated the non-integrability of a four-vortices problem of intensities  $(K_1, K_2, K_3, K_4)$  sufficiently close to the  $(1, 1, 1, 0)$  of the restricted case.

In Holmes and Marsden (1982) the authors say that “the problem treated by Ziglin possesses some difficulties that cannot be avoided without a careful study”; also, according to Aref [see Aref (1983)], Marsden in a private communication say that “the proof of Ziglin is incomplete for

some reasons, and similar criticism applies to the extension of the proof in Khanin (1982)".

In a joint work, in progress, with V. Moauro, P. Negrini and M. S. A. C. Castilla, we decided to come back to the question of the non-integrability for the problem of 4-vortices. From now on I will present some rough ideas of our work.

First of all, let  $O$  be the origin of  $\mathbf{R}^2$ ,  $M_1$  be the center of mass of  $P_i = O + \mathbf{x}_i$ ,  $i = 1, 2, 3$ , with unitary masses, and  $M_0$  be the center of mass of the four vortices, the last one  $P_4 = O + \mathbf{x}_4$  with mass  $\varepsilon > 0$ , that is,

$$M_0 - O = \left( \frac{\sum_{i=1}^3 x_i + \varepsilon x_4}{3 + \varepsilon}, \frac{\sum_{i=1}^3 y_i + \varepsilon y_4}{3 + \varepsilon} \right) = \frac{1}{3 + \varepsilon} (I_2, I_3)$$

and

$$M_1 - O = \frac{1}{3} \left( \sum_{i=1}^3 x_i, \sum_{i=1}^3 y_i \right).$$

Let us introduce now a new system of coordinates  $(\sqrt{\varepsilon} x_4, \sqrt{\varepsilon} y_4; \theta_1, \tilde{p}_1; \theta_2, \tilde{p}_2; \theta_3, \tilde{p}_3)$  by:

$$\begin{aligned} P_1 - P_2 &= \alpha \sqrt{\tilde{p}_1} e^{i\theta_1} \\ P_3 - M_1 &= \beta \sqrt{\tilde{p}_2} e^{i\theta_2} \\ P_4 - M_1 &= \gamma \sqrt{\tilde{p}_3} e^{i\theta_3}, \end{aligned}$$

where  $\alpha, \beta, \gamma$  will be determined in order that the new coordinates will be canonical.

From the definitions of  $M_1$  and  $M_0$  we have  $(P_1 - M_1) + (P_2 - M_1) + (P_3 - M_1) = 0$  and  $3(M_1 - M_0) + \varepsilon(P_4 - M_0) = 0$  that imply

$$\begin{aligned} P_1 - M_0 &= \frac{1}{2} \left[ (P_1 - P_2) - (P_3 - M_1) - \frac{2\varepsilon}{3 + \varepsilon} (P_4 - M_1) \right] \\ P_2 - M_0 &= -\frac{1}{2} \left[ (P_1 - P_2) + \frac{2\varepsilon}{3 + \varepsilon} (P_4 - M_1) + (P_3 - M_1) \right] \\ P_3 - M_0 &= (P_3 - M_1) - \frac{\varepsilon}{3 + \varepsilon} (P_4 - M_1). \end{aligned}$$

In cartesian coordinates we have then:

$$\begin{aligned} x_1 &= \frac{1}{3 + \varepsilon} I_2 + \frac{1}{2} \alpha \sqrt{\tilde{p}_1} \cos \theta_1 - \frac{1}{2} \beta \sqrt{\tilde{p}_2} \cos \theta_2 - \frac{\gamma \varepsilon}{3 + \varepsilon} \sqrt{\tilde{p}_3} \cos \theta_3 \\ y_1 &= \frac{1}{3 + \varepsilon} I_3 + \frac{1}{2} \alpha \sqrt{\tilde{p}_1} \sin \theta_1 - \frac{1}{2} \beta \sqrt{\tilde{p}_2} \sin \theta_2 - \frac{\gamma \varepsilon}{3 + \varepsilon} \sqrt{\tilde{p}_3} \sin \theta_3 \end{aligned}$$

$$\begin{aligned}
 x_2 &= \frac{1}{3+\varepsilon} I_2 - \frac{1}{2} \alpha \sqrt{\tilde{p}_1} \cos \theta_1 - \frac{1}{2} \beta \sqrt{\tilde{p}_2} \cos \theta_2 - \frac{\gamma \varepsilon}{3+\varepsilon} \sqrt{\tilde{p}_3} \cos \theta_3 \\
 y_2 &= \frac{1}{3+\varepsilon} I_3 - \frac{1}{2} \alpha \sqrt{\tilde{p}_1} \sin \theta_1 - \frac{1}{2} \beta \sqrt{\tilde{p}_2} \sin \theta_2 - \frac{\gamma \varepsilon}{3+\varepsilon} \sqrt{\tilde{p}_3} \sin \theta_3 \\
 x_3 &= \frac{1}{3+\varepsilon} I_2 + \beta \sqrt{\tilde{p}_2} \cos \theta_2 - \frac{\gamma \varepsilon}{3+\varepsilon} \sqrt{\tilde{p}_3} \cos \theta_3 \\
 y_3 &= \frac{1}{3+\varepsilon} I_3 + \beta \sqrt{\tilde{p}_2} \sin \theta_2 - \frac{\gamma \varepsilon}{3+\varepsilon} \sqrt{\tilde{p}_3} \sin \theta_3 \\
 x_4 &= \frac{1}{3+\varepsilon} I_2 + \frac{3\gamma}{3+\varepsilon} \sqrt{\tilde{p}_3} \cos \theta_3 \\
 y_4 &= \frac{1}{3+\varepsilon} I_2 + \frac{3\gamma}{3+\varepsilon} \sqrt{\tilde{p}_3} \sin \theta_3.
 \end{aligned}$$

The preservation of the symplectic form is given by the equalities

$$\omega = \varepsilon dx_4 \wedge dy_4 + \sum_{i=1}^3 dx_i \wedge dy_i = \varepsilon dx_4 \wedge dy_4 + \sum_{i=1}^3 d\tilde{p}_i \wedge d\theta_i.$$

Making the origin  $O$  to be equal to the center of mass  $M_0$  and fixing  $\alpha=2$ ,  $\beta=\frac{2\sqrt{3}}{3}$  and  $\gamma=\sqrt{\frac{2(3+\varepsilon)}{3\varepsilon}}$ , we see that the new coordinates are canonical.

Remark that  $\tilde{p}_3 = \frac{3\varepsilon}{2(3+\varepsilon)} |P_4 - M_1|^2$ , that is,  $\tilde{p}_3 = O(\varepsilon)$ .

The distances between vortices are given by:

$$\begin{aligned}
 r_{12}^2 &= 4\tilde{p}_1 \\
 r_{13}^2 &= \tilde{p}_1 + 3\tilde{p}_2 - 2\sqrt{3}\sqrt{\tilde{p}_1\tilde{p}_2} \cos(\theta_1 - \theta_2) \\
 r_{23}^2 &= \tilde{p}_1 + 3\tilde{p}_2 + 2\sqrt{3}\sqrt{\tilde{p}_1\tilde{p}_2} \cos(\theta_1 - \theta_2) \\
 r_{14}^2 &= \tilde{p}_1 + \frac{1}{3}\tilde{p}_2 + \frac{2}{3} \frac{(3+\varepsilon)}{\varepsilon} \tilde{p}_3 - \frac{2}{\sqrt{3}} \sqrt{\tilde{p}_1\tilde{p}_2} \cos(\theta_1 - \theta_2) \\
 &\quad - 2\sqrt{\frac{2}{3}} \sqrt{\frac{3+\varepsilon}{\varepsilon}} \sqrt{\tilde{p}_1\tilde{p}_3} \cos(\theta_1 - \theta_3) \\
 &\quad + \frac{2\sqrt{2}}{3} \sqrt{\frac{3+\varepsilon}{\varepsilon}} \sqrt{\tilde{p}_2\tilde{p}_3} \cos(\theta_2 - \theta_3)
 \end{aligned}$$



$$\begin{aligned}
r_{24}^2 &= \tilde{p}_1 + \frac{1}{3}\tilde{p}_2 + \frac{2}{3} \frac{(3+\varepsilon)}{\varepsilon} \tilde{p}_3 + \frac{2}{\sqrt{3}} \sqrt{\tilde{p}_1 \tilde{p}_2} \cos(\theta_1 - \theta_2) \\
&\quad + 2 \sqrt{\frac{2}{3}} \sqrt{\frac{3+\varepsilon}{\varepsilon}} \sqrt{\tilde{p}_1 \tilde{p}_3} \cos(\theta_1 - \theta_2) \\
&\quad\quad\quad + \frac{2\sqrt{2}}{3} \sqrt{\frac{3+\varepsilon}{\varepsilon}} \sqrt{\tilde{p}_2 \tilde{p}_3} \cos(\theta_2 - \theta_3) \\
r_{34}^2 &= \frac{4}{3}\tilde{p}_2 + \frac{2}{3} \frac{(3+\varepsilon)}{\varepsilon} \tilde{p}_3 - \frac{4\sqrt{2}}{3} \sqrt{\frac{3+\varepsilon}{\varepsilon}} \sqrt{\tilde{p}_2 \tilde{p}_3} \cos(\theta_2 - \theta_3)
\end{aligned}$$

The Hamiltonian function  $H$  becomes

$$\begin{aligned}
-4\pi H &= -4\pi H_0 - 4\pi\varepsilon H_1 = [\log r_{12}^2 r_{13}^2 r_{23}^2](\tilde{p}_1, \tilde{p}_2, \theta_1 - \theta_2) \\
&\quad + \varepsilon [\log r_{14}^2 r_{24}^2 r_{34}^2](\tilde{p}_1, \tilde{p}_2, \tilde{p}_3, \theta_1 - \theta_2, \theta_1 - \theta_3, \theta_2 - \theta_3, \varepsilon),
\end{aligned}$$

then, the Hamiltonian system of energy  $H$  has a first integral  $\tilde{p}_1 + \tilde{p}_2 + \tilde{p}_3$  and, of course, depends on  $\varepsilon$  as a real parameter.

Choosing a new system of coordinates:

( $\sqrt{\varepsilon}x_4, \sqrt{\varepsilon}y_4; \varphi_1, p_1; \varphi_2, p_2; \varphi_3, p_3$ ) defined by

$$\begin{aligned}
p_1 &= \tilde{p}_1, & p_2 &= \tilde{p}_1 + \tilde{p}_2 + \tilde{p}_3, & p_3 &= \tilde{p}_3 \\
\varphi_1 &= \theta_1 - \theta_2, & \varphi_2 &= \theta_2, & \varphi_3 &= \theta_3 - \theta_2,
\end{aligned}$$

and since

$$\begin{aligned}
dp_1 \wedge d\varphi_1 + dp_2 \wedge d\varphi_2 + dp_3 \wedge d\varphi_3 \\
&= d\tilde{p}_1 \wedge (d\theta_1 - d\theta_2) + (d\tilde{p}_1 + d\tilde{p}_2 + d\tilde{p}_3) \wedge d\theta_2 + d\tilde{p}_3 \wedge (d\theta_3 - d\theta_2) \\
&= d\tilde{p}_1 \wedge d\theta_1 + d\tilde{p}_2 \wedge d\theta_2 + d\tilde{p}_3 \wedge d\theta_3,
\end{aligned}$$

one sees that the new coordinates are also canonical and the new Hamiltonian  $\tilde{H}$  does not depend on  $\varphi_2$ , that is:

$$\tilde{H}(p_1, p_2, p_3, \varphi_1, \varphi_3, \varepsilon) \stackrel{\text{def}}{=} \tilde{H}_0 + \varepsilon \tilde{H}_1$$

where

$$\tilde{H}_0 = H_0(p_1, p_2 - p_1 - p_3, \varphi_1)$$

and  $\tilde{H}_1 = H_1(p_1, p_2 - p_1 - p_3, p_3, \varphi_1, \varphi_1 - \varphi_3, -\varphi_3, \varepsilon)$ . Then

$$\left. \begin{aligned} \dot{p}_1 &= \frac{\partial \tilde{H}}{\partial \varphi_1} = \frac{\partial \tilde{H}_0}{\partial \varphi_1} + \varepsilon \frac{\partial \tilde{H}_1}{\partial \varphi_1} \\ \dot{\varphi}_1 &= -\frac{\partial \tilde{H}}{\partial p_1} = -\frac{\partial \tilde{H}_0}{\partial p_1} - \varepsilon \frac{\partial \tilde{H}_1}{\partial p_1} \\ \dot{p}_2 &= 0 \\ \dot{\varphi}_2 &= -\frac{\partial \tilde{H}}{\partial p_2} = -\frac{\partial \tilde{H}_0}{\partial p_2} - \varepsilon \frac{\partial \tilde{H}_1}{\partial p_2} \\ \dot{p}_3 &= \frac{\partial \tilde{H}}{\partial \varphi_3} = \varepsilon \frac{\partial \tilde{H}_1}{\partial \varphi_3} \\ \dot{\varphi}_3 &= -\frac{\partial \tilde{H}}{\partial p_3} = -\frac{\partial \tilde{H}_0}{\partial p_3} - \varepsilon \frac{\partial \tilde{H}_1}{\partial p_3} \end{aligned} \right\} \quad (10)$$

The system above has the coordinate  $p_2$  as a first integral.

With an analogous procedure, we obtain the equations corresponding to the problem of three equal unitary vortices. But it is an easy matter to get them just making  $\varepsilon = 0$  and  $\tilde{p}_3 = O(\varepsilon) = 0$  in the first four equations of system (10):


$$\left. \begin{aligned} \dot{p}_1 &= \frac{1}{\pi} e^{4\pi H_0} [-24 p_1^2 (p_2 - p_1) \cos \varphi_1 \sin \varphi_1] \\ \dot{\varphi}_1 &= -\frac{1}{\pi} e^{4\pi H_0} [(24 p_1 p_2 - 36 p_1^2) \cos^2 \varphi_1 \\ &\quad - (3 p_2 - 2 p_1)^2 + 4 p_1 (3 p_2 - 2 p_1)] \\ &\quad p_2 = \mu = \text{const.} \\ \dot{\varphi}_2 &= -\frac{1}{\pi} e^{4\pi H_0} [12 p_1^2 \cos^2 \varphi_1 - 6 p_1 (3 p_2 - 2 p_1)] \end{aligned} \right\} \quad (11)$$

where

$$\begin{aligned} H_0 &= -\frac{1}{4\pi} \log r_{12}^2 r_{13}^2 r_{23}^2 \\ &= -\frac{1}{4\pi} \log \{ 4 p_1 [(3 p_2 - 2 p_1)^2 - 12 p_1 (p_2 - p_1) \cos^2 \varphi_1] \}. \end{aligned}$$

The system (11) is decoupled; if  $\varphi_2$  is a new time, (11) is equivalent to the system given by  $\frac{dp_1}{d\varphi_2}$  and  $\frac{d\varphi_1}{d\varphi_2}$  which has the same orbits as

$$\left. \begin{aligned} \dot{p}_1 &= -24 p_1^2 (\mu - p_1) \cos \varphi_1 \sin \varphi_1 \\ \dot{\varphi}_1 &= -[(24 p_1 \mu - 36 p_1^2) \cos^2 \varphi_1 \\ &\quad - (3 \mu - 2 p_1)^2 + 4 p_1 (3 \mu - 2 p_1)]. \end{aligned} \right\} \quad (12)$$



The phase space of (12) is given by figure for  $p_1 < \mu$ ; it can be thought as an unperturbed system and presents heteroclinic connections as well as regular tori in regions where the KAM theory can be applied.

The perturbed system (10) can be reduced to a planar time-dependent Hamiltonian system. The idea will be presented in the sequel. Consider the motions on a level of energy  $\tilde{H}(p_1, p_2 = \text{const.}, p_3, \varphi_1, \varphi_3, \varepsilon) = h$  and try to solve (locally) this equation in  $p_3$ . For this we compute the derivative  $\frac{\partial \tilde{H}}{\partial p_3} = \frac{\partial \tilde{H}_0}{\partial p_3} + \varepsilon \frac{\partial \tilde{H}_1}{\partial p_3}$  and assume it is non zero; then one obtains, by the implicit function theorem, the function

$$p_3 = F(p_1, p_2 = \text{const.}, \varphi_1, \varphi_3, \varepsilon, h). \tag{13}$$

One can show that system (10), with  $p_2 = \text{const.}$  and  $\tilde{H} = h$ , can be reduced to (13) and (14):

$$\left. \begin{aligned} \frac{dp_1}{d\varphi_3} &= \frac{\partial F}{\partial \varphi_1}(p_1, p_2 = \text{const.}, \varphi_1, \varphi_3, \varepsilon, h) \\ \frac{d\varphi_1}{d\varphi_3} &= -\frac{\partial F}{\partial p_1}(p_1, p_2 = \text{const.}, \varphi_1, \varphi_3, \varepsilon, h), \end{aligned} \right\} \tag{14}$$

where  $\varphi_3$  is considered as a new time. In fact, if we assume (13), from the identity

$$\tilde{H}(p_1, p_2 = \text{const.}, F(p_1, p_2 = \text{const.}, \varphi_1, \varphi_3, \varepsilon, h), \varphi_1, \varphi_3, \varepsilon) = h$$

we obtain, by differentiation with respect to  $p_1$  and  $\varphi_1$ :

$$\left. \begin{aligned} \frac{\partial F}{\partial \varphi_1} &= -\frac{\partial \tilde{H}}{\partial \varphi_1} \Big/ \frac{\partial \tilde{H}}{\partial p_3} = -\left( \frac{\partial \tilde{H}_0}{\partial \varphi_1} + \varepsilon \frac{\partial \tilde{H}_1}{\partial \varphi_1} \right) \Big/ \left( \frac{\partial \tilde{H}_0}{\partial p_3} + \varepsilon \frac{\partial \tilde{H}_1}{\partial p_3} \right) \\ \frac{\partial F}{\partial p_1} &= -\frac{\partial \tilde{H}}{\partial p_1} \Big/ \frac{\partial \tilde{H}}{\partial p_3} = -\left( \frac{\partial \tilde{H}_0}{\partial p_1} + \varepsilon \frac{\partial \tilde{H}_1}{\partial p_1} \right) \Big/ \left( \frac{\partial \tilde{H}_0}{\partial p_3} + \varepsilon \frac{\partial \tilde{H}_1}{\partial p_3} \right) \end{aligned} \right\} \quad (15)$$

where  $p_3$  is given by (13). This, (13) and the fact that  $\varphi_3$  can be considered in (10) as a new time, gives us the equations (14).

Remark that the system (14) is a time dependent Hamiltonian planar system with variables  $p_1$ ,  $\varphi_1$  and is periodic in time  $\varphi_3$ ; moreover, equations (15) show that for  $\varepsilon=0$  the system (14) is autonomous.

The final comment is that if we examine (14) and (13) in the extended phase space  $\{p_1, \varphi_1, \varphi_3 \pmod{2\pi}\}$ , the Poincaré-map of the plane  $\{\varphi \pmod{2\pi}\} = 0$  to itself, given by the cylindrical phase-flow, presents, for  $\varepsilon=0$ , the same orbits as the phase space of Figure. The main point now is to complete the proof using the Melnikov method, showing that the saddle connection I-V (see Fig.) of the unperturbed Poincaré-map will split, after perturbation, in two transversal separatrices.

By a classical Poincaré result (see Kozlov, 1983), one can conclude that system (10) of four vortices is not analytically integrable for  $\varepsilon$  small.

## REFERENCES

- H. AREF and N. POMPHREY, Integrable and Chaotic Motions of Four Vortices. I. The Case of Identical Vortices, *Proc. R. Soc. London Ser. A*, Vol. **380**, 1982, pp. 359-387.
- H. AREF, Integrable, Chaotic and Turbulent Vortex Motion in Two-Dimensional flows, *Ann. Rev. Fluid Mech.*, Vol. **15**, 1983, pp. 345-389.
- A. J. CHORIN and J. E. MARSDEN, *A Mathematical Introduction to Fluid Mechanics*, Springer-Verlag, 1979.
- D. DÜRR and M. PULVIRENTI, On the Vortex Flow in Bounded Domain, *Commun. Math. Phys.*, Vol. **85**, 1982, pp. 265-273.
- P. J. HOLMES, Averaging and Chaotic Motions in Forced Oscillations, *S.I.A.M. J. Appl. Math.*, Vol. **38**, (1), 1980, pp. 65-79. Errata, Vol. **40**, (1), 1981, pp. 167-168.
- P. J. HOLMES and J. E. MARSDEN, Horseshoes in Perturbations of Hamiltonian System with Two Degrees of Freedom, *Commun. Math. Phys.*, Vol. **82**, 1982, pp. 523-544.
- K. M. KHANIN, Quasi-Periodic Motions of Vortex Systems, *Physica*, Vol. **4D**, 1982, pp. 261-269.
- V. V. KOZLOV, Integrability and Non-Integrability in Hamiltonian Mechanics, *Russian Math. Surveys*, Vol. **38.1**, 1983, pp. 1-76.
- C. MARCHIORO and M. PULVIRENTI, *Vortex Methods in Two Dimensional Fluid Dynamics*, Edizioni Klim, Roma, 1983.
- V. K. MELNIKOV, On the Stability of the Center of Time-Periodic Perturbations, *Trans. Moscow Math. Soc.*, Vol. **12**, (1), 1963, pp. 1-57.
- J. MOSER, *Stable and Random Motions in Dynamical Systems*, Princeton Univ. Press, Princeton N.J., Univ. of Tokyo Press, Tokyo, 1973.

- S. SMALE, *The Mathematics of Time. Essays on Dynamical Systems, Economic Process, and Related Topics*, Springer-Verlag, 1980.
- S. L. ZIGLIN, Nonintegrability of a Problem on the Motion of Four Point Vortices, *Soviet Math. Dokl.*, Vol. **21**, No. 1, 1980, pp. 296-299.

(*Manuscript received March 15, 1911.*)