

# ANNALES DE L'I. H. P., SECTION C

B. PERTHAME

## **Some remarks on quasi-variational inequalities and the associated impulsive control problem**

*Annales de l'I. H. P., section C*, tome 2, n° 3 (1985), p. 237-260

[http://www.numdam.org/item?id=AIHPC\\_1985\\_\\_2\\_3\\_237\\_0](http://www.numdam.org/item?id=AIHPC_1985__2_3_237_0)

© Gauthier-Villars, 1985, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Some remarks on quasi-variational inequalities and the associated impulsive control problem

by

**B. PERTHAME**

E. N. S., 45, rue d'Ulm, 75230 Paris, Cedex 05

**ABSTRACT.** — We study Quasi-Variational Inequalities:

$$(1) \quad \begin{cases} \text{Max } (Au - f, u - Mu) = 0, \\ u|_{\partial\Omega} = \varphi, \end{cases}$$

where

$$(2) \quad Mu = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \bar{\Omega}}} \{ c_0(\xi) + u(x + \xi) \}.$$

In general, (1) has no solution, we prove here that (1) has a unique maximum subsolution that we characterize. Then we compare the implicit obstacle (2) and the obstacle:

$$M_+u = k + \infess_{\substack{\xi \geq 0 \\ x + \xi \in \Omega}} \{ c_0(\xi) + u(x + \xi) \}$$

and we finally show that, under general assumptions, the solution of (1) is Holder continuous.

*Key-words:* Quasi-Variational Inequalities, Implicit obstacle, maximum subsolution, Holder continuity, impulsive control.

**RÉSUMÉ.** — Nous étudions les Inéquations Quasi-Variationnelles :

$$(1) \quad \begin{cases} \text{Max } (Au - f, u - Mu) = 0 \text{ dans } \Omega, \\ u|_{\partial\Omega} = \varphi, \end{cases}$$

où

$$(2) \quad Mu = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \bar{\Omega}}} \{ c_0(\xi) + u(x + \xi) \}.$$

En général (1) n'a pas de solution, nous montrons ici que (1) admet une unique sous-solution maximale que nous caractérisons. Nous comparons ensuite l'obstacle implicite (2) et l'obstacle :

$$M_+u = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \Omega}} \{ c_0(\xi) + u(x + \xi) \},$$

et nous finissons par montrer que, sous des hypothèses générales, la solution de (1) est hölderienne.

*Mots-clefs* : Inéquations Quasi-Varationnelles, obstacle implicite, sous-solution maximale, continuité Holdérienne, contrôle impulsionnel.

We study here the Quasi-Variational Inequalities (Q.V.I.) and the associated stochastic impulsive control problem:

$$(1) \quad \begin{cases} \text{Max} (Au - f, u - Mu) = 0 & \text{on } \Omega, \\ u|_{\partial\Omega} = \varphi, \end{cases}$$

where  $\Omega$  is a regular ( $C^3$ ) connected open set of  $\mathbb{R}^N$  and  $M$  is given by:

$$(2) \quad Mu = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \Omega}} \{ c_0(\xi) + u(x + \xi) \}$$

The problem (1) was introduced in A. Bensoussan and J. L. Lions [1] (more recent results may be found in [3]). A typical result is the following if we assume  $\varphi \equiv 0$ ,  $f \geq 0$  and  $c_0$  increasing, then (1) has a unique solution which is in  $W^{1,\infty}$ .

Here we will relax these assumptions and answer the following questions. If we take general  $\varphi$  and  $f$ , (1) has in general no solution because of a difficulty involving the boundary condition: there is no reason *a priori* that  $Mu$  should be above  $\varphi$  on  $\partial\Omega$ . And if in general there is no solution of (1), the question is to determine in which sense (1) might be solved and whether the corresponding solution is the optimal cost function for the associated impulse control problem which is meaningful without any condition.

If the solution of (1) is not continuous, then formula (2) has no clear meaning and one generally defines the implicit obstacle by:

$$M_+u = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \Omega}} \{ c_0(\xi) + u(x + \xi) \}$$

But it is not clear (and to our knowledge it has never been checked before !) that  $Mu = M_+u$  (at least for  $u \in C(\overline{\Omega})$ ). We also answer here that question.

The last question that we study is to find general local regularity results,

say in  $C^{0,\alpha}$ , which do not involve the boundary data  $\varphi$ . Such a regularity has already been proved by J. Frehse and U. Mosco [9]. Here we state an analogous result but our proof is completely different from the one in [9].

In a first part we show that (1) has a unique maximal subsolution which is the solution of:

$$(3) \quad \begin{cases} \text{Max} (Au - f, u - Mu) = 0, \\ u|_{\partial\Omega} = \varphi \wedge Mu. \end{cases}$$

Here we cannot look after solutions in  $H^1$  since this space is not adapted to the obstacle  $Mu$  and, in particular,  $\varphi \wedge Mu \notin H^{1/2}(\partial\Omega)$ . But under a general assumption introduced in B. Perthame [16],  $\varphi \wedge Mu$  and  $Mu$  are continuous so that we can deal with continuous solution of (3) called viscosity solution. This kind of solution (a particular case of the notion introduced in M. G. Crandall and P. L. Lions [7], P. L. Lions [13] [14]) allows us to solve (3) with the classical argument of B. Hanouzet and J. L. Joly [10]. Remark that in Appendix 2 we prove equivalence between different notions of solution of the obstacle problem. In particular it appears that the viscosity solution is also the classical solution in  $H_{loc}^1$ .

We prove in a second part that this analytical solution is the one we should expect in view of the associated impulsive control problem. This is a verification analogous to the one in [3] [17] which clearly shows that the above notion is the correct one.

We give in a third part some comparison results on  $Mu$  and  $M_+u$ . There are two main results: if  $\Omega$  is sufficiently smooth ( $C^N$ ) then  $Mu = M_+u$  a. e. and if  $Mu$  or  $M_+u$  is continuous then  $Mu = M_+u$  everywhere (remark  $Mu$  and  $M_+u$  are defined pointwise and not almost everywhere).

Finally we prove that when  $c_0$  is continuous and  $c_0(\xi) \leq C|\xi|^\alpha$ ,  $\alpha$  small enough, then the solution of (1) is in  $C_{loc}^{0,\alpha}$ . The method is to replace  $Mu$  by an other obstacle which is in  $C_{loc}^{0,\alpha}$  and which is built locally with the help of general properties of  $\bar{M}u$ . Remark that we can give an example with very regular data where  $u$  is only  $W_{loc}^{1,\infty}$  (cf. B. Perthame [18]), and that in general no regularity up to the boundary holds.

## I. Q. V. I. WITHOUT THE EXISTENCE OF A SUBSOLUTION

### 1. Assumptions and main results.

This section is devoted to the Q. V. I.:

$$(1) \quad \begin{cases} \text{Max} (Au - f, u - Mu) = 0, \\ u|_{\partial\Omega} = \varphi, \end{cases}$$

We prove (1) has a unique maximum subsolution and that it is the solution of:

$$(3) \quad \begin{cases} \text{Max } (Au - f, u - Mu) = 0, \\ u|_{\partial\Omega} = \varphi \wedge Mu. \end{cases}$$

Here we call subsolution of (1) any  $u \in C(\overline{\Omega})$ , satisfying:

$$\begin{cases} Au \leq f & \text{in } \mathcal{D}'(\Omega), \\ u \leq Mu & \text{in } \overline{\Omega}, \\ u \leq \varphi & \text{on } \partial\Omega, \end{cases}$$

where we have set:

$$Au = -a_{ij} \frac{\partial^2 u}{\partial X_i \partial X_j} + b_i \frac{\partial u}{\partial X_i} + cu.$$

We assume:

$$(5) \quad a_{ij}, b_i, c, f \in W^{2,\infty}(\mathbb{R}^N)$$

$$(6) \quad \exists v > 0, \quad a_{ij} \xi_i \xi_j \geq v |\xi|^2 \quad \forall \xi = (\xi_1, \dots, \xi_N) \in \mathbb{R}^N,$$

$$(7) \quad \exists \lambda > 0, \quad c \geq \lambda,$$

and

$$(2) \quad Mu = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \overline{\Omega}}} \{ c_0(\xi) + u(x + \xi) \},$$

where  $k$  is a positive constant,  $c_0: (\mathbb{R}^+)^N \rightarrow \mathbb{R}^+$  is a lower semi-continuous sub-additive function with  $c_0(0) = 0$  and  $\xi \geq 0$  means  $\xi = (\xi_1, \dots, \xi_N)$  with  $\xi_i \geq 0$ .

We will also assume that the boundary data satisfies:

$$(8) \quad \varphi \in C(\partial\Omega),$$

$$(9) \quad M_0\varphi = k + \inf_{\xi \geq 0, x + \xi \in \partial\Omega} \{ c_0(\xi) + \varphi(x + \xi) \} \in C(\overline{\Omega}),$$

These assumptions will allow us to deal with continuous solutions of (1) or (3) which are called viscosity solutions. Recall that P. L. Lions [13] [14] has introduced the notion of viscosity solution of general second order equations adapting to second order equations the notion introduced by M. G. Grandall and P. L. Lions [7] for first order equations. A very particular application of this notion yield that the following obstacle problem:

$$(10) \quad \begin{cases} \text{Max } (Au - f, u - \Psi) = 0, \\ u|_{\partial\Omega} = \varphi \wedge \Psi, \end{cases}$$

has a unique viscosity solution  $u \in C(\overline{\Omega})$  if the obstacle  $\Psi \in C(\overline{\Omega})$ . In this particular setting  $u$  is nothing else that the limit in  $C(\overline{\Omega})$  of all regularized obstacle problems with nice standard solutions. (The different notions of solution of (10) are collected in Appendix 2 where we prove the equivalence of these notions).

In the following we will call viscosity solution of (3) a continuous function  $u$  such that  $u|_{\partial\Omega} = \varphi \wedge Mu$ ,  $Mu \in C(\overline{\Omega})$  and  $u$  is a viscosity solution of (10) with  $\Psi = Mu$ .

*Remarks.* — 1) Below it will be proved that, under assumption (9) and with the definition of  $Mu$  in (2), we always have  $Mu \in C(\overline{\Omega})$  when  $u \in C(\overline{\Omega})$  and  $u|_{\partial\Omega} = \varphi \wedge Mu$ .

2) In the particular case treated here, the notion of viscosity solution is equivalent to  $u \in C(\overline{\Omega})$ ,  $Mu \in C(\overline{\Omega})$  and:

$$\begin{cases} Au \leq f & \text{in } \mathcal{D}'(\Omega), \quad u \leq Mu & \text{on } \Omega, \\ & \text{on the open set } \mathcal{O} = \{u < Mu\} \cap \Omega, \\ u \in W_{loc}^{2,\infty}(\mathcal{O}) & \text{and } Au = f \text{ a. e.,} \end{cases}$$

moreover in Appendix 2 it is proved that  $u \in H_{loc}^1$ .

3) Exactly as in B. Perthame [16] where the assumption has been introduced to get the continuity of the solution, one easily checks that (9) may be replaced by:

$$(9') \quad Mw \in C(\overline{\Omega}),$$

where  $w$  is any continuous super solution of (1) such that  $w|_{\partial\Omega} = \varphi$ .

Then we can state the:

**THEOREM 1.** — *Under assumptions (5)-(9), equation (1) has a unique maximal subsolution  $u$  and it is the unique solution of (3).*

*If  $\varphi \in W^{3,\infty}(\partial\Omega)$  and  $M_0\varphi \in W_{loc}^{1,\infty}(\Omega)$  (resp. is locally semi-concave) then  $u \in W_{loc}^{1,\infty}(\Omega)$  (resp.  $W_{loc}^{2,\infty}$ ).*

Let us recall that a function  $u \in C(\overline{\Omega})$  is said to be locally semi-concave if for each open set  $\mathcal{O} \subset \overline{\mathcal{O}} \subset \Omega$  there exist a constant  $C$  such that:

$$\frac{\partial^2 u}{\partial \chi^2} \leq C \quad \text{in } \mathcal{D}'(\mathcal{O}), \quad \forall \chi, \quad |\chi| = 1,$$

or, in other words:  $u(x) - \frac{1}{2}C|x|^2$  is concave on convex subsets of  $\mathcal{O}$ .

The end of this section is devoted to the proof of Theorem 1, it is divided in three parts: first we build a decreasing process that is uniformly converging to a solution of (3) (this is a variant of the « usual » proof due to B. Hanouzet and J. L. Joly [10]). Here the difficulty comes from the fact that the boundary value of the process changes at each step. The concavity of the operator which associates  $\varphi \wedge Mu$  to  $u$  enables us to conclude. In the second part we state the uniqueness result and finally we prove the regularity.

## 2. Existence of a solution.

Throughout section I we will assume that  $f \geq 0$ ,  $\varphi > 0$ . Indeed (7) enables us to make such an assumption by adding constants to  $u$  and  $\varphi$  and a positive function to  $f$ . We can now define the following process:  $u_0 \in C(\bar{\Omega})$  is the solution of:

$$(11) \quad \begin{cases} Au_0 = f \\ u_0|_{\partial\Omega} = \varphi, \end{cases}$$

then we define  $u_1$  as the solution of:

$$\begin{cases} \text{Max} (Au_1 - f, u_1 - Mu_0) = 0, \\ u_1|_{\partial\Omega} = \varphi \wedge Mu_0, \end{cases}$$

and by induction,

$$\begin{cases} \text{Max} (Au_n - f, u_n - Mu_{n-1}) = 0, \\ u_n|_{\partial\Omega} = \varphi \wedge Mu_{n-1}. \end{cases}$$

The existence of this sequence is justified by the:

LEMMA 1. — For each  $n \geq 0$   $Mu_n \in C(\bar{\Omega})$ .

*Proof.* — It is easily checked (cf. [16]) that under assumption (9)  $Mu_0$  is continuous on  $\bar{\Omega}$ . So, we assume by induction that for  $n \geq 0$ ,  $Mu_n$  is continuous and we prove that  $Mu_{n+1}$  is also continuous. We know (see [16]) that  $Mu_{n+1}$  is lower semi-continuous. Let us prove it is upper semi-continuous.

Let:  $Mu_{n+1}(x_0) = k + c_0(\xi_0) + u_{n+1}(x_0 + \xi_0)$ ,  $\xi_0 \geq 0$ ; if  $x_0 + \xi_0 \in \Omega$ , then, on a neighbourhood of  $x_0$  we have:

$$Mu_{n+1}(x) \leq k + c_0(\xi_0) + u_{n+1}(x + \xi_0)$$

and the result is proved. If  $x_0 + \xi_0 \in \partial\Omega$ , we will prove that:

$$(12) \quad u_{n+1}(x_0 + \xi_0) = \varphi(x_0 + \xi_0) < Mu_n(x_0 + \xi_0),$$

and this prove that  $Mu_{n+1}$  is upper semi-continuous at  $x_0$  since:

$$\begin{cases} Mu_n(x) \leq M_0\varphi(x) \quad \text{on } \bar{\Omega}, \\ Mu_n(x_0) = M_0\varphi(x_0). \end{cases}$$

But (12) is deduced from the:

LEMMA 2. — If  $v \in C(\bar{\Omega})$  and  $Mv(x_0) = k + c_0(\xi_0) + v(x_0 + \xi_0)$ ,  $\xi_0 \geq 0$ , then  $v(x_0 + \xi_0) = Mv(x_0 + \xi_0) - k$ .

Indeed take  $v = u_{n+1}$  in lemma 2, then:

$$u_{n+1}(x_0 + \xi_0) < Mu_{n+1}(x_0 + \xi_0) \leq Mu_n(x_0 + \xi_0),$$

since  $u_n$  is a decreasing sequence). But  $x_0 + \xi \in \Gamma$  and thus we have

$$u_{n+1}(x_0 + \xi_0) = \varphi(x_0 + \xi_0) < Mu_n(x_0 + \xi_0),$$

and (12) is proved, concluding the proof of Lemma 1.

*Proof of Lemma 2.* — Notice that, if  $\eta \geq 0$ ,  $x_0 + \xi_0 + \eta \in \bar{\Omega}$ , then:

$$\begin{aligned} Mv(x_0) &= k + c_0(\xi_0) + v(x_0 + \xi_0) \\ &\leq k + c_0(\xi_0 + \eta) + v(x_0 + \xi_0 + \eta) \\ &\leq k + c_0(\xi_0) + c_0(\eta) + v(x_0 + \xi_0 + \eta) \end{aligned}$$

and, taking the infimum over all  $\eta \geq 0$ , we find:

$$v(x_0 + \xi_0) \leq Mv(x_0 + \xi_0) - k,$$

since equality holds for  $\eta = 0$ , Lemma 2 is proved.

*Remarks.* — 1) This also proves Remark 1 in I.1: take  $u_{n+1} \equiv u_n \equiv u$  in the proof of Lemma 1.

2) The argument used above is very similar to the one introduced in L. Caffarelli and A. Friedman [5] [6].

As mentioned in the proof of Lemma 1, the sequence  $u_n$  is decreasing, but  $u_n \geq 0$  since  $f \geq 0$ ,  $\varphi \geq 0$ , so that  $u_n$  converges to some function  $u$ ; moreover we have:

**PROPOSITION 1.** — *The sequence  $u_n$  converges uniformly to  $u \in C(\bar{\Omega})$  which is a viscosity solution of (3) and  $0 \leq u \leq u_0$ .*

*Proof.* — The proof below is adapted from B. Hanouzet and J. L. Joly [10]. Notice that (3) is for example deduced from the « stability » of viscosity solutions of second order equations by uniform convergence. Let us prove the uniform convergence of  $u_n$ : choose  $\mu \in ]0, 1[$  such that  $u \leq \frac{k \wedge \varphi}{\varphi}$ ,  $\mu \leq \frac{k}{\|u_0\|_{L^\infty}}$  and assume that for some  $\theta \in [0, 1]$  and some  $n \geq 0$  we have:

$$u_n - u_{n+1} \leq \mathcal{O}u_n,$$

then, since  $M$  is a concave mapping:

$$(1 - \mathcal{O})Mu_n + \mathcal{O}.M0 = (1 - \mathcal{O})Mu_n + \mathcal{O}k \leq Mu_{n+1}$$

Let us call  $z$  the solution of the obstacle problem (10) with

$$\varphi = (1 - \mathcal{O})Mu_n + \mathcal{O}k$$

and  $v_0$  the solution of

$$(13) \quad \begin{cases} \text{Max}(Av_0 - f, v_0 - k) = 0, \\ v_0|_{\partial\Omega} = \varphi \wedge k, \end{cases}$$

then, applying the maximum principle (regularizing  $Mu_n$  if necessary) and noticing that:

$$(1 - \mathcal{O})\varphi \wedge Mu_n + \mathcal{O}\varphi \wedge k \leq \varphi \wedge \{(1 - \mathcal{O})Mu_n + \mathcal{O}.k\}$$

one easily checks that:

$$(1 - \mathcal{O})u_{n+1} + \mathcal{O}.v_0 \leq z \leq u_{n+2}.$$

But the choice of  $\mu$  shows that:  $\mu u_{n+1} \leq \mu u_0 \leq v_0$  and thus we obtain:  $u_{n+1} - u_{n+2} \leq \mathcal{O}(1 - \mu)u_{n+1}$ .

Since  $u_0 - u_1 \leq u_0$ , iterating the above inequalities we obtain:

$$u_n - u_{n+1} \leq (1 - \mu)^n \|u_0\|_{L^\infty},$$

and this proves Proposition 1.

*Remark.* — Of course this also proves that  $Mu_n$  converges uniformly to  $Mu$  hence  $u$  is a viscosity solution of (3).

### 3. Uniqueness.

We must prove two results: that the solution of (3) is unique and also that it is the maximum subsolution. This last point is clear, if a function  $w \in C(\bar{\Omega})$  satisfies:

$$\begin{aligned} Aw &\leq f && \text{in } \mathcal{D}'(\Omega), \\ w &\leq Mw && \text{in } \bar{\Omega}, \\ w &\leq \varphi && \text{on } \Gamma, \end{aligned}$$

(this actually means that  $w$  is a viscosity subsolution of (1) if  $Mw \in C(\Omega)$ ) then, recalling the result of P. L. Lions [13],  $w \leq u_0$  since for example  $w$  is a viscosity subsolution of (11). We deduce that:

$$\begin{aligned} w &\leq Mw \leq Mu_0, \\ w &\leq \varphi \wedge Mw \leq \varphi \wedge Mu_0 \quad \text{on } \Gamma, \end{aligned}$$

and so  $w \leq u_1$ . An easy induction proves that  $w \leq u_n$  for each  $n \geq 0$  and so  $w \leq u$ .

**PROPOSITION 2.** — *Under the assumptions of theorem 1, if  $w \in C(\bar{\Omega})$  is a viscosity solution of (3), then  $w = u$ .*

*Proof.* — We already know that  $w \leq u$ . First we show that  $w \geq 0$ : we know (by the remark in I 2°) that  $Mw \in C(\bar{\Omega})$  so we can build a sequence  $\Psi_n \in C^\infty(\bar{\Omega})$  such that  $\Psi_n \xrightarrow{n \rightarrow \infty} Mw$  in  $C(\bar{\Omega})$ . Let us denote by  $w_n$  the solu-

tion of (10) for  $\Psi = \Psi_n$  so that  $w_n \xrightarrow{n \rightarrow \infty} w$  in  $C(\bar{\Omega})$  and  $w_n \in W_{loc}^{2,\infty}(\Omega)$  (see [16]). Take  $x_n \in \bar{\Omega}$  so that:  $\text{Min}_{x \in \Omega} w_n(x) = w_n(x_n)$ ; as

$$w_n(x_n) \xrightarrow{n \rightarrow \infty} \text{Min}_{x \in \Omega} w(x) < \text{Min}_{x \in \Omega} Mw(x),$$

we may assume (at least if  $n$  is large enough) that  $w_n(x_n) < Mw_n(x_n)$ .

If  $x_n \in \Gamma$  we deduce that  $w_n(x_n) = \varphi(x_n) \geq 0$ , if not the maximum principle shows that  $w_n(x_n) \geq 0$  and in both cases  $w_n \geq 0$  and thus  $w \geq 0$ .

We now prove Proposition 2 with the help of B. Hanouzet and J. L. Joly method. We have:

$$u - w \leq u,$$

by the same arguments as in Proposition 1 we get

$$u - w \leq (1 - \mu)^n u, \quad \forall n \geq 0,$$

and so  $u \leq w$  and Proposition 2 is proved.

#### 4. Regularity.

Here we assume that  $M_0\varphi \in W_{loc}^{1,\infty}(\Omega) \cap C(\bar{\Omega})$  (Resp. that  $M_0\varphi$  is locally semi-concave) and we prove that  $u \in W_{loc}^{1,\infty}(\Omega)$  (resp.  $W_{loc}^{2,\infty}(\Omega)$ ).

We only sketch the proof since it is nearly the same as in [16]. One introduces:

$$F = \{ y \in \Omega \mid \exists x \leq y, Mu(x) = k + c_0(y - x) + u(y) \},$$

and one easily proves that there exists some open set  $G$  (for the topology of  $\bar{\Omega}$ ) such that:

$$F \subset G \subset \bar{G} \subset \{ u < Mu \}$$

and so  $u \in W^{1,\infty}(\bar{G})$  (resp.  $u \in W^{2,\infty}(\bar{G})$ ). Now, as  $Mu \ll$  takes its values  $\gg$  in  $F$  and with the regularity of  $M_0\varphi$  one can prove that  $Mu$  is in  $W_{loc}^{1,\infty}(\Omega)$  (resp. locally semi-concave) and applying classical results on the obstacle problem this proves that  $u \in W_{loc}^{1,\infty}(\Omega)$  (Resp.  $u \in W_{loc}^{2,\infty}(\Omega)$ ), and the proof of Theorem 1 is completed.

*Remarks.* — 1) In particular if  $\Omega$  is convex the solution belongs to  $W_{loc}^{2,\gamma}$  when  $\varphi \in W^{3,\infty}(\partial\Omega)$ . Indeed  $Mu_0 \in C(\Omega)$  and is locally semi-concave and this is enough by remark 3 in I. 1 (cf. [16], [18]).

2) Of course these results extend to Hamilton-Jacobi-Bellman equations (see P. L. Lions [12], L. C. Evans and P. L. Lions [8]). The associated Q. V. I. is:

$$\begin{cases} \text{Max}_{1 \leq i \leq m} (A^i u - f^i), u - Mu = 0, \\ u|_{\partial\Omega} = \varphi. \end{cases}$$

(For results on this problem see S. Lehnart [11], B. Perthame [16]).

## II. THE IMPULSIVE CONTROL PROBLEM

Our purpose, here, is to give the stochastic interpretation of  $u$ , the solution of (3), in terms of control of diffusion processes. Two remarks are to be developed below: first, despite of the jumps the process is constrained to stay in  $\bar{\Omega}$ ; next, the change of boundary data (from  $\varphi$  to  $\varphi \wedge Mu$ ) does not induce any change on the optimal cost function.

### 1. The optimal cost function.

Let  $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P}, w_t)$  be a standard space composed by a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  with a right-continuous increasing filtration of complete sub- $\sigma$  algebra  $\mathcal{F}_t$  and a Brownian motion  $w_t$  in  $\mathbb{R}^N$ ,  $\mathcal{F}_t$  adapted. For any sequences  $\mathcal{O}^1 < \mathcal{O}^2 < \dots < \mathcal{O}^n < \dots$ ,  $\mathcal{O}^n \xrightarrow[n \rightarrow \infty]{} \infty$ , of stopping times and  $\xi^1, \dots, \xi^n, \dots$  of  $\mathcal{F}_{\mathcal{O}^n}$ -random variables in  $(\mathbb{R}^+)^n$  we can define by induction:

$$(14) \quad \begin{cases} dy_x^n(t) = \sigma(y_x^n(t))dw_t - b(y_x^n(t))dt, & t \geq \mathcal{O}^n \\ y_x^n(\mathcal{O}^n) = y_x^{n-1}(\mathcal{O}^n) + \xi^n, \end{cases}$$

and:

$$y_x^0(0) = x$$

we set:

$$y_x(t) = y_t = y_x^n(t), \quad \mathcal{O}^n \leq t < \mathcal{O}^{n+1}.$$

We will say that such a system  $\mathcal{A}$  is admissible if  $y_x(\mathcal{O}^n) \in \bar{\Omega}$  whenever  $y_x^{n-1}(\mathcal{O}^n) \in \bar{\Omega}$ . Then we define

$$\tau = \inf \{ t \geq 0, y_x(t) \notin \bar{\Omega} \},$$

and the cost function of this system is:

$$(15) \quad j(x, \mathcal{A}) = \mathbb{E} \left\{ \int_0^\tau f(y_s) e^{-\lambda s} ds + \sum_{n=1}^{\infty} (k + c_0(\xi^n)) e^{-\lambda \mathcal{O}^n} + \varphi(y_x(\tau)) e^{-\lambda \tau} \right\}$$

and the optimal cost function is:

$$(16) \quad v(x) = \inf_{\mathcal{A} \text{ adm.}} j(x, \mathcal{A}).$$

*Remark.* — 1) With our definition of admissible systems we know that  $y_x(\tau) \in \partial\Omega$ , and so (15) is meaningful.

2) Here we have taken  $c(x) \equiv \lambda$  to simplify notations but the following results still hold with any  $c(x) > 0$ .

**THEOREM 2.** — *Under the assumptions of Theorem 1, the solution  $u$  of (3) is the optimal cost function given by (16)*

*Proof.* — *i)  $u \leq v$ :* This is clear in view of A. Bensoussan and J. L. Lions [4] or P. L. Lions [13]:  $u$  is solution of the Q. V. I. (1) with  $\tilde{\varphi} = \varphi \wedge \text{Mu}$  so that it is the optimal cost function for a cost function defined by (15) with  $\tilde{\varphi}$  in place of  $\varphi$ . These cost functions are less than  $j(x, \mathcal{A})$  since  $\tilde{\varphi} \leq \varphi$  so that  $u \leq v$ .

*ii)  $u \geq v$ :* Here we give an optimal admissible system as in [3] [17]. Let  $\hat{\xi}$  be measurable function in  $(\mathbb{R}^+)^N$  such that:

$$\text{Mu}(x) = k + c_0(\hat{\xi}(x)) + u(x + \hat{\xi}(x)) \quad \forall x \in \bar{\Omega}.$$

we can find a standard space  $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P}, w_t)$  where we solve:

$$\begin{cases} dy_x^0(t) = \sigma(y_x^0(t))dw_t + b(y_x^0(t))dt, \\ y_x^0(0) = x, \end{cases}$$

let  $\hat{\mathcal{O}}_1$  be the first time when  $y_x^0$  belongs to  $\mathcal{C} = \{u = \text{Mu}\}$  ( $\hat{\mathcal{O}}_1 = +\infty$  if  $\mathcal{C}$  is not reached),  $\hat{\xi}_1 = \hat{\xi}(y_x^0(\hat{\mathcal{O}}_1))$  ( $\hat{\xi}_1$  is anything measurable if  $\hat{\mathcal{O}}_1 = +\infty$ ) and by induction

$$\begin{cases} dy_x^n(t) = \sigma(y_x^n(t))dw(t) + b(y_x^n(t))dt, \\ y_x^n(\hat{\mathcal{O}}^n) = y_x^{n-1}(\hat{\mathcal{O}}^n) + \hat{\xi}^n, \end{cases}$$

and  $\hat{\mathcal{O}}_{n+1}$  is the first time  $y_x^n$  enters  $\mathcal{C}$  and  $\hat{\xi}^{n+1} = \hat{\xi}(y_x^n(\hat{\mathcal{O}}^{n+1}))$ .

From [2] we know the:

LEMMA 3. — For any  $n \geq 0$ , one has:

$$(17) \quad u(y_x(\hat{\mathcal{O}}^n \wedge \tau))e^{-\lambda \hat{\mathcal{O}}^n \wedge \tau} = \mathbb{E} \left\{ \int_{\hat{\mathcal{O}}^n \wedge \tau}^{\hat{\mathcal{O}}^{n+1} \wedge \tau} f(y_x(s))e^{-\lambda s} ds + u(y_x^-(\hat{\mathcal{O}}^{n+1} \wedge \tau))e^{-\lambda \hat{\mathcal{O}}^{n+1} \wedge \tau} \mid \mathcal{F}_{\hat{\mathcal{O}}^n} \right\} \quad \text{a. e.}$$

*Remarks.* — 1°)  $y_x(\hat{\mathcal{O}}^n)$  and  $y_x(\tau)$  are in the set  $\{u < \text{Mu}\}$ , indeed whenever  $y_x(s)$  reaches  $\{u = \text{Mu}\}$  it jumps to come back in  $\{u < \text{Mu}\}$ . In particular  $u(y_x(\tau))1_{(\tau < \infty)} = \varphi(y_x(\tau))1_{(\tau < \infty)}$ .

2°) One checks as in [3] that  $\hat{\mathcal{O}}^n \xrightarrow{n \rightarrow \infty} +\infty$ ; indeed for each  $n$ ,  $u(y_x^{n-1}(\hat{\mathcal{O}}^n)) - u(y_x^{n-1}(\hat{\mathcal{O}}^n) + \hat{\xi}_n) \geq k$ , and, since  $u$  is continuous,  $|\hat{\xi}^n| \geq L > 0$ . Since  $\hat{\xi}^n \geq 0$  we obtain  $\hat{\xi}^1 + \dots + \hat{\xi}^n \geq L\sqrt{n}$  and the formula:

$$y_x(\hat{\mathcal{O}}^n) = x + \int_0^{\hat{\mathcal{O}}^n} \sigma(y_x(s))dw_s - \int_0^{\hat{\mathcal{O}}^n} b(y_x(s))ds + \hat{\xi}^1 + \dots + \hat{\xi}^n$$

on  $\{\hat{\mathcal{O}}^n < \infty\}$  shows that  $\hat{\mathcal{O}}^n \xrightarrow{n \rightarrow \infty} +\infty$ .

In (17) we remark that:

$$u[y_x^-(\hat{\mathcal{O}}^{n+1} \wedge \tau)]1_{(\hat{\mathcal{O}}^n \leq \tau)} = u[y_x(\hat{\mathcal{O}}^{n+1})]1_{(\hat{\mathcal{O}}^{n+1} \leq \tau)} + [k + c_0(\hat{\xi}^{n+1})]1_{(\hat{\mathcal{O}}^{n+1} \leq \tau)} + u[y_x(\tau)]1_{(\hat{\mathcal{O}}^n \leq \tau < \hat{\mathcal{O}}^{n+1})},$$

but we have:

$$e^{-\lambda \hat{\theta}^{n+1}} 1_{\hat{\theta}^{n+1} \leq \tau} = e^{-\lambda \hat{\theta}^{n+1}} 1_{\hat{\theta}^{n+1} \leq \tau < \infty} = e^{-\lambda \hat{\theta}^{n+1}} 1_{\hat{\theta}^{n+1} < \infty} = e^{-\lambda \hat{\theta}^{n+1}}$$

since  $\hat{\theta}^n \leq \tau$  whenever  $\hat{\theta}^n < \infty$ . Thus adding (17) for each  $n \geq 0$  and using remark 2, we get

$$u(x) = E \left\{ \int_0^\infty f(y_x(s)) e^{-\lambda s} ds + \sum_{n=1}^\infty [k + c_0(\hat{\xi}^n)] e^{-\lambda \hat{\theta}^n} + \varphi(y_x(\tau)) e^{-\lambda \tau} \right\},$$

and this means that  $u \geq w$  and theorem 2 is proved.

### III. SOME REMARKS ON THE OPERATOR M

The operator M which defines the Q. V. I. is not always given by (2). Indeed when one deals with discontinuous solution the operator Mu has no clear meaning and the Q. V. I. is generally defined with an operator  $M_+$ :

$$M_+u = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \Omega}} \{ c_0(\xi) + u(x + \xi) \}.$$

From a stochastical viewpoint it is also natural to consider admissible systems such that the jumps  $\xi$  at a point  $x$  are constrained to satisfy  $x + \xi \in \Omega$  and no longer  $x + \xi \in \bar{\Omega}$ . It will rise an other operator, defined for  $x \in \Omega$ :

$$M_-u = k + \inf_{\substack{\xi \geq 0 \\ x + \xi \in \Omega}} \{ c_0(\xi) + u(x + \xi) \}.$$

$M_+$  is defined for any bounded from below measurable function and  $M_-$  for  $u \in C(\Omega)$ . Here we give the relation between M,  $M_-$ ,  $M_+$ .

#### 1. Regularity of M, $M_-$ , $M_+$ .

PROPOSITION 3. — *Let  $u \in C(\bar{\Omega})$  then Mu is lower semi-continuous on  $\bar{\Omega}$  (l.s.c.),  $M_-u$  is upper semi-continuous (u. s. c.) and  $Mu \leq M_-u$  on  $\Omega$ .*

*Proof.* — The regularity of Mu and the inequality are clear. The other results are based on the following remark: take  $u_n \in C(\bar{\Omega})$  a decreasing sequence such that  $u_n \xrightarrow{n \rightarrow \infty} u$  pointwise on  $\Omega$ , then  $M_-u_n$  decreases to  $M_-u$ . Now if we choose  $u_n$  such that  $u_n|_{\partial\Omega} = \|u_n\|_{L^\infty} > \|u\|_{L^\infty}$  and  $u_n < \|u_n\|_{L^\infty}$  on  $\Omega$ , then it is easily checked that  $M_-u_n$  is continuous and so, that Mu is u. s. c.

PROPOSITION 4. — *If u is u. s. c. then  $M_-u$  is u. s. c. moreover, if  $c_0$  is continuous and u is u. s. c. then  $M_-u = M_+u$  on  $\Omega$ .*

*Proof.* — The proof of Proposition 3 directly proves the first part of

Proposition 4. Moreover, if  $c_0$  is continuous and  $u_n$  is chosen as above it is clear that  $M_+u_n = M_-u_n$  so that:

$M_-u \leq M_+u \leq M_+u_n \xrightarrow{n \rightarrow \infty} M_+u$  on  $\Omega$ , and Proposition 4 is proved.

## 2. Relation between $M$ and $M_-$ (or $M_+$ ).

PROPOSITION 5. — If  $c_0$  is continuous,  $u \in C(\bar{\Omega})$  and  $\Omega$  is of class  $C^N$  then  $Mu = M_+u$  a. e.

*Proof of Proposition 5.* — Let us introduce some notations:  $F$  will be the set of points  $x \in \Omega$  where possibly  $Mu(x) \neq M_+u(x)$ , and will denote by  $T\Gamma(y)$  the tangent hyperplane to  $\partial\Omega$  at  $y$  and  $v(y)$  the unit outward normal to  $\partial\Omega$  at  $y$ . Then we have:

$F \subset G = \{x \in \Omega, \exists y > x, y \in \partial\Omega, (e_i, y-x) = 0 \text{ for some } i \text{ and } x \in T\Gamma(y)\}$ ,

where  $(e_i)_{1 \leq i \leq N}$  is the canonical basis of  $\mathbb{R}^N$ . This result has been proved in [15], but it is not precise enough. In fact we have  $F \subset \bigcup_{n < N} G_n$  where:

$G_1 = \{x \in \Omega, \exists y \in \partial\Omega, y > x, \exists i \leq N \text{ such that:}$

$$(e_i, y-x) = 0, v(y) \parallel e_i, (e_j, y-x) \neq 0 \quad \forall j \neq i\}$$

$G_k = \{x \in \Omega, \exists y \in \partial\Omega, y > x, \exists i_1 < i_2 < \dots < i_k \text{ such that:}$

$$(e_{i_j}, y-x) = 0, (e_i, y-x) \neq 0 \quad i \neq i_j, 1 \leq j \leq k; \quad v(y) \in \text{vect}(e_{i_1}, \dots, e_{i_k})\}.$$

Let us prove that  $F \subset \bigcup_{n < N} G_n$ : take some  $x_0 \in F$ , as  $x_0 \in G$  there exists

$y > x, y \in \partial\Omega$  and some  $e_{i_1}$  such that  $(e_{i_1}, y-x) = 0$ . We may always choose  $e_{i_2} \dots e_{i_k} \dots e_{i_n}$  such that:  $(e_{i_j}, y-x) = 0$  for  $1 \leq j \leq k$ ,  $(e_{i_j}, y-x) \neq 0$  for  $j > k$ . Now assume that  $x \notin G_k$ , it means that for some  $1 > k$ ,  $(e_{i_1}, v(y)) \neq 0$ . Since  $y > x$  then  $y \pm \varepsilon e_{i_1} > x$  for  $\varepsilon$  small enough, and either  $y - \varepsilon e_{i_1} \in \Omega$ , or  $y + \varepsilon e_{i_1} \in \Omega$  for  $\varepsilon$  small enough. In both cases we get that  $M_+u(x_0) = Mu(x_0)$ , indeed we know that  $M_+u(x_0) \geq Mu(x_0)$  and

$$\begin{aligned} M_+u(x_0) &\leq k + c_0(y + \varepsilon e_{i_1} - x_0) + u(y + \varepsilon e_{i_1}) \\ &\leq k + c_0(y - x_0) + u(y) \quad (\text{as } \varepsilon \rightarrow 0) \\ &\leq Mu(x_0). \end{aligned}$$

This yields a contradiction and thus  $x \in G_k$ . We now conclude with the:

LEMMA 4. — For  $1 \leq k \leq N$ ,  $\text{meas}(G_k) = 0$ .

STEP 1. — We prove that  $G_1$  is of zero measure and it is enough to prove that

$$H_1 = \{x \in \Omega, \exists y \in \partial\Omega, y > x,$$

$v(y)$  is parallel to  $e_N$  and  $(e_N, y - x) = 0$  } is of zero measure. Since the set of  $y \in \partial\Omega$  such that  $v(y)$  is parallel to  $e_N$  is compact, it can be covered by a finite number of open sets  $(\Gamma_i)_{i \in I}$

$$\Gamma_i = \{ y, y_N = f_i(y_1 \dots y_{N-1}), \quad (y_1 \dots y_{N-1}) \in \mathcal{O}_i \}$$

where  $f_i \in C^N$  and  $\mathcal{O}_i$  is an open set.

Denoting  $y' = (y_1 \dots y_{N-1})$  it remains to prove that for these functions  $f_i \in C^N(\theta_i)$  the sets:

$$H = \{ x \in \Omega, \exists y = (y', y_N), y > x, y_N = f_i(y') \\ Df_i(y') = 0 \quad \text{and} \quad (e_N, y - x) = 0 \}$$

is of zero measure. But:

$$H \subset \{ y + (\alpha', 0), |\alpha'| \leq \text{diam}(\Omega), \quad y \in \partial\Omega$$

and

$$y_N = f_i(y') \quad \text{and} \quad v(y) = \pm e_N \}.$$

Denoting by  $m_{\mathbb{R}}$  the Lebesgue measure on  $\mathbb{R}$  we have, applying Sard's Theorem:

$$m_{\mathbb{R}} \{ z = f_i(y'), y' \in \mathcal{O}_i, Df_i(y') = 0 \} = 0,$$

thus  $H$  is of zero measure and we conclude the first step:  $\text{meas}(\mathbf{G}_1) = 0$ .

STEP 2. — In the same way we prove that  $\mathbf{G}_k$  is of zero measure for  $k > 1$ : it is enough to prove that  $H_k$  is of zero measure where:

$$H_k = \{ x \in \Omega, \exists y \in \partial\Omega, y > x, v(y) \in \text{Vect}(e_N, \dots, e_{N-k+1})$$

and

$$(e_j, y - x) = 0 \quad \forall j, \quad N - k + 1 \leq j \leq N \}.$$

By the implicit function Theorem and changing  $e_N$  into some  $e_j, j \geq N - k + 1$ , if necessary we have locally:

$$\Gamma_i = \{ y, y_N = f_i(y'), y' \in \mathcal{O}_i \},$$

and it is enough to study the set:

$$H = \{ x \in \Omega, \exists y > x, y_N = f_i(y'), \frac{\partial f_i}{\partial y_j} = 0 \\ \forall j \leq N - k; (e_j, y - x) = 0 \quad \forall j > N - k \}.$$

But its measure is less than:

$$(2 \text{ diam } \Omega)^{N-k} m_{\mathbb{R}^k} \left\{ (f_i(y'), y_{N-1}, \dots, y_{N-k+1}), y' \in \mathcal{O}_i \right. \\ \left. \text{and} \quad \frac{\partial f_i}{\partial y_j} = 0 \quad \forall j \leq N - k \right\},$$

this last term is the measure of the critical values of the function

$$F(y_1 \dots y_{N-1}) = (f_i(y_1 \dots y_{N-1}), y_{N-1}, \dots, y_{N-k+1})$$

and so is zero by Sard's theorem. This concludes the proof of Lemma 4 and of Proposition 5.

It is quite easier to get the following comparison Proposition in which we consider  $M_-u$  and  $Mu$  are defined for  $x \in \Omega$ :

PROPOSITION 6. — *If  $u \in C(\bar{\Omega})$ , and  $c_0$  is a non-negative, sub-additive function such that  $c_0(0) \equiv 0$ , then the set of discontinuity points in  $\Omega$  of  $M_-u$  is also the set of discontinuity points in  $\Omega$  of  $Mu$  and it is also the set of points where  $M_-u(x) \neq Mu(x)$ .*

COROLLARY. — *If  $u \in C(\bar{\Omega})$ ,  $M_-u = Mu$  a. e. in Baire Sense.*

This is because we know that the set of discontinuity of a l. s. c. (or u. s. c.) function is rare in Baire sense.

Proposition 6 is clearly deduced from the following:

LEMMA 5. — *Under the assumptions of Proposition 6:*

$$\liminf_{x \rightarrow x_0} M_-u(x) = Mu(x_0), \quad \limsup_{x \rightarrow x_0} Mu(x) = M_-u(x_0).$$

Let us prove the second equality which is the most difficult one: we know that  $\limsup_{x \rightarrow x_0} Mu(x) \leq M_-u(x_0)$ , and we shall prove that:

$$\liminf_{\substack{x \rightarrow x_0 \\ x > x_0}} Mu(x) \geq M_-u(x_0).$$

Take  $x_n > x_0$  and  $x_n \xrightarrow{n \rightarrow \infty} x_0$ ,  $Mu(x_n) = k + c_0(\xi_n) + u(x_n + \xi_n)$ ,  $\xi_n \geq 0$ . Extracting a subsequence if necessary we assume:  $\xi_n \xrightarrow{n \rightarrow \infty} \xi$ . Since  $x_n + \xi_n - x_0 \geq x_n - x_0 > 0$ , we can find an  $\varepsilon_n \in \mathbb{R}^N$ ,  $\varepsilon_n \xrightarrow{n \rightarrow \infty} 0$ ,  $x_n - x_0 + \xi_n + \varepsilon_n \geq 0$  and  $x_0 + \xi_n + \varepsilon_n \in \Omega$ . We have then:

$$\begin{aligned} M_-u(x_0) &\leq c_0(x_n - x_0 + \xi_n + \varepsilon_n) + u(x_n + \xi_n + \varepsilon_n) \\ &\leq \liminf_{n \rightarrow \infty} \{ c_0(x_n - x_0 + \varepsilon_n) + c_0(\xi_n) + u(x_n + \xi_n) \} \\ &\leq \liminf_{n \rightarrow \infty} Mu(x_n) \end{aligned}$$

Remark. — 1) Here we have used the three properties of  $c_0$ :  $c_0$  is subadditive,  $c_0(\xi) \xrightarrow{\xi \rightarrow 0} 0$  and  $c_0$  is l. s. c. To prove the first equality of Proposition 6 we only need  $c_0$  to be l. s. c.

2) A useful consequence of Proposition 6 is that the solution of (3) is also the solution of:

$$\begin{cases} \text{Max } (Au - f, u - M_+u) = 0 \\ u|_{\partial\Omega} = \varphi \wedge M_+u, \end{cases}$$

in other words  $u$  is also the solution of the impulsive control problem where the admissible systems are defined in such away that the process  $y_x(t)$  stays in  $\Omega$  (and no longer in  $\bar{\Omega}$ ) and the stopping time  $\tau$  is the first

exit time from  $\Omega$  (and no longer from  $\bar{\Omega}$ ). This remark also applies to the results proved in [16] for Q. V. I. associated to Hamilton-Jacobi-Bellman equations.

**IV. HOLDER CONTINUITY OF THE SOLUTION OF Q. V. I.**

In this section we prove a general Holder continuity result for the Q. V. I.:

$$(18) \quad \begin{cases} a(u, v - u) \geq (f, v - u), \\ \forall v \in H_0^1, v \leq M_+u; u \in H_0^1, u \leq M_+u, \end{cases}$$

where  $f \geq 0$  and  $a(., .)$  is the bilinear form associated with A:

$$(Au, v) = a(u, v)$$

Indeed if we do not assume (9) the general theory developed in A. Bensoussan and J. L. Lions [3] shows that (18) has a unique solution in  $H_0^1$ , but it is not continuous in general since our assumptions on  $c_0$  (see (2)) are too weak. Here we give a proof of the Holder continuity of  $u$  under assumption:

$$(19) \quad c_0 \text{ is continuous on } (\mathbb{R}^+)^N, \quad c_0(\xi) \leq C |\xi|^\alpha, \quad \forall \xi \geq 0,$$

where  $\alpha$  is an exponent to be precised in Theorems 3 and 4. Such an assumption has already been introduced in A. Bensoussan, J. Frehse and U. Mosco [4], or J. Frehse and U. Mosco [9] to prove the continuity of Q. V. I. with quadratic growth. The method used here is completely different of the one in [4] [9]. It consists in interpreting  $u$  as the solution of an obstacle problem where the obstacle  $\Psi$  has some special property. Using a regularity result for linear second order elliptic equations on open sets satisfying an exterior cone condition we prove that the solution is Holder continuous. The precise result is the following:

**THEOREM 3.** — *Under assumptions (5)-(7) there exist some  $\alpha_0, 0 < \alpha_0 < 1$ , such that if (19) holds with  $0 < \alpha < \alpha_0$  then the solution of (18) belongs to  $C_{loc}^{0,\alpha}(\Omega)$ .*

The proof of this Theorem is divided in two parts: first we prove that the solution  $u$  of (18) is u. s. c., then we prove the Hölder continuity of  $u$ .

*Remark.* — Of course this result extends to the more general case where  $f$  is eventually negative but (18) has a subsolution. It also extends to any boundary data.

**1. Selection of an u. s. c. solution.**

Here we make the assumptions of Theorem 3. We know that  $u$  is obtained as the limit of the decreasing process as in I:  $u_0$  is the solution of  $Au_0 = f, u_0 \in C(\bar{\Omega})$ . Then  $u_n$  is the solution of:

$$\begin{cases} a(u_n, v - u_n) \geq (f, v - u_n), \\ u_n \in H_0^1, u_n \leq M_+u_{n-1}; \quad \forall v \in H_0^1 \quad v \leq M_+u_{n-1}. \end{cases}$$

At each step we can select a u. s. c. solution  $u_n$ . Indeed if  $u_{n-1}$  is u. s. c. then  $M_-u_{n-1} = M_+u_{n-1}$  by Proposition 5 since  $c_0$  is continuous (in particular it doesn't depend on the choice of the u. s. c. selection) and we know  $M_-u_{n-1}$  is u. s. c. Now take a decreasing sequence of obstacles  $\Psi_p \xrightarrow{p \rightarrow \infty} Mu_{n-1}$  the solution  $u^p$  of the obstacle problem for  $\Psi_p$  is continuous and decreases with  $p$  to  $u_n$  (moreover  $u^p \xrightarrow{p \rightarrow \infty} u_n$  in  $H_0^1$  by the general theory in [3]) so that we can select an u. s. c. solution  $u_n$  at each step.

Finally since  $u_n$  is decreasing to  $u$ ,  $u$  has a u. s. c. representant.

**2. Holder continuity of  $u$  in theorem 3.**

Take a sequence  $u^n \searrow u$  in  $\Omega$ ,  $u^n \in C(\bar{\Omega})$  and assume  $u^n|_{\Gamma} = \|u^n\|_{L^\infty}$  is large enough so that  $M_-u^n$  is continuous and  $M_-u^n \rightsquigarrow M_-u$ . We prove in what follows that each solution  $v_n$  of the obstacle problem for  $\Psi = M_-u^n$  belongs to  $C_{loc}^{0,\alpha}$  with uniform bounds on compact subsets of  $\Omega$ .

For any  $x_0 \in \Omega$  we denote by  $d(x_0)$  the function  $d(x_0) = \inf_{y \in \partial\Omega} d(x_0, y)$ . Since  $c_0$  is subadditive and by (19) we have for all  $y \in \Omega$ ,  $y \leq x_0$ :

$$\begin{aligned} M_-u^n(y) &\leq M_-u^n(x_0) + c_0(x_0 - y) \\ &\leq M_-u^n(x_0) + C|x_0 - y|^\alpha \\ &\leq M_-u^n(x_0) + \text{Max} \left( C, \frac{\|u_0\|_{L^\infty}}{d^\alpha(x_0)} \right) |x_0 - y|^\alpha. \end{aligned}$$

In the following we will denote by:

$$C(x_0) = \text{Max} \left( C, \frac{\|u_0\|_{L^\infty}}{d^\alpha(x_0)} \right)$$

and we define two open sets:

$$\begin{aligned} Q(x_0) &= \{ y \in \Omega, y < x_0, |y - x_0| < d(x_0) \} \\ K(x_0) &= \{ y \in \Omega, |y - x_0| < d(x_0), y \notin \bar{Q}(x_0) \}. \end{aligned}$$

By the appendix 1 if  $\alpha$  is small enough  $\left( \alpha \leq \alpha \left( \frac{\sqrt{2}}{2} \right) \right)$  in Theorem 4) we can solve:

$$\begin{cases} Av^1 = f, v^1 \in C^{0,\alpha}(\bar{Q}(x_0)) \cap C^\infty(Q(x_0)), \\ v^1|_{\partial Q(x_0)} = Mu^n(x_0) + C(x_0)|x_0 - y|^\alpha, \\ Av^2 = f, v^2 \in C^{0,\alpha}(K(x_0)) \cap C^\infty(K(x_0)), \\ v^2|_{\partial K(x_0)} = Mu^n(x_0) + c(x_0)|x_0 - y|^\alpha, \end{cases}$$

then we set:

$$v_{x_0} = \begin{cases} v^1 & \text{on } \overline{Q(x_0)} \\ v^2 & \text{on } \overline{K(x_0)}, \\ Mu^n(x_0) + C(x_0)d^\alpha(x_0) & \text{on } \Omega \setminus (Q(x_0) \cup K(x_0)). \end{cases}$$

The Appendix 1 shows that  $v_{x_0}(y)$  is in  $C^{0,\alpha}$  if  $\alpha$  is small enough and  $\|v_{x_0}\|_{C^{0,\alpha}} \leq D(x_0)$ , where  $D(x_0)$  only depends on  $d(x_0)$ , and so the obstacle  $\Psi$  defined below is in  $C_{loc}^{0,\alpha} \cap C(\Omega)$ :

$$\Psi = \left( \inf_{x_0 \in \Omega} v_{x_0} \right) \wedge u_0,$$

and we conclude by the:

**LEMMA 6.** — *The solution  $v_n$  of the obstacle problem for the obstacle  $M - u^n$  is the solution of:*

$$\begin{cases} \text{Max} (Aw - f, w - \Psi) = 0, & w \in C(\overline{\Omega}) \cap C^{0,\alpha}(\Omega), \\ w|_{\partial\Omega} = 0. \end{cases}$$

*Proof.* — For any  $x \in \Omega$  we have:

$$w(x) \leq \Psi(x) \leq v_x(x) = Mu^n(x),$$

hence  $w$  is a subsolution of (18) and  $w \leq v_n$ . On the other hand for any  $x_0 \in \Omega$  we have:

$$\begin{cases} Aw_n \leq f & \text{on } Q(x_0) \\ v_n|_{\partial Q(x_0)} \leq Mu^n|_{\partial Q(x_0)} \leq Mu^n(x_0) + C(x_0)|x_0 - y|^\alpha, \\ \\ \begin{cases} Av_n \leq f & \text{on } K(x_0), \\ v_n|_{\partial K(x_0)} \leq Mu^n(y) \leq Mu^n(x_0) + C(x_0)|x_0 - y|^\alpha, & \text{if } y \leq x_0, \\ v_n|_{\partial K(x_0)} \leq \|u_0\|_{L^\infty} \leq C(x_0)|x_0 - y|^\alpha & \text{if } y > x_0. \end{cases} \end{cases}$$

This implies  $v_n \leq v_{x_0}$ ,  $\forall x_0$  i.e.  $v_n \leq \Psi$  and then  $v_n \leq w$ ; and Lemma 6 is proved.

Now we deduce Theorem 3 from Lemma 6: we notice that the Hölder estimates on  $v_n$  do not depend on  $u^n$  and since  $v_n$  converges to  $u$ , Theorem 3 is proved.

## APPENDIX I

### CONSTRUCTION OF A BARRIER FUNCTION FOR AN OPEN SET SATISFYING THE EXTERIOR CONE CONDITION

In section IV we need to solve linear elliptic second order equations for open sets satisfying only the exterior cone condition. The purpose of this appendix is to show how to construct a barrier function in such cases (which do not seem to be classical). The results below are due to K. Miller [15] where one can find another way to prove them.

In this section we are given an operator  $A$  and a function  $f$  satisfying assumptions (and notations) of section I. We denote  $B = \|bi\|_{L^\infty(\mathbb{R}^N)}$  and by  $C_\mu$  the cone:

$$C_\mu = \left\{ x, \frac{x_N}{r} < -\mu \right\}, \quad 0 \leq \mu < 1,$$

where  $r = |x|$ , and we construct a barrier function at the point 0 for the open set:

$$\mathcal{O} = \{ x, |x| < R_0 \} \setminus \overline{C}_\mu.$$

The existence of a barrier function and its regularity are given in the:

**THEOREM 4.** — *Let  $\mu \in [0, 1[$  then there exist  $\alpha \in ]0, 1[$  such that for  $h \in C^{0,\alpha}(\mathbb{R}^N)$  there is a function  $v \in C^\alpha(\mathcal{O}) \cap C^{0,\alpha}(\overline{\mathcal{O}})$  such that:*

$$(20) \quad \begin{cases} Av \geq f, \\ v|_{\partial\mathcal{O}} \geq h|_{\partial\mathcal{O}}, \quad v(0) = h(0), \end{cases}$$

moreover  $\alpha$  and  $\|v\|_{C^{0,\alpha}}$  only depends on  $R_0, \mu, A, f$  and  $\|h\|_{C^{0,\alpha}}$ .

*Proof.* — We may assume  $h(0) = 0$  and we look for  $v$  satisfying:

$$v(x) = r^\alpha g\left(\frac{x_N}{r}\right), \quad 0 < \alpha < 1.$$

We will divide the proof in three parts: computation of  $Av$ , lower bound for  $Av$ , existence of  $g$  and  $\alpha$ .

#### 1. Computation of $Av$ .

We assume  $g \in C^\infty([-\mu, +1], \mathbb{R}^+)$  and we compute:

$$\begin{aligned} \frac{\partial v}{\partial x_i} &= \alpha r^{\alpha-2} g\left(\frac{x_N}{r}\right) x_i + r^{\alpha-3} g'\left(\frac{x_N}{r}\right) \{ \delta_{Ni} r^2 - x_N x_i \} \\ \frac{\partial^2 v}{\partial x_i \partial x_j} &= \alpha(\alpha-2) r^{\alpha-4} x_i x_j g\left(\frac{x_N}{r}\right) + \alpha r^{\alpha-2} \partial_{ij} g\left(\frac{x_N}{r}\right) \\ &+ g'\left(\frac{x_N}{r}\right) \{ \alpha r^{\alpha-5} x_i (\partial_{Ni} r^2 - x_N x_j) \\ &\quad + (\alpha-3) r^{\alpha-5} x_j (\delta_{Ni} r^2 - x_N x_i) + r^{\alpha-3} (2\delta_{Ni} x_j - x_N \delta_{ij} - x_i \delta_{jN}) \} \\ &\quad + r^{\alpha-6} g''\left(\frac{x_N}{r}\right) \{ \delta_{Ni} r^2 - x_N x_i \} \{ \delta_{Nj} r^2 - x_N x_j \}. \end{aligned}$$

Introducing  $y = \left( -\frac{x_N x_1}{r^2}, \dots, -\frac{x_N x_{N-1}}{r^2}, 1 - \frac{x_N}{r} \right)$  we get:

$$Av = g\left(\frac{x_N}{r}\right)n^\alpha c(x) + \alpha g\left(\frac{x_N}{r}\right)^{\alpha-2} \left\{ b_i x_i - (\alpha-2)a_{ij} \frac{x_i x_j}{r^2} - a_{ij} \delta_{ij} \right\} \\ + r^{\alpha-2} g'\left(\frac{x_N}{r}\right) \left\{ r b_i \cdot y_i - \alpha a_{ij} \frac{x_i}{r} y_j - (\alpha-3)a_{ij} \frac{x_j}{r} y_i - a_{ij} \left( 2\delta_{Ni} \frac{x_j}{r} - \frac{x_N}{r} \delta_{ij} \right. \right. \\ \left. \left. - \frac{x_i}{r} \delta_{jN} \right) \right\} - r^{\alpha-2} g''\left(\frac{x_N}{r}\right) a_{ij} y_i y_j$$

2. A lower bound for Av.

We assume now  $g \geq 0, g' \geq 0, g'' \leq 0$  and we put  $t = \frac{x_N}{r} \in [-1, 1]$ , then we get:

$$Av \geq -\alpha C_0 g(t) r^{\alpha-2} + r^{\alpha-2} g'(t) \left\{ -2 BR_0 + (2\alpha - 3) a_{ij} \frac{x_i x_j}{r^2} t \right. \\ \left. + (1 - \alpha) a_{Nj} \frac{x_j}{r} + (1 - \alpha) a_{iN} \frac{x_i}{r} + tr(a_{ij})t - r^{\alpha-2} g''(t) v(1 - t^2) \right\},$$

now we put  $x = (x', x_N)$  and  $d = (a_{ij})_{1 \leq i, j \leq N-1}$  so that the coefficient of  $r^{\alpha-2} g'(t)$  is equal to:

$$-2 BR_0 + (2\alpha - 3) \left( d \frac{x'}{r}, \frac{x'}{r} \right) t + \sum_{i=1}^{N-1} (a_{Ni} + a_{iN}) \left[ \frac{x_i}{r} (1 - \alpha) \right. \\ \left. + t^2 \frac{x_i}{r} (2\alpha - 3) \right] + tr(a_{ij})t + a_{NN} [2(1 - \alpha)t + (2\alpha - 3)t^3] \\ \geq -2 BR_0 - Ct^+(1 - t) - C\sqrt{1 - t} + tr(a_{ij})t + a_{NN}t [2 - 2\alpha + (2\alpha - 3)t^2] \\ \geq -2 BR_0 - C\sqrt{1 - t} + tr(a_{ij})t^+ - a_{NN}t^+ \\ \geq -2 BR_0 - C\sqrt{1 - t} + vt^+,$$

here C denotes different constants depending only on  $(a_{ij})$  and we assume  $N \geq 2$ .

Finally we get:

$$Av \geq -\alpha C_0 r^{\alpha-2} g(t) + r^{\alpha-2} g'(t) \left\{ -\frac{v}{2} + vt^+ - C\sqrt{1 - t} \right\} - r^{\alpha-2} g''(t)(1 - t^2).$$

It remains to find some g for which this expression is large enough.

3. Existence of g.

LEMMA 7. — For  $\mu \in [0, 1[$  there exist  $\alpha \in ]0, 1[$  and  $g_\mu \in C^\infty([- \mu, 1], \mathbb{R}^+)$ ,  $g_\mu \geq 1$ ,  $g'_\mu \geq 0, g''_\mu \geq 0$  such that:

$$(21) \quad -\alpha C_0 g_\mu(t) + g'_\mu(t) \left\{ -\frac{v}{2} + vt^+ - C\sqrt{1 - t} \right\} - g''_\mu(t)(1 - t^2) > 0.$$

Proof of lemma 7. — We look for  $g_\mu$  such that:

$$g_\mu(t) = 1 + e^{+K\mu} - e^{-Kt}$$

then (21) becomes:

$$-\alpha C_0 g_\mu(t) + g'_\mu(t) \left\{ -\frac{v}{2} + vt^+ - C\sqrt{1 - t} \right\} - g''_\mu(t)(1 - t^2) \\ \geq -\alpha C_0 g_\mu(t) + Ke^{-Kt} \left\{ -\frac{v}{2} + vt^+ - C\sqrt{1 - t} + K(1 - t)^2 \right\}$$

and a straightforward study of the function between the brackets shows that, if  $K$  is large enough (depending on  $\mu$  and  $(a_{ij})$ ) it is positive. Choosing  $\alpha$  small enough we get lemma 7.

We can now conclude the proof of Theorem 4; by lemma 7 and choosing  $g = Cg_\mu$  we have:  $\Delta v \geq Cr^{\alpha-2}C(\mu)$  where  $C(\mu)$  is the minimum in (21),  $C(\mu) > 0$ ; for  $C$  large enough (20) is satisfied and the dependance of  $\|v\|_{C^{0,\alpha}}$  on  $\mu, A, f$  and  $\|h\|_{C^{0,\alpha}}$  is clear.

*Remark.* — Of course for each  $\beta$ ,  $0 < \beta \leq \alpha$ , and  $h \in C^{0,\beta}(\mathbb{R}^N)$  we can find a barrier function in  $C^{0,\beta}(\bar{C})$ .

## APPENDIX 2

### DIFFERENT NOTIONS OF SOLUTION FOR THE OBSTACLE PROBLEM

Throughout this paper we have used the notion of viscosity solution for the obstacle problem. An other definition is given in a remark of Section I. On the other hand we could use the usual definition of variational inequalities. The purpose of this Appendix is to prove the equivalence between these three notions. Namely, we have, with the notations and assumptions of section I and IV:

**PROPOSITION 7.** — *Let  $u$  and  $\Psi$  belong to  $C(\bar{\Omega})$  then the following are equivalent:*

i)  $u$  is a viscosity solution of:

$$\begin{aligned} & \text{Max } (Au - f, u - \psi) = 0, \\ \text{ii)} \quad & \begin{cases} Au \leq f \text{ in } \mathcal{D}'(\Omega), \quad u \leq \Psi \text{ on } \bar{\Omega}, \\ \text{on the open set } \mathcal{O} = \{u < \Psi\}, \quad u \in W_{\text{loc}}^{2,\infty}(\mathcal{O}), \\ \text{and } Au = f, \end{cases} \end{aligned}$$

iii)  $u \in H_{\text{loc}}^1(\Omega)$  and  $u$  is a solution of the variational inequality associated to  $\Psi$  i.e.:

$$(22) \quad \begin{cases} a(u, v - u) \geq (f, v - u) \\ \forall v \in H_{\text{loc}}^1(\Omega), \quad v \leq \Psi, \quad u \leq \Psi \quad \text{and} \quad v = u \quad \text{on a neighbourhood on } \partial\Omega. \end{cases}$$

iv)  $u$  is the limit of solutions of all regularized obstacle problem with nice standard solution (actually this limit holds in  $C(\bar{\Omega})$  and weakly in  $H^1(\Omega')$  for all  $\Omega' \subset \bar{\Omega}' \subset \Omega$ ). In other words if we consider  $u_\varepsilon$  solution of:

$$(23) \quad \begin{cases} \text{Max } (Au_\varepsilon - f, u_\varepsilon - \Psi_\varepsilon) = 0, & u_\varepsilon \in C(\bar{\Omega}) \cap W_{\text{loc}}^{2,\infty}(\Omega), \\ u_\varepsilon|_{\partial\Omega} = \varphi_\varepsilon \end{cases}$$

with  $\Psi_\varepsilon \geq \varphi_\varepsilon$  on  $\partial\Omega$ ,  $\Psi_\varepsilon \rightarrow \Psi$  in  $C(\Omega)$  and  $\varphi_\varepsilon \rightarrow \varphi$  in  $C(\partial\Omega)$ , then  $u_\varepsilon \rightarrow u$  in  $C(\bar{\Omega})$  and weakly in  $H^1(\Omega')$ .

*Proof.* — The equivalence between (i) and (ii) is a simple consequence of the notion of viscosity solution as defined in P. L. Lions [13].

The equivalence between (i) and (iv) is easily deduced from the stability (under convergence in  $C(\bar{\Omega})$ ) of the notion of viscosity solution and the existence results for (23) which are proved in [2] and [16]. Remark that the family  $(u_\varepsilon)$  is bounded in  $H^1(\Omega')$  (and thus the weak convergence holds) since we have:

$$Au_\varepsilon \leq f,$$

and for  $\xi \in \mathcal{D}_+(\Omega)$  and  $C$  large enough:

$$\begin{aligned} & \int_{\Omega} Au_\varepsilon \xi^2 (u_\varepsilon + C) \leq \int_{\Omega} \xi^2 f (u_\varepsilon + C), \\ \int_{\Omega} |Du_\varepsilon|^2 \xi^2 - \int_{\Omega} C |Du_\varepsilon|^2 \xi^2 (u_\varepsilon + C) & - \int_{\Omega} C \xi |D\xi| \cdot |Du_\varepsilon| (u_\varepsilon + C) \\ & \leq \int_{\Omega} f \xi^2 (u_\varepsilon + C) - \int_{\Omega} c(x) u_\varepsilon (u_\varepsilon + C) \end{aligned}$$

and a simple minoration yields:

$$\frac{\nu}{2} \int_{\Omega} |Du_{\varepsilon}|^2 \xi^2 \leq C,$$

(here  $C$  denotes different constants independant of  $\varepsilon$ ). This concludes the equivalence between (i) and (iv).

Using the estimates proved above the equivalence (iii)  $\Leftrightarrow$  (iv) is clear since the solution of the variational inequality (22) is stable under uniform convergence of the obstacle (see [2]).

*Remark.* — The regularity stated in (ii) for  $u$  on the set  $\mathcal{O} = \{u < \Psi\}$  is not optimal. It depends of course on the regularity of  $f$ : if  $f \in L^2$ ,  $u \in H_{loc}^2$ ; if  $f \in L^{\infty}$   $u \in W_{loc}^{2,p} \forall p < \infty \dots$  etc...

## REFERENCES

- [1] A. BENSOUSSAN and J. L. LIONS, Nouvelle formulation de problèmes de contrôle impulsif et applications. *Comptes Rendus, Paris*, t. **276**, 1973, p. 1189-1192.
- [2] A. BENSOUSSAN and J. L. LIONS, *Application des Inéquations variationnelles en contrôle stochastique*. Dunod, 1978.
- [3] A. BENSOUSSAN and J. L. LIONS, *Contrôle impulsif et Inéquations Quasi-Variationnelles*. Dunod, 1982.
- [4] A. BENSOUSSAN, J. FREHSE, U. MOSCO, A stochastic impulse control problem with quadratic growth Hamiltonian and the corresponding quasi-variational Inequality. *J. Reine Angew. Math.*, t. **331**, 1982, p. 124-245.
- [5] L. A. CAFFARELLI and A. FRIEDMAN, Regularity of the solution of the quasi-variational inequality for the impulse control problem. *Comm. In P. D. E.*, t. **3**, (8), 1978, p. 745-753.
- [6] L. A. CAFFARELLI and A. FRIEDMAN, Regularity of the solution of the Quasi-variational inequality for the impulse control problem. *Comm. in P. D. E.*, t. **4** (3), 1979, p. 279-291.
- [7] M. G. CRANDALL and P. L. LIONS, *Viscosity solutions of Hamilton-Jacobi equations*.
- [8] L. C. EVANS and P. L. LIONS, Résolution des équations de Hamilton-Jacobi-Bellman pour les équations uniformément elliptiques. *C. R. A. S.*, t. **290**, 1980, p. 1049-1052.
- [9] J. FREHSE and U. MOSCO, Irregular obstacles and quasi-Variational Inequalities of Stochastic Impulse Control. *Annali di Pisa*, t. **4**, 91, 1982, p. 105-157.
- [10] B. HANOZET and J. L. JOLY, Convergence uniforme des itérés définissant la solution faible d'une Inéquation Quasi-variationnelle. *C. R. A. S.*, t. **286**, 1978, p. 735-738.
- [11] S. LEHNART, Bellman equations for Optimal stopping time problems. *Indiana U. Math. J.*, t. **32**, 3, 1983, p. 363-375.
- [12] P. L. LIONS, Résolution Analytique des Problèmes des Bellman Dirichlet. *Acta Mathematica*, t. **146**, 1981, p. 151-166.
- [13] P. L. LIONS, Optimal control and Hamilton-Jacobi-Bellman equations. Part 2, Viscosity solutions and uniqueness. *Comm. in P. D. E.*, t. **8**, 1983, p. 1101-1174.
- [14] P. L. LIONS, *Optimal control and Hamilton-Jacobi-Bellman equations. Part. 3*. Regularity of the optimal cost function. In Non-linear partial differential equations and applications, Collège de France, Seminar, Vol. V, Pitman, London, 1983.
- [15] K. MILLER, Barriers on cones for uniformly elliptic operators. *Annali di Matematica*, t. **2**, 76, 1967, p. 93-105.

- [16] B. PERTHAME, Quasi-Variational Inequalities and Hamilton-Jacobi-Bellman Equations in a bounded region. *Comm. in P. D. E.*, t. 9, 6, 1984, p. 561-595.
- [17] B. PERTHAME, Continuous and Impulsive control of diffusion in  $\mathbb{R}^N$ . *Non Linear Analysis T. M. A.*, t. 8, 10, 1984, p. 1227-1239.
- [18] B. PERTHAME, On the regularity of the solutions of Quasi-Variational Inequalities. To appear.

(Manuscrit reçu le 12 septembre 1984)

---

#### COPYRIGHT

Reproduction in whole or in part without the permission of the author or his representative is prohibited (law of March 11, 1957, Article 40, line 1). Such reproduction by whatever means, constitutes an infringement forbidden by Article 425 and those following it, of the Penal Code. The law of March 11, 1957, lines 2 and 3 of Article 41, authorizes only those copies or reproductions made for the exclusive use of the copyst, and not intended for collective use and such analyses and short quotations as are made for the purposes of an example or illustration.

Toute représentation ou reproduction intégrale ou partielle, faite sans le consentement de l'auteur ou de ses ayants droit ou ayants cause, est illicite (loi du 11 mars 1957, alinéa 1<sup>er</sup> de l'article 40). Cette représentation ou reproduction, par quelque procédé que ce soit, constituerait une contrefaçon sanctionnée par les articles 425 et suivants du Code pénal. La loi du 11 mars 1957 n'autorise, aux termes des alinéas 2 et 3 de l'article 41, que les copies ou reproductions strictement réservées à l'usage privé du copiste et non destinées à une utilisation collective d'une part et d'autre part que les analyses et les courtes citations dans un but d'exemple et d'illustration.

The appearance of the code at the bottom of the first page of an article in this journal indicates the copyright owner's consent that copies of the article may be made for personal or internal use, or for the personal or internal use of specific clients. This consent is given on the condition, however, that the copier pay the stated per-copy fee through the Copyright Clearance Center, Inc., Operations Center, 21 Congress St., Salem, Mass. 01970, U.S.A. for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purpose, for creating new collective works, or for resale.

© GAUTHIER-VILLARS 1985

*Annales de l'Institut Henri Poincaré - Analyse non linéaire*