

ANNALES DE L'I. H. P., SECTION C

ARRIGO CELLINA

GIOVANNI COLOMBO

ALESSANDRO FONDA

A continuous version of Liapunov's convexity theorem

Annales de l'I. H. P., section C, tome 5, n° 1 (1988), p. 23-36

http://www.numdam.org/item?id=AIHPC_1988__5_1_23_0

© Gauthier-Villars, 1988, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

A continuous version of Liapunov's convexity theorem

by

Arrigo CELLINA, Giovanni COLOMBO and Alessandro FONDA

International School for Advanced Studies (S.I.S.S.A.),
Trieste, Italy

ABSTRACT. — Given a continuous map $s \mapsto \mu_s$, from a compact metric space into the space of nonatomic measures on T , we show the existence of a family $(A_\alpha^s)_{\alpha \in [0, 1]}$, increasing in α and continuous in s , such that

$$\mu_s(A_\alpha^s) = \alpha \mu_s(T) \quad (\alpha \in [0, 1]).$$

Key words : Liapunov's convexity theorem - Measure theory - Selections.

RÉSUMÉ. — Étant donnée une application continue $s \mapsto \mu_s$, d'un espace métrique compact dans l'espace des mesures nonatomicques sur T , nous montrons l'existence d'une famille $(A_\alpha^s)_{\alpha \in [0, 1]}$, croissante avec α et continue en s , telle que

$$\mu_s(A_\alpha^s) = \alpha \mu_s(T) \quad (\alpha \in [0, 1]).$$

1. INTRODUCTION

Let μ be a non-atomic finite measure on a measurable space T . A result of measure theory states the existence of a family $(A_\alpha)_\alpha$ of subsets of T , increasing with α in $[0, 1]$ and such that

$$\mu(A_\alpha) = \alpha \mu(T).$$

Classification A.M.S. : 28 A 10, 54 C 65.

According to Liapunov's Convexity Theorem on the range of vector measures (see Halmos [2], [3] and Liapunov [4]) the above result holds for a finite family of nonatomic measures μ_i , $i = 1, \dots, n$: there exists an increasing family $(A_\alpha)_\alpha$ such that

$$\mu_i(A_\alpha) = \alpha \mu_i(T), \quad i = 1, \dots, n.$$

In general, the above is not true for an infinite family $(\mu_s)_s$ of measures (see Liapunov [5]). In this paper we consider a map $s \rightarrow \mu_s$, continuous for s in a compact metric space S . Denoting by $\mathcal{A}(\mu_s)$ the set of increasing families $(A_\alpha^s)_\alpha$ satisfying

$$\mu_s(A_\alpha^s) = \alpha \mu_s(T),$$

we show the existence of a selection $(\tilde{A}_\alpha^s)_\alpha$ of the multivalued map $\mathcal{A}(\mu_s)$ continuously depending on s in the sense of Definition 2 of the following section.

2. NOTATIONS AND PRELIMINARY RESULTS

We consider a measure space (T, \mathcal{F}, μ_0) where μ_0 is a non-atomic positive measure on a σ -algebra \mathcal{F} and $\mu_0(T) = 1$. Denote by \mathcal{M} the set of positive finite measures μ on T which are absolutely continuous with respect to μ_0 , hence non-atomic. The metric in \mathcal{M} is induced by the norm $\|\mu\|$ given by the variation of μ .

DEFINITION 1. — A family $(A_\alpha)_{\alpha \in [0, 1]}$, $A_\alpha \in \mathcal{F}$, is called *increasing* if

$$A_\alpha \subseteq A_\beta \quad \text{when } \alpha \leq \beta.$$

An increasing family is called *refining* $A \in \mathcal{F}$ with respect to the measure $\mu = (\mu_1, \dots, \mu_n) \in \mathcal{M}^n$ if $A_0 = \emptyset$, $A_1 = A$ and

$$\mu(A_\alpha) = \alpha \mu(A) \quad (\alpha \in [0, 1]).$$

The set of the families refining T with respect to μ is denoted by $\mathcal{A}(\mu)$.

The proofs of Lemmas 1 and 2 are based on Liapunov's theorem (see Fryszkowski [1]).

LEMMA 1. — Consider a vector measure $\mu \in \mathcal{M}^n$. For each $A \in \mathcal{F}$ there exists a family $(A_\alpha)_{\alpha \in [0, 1]}$ refining A with respect to μ . In particular, the set $\mathcal{A}(\mu)$ is nonempty.

In what follows, S is a compact metric space with distance d .

LEMMA 2. — Let $s \rightarrow \mu_s$ be a continuous map from S into \mathcal{M}^n . Then for every $\varepsilon > 0$ there exists an increasing family $(A_\alpha)_\alpha$ satisfying

- (i) $\mu_0(A_\alpha) = \alpha$ ($\alpha \in [0, 1]$);
- (ii) $|\mu_s(A_\alpha) - \alpha \mu_s(T)| < \varepsilon$ ($\alpha \in [0, 1], s \in S$).

DEFINITION 2. — A map $s \rightarrow (A_\alpha^s)_\alpha$ is called *continuous* on S if for every $s^0 \in S$ and $\varepsilon > 0$ there exists a $\delta > 0$ such that: s, s' and s'' in $B(s^0, \delta)$ implies

$$\sup_{\alpha \in [0, 1]} \mu_s(A_\alpha^{s'} \Delta A_\alpha^{s''}) < \varepsilon.$$

Analogously we set

DEFINITION 3. — The set valued map $s \rightarrow \mathcal{A}(\mu_s)$ is called *continuous* if for every $s^0 \in S$ and $\varepsilon > 0$ there exists a $\delta > 0$ such that: s, s' and s'' in $B(s^0, \delta)$ implies $\forall (A'_\alpha) \in \mathcal{A}(\mu_{s'}), \exists (A''_\alpha) \in \mathcal{A}(\mu_{s''})$ such that

$$\sup_{\alpha \in [0, 1]} \mu_s(A'_\alpha \Delta A''_\alpha) < \varepsilon.$$

We will use the symbol \cup to denote the union of disjoint sets. Finally, we recall that $\rho(\dots)$ defined as $\rho(A, B) = \mu(A \Delta B)$ ($\mu \in \mathcal{M}$) is a pseudometric on \mathcal{F} .

Remarks. — (a) In [5], Liapunov considers a sequence μ_n of measures on $[0, 2\pi]$ defined by a family of densities f_n converging strongly in L^1 to zero. He shows that there cannot exist any Borel subset A of $[0, 2\pi]$ such that for every $n, \mu_n(A) = \frac{1}{2} \mu_n([0, 2\pi])$. By associating μ_n to the point $1/n$ and $\mu_\infty = 0$ to the point 0, we have a map $s \mapsto \mu_s$ from the compact metric space $S = \{1/n : n \in \mathbb{N}\} \cup \{0\}$ into the space of nonatomic measures. The continuity at 0 follows from the strong convergence of (f_n) . This example shows that the assumptions of Theorem 1 below do not guarantee the existence of a constant selection.

(b) A further example is taken from Valadier [7]. Let S and T be the real interval $[0, 1]$, and set $\mu_s(A) = \int_A e^{-st} dt$. Assume there exists a set $\bar{A} \subseteq T$ such that

$$\forall s, \mu_s(A) = \frac{1}{2} \mu_s(T).$$

Then

$$\int_{-\infty}^{+\infty} \chi_A(t) e^{-st} dt = \int_{-\infty}^{+\infty} \frac{1}{2} \chi_T(t) e^{-st} dt.$$

Since the Laplace transformations of χ_A and $\frac{1}{2}\chi_T$, both of compact support, are analytic and coincide on $[0,1]$, they are identical. By the injectivity of the Laplace Transformation, we have

$$\chi_A = \frac{1}{2}\chi_T,$$

a contradiction. Hence again we have an example where there exist no constant selections.

(c) It seems more natural to express the continuity in terms of the pseudometric $\rho(A, B) = \mu_0(A, B)$. However, Definition 2 is not necessarily equivalent to the continuity with respect to this pseudometric when μ_0 is not absolutely continuous with respect to μ_s^0 .

3. MAIN RESULTS

In order to prove our main theorem we need three additional Lemmas.

LEMMA 3. — Consider a 1-dimensional measure $\mu \in \mathcal{M}$ and an increasing family $(A_\alpha^1)_\alpha$ such that for some $\varepsilon > 0$,

$$|\mu(A_\alpha^1) - \alpha\mu(T)| < \varepsilon \quad (\alpha \in [0, 1]).$$

There exists an increasing family $(A_\alpha^2)_\alpha$ such that

- (i) $\mu(A_\alpha^2) = \alpha\mu(T) \quad (\alpha \in [0, 1])$
- (ii) $\mu(A_\alpha^1 \triangle A_\alpha^2) < 6\varepsilon \quad (\alpha \in [0, 1]).$

Proof. — Fix M so that $\frac{1}{M} \geq \frac{\varepsilon}{\mu(T)} \geq \frac{1}{M+1}$. We begin by defining recursively an increasing family $(A_\alpha^2)_\alpha$ for $\alpha = i/M, i=0, \dots, M$, such that (i) holds and $A_{i/M}^2 \subseteq A_{(i+1)/M}^1$. Set $A_0^2 = \emptyset$ and assume $A_{i/M}^2$ has been defined for $i=0, \dots, n < M$.

Case 1. — When $\mu(A_{(n+1)/M}^1) \geq \frac{n+1}{M}\mu(T)$, define $A_{(n+1)/M}^2$ by Lemma 1, as a set such that $A_{n/M}^2 \subseteq A_{(n+1)/M}^2 \subseteq A_{(n+1)/M}^1$ and

$$\mu(A_{(n+1)/M}^2) = \frac{n+1}{M}\mu(T).$$

Case 2. — When $\mu(A_{(n+1)/M}^1) < \frac{n+1}{M}\mu(T)$, we first notice that by the

choice of M we have that $\mu(A_{(n+2)/M}^1) \geq \frac{n+1}{M} \mu(T)$; hence we can define

$A_{(n+1)/M}^2$ as a set such that $A_{(n+1)/M}^1 \subseteq A_{(n+1)/M}^2 \subseteq A_{(n+2)/M}^1$ and

$$\mu(A_{(n+1)/M}^2) = \frac{n+1}{M} \mu(T).$$

Notice that $A_{(n+1)/M}^2 \supseteq A_{n/M}^2$, since $A_{(n+1)/M}^1 \supseteq A_{n/M}^2$ by the inductive hypothesis.

In either case, we have

$$\begin{aligned} \mu(A_{(n+1)/M}^2 \triangle A_{(n+1)/M}^1) &= \left| \mu(A_{(n+1)/M}^2) - \mu(A_{(n+1)/M}^1) \right| \\ &\leq \left| \mu(A_{(n+1)/M}^2) - \frac{n+1}{M} \mu(T) \right| \\ &\quad + \left| \mu(A_{(n+1)/M}^1) - \frac{n+1}{M} \mu(T) \right| \\ &< \varepsilon. \end{aligned}$$

By Lemma 1 it is now easy to define a family $(A_\alpha^2)_{\alpha \in [0, 1]}$ such that

- (a) $A_{i/M}^2 \subseteq A_\alpha^2 \subseteq A_\beta^2 \subseteq A_{(i+1)/M}^2$ for $\frac{i}{M} \leq \alpha \leq \beta \leq \frac{i+1}{M}$;
 (b) $\mu(A_\alpha^2) = \alpha \mu(T)$.

Now we check that (ii) holds for $\frac{i}{M} \leq \alpha \leq \frac{i+1}{M}$. We can as well assume that $\mu(T) \geq 6\varepsilon$ otherwise (ii) trivially holds.

$$\begin{aligned} \mu(A_\alpha^1 \triangle A_\alpha^2) &= \mu(A_\alpha^1 \setminus A_\alpha^2) + \mu(A_\alpha^2 \setminus A_\alpha^1) \\ &\leq \mu(A_{(i+1)/M}^1 \setminus A_{i/M}^1) + \mu(A_{i/M}^1 \setminus A_{i/M}^2) \\ &\quad + \mu(A_{(i+1)/M}^2 \setminus A_{i/M}^2) + \mu(A_{i/M}^2 \setminus A_{i/M}^1) \\ &\leq \frac{1}{M} \mu(T) + 2\varepsilon + \frac{1}{M} \mu(T) + \varepsilon \\ &\leq 2 \frac{\varepsilon \mu(T)}{\mu(T) - \varepsilon} + 3\varepsilon = \frac{2\varepsilon}{1 - (\varepsilon/\mu(T))} + 3\varepsilon \\ &\leq \left(\frac{12}{5} + 3 \right) \varepsilon < 6\varepsilon. \quad \blacksquare \end{aligned}$$

COROLLARY. — *The set-valued map $s \rightarrow \mathcal{A}(\mu_s)$ is continuous.*

Proof. — Choose s^0 and $\varepsilon > 0$. Let $\delta > 0$ be such that $d(s, s^0) < \delta$ implies $\|\mu_s - \mu_{s^0}\| < \varepsilon/26$. Fix s, s' and s'' in $B(s^0, \delta)$ and $A'_\alpha \in \mathcal{A}(\mu_s)$. Since

$$\begin{aligned} |\mu_{s^0}(A'_\alpha) - \alpha\mu_{s^0}(T)| &= |\mu_{s^0}(A'_\alpha) - \mu_{s'}(A'_\alpha) + \mu_{s'}(A'_\alpha) \\ &\quad - \alpha\mu_{s'}(T) + \alpha\mu_{s'}(T) - \alpha\mu_{s^0}(T)| \\ &\leq 2\|\mu_{s'} - \mu_{s^0}\| < \varepsilon/13, \end{aligned}$$

by Lemma 3 there exists $A_\alpha^0 \in \mathcal{A}(\mu_{s^0})$ such that $\mu_{s^0}(A'_\alpha \triangle A_\alpha^0) \leq 6\varepsilon/13$. Analogously, given A_α^0 , there exists $A''_\alpha \in \mathcal{A}(\mu_{s''})$ such that $\mu_{s''}(A_\alpha^0 \triangle A''_\alpha) \leq 6\varepsilon/13$.

Hence

$$\begin{aligned} \mu_s(A'_\alpha \triangle A''_\alpha) &\leq |\mu_s(A'_\alpha \triangle A''_\alpha) - \mu_{s^0}(A'_\alpha \triangle A''_\alpha)| + \mu_{s^0}(A'_\alpha \triangle A''_\alpha) \\ &\leq \|\mu_s - \mu_{s^0}\| + \mu_{s^0}(A'_\alpha \triangle A_\alpha^0) + \mu_{s^0}(A_\alpha^0 \triangle A''_\alpha) \\ &\leq \varepsilon/26 + 6\varepsilon/13 + \|\mu_{s^0} - \mu_{s''}\| + \mu_{s''}(A_\alpha^0 \triangle A''_\alpha) \\ &\leq \varepsilon. \quad \blacksquare \end{aligned}$$

In the following Lemmas, the symbol $\sup_{\lambda_j(s) > 0}$ is a shorthand notation

for $\sup_{\{j \in \mathbb{N} : \lambda_j(s) > 0\}}$.

LEMMA 4. — Let $s \rightarrow \mu_s$ be a continuous map from a metric space S into the space \mathcal{M} and let $(B(s_j, \eta_j))_{j=1, \dots, N}$ be a finite open covering of S . Let $(\lambda_j(\cdot))_{j=1, \dots, N}$ be a continuous partition of unity subordinate to it such that $\lambda_j(s_j) = 1$.

For any center $s_j, j=1, \dots, N$, let be defined a finite increasing family $(\bar{A}_{i/M}^{s_j})_{i=0, \dots, M}$ such that

$$\mu_{s_j}(\bar{A}_{i/M}^{s_j}) = \frac{i}{M} \mu_{s_j}(T) \quad (i \in \{0, \dots, M\}).$$

Then for each $s \in S$ there exists an increasing family $(A_\alpha^s)_\alpha$ that extends the family $(\bar{A}_{i/M}^{s_j})_i$ in the sense that $A_{i/M}^{s_j} = \bar{A}_{i/M}^{s_j}$ for every i and j , and such that the following properties hold:

$$(i) \quad |\mu_s(A_\alpha^s) - \alpha\mu_s(T)| \leq 6 \sup_{\lambda_j(s) > 0} \|\mu_s - \mu_{s_j}\| \quad (\alpha \in [0, 1]);$$

$$(ii) \quad \text{for } \alpha \in \left[\frac{i}{M}, \frac{i+1}{M} \right] \text{ and any center } s_j,$$

$$\mu_{s_j}(A_\alpha^s \triangle A_\alpha^{s_j}) \leq \sup_{\lambda_k(s) > 0} \mu_{s_j}(\bar{A}_{(i+1)/M}^{s_k} \triangle \bar{A}_{(i+1)/M}^{s_j})$$

$$+ \sup_{\lambda_k(s) > 0} \|\mu_{s_j} - \mu_{s_k}\| + \frac{1}{M} (\sup_{\lambda_k(s) > 0} \mu_{s_k}(T) + \mu_{s_j}(T));$$

(iii) $\lim_{s \rightarrow s^*} \sup_{\alpha \in [0,1]} \mu_0(A_\alpha^s \triangle A_\alpha^{s^*}) = 0.$

Proof. — For each $s \in S$, first we will define the sets $(A_{i/M}^i)_i$ by interpolating among the given families $(\bar{A}_{i/M}^{s_j})_i$, taking from each set a subset having measure proportional to the corresponding $\lambda_i(s)$. Then we extend the construction for $\alpha \in]i/M, (i+1)/M[$. Finally we check that (i)-(iii) hold.

I. For any set $A \subseteq T$, we define $A^1 = A$ and $A^0 = T \setminus A$. We denote by \mathcal{X} the set of all $N \times (M-1)$ matrices $\Gamma = (\gamma_{ij})$ whose elements are in $\{0,1\}$.

Now we define

$$\begin{aligned} A(\Gamma) = & (\bar{A}_{1/M}^{s_1})^{\gamma_{11}} \cap \dots \cap (\bar{A}_{1/M}^{s_N})^{\gamma_{1N}} \\ & \cap (\bar{A}_{2/M}^{s_1})^{\gamma_{21}} \cap \dots \cap (\bar{A}_{2/M}^{s_N})^{\gamma_{2N}} \\ & \dots \dots \dots \\ & \cap (\bar{A}_{(M-1)/M}^{s_1})^{\gamma_{M-1,1}} \cap \dots \cap (\bar{A}_{(M-1)/M}^{s_N})^{\gamma_{M-1,N}}. \end{aligned}$$

Note that:

- (a) since the family $(\bar{A}_{i/M}^{s_j})_i$ is increasing in i , $A(\Gamma) = \emptyset$ if $\exists i, j: \gamma_{ij} = 1, \gamma_{i+1,j} = 0$; moreover, if $\Gamma_1 \neq \Gamma_2$, then $A(\Gamma_1) \cap A(\Gamma_2) = \emptyset$;
- (b) for any i, j

$$\bar{A}_{i/M}^{s_j} = \bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij} = 1}} A(\Gamma),$$

i. e. the family at the r. h. s. is a partition of $\bar{A}_{i/M}^{s_j}$;

(c) $\bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij} = 0, \gamma_{ik} = 1}} A(\Gamma) = A_{i/M}^{s_k} \setminus A_{i/M}^{s_j}, \quad \bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij} = 1, \gamma_{ik} = 1}} A(\Gamma) = A_{i/M}^{s_j} \cap A_{i/M}^{s_k}.$


By lemma 1, for each $\Gamma \in \mathcal{X}$ there exists a family $(A(\Gamma)_\alpha)_{\alpha \in [0,1]}$ refining $A(\Gamma)$ with respect to the measure $(\mu_0, \mu_{s_1}, \dots, \mu_{s_N})$. Define

$$\beta_\Gamma^i(s) = \sum_{k=1}^N \gamma_{ik} \lambda_k(s)$$

and

$$A_{i/M}^s = \bigcup_{\Gamma \in \mathcal{X}} A(\Gamma)_{\beta_\Gamma^i(s)} \tag{1}$$

(see Fig., where the case $N = M = 3$ is described).



The family $(A_{i/M}^s)_i$ coincides with $(\bar{A}_{i/M}^{s_j})_i$ for $s=s_j$; in fact we have $\beta_\Gamma^i(s_j) = \gamma_{ij}$ so that, by (b),

$$A_{i/M}^{s_j} = \bigcup_{\Gamma \in \mathcal{X}} A(\Gamma)_{\gamma_{ij}} = \bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij}=1}} A(\Gamma) = \bar{A}_{i/M}^{s_j}$$

Next we have:

$$\begin{aligned} \mu_{s_j}(A_{i/M}^s) &= \sum_{\Gamma \in \mathcal{X}} \mu_{s_j}(A(\Gamma)_{\beta_\Gamma^i(s)}) = \sum_{\Gamma \in \mathcal{X}} \beta_\Gamma^i(s) \mu_{s_j}(A(\Gamma)) \\ &= \sum_{\Gamma \in \mathcal{X}} \left(\sum_{k=1}^N \gamma_{ik} \lambda_k(s) \right) \mu_{s_j}(A(\Gamma)) \\ &= \sum_{k=1}^N \lambda_k(s) \sum_{\Gamma \in \mathcal{X}} \gamma_{ik} \mu_{s_j}(A(\Gamma)) \end{aligned} \tag{2}$$

$$\begin{aligned}
&= \sum_{k=1}^N \lambda_k(s) \mu_{s_j} \left(\bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ik}=1}} A(\Gamma) \right) \\
&= \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{i/M}^{s_k}).
\end{aligned}$$

II. Set, for $\alpha = (1-t)i/M + t(i+1)/M$ ($t \in [0, 1]$) and $s \in S$,

$$A_\alpha^s = \bigcup_{\Gamma \in \mathcal{X}} A(\Gamma)_{(1-t)\beta_\Gamma^i(s) + t\beta_\Gamma^{i+1}(s)}.$$

Remark that by the above definition and (1), it follows that

$$\mu_{s_j}(A_\alpha^s) = (1-t)\mu_{s_j}(A_{i/M}^s) + t\mu_{s_j}(A_{(i+1)/M}^s).$$

We claim that

$$\mu_{s_j}(A_\alpha^s) = \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_\alpha^{s_k}) \quad (j=1, \dots, N; \alpha \in [0, 1]; s \in S).$$

In fact, for α as above, we have:

$$\begin{aligned}
\mu_{s_j}(A_\alpha^s) &= \sum_{\Gamma \in \mathcal{X}} [(1-t)\beta_\Gamma^i(s) + t\beta_\Gamma^{i+1}(s)] \mu_{s_j}(A(\Gamma)) \\
&= (1-t) \sum_{k=1}^N \lambda_k(s) \sum_{\Gamma \in \mathcal{X}} \gamma_{ik} \mu_{s_j}(A(\Gamma)) \\
&\quad + t \sum_{k=1}^N \lambda_k(s) \sum_{\Gamma \in \mathcal{X}} \gamma_{i+1,k} \mu_{s_j}(A(\Gamma)) \\
&= (1-t) \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{i/M}^{s_k}) + t \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{(i+1)/M}^{s_k}) \\
&= \sum_{k=1}^N \lambda_k(s) [(1-t)\mu_{s_j}(A_{i/M}^{s_k}) + t\mu_{s_j}(A_{(i+1)/M}^{s_k})] \\
&= \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_\alpha^{s_k}).
\end{aligned}$$

III. We are now in the position of proving (i). Fix $s \in S$ and $\alpha \in [0, 1]$ and set $\omega_s = \sup \{ \|\mu_s - \mu_{s_j}\| : \lambda_j(s) > 0 \}$. We have:

$$\begin{aligned}
|\mu_s(A_\alpha^s) - \alpha\mu_s(T)| &\leq |\mu_s(A_\alpha^s) - \mu_{s_j}(A_\alpha^s)| \\
&\quad + |\mu_{s_j}(A_\alpha^s) - \alpha\mu_{s_j}(T)| + \alpha |\mu_{s_j}(T) - \mu_s(T)|
\end{aligned}$$

$$\begin{aligned}
&\leq 2\omega_s + \left| \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_\alpha^{s_k}) - \alpha \mu_{s_j}(T) \right| \\
&\leq 2\omega_s + \sum_{k=1}^N \lambda_k(s) [|\mu_{s_j}(A_\alpha^{s_k}) - \mu_{s_k}(A_\alpha^{s_k})| \\
&\quad + \alpha |\mu_{s_k}(T) - \mu_{s_j}(T)|] \\
&\leq 6\omega_s.
\end{aligned}$$

In order to prove (ii), note first that

$$\begin{aligned}
A_{i/M}^s \triangle A_{i/M}^{s_j} &= \left(\bigcup_{\Gamma \in \mathcal{X}} A(\Gamma)_{\beta_\Gamma^i(s)} \right) \triangle \left(\bigcup_{\Gamma \in \mathcal{X}} A(\Gamma)_{\beta_\Gamma^i(s_j)} \right) \\
&= \bigcup_{\Gamma \in \mathcal{X}} (A(\Gamma)_{\beta_\Gamma^i(s)} \triangle A(\Gamma)_{\beta_\Gamma^i(s_j)})
\end{aligned} \tag{3}$$

and that, by a calculation similar to (2) and by (c),

$$\mu_{s_j} \left(\bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij}=0}} A(\Gamma)_{\beta_\Gamma^i(s)} \right) = \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{i/M}^{s_k} \setminus A_{i/M}^{s_j}), \tag{4}$$

$$\mu_{s_j} \left(\bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij}=1}} (A(\Gamma) \setminus A(\Gamma)_{\beta_\Gamma^i(s)}) \right) = \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{i/M}^{s_j} \setminus A_{i/M}^{s_k}). \tag{5}$$

Therefore, for any i, j , from (3) and recalling that $\beta_\Gamma^i(s_j) = \gamma_{ij}$, we have

$$\mu_{s_j}(A_{i/M}^s \triangle A_{i/M}^{s_j}) = \mu_{s_j} \left(\bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij}=0}} A(\Gamma)_{\beta_\Gamma^i(s)} \right) + \mu_{s_j} \left(\bigcup_{\substack{\Gamma \in \mathcal{X} \\ \gamma_{ij}=1}} (A(\Gamma) \setminus A(\Gamma)_{\beta_\Gamma^i(s)}) \right)$$

and from (4), (5) this last expression is

$$\begin{aligned}
&\sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{i/M}^{s_k} \setminus A_{i/M}^{s_j}) + \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{i/M}^{s_j} \setminus A_{i/M}^{s_k}) \\
&= \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{i/M}^{s_k} \triangle A_{i/M}^{s_j}) \\
&\leq \sup \{ \mu_{s_j}(A_{i/M}^{s_k} \triangle A_{i/M}^{s_j}) : \lambda_k(s) > 0 \}.
\end{aligned}$$

Hence (ii) holds for $\alpha = i/M$.

In order to prove (ii) for α in $]i/M, (i+1)/M[$, let us note that

$$\begin{aligned}
A_\alpha^s \setminus A_\alpha^{s'} &\subseteq [(A_{(i+1)/M}^s \setminus A_{(i+1)/M}^{s_j}) \setminus A_\alpha^{s'}] \cup [A_{(i+1)/M}^{s_j} \setminus A_\alpha^{s'}] \\
&\subseteq (A_{(i+1)/M}^s \setminus A_{(i+1)/M}^{s_j}) \cup (A_{(i+1)/M}^{s_j} \setminus A_{i/M}^{s'}),
\end{aligned}$$

so that

$$\mu_{s_j}(A_\alpha^s \setminus A_\alpha^{s_j}) \leq \mu_{s_j}(A_{(i+1)/M}^s \setminus A_{(i+1)/M}^{s_j}) + \mu_{s_j}(A_{(i+1)/M}^{s_j} \setminus A_{i/M}^{s_j})$$

and

$$\mu_{s_j}(A_\alpha^{s_j} \setminus A_\alpha^s) \leq \mu_{s_j}(A_{(i+1)/M}^{s_j} \setminus A_{(i+1)/M}^s) + \mu_{s_j}(A_{(i+1)/M}^s \setminus A_{i/M}^s).$$

Hence

$$\begin{aligned} \mu_{s_j}(A_\alpha^s \triangle A_\alpha^{s_j}) &\leq \mu_{s_j}(A_{(i+1)/M}^s \triangle A_{(i+1)/M}^{s_j}) \\ &\quad + \mu_{s_j}(A_{(i+1)/M}^{s_j} \setminus A_{i/M}^{s_j}) + \mu_{s_j}(A_{(i+1)/M}^s \setminus A_{i/M}^s) \\ &\leq \sup \{ \mu_{s_j}(A_{(i+1)/M}^{s_k} \triangle A_{(i+1)/M}^{s_j}) : \lambda_k(s) > 0 \} \\ &\quad + (1/M) \mu_{s_j}(T) + \sum_{k=1}^N \lambda_k(s) \mu_{s_j}(A_{(i+1)/M}^{s_k} \setminus A_{i/M}^{s_k}) \\ &\leq \sup_{\lambda_k(s) > 0} \mu_{s_j}(A_{(i+1)/M}^{s_k} \triangle A_{(i+1)/M}^{s_j}) + (1/M) \mu_{s_j}(T) \\ &\quad + \sum_{k=1}^N \lambda_k(s) | \mu_{s_j}(A_{(i+1)/M}^{s_k} \setminus A_{i/M}^{s_k}) - \mu_{s_k}(A_{(i+1)/M}^{s_k} \setminus A_{i/M}^{s_k}) | \\ &\quad + \sum_{k=1}^N \lambda_k(s) \mu_{s_k}(A_{(i+1)/M}^{s_k} \setminus A_{i/M}^{s_k}) \\ &\leq \sup_{\lambda_k(s) > 0} \mu_{s_j}(\bar{A}_{(i+1)/M}^{s_k} \triangle A_{(i+1)/M}^{s_j}) \\ &\quad + \sup_{\lambda_k(s) > 0} \| \mu_{s_j} - \mu_{s_k} \| \\ &\quad + (1/M) \sup_{k=1, \dots, N} \mu_{s_k}(T). \end{aligned}$$

This proves (ii).

Finally we prove (iii); for $\alpha = (1-t) i/M + t (i+1)/M$ we have

$$\begin{aligned} \mu_0(A_\alpha^s \triangle A_\alpha^{s^*}) &= \sum_{\Gamma \in \mathcal{X}} \mu_0(A(\Gamma)_{(1-t)\beta_\Gamma^i(s) + t\beta_\Gamma^{i+1}(s)} \\ &\quad \triangle A(\Gamma)_{(1-t)\beta_\Gamma^i(s^*) + t\beta_\Gamma^{i+1}(s^*)}) \\ &= \sum_{\Gamma \in \mathcal{X}} \{ |[(1-t)\beta_\Gamma^i(s) + t\beta_\Gamma^{i+1}(s)] \\ &\quad - [(1-t)\beta_\Gamma^i(s^*) + t\beta_\Gamma^{i+1}(s^*)]| \mu_0(A(\Gamma)) \} \\ &\leq (1-t) \sum_{\Gamma \in \mathcal{X}} | \beta_\Gamma^i(s) - \beta_\Gamma^i(s^*) | \mu_0(A(\Gamma)) \\ &\quad + t \sum_{\Gamma \in \mathcal{X}} | \beta_\Gamma^{i+1}(s) - \beta_\Gamma^{i+1}(s^*) | \mu_0(A(\Gamma)). \end{aligned}$$

By taking the limit as s tends to s^* we conclude the proof. ■

LEMMA 5. — Let $s \rightarrow \mu_s$ be a continuous map from a compact metric space S into the space \mathcal{M} and, for each $s \in S$, let $(\bar{A}_\alpha^s)_\alpha$ be an increasing family, continuous with respect to s and such that, for some $\varepsilon > 0$,

$$|\mu_s(\bar{A}_\alpha^s) - \alpha \mu_s(T)| < \varepsilon \quad (\alpha \in [0, 1], s \in S).$$

For every $s \in S$ there exists an increasing family $(A_\alpha^s)_\alpha$ continuous with respect to s and such that

- (i) $|\mu_s(A_\alpha^s) - \alpha \mu_s(T)| < \varepsilon/10 \quad (\alpha \in [0, 1]);$
- (ii) $\sup_{\alpha \in [0, 1]} \mu_s(\bar{A}_\alpha^s \triangle A_\alpha^s) < 10\varepsilon.$

Proof. — By continuity, for each $s \in S$ there is a $\eta_s > 0$ such that $d(s, s') < 2\eta_s$ implies $\|\mu_s - \mu_{s'}\| < \varepsilon/60$ and $\mu_{s'}(\bar{A}_\alpha^s \triangle \bar{A}_\alpha^{s'}) < \varepsilon$. The open balls $B(s, \eta_s)$ cover S . Let $\{B(s_j, \eta_j) : j = 1, \dots, N\}$ be a finite sub-covering and $\{\lambda_j : j = 1, \dots, N\}$ be a continuous partition of unity subordinate to it and such that $\lambda_j(s_j) = 1, j = 1, \dots, N$.

Let $(A_\alpha^{s_j})_\alpha$ be the families defined by Lemma 3 by taking $\mu = \mu_{s_j}$.

Fix j such that $\mu_{s_j}(T) = \max\{\mu_{s_k}(T) : k = 1, \dots, N\}$ and choose $M \geq 2\mu_{s_j}(T)/\varepsilon$. By Lemma 4, extend the collection $(A_{i/M}^{s_k})_{i=0, \dots, M}$ ($k = 1, \dots, N$) to the family $(A_\alpha^s)_{\alpha \in [0, 1]} (s \in S)$.

The continuity of $s \rightarrow (A_\alpha^s)_{\alpha \in [0, 1]}$ follows from (iii) of Lemma 4, recalling that $\mu_s \leq \mu_0$ for each $s \in S$.

The choice of η_s and (i) of Lemma 4 imply that (i) holds. Moreover

$$\mu_{s_j}(\bar{A}_\alpha^s \triangle A_\alpha^s) \leq \mu_{s_j}(\bar{A}_\alpha^s \triangle \bar{A}_\alpha^{s_j}) + \mu_{s_j}(\bar{A}_\alpha^{s_j} \triangle A_\alpha^{s_j}) + \mu_{s_j}(A_\alpha^{s_j} \triangle A_\alpha^s).$$

By the choice of η_s and (ii) of Lemma 3, the r. h. s. is bounded by

$$\varepsilon + 6\varepsilon + \mu_{s_j}(A_\alpha^{s_j} \triangle A_\alpha^s),$$

which, by (ii) of Lemma 4 and the choice of M , yields

$$\mu_{s_j}(\bar{A}_\alpha^s \triangle A_\alpha^s) \leq \left(9 + \frac{1}{60}\right)\varepsilon.$$

Since $\|\mu_{s_j} - \mu_s\| < \varepsilon/60$, (ii) follows. ■

The following theorem shows the existence of a selection (\tilde{A}_α^s) from $\mathcal{A}(\mu_s)$, continuously depending on s .

THEOREM 1. — Let $s \rightarrow \mu_s$ be a continuous map from a compact metric space S into the space \mathcal{M} . For every $s \in S$ there an increasing family $(\tilde{A}_\alpha^s)_\alpha$ of measurable subsets of T satisfying

$$\mu_s(\tilde{A}_\alpha^s) = \alpha \mu_s(T) \quad (\alpha \in [0, 1]) \quad (6)$$

and such that the map $s \rightarrow (\tilde{A}_\alpha^s)_\alpha$ is continuous.

Proof. — We assume that we have defined for s in S an increasing family $(A_\alpha^{s,n})_\alpha$ which is continuous with respect to s and satisfies

$$|\mu_s(A_\alpha^{s,n}) - \alpha\mu_s(T)| < 10^{-n}.$$

By Lemma 2, the above is true for $n=1$ taking a family $(A_\alpha^{s,1})_\alpha$ constant with respect to s .

We obtain the existence of an increasing family $(A_\alpha^{s,n+1})_\alpha$ continuous with respect to s and such that

$$|\mu_s(A_\alpha^{s,n+1}) - \alpha\mu_s(T)| < 10^{-(n+1)} \quad (7)$$

and

$$\mu_s(A_\alpha^{s,n+1} \triangle A_\alpha^{s,n}) < 10^{-(n-1)}. \quad (8)$$

In fact, set in Lemma 5 \bar{A}_α^s to be $A_\alpha^{s,n}$ and ε to be 10^{-n} to infer the existence of a family, denoted by $(A_\alpha^{s,n+1})_\alpha$, satisfying (7) and (8).

Consider now the sequence $((A_\alpha^{s,n})_{n \in \mathbb{N}})$ defined by the above recursive procedure: we wish to show that it converges to a family $(\tilde{A}_\alpha^s)_\alpha$ which is continuous with respect to s and satisfies (6).

Property (8) implies that the sequence $(A_\alpha^{s,n})_n$ (s and α fixed) is a Cauchy sequence in \mathcal{F} supplied with the pseudometric $\rho_s(A, B) = \mu_s(A \triangle B)$. The procedure in Oxtoby [6], Chap. 10, defines a limit family $(\tilde{A}_\alpha^s)_\alpha$, which is increasing: $\tilde{A}_\alpha^s = \bigcup_{n \in \mathbb{N}} \bigcap_{m \geq n} A_\alpha^{s,m}$.

By the inequality

$$|\mu_s(A) - \mu_s(B)| \leq \mu_s(A \triangle B)$$

and (7) we have

$$\mu_s(\tilde{A}_\alpha^s) = \lim_{n \rightarrow \infty} \mu_s(A_\alpha^{s,n}) = \alpha\mu_s(T).$$

In order to check the continuity of the map $s \rightarrow (\tilde{A}_\alpha^s)_\alpha$, fix $\varepsilon > 0$ and $s^0 \in S$. Since the inequality (8) is uniform with respect to s and α , there exists an \bar{n} such that $\mu_s(A_\alpha^{s,\bar{n}} \triangle \tilde{A}_\alpha^s) < \varepsilon/5$ for every s in S and α in $[0, 1]$. Let $\delta > 0$ be such that

$$\|\mu_s - \mu_{s^0}\| < \varepsilon/10 \quad [s \text{ in } B(s^0, \delta)]$$

and

$$\sup_{\alpha \in [0, 1]} \mu_s(A_\alpha^{s',\bar{n}} \triangle A_\alpha^{s'',\bar{n}}) < \varepsilon/5 \quad [s, s' \text{ and } s'' \text{ in } B(s^0, \delta)].$$

Then for every $\alpha \in [0, 1]$, s, s' and s'' in $B(s^0, \delta)$, we have:

$$\begin{aligned} \mu_s(\tilde{A}_\alpha^{s'} \triangle \tilde{A}_\alpha^{s''}) &\leq \mu_s(\tilde{A}_\alpha^{s'} \triangle A_\alpha^{s',\bar{n}}) + \mu_s(A_\alpha^{s',\bar{n}} \triangle \tilde{A}_\alpha^{s''}) \\ &\leq \mu_s(\tilde{A}_\alpha^{s'} \triangle A_\alpha^{s',\bar{n}}) + \mu_s(A_\alpha^{s',\bar{n}} \triangle A_\alpha^{s'',\bar{n}}) + \mu_s(A_\alpha^{s'',\bar{n}} \triangle \tilde{A}_\alpha^{s''}) \end{aligned}$$

$$\begin{aligned} &\leq \|\mu_s - \mu_{s'}\| + \mu_{s'}(\tilde{A}_\alpha^{s'} \triangle A_\alpha^{s'}, \bar{n}) + \mu_s(A_\alpha^{s'}, \bar{n} \triangle A_\alpha^{s'}, \bar{n}) \\ &\quad + \|\mu_s - \mu_{s''}\| + \mu_{s''}(A_\alpha^{s'}, \bar{n} \triangle \tilde{A}_\alpha^{s''}) \\ &< \varepsilon. \quad \blacksquare \end{aligned}$$

COROLLARY. — Under the same assumptions, for every $\eta > 0$ and for every increasing family $(A_\alpha)_\alpha$ satisfying

$$|\mu_s(A) - \alpha \cdot \mu_s(T)| < \eta \quad (\alpha \in [0, 1], s \in S),$$

the family $(\tilde{A}_\alpha^s)_\alpha$ of Theorem 1 can be chosen as to satisfy, in addition,

$$\mu_s(\tilde{A}_\alpha^s \triangle A_\alpha) < \eta \quad (\alpha \in [0, 1], s \in S).$$

Proof. — Set $A_\alpha^{s,1}$ to be A_α in the proof of Theorem 1. \blacksquare

REFERENCES

- [1] A. FRYSZKOWSKI, Continuous Selections for a Class of Non-Convex Multivalued Maps, *Studia Mathematica*, T. LXXVI, 1983, pp. 163-174.
- [2] P. HALMOS, The Range of a Vector Measure, *Bull. Am. Math. Soc.*, Vol. 54, 1948, pp. 416-421.
- [3] P. HALMOS, *Measure Theory*, Van Nostrand, Princeton, 1950.
- [4] A. LIAPUNOV, Sur les fonctions-vecteurs complètement additives, *Bull. Acad. Sci. U.R.S.S., Ser. Math.*, Vol. 4, 1940, pp. 465-478 (Russian).
- [5] A. LIAPUNOV, same as above, Vol. 10, 1946, pp. 277-279 (Russian); for an English version, see J. DIESTEL and J. J. UHL, *Vector Measures*, 1977, A.M.S., Providence, Rhode Island, p. 262.
- [6] J. OXTOBY, *Measure and Category*, Second Edition, Springer Verlag, New York, 1980.
- [7] M. VALADIER, Une mesure vectorielle sans atome dont l'ensemble des valeurs est non convexe, *Boll. U.M.I., Serie VI, Vol. II-C, No. 1*, 1983, pp. 293-296.

(Manuscrit reçu le 5 septembre 1986)

(corrigé le 17 décembre 1986.)