

# ANNALES DE L'I. H. P., SECTION C

SISTO BALDO

## **Minimal interface criterion for phase transitions in mixtures of Cahn-Hilliard fluids**

*Annales de l'I. H. P., section C*, tome 7, n° 2 (1990), p. 67-90

[http://www.numdam.org/item?id=AIHPC\\_1990\\_\\_7\\_2\\_67\\_0](http://www.numdam.org/item?id=AIHPC_1990__7_2_67_0)

© Gauthier-Villars, 1990, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section C » (<http://www.elsevier.com/locate/anihpc>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Minimal interface criterion for phase transitions in mixtures of Cahn-Hilliard fluids

by

**Sisto BALDO**

Dipartimento di Matematica, Università di Pisa,  
Via Buonarroti 2, Pisa-Italia

---

**ABSTRACT.** — In this paper we extend the Van der Waals-Cahn-Hilliard theory of phase transitions to the case of a mixture of  $n$  non-interacting fluids. By describing the state of the mixture as given by a vector density function  $u = (u_1, \dots, u_n)$ , the problem consists in studying the asymptotic behaviour as  $\varepsilon \rightarrow 0^+$  of minimizers of the energy functionals:

$$E_\varepsilon(u) = \int_{\Omega} |\varepsilon^2 |Du|^2 + W(u)| dx$$

under the volume constraint  $\int_{\Omega} u(x) dx = m$ , with  $m \in \mathbf{R}^n$  fixed. The function  $W$ , which represents the Gibbs free energy, is non-negative and vanishes only in a finite number of points  $\alpha_1, \dots, \alpha_k \in \mathbf{R}^n$ . The result is that the minimizers asymptotically approach a configuration which corresponds to a partition of the container  $\Omega$  into  $k$  subsets whose boundaries satisfy a minimality condition.

*Key words* : Calculus of variations,  $\Gamma$ -convergence, relaxation, fluids, phase transitions.

**RÉSUMÉ.** — Dans cet article nous étendons la théorie des transitions de phase de Van der Waals-Cahn-Hilliard au cas d'un mélange de  $n$  fluides non interagés. En supposant l'état du mélange décrit par une fonction

---

*Classification A.M.S.* : 49 A 50.

vectorielle de densité  $u = (u_1, \dots, u_n)$ , le problème consistera dans l'étude du comportement asymptotique par  $\varepsilon \rightarrow 0^+$  des minimisants des énergies :

$$E_\varepsilon(u) = \int_{\Omega} |\varepsilon^2 |Du|^2 + W(u)| dx$$

sous la contrainte de volume  $\int_{\Omega} u(x) dx = m$ , avec  $m \in \mathbf{R}^n$  fixé. La fonction

$W$  représente l'énergie libre de Gibbs, à valeurs non négatives et qui est nulle sur un nombre fini de points  $\alpha_1, \dots, \alpha_k \in \mathbf{R}^n$ . Nous obtenons alors que les minimisants approchent asymptotiquement une configuration qui correspond à une partition du container  $\Omega$  en  $k$  sous-ensembles dont les bords satisfont à une certaine condition de minimalité.

## 1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

We briefly recall the theory for a single fluid (*see*, for example, [GM]). Suppose we have a fluid, contained in an open set  $\Omega \subset \mathbf{R}^N$ , of total mass  $m$  and density given by a function  $u$  defined on  $\Omega$ .

Under isothermal conditions, the stable configurations of the fluid are given by the functions  $u$  which minimize the total energy:

$$E_\varepsilon(u) = \int_{\Omega} [\varepsilon^2 |Du|^2 + W(u)] dx \rightarrow \min \quad (1.1)$$

among all  $u$  such that  $\int_{\Omega} u(x) dx = m$  and  $u(x) \geq 0$ . Here  $W$  is a nonnegative function that represents the Gibbs free energy relative to an ideal fluid, and  $\varepsilon$  is a small parameter that takes into account the energy connected with the formation of interfaces.

Assume that  $W$  acts as in Figure 1 and that  $\alpha|\Omega| < m < \beta|\Omega|$ , where  $|\Omega|$  denotes the volume of  $\Omega$ .

Note that the solutions of the variational problem (1.1) have a certain regularity, at least  $u \in H^1(\Omega)$ , while it is physically reasonable that the density  $u$  should be a piecewise constant function admitting only the values  $\alpha$  and  $\beta$  (*i. e.* the fluid splits into two phases). On the other hand, experience indicates that stable configurations must minimize the surface area of the interfaces between the phases, and so we cannot simply neglect the gradient term in (1.1).




FIG.

The idea to overcome such difficulties is to study the asymptotic behaviour of the variational problem (1.1) as  $\varepsilon \rightarrow 0^+$ . The problem was recently solved in full generality by Modica [M1]. He proved that, if  $u_\varepsilon$  is a solution of the variational problem (1.1) and  $u_\varepsilon$  converges in  $L^1(\Omega)$  as  $\varepsilon \rightarrow 0^+$  to a function  $u_0$ , then  $u_0$  takes only the values  $\alpha$  and  $\beta$  and the set  $E = \{x \in \Omega : u(x) = \alpha\}$  minimizes the surface area of its boundary among all subsets of  $\Omega$  having the same volume.

It should be remarked that Modica's result does not work for more than two phases. For instance, if  $W(\alpha) = W(\beta) = W(\gamma) = 0$  with  $\alpha < \beta < \gamma$ , the limit density  $u_0$  takes again only two values: either  $\alpha$  and  $\beta$ , or  $\beta$  and  $\gamma$ , depending on  $m$ . The multiphases systems, as well as the mixtures of non-interacting fluids, can be studied by passing from the scalar case  $u \in \mathbb{R}$  to the vector case  $u \in \mathbb{R}^n$ . This is the aim of the present paper.

Let  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 2$  be open and bounded, with lipschitz-continuous boundary. Suppose we have  $n$  fluids that we can describe with a vector-valued density function  $u = (u^1, \dots, u^n) \in L^1(\Omega; \mathbb{R}^n)$ : each scalar component of  $u$  is the density of an ingredient of the mixture. The appropriate constraint are again, with obvious meaning of the symbols,

$$\int_{\Omega} u(x) dx = m, \quad u > 0$$

where  $m = (m^1, \dots, m^n) \in \mathbb{R}^n$  is given and satisfies the condition  $\min \{ \alpha_1^i, \dots, \alpha_k^i \} \leq \frac{m^i}{|\Omega|} \leq \max \{ \alpha_1^i, \dots, \alpha_k^i \}$  for every  $i = 1, \dots, n$ . By physical considerations, it seems to be reasonable that the free energy of the mixture is the sum of the free energies of the components. So we assume that the free energy of the fluid is a nonnegative, continuous function  $W(u)$ , defined for  $u \in \mathbb{R}^n$ ,  $u > 0$ , which vanishes in a finite number of points  $\alpha_1, \dots, \alpha_k \in \mathbb{R}^n$ .

For example, if  $n=2$  and  $W(u^1, u^2) = W_1(u^1) + W_2(u^2)$  with  $W_1$  and  $W_2$  acting as in Figure 1 with zeros  $\alpha_1, \beta_1$  and  $\alpha_2, \beta_2$ , then  $k=4$  and  $\alpha^1 = (\alpha_1, \alpha_2)$ ,  $\alpha^2 = (\alpha_1, \beta_2)$ ,  $\alpha^3 = (\beta_1, \alpha_2)$ ,  $\alpha^4 = (\beta_1, \beta_2)$ .

We need also the following technical assumption on  $W$ : there exist  $K_1, K_2 \in \mathbf{R}$  with  $0 < K_1 < K_2$ , such that:

$$W(u) \geq \sup \{ W(v) : v \in [K_1, K_2]^n \} \quad (1.2)$$

for every  $u \notin [K_1, K_2]^n$ .

We can now state our main result. For every  $A \subset \mathbf{R}^N$  denote by  $\mathbf{1}_A$  its characteristic function, by  $\partial^* A$  its reduced boundary (cf. Giusti [Gi]), by  $|A|$  its Lebesgue measure, by  $H_{N-1}(A)$  its  $(N-1)$ -dimensional Hausdorff measure. For every  $i, j \in \{1, \dots, k\}$  define

$$d(\alpha_i, \alpha_j) = \inf \left\{ \int_0^1 W^{1/2}(\gamma(t)) |\gamma'(t)| dt : \gamma \in C^1([0, 1]; \mathbf{R}^n), \right. \\ \left. \gamma > 0, \gamma(0) = \alpha_i, \gamma(1) = \alpha_j \right\}.$$

**THEOREM.** — For every  $\varepsilon > 0$ , let  $u_\varepsilon$  be a solution of the minimization problem:

$$\min \left\{ \int_{\Omega} [\varepsilon^2 |Du|^2 + W(u)] dx : u \in H^1(\Omega; \mathbf{R}^n), \int_{\Omega} u(x) dx = m, u > 0 \right\}.$$

Assume that the functions  $u_\varepsilon$  converge in  $L^1(\Omega, \mathbf{R}^n)$ , as  $\varepsilon \rightarrow 0^+$ , to a function  $u_0$ . Then

$$u_0(x) = \sum_{i=1}^k \alpha_i \mathbf{1}_{S_i}(x)$$

where  $S_1, \dots, S_k$  is a partition of  $\Omega$  which minimizes the quantity

$$\sum_{i, j=1}^k d(\alpha_i, \alpha_j) H_{N-1}(\partial^* S_i \cap \partial^* S_j)$$

among all other partitions of  $\Omega$  such that  $\sum_{i=1}^k |S_i| \alpha_i = m$ .

We mention that results of this type in the vector case have been recently obtained by P. Sternberg [S] and I. Fonseca & L. Tartar [FT]. In [S] the problem is studied by taking  $n=2$  and  $W$  which vanishes on two simple smooth curves in the plane and without assuming any constraint. In [FT] the problem is studied with  $n$  arbitrary, but  $k=2$ , and with the same volume constraint as in the present paper. Furthermore, Ambrosio [A] has announced general results in the case in which the set of zeros of  $W$  is essentially any closed subset of  $\mathbf{R}^n$ . All the mentioned papers, as well as the present paper, rely on the  $\Gamma$ -convergence theory.

I wish to thank Professor L. Modica for suggesting me this research as subject of my thesis for the “Laurea in Matematica” at the university of Pisa.

2. PRELIMINARY RESULTS

For any  $v \in L^1(\Omega)$ , it is usual to define

$$\int_A |Dv| = \sup \left\{ \int_A v \operatorname{div} g \, dx : g \in C_0^1(\Omega; \mathbf{R}^N), |g| \leq 1 \right\}$$

for any open subset  $A$  of  $\Omega$ . If  $\int_\Omega |Dv| < +\infty$ , then  $u \in BV(\Omega)$ ; in this case the set function  $A \mapsto \int_A |Dv|$  is the trace on the open subsets of  $\Omega$  of a borel measure on  $\Omega$ , which will be denoted by  $\int_E |Dv|$  for any borel subset  $E$  of  $\Omega$ . If  $v$  is the characteristic function  $\mathbf{1}_S$  of a measurable subset  $S$  of  $\Omega$ , then we let

$$\int_\Omega |Dv| = P_\Omega(S)$$

and  $P_\Omega(S)$  is called perimeter of  $S$  in  $\Omega$ , because  $P_\Omega(S) = H_{N-1}(\partial S \cap \Omega)$  in the case that  $S$  has a smooth boundary. For every  $S$ , it is possible to construct a subset  $\partial^* S \subset \partial S$ , called reduced boundary of  $S$ , such that

$$P_\Omega(S) = H_{N-1}(\partial^* S \cap \Omega).$$

Denote by  $\mathbf{R}_+^n$  the set of all  $u \in \mathbf{R}^n$  such that  $u \geq 0$ , that is  $u_1 \geq 0, \dots, u_n \geq 0$ . On  $\mathbf{R}_+^n$  we define the following metric:

$$d(\zeta_1, \zeta_2) = \inf \left\{ \int_0^1 W^{1/2}(\gamma(t)) |\gamma'(t)| \, dt : \gamma \in C^1([0, 1]; \mathbf{R}_+^n), \gamma(0) = \zeta_1, \gamma(1) = \zeta_2 \right\}$$

which is the riemannian metric derived from  $W^{1/2}$ .

For  $\zeta \in \mathbf{R}_+^n$ , let  $\varphi_i(\zeta) = d(\zeta, \alpha_i)$ ,  $i = 1, \dots, k$ , where  $\alpha_i$  are the zeros of  $W$ . We begin with a simple result on the metric  $d$ .

PROPOSITION 2.1. — *The function  $\varphi_i$  is locally lipschitz-continuous. Moreover, if  $u \in H^1(\Omega) \cap L^\infty(\Omega)$ , then  $\varphi_i \circ u \in W^{1,1}(\Omega)$  and the following inequality holds:*

$$\int_\Omega |D(\varphi_i \circ u)| \, dx \leq \int_\Omega W^{1/2}(u(x)) |Du(x)| \, dx. \tag{2.1}$$

*Proof.* — We omit the very easy proof that  $\varphi_i$  is locally lipschitz-continuous. If the following inequality holds:

$$\int_{\Omega'} |D(\varphi_i \circ u)| \leq \int_{\Omega'} W^{1/2}(u(x)) |Du(x)| dx \quad (2.2)$$

for any  $\Omega'$  open set in  $\Omega$ , then, by the Radon-Nikodym Theorem,  $D(\varphi_i \circ u)$  is a  $L^1$  function and (2.1) immediately follows. Let us prove (2.2).

Consider the case  $u \in C^1(\Omega)$ . The function  $\varphi_i \circ u$  is locally lipschitz-continuous, and then differentiable almost everywhere in  $\Omega$ . Let  $x$  be a differentiability point for  $\varphi_i \circ u$ ,  $\{x_h\}$  be a sequence converging to  $x$  in  $\Omega$  and  $\sigma_h$  be the segment  $\sigma_h(t) = (1-t)x + tx_h$ . We have, by definition of the metric:

$$\begin{aligned} |\varphi_i(u(x)) - \varphi_i(u(x_h))| &\leq d(u(x), u(x_h)) \\ &\leq \int_0^1 W^{1/2}(u(\sigma_h(t)) |Du(\sigma_h(t))| |x_h - x| dt. \end{aligned}$$

By applying the Mean Value Theorem to the right-hand side integral, dividing by  $|x_h - x|$ , and passing to the limit we obtain

$$|D(\varphi_i(u(x)))| \leq W^{1/2}(u(x)) |Du(x)|,$$

and (2.2) follows.

Suppose now  $u \in H^1(\Omega) \cap L^\infty(\Omega)$ , and let  $\{u_h\} \subset C^1(\Omega)$  be a sequence such that  $u_h \rightarrow u$  in  $H^1(\Omega)$ . Suppose also that  $u_h(x) \rightarrow u(x)$  and  $Du_h(x) \rightarrow Du(x)$  almost everywhere in  $\Omega$ .

By using the Dominated Convergence Theorem and the Fatou's Lemma we obtain for any  $g \in C_0^1(\Omega'; \mathbf{R}^N)$ ,  $|g| \leq 1$  in  $\Omega'$ :

$$\begin{aligned} \int_{\Omega'} (\varphi_i \circ u) \operatorname{div} g \, dx &= \lim_{h \rightarrow +\infty} \int_{\Omega'} (\varphi_i \circ u_h) \operatorname{div} g \, dx \\ &- \lim_{h \rightarrow +\infty} \int_{\Omega'} \langle D(\varphi_i \circ u_h), g \rangle \, dx \leq \lim_{h \rightarrow +\infty} \int_{\Omega'} |D(\varphi_i \circ u_h)| \, dx \\ &\leq \limsup_{h \rightarrow +\infty} \int_{\Omega'} W^{1/2}(u_h(x)) |Du_h(x)| \, dx \leq \int_{\Omega'} W^{1/2}(u(x)) |Du(x)| \, dx \end{aligned}$$

and the proposition is proved. ■

Let  $\mu$  and  $\nu$  be two regular positive Borel measures on  $\Omega$ . We define the supremum  $\mu \vee \nu$  of  $\mu$  and  $\nu$  as the smallest regular positive measure which is greater than or equal to  $\mu$  and  $\nu$  on all borel subsets of  $\Omega$ . We

have:

$$(\mu \vee \nu)(A) = \sup \left\{ \mu(A') + \nu(A'') : A' \cap A'' = \emptyset, \right. \\ \left. A' \cup A'' \subset A, A' \text{ and } A'' \text{ are open sets in } \Omega \right\}$$

for any open subset A of  $\Omega$ .

Let  $u$  be a function such that  $W(u(x))=0$  almost everywhere and  $\varphi_i \circ u \in BV(\Omega)$ ; hence

$$u(x) = \sum_{i=1}^k \alpha_i \mathbf{1}_{S_i}(x) \tag{2.3}$$

where  $S_1, \dots, S_k$  are pairwise disjoint sets in  $\Omega$  such that  $\left| \Omega \setminus \bigcup_{i=1}^k S_i \right| = 0$ .

PROPOSITION 2.2. — Denote  $\mu_i$  the Borel measure  $\mu_i : E \mapsto \int_E |D(\varphi_i \circ u)|$ . Then  $P_\Omega(S_i) < +\infty$  for  $i=1, \dots, k$  and

$$\left( \bigvee_{i=1}^k \mu_i \right) (\Omega) = 1/2 \sum_{i,j=1}^k d(\alpha_i, \alpha_j) H_{N-1}(\partial^* S_i \cap \partial^* S_j \cap \Omega). \tag{2.4}$$

Proof. — Let us prove that the perimeters of the sets  $S_i$  in  $\Omega$  are bounded. By applying the coarea-type Fleming-Rishel formula:

$$\int_\Omega |D(\varphi_i \circ u)| = \int_{-\infty}^{+\infty} P_\Omega(\{x : \varphi_i(u(x)) \leq t\}) dt \\ \geq \int_0^{d_i} P_\Omega(\{x : \varphi_i(u(x)) \leq t\}) dt = d_i P_\Omega(S_i)$$

where  $d_i = \min_{\substack{j=1, \dots, k \\ j \neq i}} (d(\alpha_i, \alpha_j))$ ; hence  $P_\Omega(S_i) < +\infty$ .

Suppose we have proved that for every index  $i=1, \dots, k$  and for every open set  $\Omega' \subset \Omega$  we have:

$$2 \int_{\Omega'} |D(\varphi_i \circ u)| = \sum_{j,h=1}^k H_{N-1}(\partial^* S_h \cap \partial^* S_j \cap \Omega') |\varphi_i(\alpha_h) - \varphi_i(\alpha_j)|. \tag{2.5}$$

Since we have  $|\varphi_l(\alpha_i) - \varphi_l(\alpha_j)| \leq d(\alpha_i, \alpha_j)$  for any  $i, j, l=1, \dots, k$ , and the equality holds for  $l=i$  or  $l=j$ , the result follows from the lemma below, by passing to the measure theoretical supremum in (2.5).

We now prove (2.5). Note that for any  $J \subset \{1, \dots, k\}$  we have

$$P_{\Omega'}(\bigcup_{j \in J} S_j) = \sum_{j \in J} \sum_{\substack{i=1 \\ i \notin J}}^k H_{N-1}(\partial^* S_i \cap \partial^* S_j \cap \Omega') \tag{2.6}$$



In fact, reasoning as in Vol'pert [Vo]:

$$\begin{aligned} P_{\Omega'}(S_i \cup S_j) &= H_{N-1}(\partial^* S_i \Delta \partial^* S_j) \quad (i \neq j) \\ \partial^* S_i &= \bigcup_{\substack{j=1 \\ j \neq i}}^k (\partial^* S_i \cap \partial^* S_j) \cup N \quad \text{where } H_{N-1}(N) = 0. \end{aligned}$$

An inductive argument on the cardinality of  $J$  proves (2.6). We now return to the proof of (2.5). Suppose for simplicity that  $i=1$  and that the indices are numbered in such a way that

$$0 = \varphi_1(\alpha_1) < \varphi_1(\alpha_2) < \dots < \varphi_1(\alpha_k).$$

(It's possible that there are indices  $j$  such that  $\varphi_1(\alpha_j) = \varphi_1(\alpha_{j+1})$ , but the changes in the proof are trivial.) By the Fleming-Rishel formula we obtain:

$$\begin{aligned} \int_{\Omega'} |D(\varphi_1 \circ u)| &= \int_0^{\varphi_1(\alpha_1)} P_{\Omega'}(\{x \in \Omega' : \varphi_1(u(x)) \leq t\}) dt \\ &= \sum_{j=1}^{k-1} [\varphi_1(\alpha_{j+1}) - \varphi_1(\alpha_j)] P_{\Omega'}\left(\bigcup_{l=1}^j S_l\right) \\ &= \sum_{j=1}^{k-1} \sum_{l=1}^j \sum_{m=j+1}^k [\varphi_1(\alpha_{j+1}) - \varphi_1(\alpha_j)] H_{N-1}(\partial^* S_l \cap \partial^* S_m). \end{aligned}$$

Reordering the terms in the sum, we finally obtain (2.5). ■

LEMMA 2.3. — *Let  $\mu$  be a regular positive borel measure on  $\Omega$ ,  $B_1, \dots, B_m$  be disjoint borel subsets of  $\Omega$  with finite  $\mu$ -measure and  $c_i^h$ ,  $i=1, \dots, m$ ,  $h=1, \dots, k$  be positive coefficients. Define*

$$\mu_h(A) = \sum_{i=1}^m c_i^h \mu(A \cap B_i); \quad \nu(A) = \sum_{i=1}^m (\max_h c_i^h) \mu(A \cap B_i).$$

Then  $\nu = \bigvee_{i=1}^k \mu^h$ .

*Proof.* — We omit the easy proof. ■

On the basis of the previous proposition, we define, for any  $u \in L^1(\Omega)$ :

$$F_0(u) = \begin{cases} 2 \bigvee_{i=1}^k \mu_i = 2 \bigvee_{i=1}^k \int_{\Omega} |D(\varphi_i \circ u)| & \text{if } \varphi_i \circ u \in \text{BV}(\Omega), \quad W(u(x)) = 0 \text{ a. e.}, \quad \int_{\Omega} u dx = m \\ +\infty & \text{otherwise} \end{cases}$$

and we obtain that, if  $F_0(u) < +\infty$ , then

$$F_0(u) = \sum_{i,j=1}^k d(\alpha_i, \alpha_j) H_{N-1}(\partial^* S_i \cap \partial^* S_j \cap \Omega).$$

We now recall the definition of  $\Gamma$ -convergence [DGF]:

Let  $\{G_h\}$  be a sequence of real-extended functionals on  $L^1(\Omega)$ ,  $G_0$  a functional of the same kind. We say that  $G_h$   $\Gamma$ -converges to  $G_0$  as  $h \rightarrow +\infty$  at a point  $u \in L^1(\Omega)$ , and we write:

$$\Gamma\text{-}\lim_{h \rightarrow +\infty} G_h(u) = G_0(u)$$

if and only if the following relations hold:

$$\forall u_h \rightarrow u \text{ in } L^1(\Omega) \text{ we have } G_0(u) \leq \liminf_{h \rightarrow +\infty} G_h(u_h) \tag{2.8}$$

$$\exists u_h \rightarrow u \text{ in } L^1(\Omega) \text{ such that } G_0(u) \geq \limsup_{h \rightarrow +\infty} G_h(u_h) \tag{2.9}$$

As immediate consequence, we obtain the following result (see [DGF]):

**PROPOSITION 2.4.** - Assume  $G_0(u) = \Gamma\text{-}\lim_{h \rightarrow +\infty} G_h(u)$  for every  $u \in L^1(\Omega)$ . Let  $u_h$  be such that  $G_h(u_h) = \min \{ G_h(u) : u \in L^1(\Omega) \}$  for every  $h \in \mathbb{N}$ . If  $u_h \rightarrow u_0$  in  $L^1(\Omega)$ , then:

$$\lim_{h \rightarrow +\infty} G_h(u_h) = G_0(u_0) = \min \{ G_0(u) : u \in L^1(\Omega) \}.$$

The result we shall prove in the next sections is the following. Define, for any  $u \in L^1(\Omega)$ ,

$$F_\varepsilon(u) = \begin{cases} \int_\Omega \left[ \varepsilon |Du|^2 + \frac{1}{\varepsilon} W(u) \right] dx & \text{if } u \in H^1(\Omega; \mathbb{R}^n_+) \text{ and } \int_\Omega u(x) dx = m \\ +\infty & \text{otherwise.} \end{cases}$$

**THEOREM 2.5.** - For every  $\varepsilon_h \downarrow 0$ ,

$$\Gamma\text{-}\lim_{h \rightarrow +\infty} F_{\varepsilon_h}(u) = F_0(u), \quad \forall u \in L^1(\Omega).$$

Our main result is now a consequence of Theorem 2.5. Indeed, by the hypothesis  $u_\varepsilon \rightarrow u_0$ . Proposition 2.4 yields that  $F_0$  admits a minimum value and  $u_0$  is a minimum point of  $F_0$ . Furthermore, the fact that this minimum value is obviously finite gives the formula for the asymptotic value of the energies  $F_\varepsilon(u_\varepsilon)$ .

### 3. THE PROOF OF THE MAIN RESULT

To prove our main result in the form of Theorem 2.5 we only need to prove the statements (2.8) and (2.9) in the definition of  $\Gamma$ -convergence.

*Proof of (2.8).* — Let  $\varepsilon_h \downarrow 0$  be a fixed sequence. It is not restrictive to assume  $\lim_{h \rightarrow +\infty} F_{\varepsilon_h}(u_h)$  exists and it is finite, being trivial the other cases.

By choosing a subsequence  $u_{h_k}$  that converges to  $u$  pointwise almost everywhere in  $\Omega$ , and by Fatou's Lemma, we obtain:

$$\int_{\Omega} W(u) dx \leq \liminf_{k \rightarrow +\infty} \int_{\Omega} W(u_{h_k}) dx \leq \liminf_{k \rightarrow +\infty} \varepsilon_{h_k} F_{\varepsilon_{h_k}}(u_{h_k}) = 0;$$

hence  $W(u(x)) = 0$  almost everywhere in  $\Omega$  because  $W$  is a continuous and nonnegative function.

We now need to prove that

$$2 \bigvee_{i=1}^k \int_{\Omega} |D(\varphi_i \circ u)| \leq \lim_{h \rightarrow +\infty} \int_{\Omega} \left[ \varepsilon_h |Du_h|^2 + \frac{1}{\varepsilon_h} W(u_h) \right] dx \quad (3.1)$$

We further reduce the problem by making the sequence  $u_h$  to be equibounded. If this is not true, by using the technical assumption (1.2) on  $W$ , we can replace  $u_h$  by the  $\tilde{u}_h$  obtained through truncation of each scalar component by  $K_1$  and  $K_2$ . Note that  $\tilde{u}_h \rightarrow u$  in  $L^1(\Omega)$  and the integrals in the right-hand side of (3.1) decrease. Then,  $(\varphi_i \circ \tilde{u}_h) \rightarrow (\varphi_i \circ u)$  in  $L^1(\Omega)$  for every  $i = 1, \dots, k$ , because of the continuity of  $\varphi_i$ , and lower semicontinuity yields:

$$|D(\varphi_i \circ u)|(A) \leq \liminf_{h \rightarrow +\infty} |D(\varphi_i \circ u_h)|(A)$$

for every  $i = 1, \dots, k$ ,  $A$  open set in  $\Omega$ .

Thus:

$$\begin{aligned} \bigvee_{i=1}^k |D(\varphi_i \circ u)|(\Omega) &= \sup \left\{ \sum_{i=1}^k |D(\varphi_i \circ u)|(A_i) : \bigcup_{i=1}^k A_i \subset \Omega, \right. \\ &\quad \left. A_i \cap A_j = \emptyset \text{ if } i \neq j, A_i \text{ open sets in } \Omega \right\} \\ &\leq \liminf_{h \rightarrow +\infty} \bigvee_{i=1}^k |D(\varphi_i \circ u_h)|(\Omega) \end{aligned}$$

Recalling Proposition 2.1 we finally obtain:

$$\begin{aligned} \liminf_{h \rightarrow +\infty} \bigvee_{i=1}^k |D(\varphi_i \circ u_h)|(\Omega) &\leq \liminf_{h \rightarrow +\infty} \int_{\Omega} W^{1/2}(u_h) |Du_h(x)| dx \\ &\leq \frac{1}{2} \liminf_{h \rightarrow +\infty} \int_{\Omega} \left[ \varepsilon_h |Du_h|^2 + \frac{1}{\varepsilon_h} W(u_h) \right] dx. \quad \blacksquare \end{aligned}$$

*Proof of (2.9).* — Fix  $\varepsilon_h \downarrow 0$ . If  $u \in L^1(\Omega)$  is such that  $F_0(u) = +\infty$ , the construction of the sequence  $u_{\varepsilon_h}$  is trivial. Therefore assume

$u(x) = \sum_{i=1}^k \alpha_i \mathbf{1}_{S_i}(x)$ , with  $S_1, \dots, S_k$  pairwise disjoint sets with finite perimeter in  $\Omega$  such that  $\left| \Omega \setminus \bigcup_{i=1}^k S_i \right| = 0$ .

The following lemma, whose proof is given in appendix, and a diagonal argument allows us to consider partitions such that the sets  $S_1, \dots, S_k$  are polygonal domains in  $\mathbf{R}^N$ , with  $H_{N-1}(\partial S_i \cap \partial \Omega) = 0$ :

LEMMA 3.1. — *Let  $\{S_1, \dots, S_k\}$  as before. Then there exists a sequence  $\{S_1^h, \dots, S_k^h\}_{h \in \mathbf{N}}$  of partitions of  $\Omega$  such that:*

(i)  $S_i^h$  is a polygonal domain and  $H_{N-1}(\partial S_i^h \cap \partial \Omega) = 0$  for any  $i = 1, \dots, k, h \in \mathbf{N}$ ;

(ii) if  $u_h(x) = \sum_{i=1}^k \alpha_i \mathbf{1}_{S_i^h}(x), u(x) = \sum_{i=1}^k \alpha_i \mathbf{1}_{S_i}(x)$ , then  $u_h \rightarrow u$  in  $L^1(\Omega)$  as  $h \rightarrow +\infty$ ;

(iii)  $\int_{\Omega} u_h(x) dx = \int_{\Omega} u(x) dx = m$  for any  $h \in \mathbf{N}$ ;

(iv)  $\lim_{h \rightarrow +\infty} \bigvee_{i=1}^k \int_{\Omega} |D(\varphi_i \circ u_h)| = \bigvee_{i=1}^k \int_{\Omega} |D(\varphi_i \circ u)|$ .

The following lemma, that will be essential in the construction of our sequence  $u_h$ , generalizes an idea of Modica [M2].

We suppose for simplicity that for any  $i, j = 1, \dots, k, i \neq j$ , there exists the distance-minimizing geodesic connecting  $\alpha_i$  and  $\alpha_j$ , i.e. we suppose that there exists a  $C^1$ -path  $\gamma_{ij}$  such that  $\gamma_{ij}(0) = \alpha_i, \gamma_{ij}(1) = \alpha_j$  and

$d(\alpha_i, \alpha_j) = \int_0^1 W^{1/2}(\gamma_{ij}(t)) |\gamma'_{ij}(t)| dt$ . We shall see later how to change the

proof if such geodesics does not exist. Note that it is not restrictive to assume  $|\gamma'_{ij}(t)| \neq 0, \forall t \in (0, 1)$ .

LEMMA 3.2. — *Consider the following ordinary differential equations:*

$$(y_\varepsilon^{ij})' = \frac{\delta + W(\gamma_{ij}(y_\varepsilon^{ij}))}{\varepsilon^2 |\gamma'_{ij}(y_\varepsilon^{ij})|^2} \tag{3.2}$$

where  $i, j = 1, \dots, k, i \neq j$  and  $\delta > 0$  is a fixed constant.

Thus, for every  $\varepsilon > 0$ , there exist a lipschitz-continuous function  $\chi_\varepsilon: \mathbf{R}^{k-1} \rightarrow \mathbf{R}^n$  and three constants  $C_1, C_2, C_3$  (depending only on  $\delta$ ) such that:

(i)  $\chi_\varepsilon(t_1, \dots, t_{k-1}) = \alpha_1$  if  $t_1 < 0$ .

$\chi_\varepsilon(t) = \alpha_i$  if  $t_1 > C_1 \varepsilon, \dots, t_{i-1} > C_1 \varepsilon, t_i < 0$  for any  $i = 2, \dots, k-1$ .

$\chi_\varepsilon(t) = \alpha_k$  if  $t_1 > C_1 \varepsilon, \dots, t_k > C_1 \varepsilon$ .

(ii)  $0 < |\chi_\varepsilon| < C_2, |D_{\chi_\varepsilon}| < C_3/\varepsilon$  a.e. in  $\mathbf{R}^{k-1}$

(iii) If  $j > i$ , on the set  $\{t \in \mathbf{R}^{k-1} : 0 < t_i < C_1 \varepsilon; t_j < 0; t_h > C_1 \varepsilon \text{ for any } h \neq i, j\}$   $\chi_\varepsilon$  depends only on  $t_i$ , and we can write:

$\chi_\varepsilon(t_i) = \gamma_{ij}(y_\varepsilon^{ij}(t_i))$  [where  $y_\varepsilon^{ij}$  solves (3.2)] for any  $t_i$  such that  $\chi_\varepsilon(t_i) \neq \alpha_i$   
 (Note: if  $j = k$ , ignore the condition  $t_j < 0$ , that makes no sense.)

*Proof.* — To prove the lemma, we only need to find the constants  $C_1, C_2, C_3$  and to define  $\chi_\varepsilon$  at the points different from those considered in (i). First we search the solutions of (3.1), for fixed  $i, j$ . The function

$$\psi_\varepsilon(t) = \int_0^t \frac{\varepsilon |\gamma'_{ij}(s)|}{[\delta + W(\gamma_{ij}(s))]^{1/2}} ds \quad (t \in [0, 1])$$

is obviously increasing and, if  $\eta_\varepsilon = \psi_\varepsilon(1)$ , we immediately obtain:  $\eta_\varepsilon \leq \varepsilon \delta^{-1/2} \text{ length}(\gamma_{ij})$ . Now, the inverse function  $\tilde{y}_\varepsilon : [0, \eta_\varepsilon] \rightarrow [0, 1]$  of  $\psi_\varepsilon$  satisfies the differential equation (3.1). We extend the function to the whole  $\mathbf{R}$  by putting:

$$y_\varepsilon(t) = \begin{cases} 0 & t \leq 0 \\ \tilde{y}_\varepsilon & 0 \leq t \leq \eta_\varepsilon \\ 1 & t \geq \eta_\varepsilon \end{cases}$$

Now  $y_\varepsilon$  is a lipschitz-continuous function satisfying (3.1) in all the points where  $y_\varepsilon \neq 1$ . Putting:

$$C_1 = \max_{i, j=1, \dots, k} \{ \delta^{-1/2} \text{ length}(\gamma_{ij}) \}$$

we can define  $\chi_\varepsilon$  on the strips as in (iii). We choose

$$C_2 > \sup \left\{ |y| : y \in \bigcup_{i, j=1}^k \gamma_{ij}([0, 1]) \right\}$$

If

$$K = \sup_{\substack{i, j=1, \dots, k \\ i \neq j, t \in [0, 1]}} \left[ \frac{[\delta + W(\gamma_{ij}(t))]^{1/2}}{|\gamma'_{ij}(t)|} \right]$$

we have  $|D_{\chi_\varepsilon}| \leq K/\varepsilon$ . Standard extension results for lipschitz-continuous functions allow one to define  $\chi_\varepsilon$  on the whole  $\mathbf{R}^{k-1}$ , with  $C_3 > K$  suitably chosen. ■

The last lemma has to do with tubular neighborhood of polygonal domains, and we omit the standard proof.

LEMMA 3.3. — Let  $\Omega$  be an open set in  $\mathbf{R}^N$ ,  $A$  a polygonal domain in  $\mathbf{R}^N$  with  $\partial A$  compact and such that  $H_{N-1}(\partial A \cap \partial \Omega) = 0$ . Put:

$$h(x) = \begin{cases} \text{dist}(x, \partial A) & \text{if } x \notin A \\ -\text{dist}(x, \partial A) & \text{if } x \in A. \end{cases}$$

Then there exists a constant  $\eta > 0$  such that  $h$  is lipschitz-continuous on  $H_\eta = \{x \in \mathbb{R}^N : |h(x)| < \eta\}$ , and  $|Dh(\alpha)| = 1$  for almost all  $x \in H_\eta$ .

Finally, if  $S_t$  denotes the set  $\{x \in \mathbb{R}^N : h(x) = t\}$  we have:

$$\lim_{t \rightarrow 0^\pm} H_{N-1}(S_t \cap \Omega) = H_{N-1}(\partial A \cap \Omega).$$

Let us return to the proof of (2.9) for polygonal domains. We define, for  $i \in \{1, \dots, k\}$ :

$$h_i(x) = \begin{cases} \text{dist}(x, \partial S_i) & \text{if } x \notin S_i \\ -\text{dist}(x, \partial S_i) & \text{if } x \in S_i \end{cases}$$

Fix  $\delta > 0$ . For  $\varepsilon$  small enough we have  $|Dh_i(x)| = 1$  a.e. on the set  $\{x \in \Omega : |h_i(x)| < C_1 \varepsilon\}$ , for all  $i = 1, \dots, k$ .

Consider the sequence of functions:

$$\tilde{u}_\varepsilon(x) = \chi_\varepsilon(h_1(x), \dots, h_{k-1}(x)).$$

Putting  $\Sigma_i^t = \{x \in \Omega : h_i(x) = t\}$ ,  $t > 0$ ,  $i = 1, \dots, k-1$ , we obtain:

$$\begin{aligned} \int_\Omega |\tilde{u}_\varepsilon - u| &\leq \sum_{i=1}^{k-1} \int_{\{x \in \Omega : 0 < h_i(x) < C_2 \varepsilon\}} |\tilde{u}_\varepsilon - u| dx \\ &\leq 2 C_2 \sum_{i=1}^{k-1} |\{x \in \Omega : 0 \leq h_i(x) \leq C_1 \varepsilon\}| \\ &= 2 C_2 \sum_{i=1}^{k-1} \int_{\{x \in \Omega : 0 \leq h_i(x) \leq C_1 \varepsilon\}} |Dh_i(x)| dx = 2 C_2 \sum_{i=1}^{k-1} \int_0^{C_1 \varepsilon} H_{N-1}(\Sigma_i^t) dt \end{aligned}$$

where the last equality follows from the coarea formula. By Lemma 3.3 we have  $\int_\Omega |\tilde{u}_\varepsilon - u| dx \leq C_1 \varepsilon$  and we conclude that  $\tilde{u}_\varepsilon \rightarrow u$  in  $L^1(\Omega)$ . If

$\int_\Omega \tilde{u}_\varepsilon = m$  we define  $u_\varepsilon$ . Otherwise we put:

$$\eta_\varepsilon = \int_\Omega \tilde{u}_\varepsilon(x) dx - \int_\Omega u(x) dx$$

Each scalar component of the vector  $\eta_\varepsilon$  is (in absolute value) less than or equal to  $C_1 \varepsilon$ . Let  $x_0$  be a point in the interior of the set  $\Omega \cap S_1$  (if it is empty, the changes in the proof are trivial). If  $\varepsilon$  is small enough, the open ball  $B_\varepsilon = B(x_0, \varepsilon^{1/N})$  is contained in  $\{x : \tilde{u}_\varepsilon(x) = \alpha_1\}$ , by definition of  $\tilde{u}_\varepsilon$ . Define:

$$u_\varepsilon(x) = \begin{cases} \tilde{u}_\varepsilon(x) & \text{if } x \in \Omega \setminus B_\varepsilon \\ \alpha_1 + h_\varepsilon(1 - \varepsilon^{-1/N} |x - x_0|) & \text{if } x \in B_\varepsilon \end{cases}$$

where  $h_\varepsilon = -N \omega_{N-1}^{-1} \eta_\varepsilon \varepsilon^{(1-N)/N}$  and  $\omega_{N-1}$  is the volume of the  $(N-1)$ -dimensional unit ball. The functions  $u_\varepsilon$  satisfy the constraint  $\int u_\varepsilon = m$  for  $\varepsilon$  small enough, and converge in  $L^1(\Omega)$  to  $u$ . It remains to give a sharp estimate of  $\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon)$ . Let us consider the following partition of  $\Omega$ :

- (i)  $\Omega_1^\varepsilon = S_1 \setminus B_\varepsilon$ .  
(ii)  $\Omega_i^\varepsilon = \{x \in S_i : h_j(x) > C_1 \varepsilon, j = 1, \dots, i-1\}$  for  $i = 2, \dots, k$ .  
(iii)  $\Omega_{ij}^\varepsilon = \{x \in \Omega : 0 < h_i(x) < C_1 \varepsilon, h_j(x) < 0, h_l(x) > C_1 \varepsilon \text{ for } l \in \{1, \dots, k-1\} \setminus \{i, j\}\}$

for  $i, j \in \{1, \dots, k\}$  and  $i \neq j$ .

$$(iv) \Omega_0^\varepsilon = \Omega \setminus \left( B_\varepsilon \cup \left( \bigcup_{i=1}^k \Omega_i^\varepsilon \right) \cup \bigcup_{\substack{i, j=1 \\ i \neq j}}^k \Omega_{ij}^\varepsilon \right).$$

If we put:

$$F_\varepsilon(u, \Omega') = \int_{\Omega'} [\varepsilon |Du|^2 + \frac{1}{\varepsilon} W(u)] dx$$

we obviously have:

$$\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) \leq \sum_{i=1}^k \limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, \Omega_i^\varepsilon) + \sum_{i, j} \limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, \Omega_{ij}^\varepsilon) + \limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, B_\varepsilon) + \limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, \Omega_0^\varepsilon). \quad (3.3)$$

The first sum of the right-hand side of (3.3) vanishes on  $\Omega_i^\varepsilon$  because  $u_\varepsilon \equiv \alpha_i$ .

Since:

$$F_\varepsilon(u_\varepsilon, B_\varepsilon) = \varepsilon |h_\varepsilon|^2 \varepsilon^{-2/N} |B_\varepsilon| + \varepsilon^{-1} \int_{B_\varepsilon} W(\alpha_1 + h_\varepsilon(1 - \varepsilon^{-1/N} |x - x_0|)) dx \\ \leq K \varepsilon^2 + \int_0^1 W(\alpha_1 + h_\varepsilon(1-r)) r^{N-1} dr$$

and  $h_\varepsilon \rightarrow 0$  for  $\varepsilon \rightarrow 0$ , we have

$$\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, B_\varepsilon) = 0.$$

Now define:  $K_{ij}^\varepsilon = \{x \in \Omega : 0 < h_i(x) < C_1 \varepsilon, 0 < h_j(x) < C_1 \varepsilon\}$ ,  $i > j$ . Since  $\Omega_0^\varepsilon \subset \bigcup_{ij} K_{ij}^\varepsilon$ , by using Lemma 3.2 we obtain:

$$F_\varepsilon(u_\varepsilon, K_{ij}^\varepsilon) = \int_{K_{ij}^\varepsilon} [\varepsilon |Du_\varepsilon|^2 + \frac{1}{\varepsilon} W(u_\varepsilon)] dx \leq C_5 \varepsilon^{-1} |K_{ij}^\varepsilon|.$$

Putting now  $S'_j = \{x \in \Omega : h_j(x) > t \text{ or } h_j(x) > C_1 \varepsilon\}$  and employing again the coarea formula we have:

$$\begin{aligned} |\mathbf{K}_{ij}^\varepsilon| &= \int_0^{C_1 \varepsilon} \mathbf{H}_{N-1}(\{x \in \Omega : h_i(x) = s, 0 < h_j(x) < C_1 \varepsilon\}) ds \\ &\leq C_1 \varepsilon \sup_{0 \leq s \leq C_1 \varepsilon} \mathbf{H}_{N-1}(\{x \in \Omega : h_i(x) = s, 0 < h_j(x) < C_1 \varepsilon\}) \\ &= C_1 \varepsilon \sup_{0 \leq s \leq C_1 \varepsilon} \mathbf{H}_{N-1}\left(\sum_i^s S_j^{C_1 \varepsilon}\right). \end{aligned}$$

Remark that, for almost all  $\rho > 0$ , we have  $\mathbf{H}_{N-1}(\partial S_i \cap \partial S_j^\rho) = 0$ . Thus, recalling Lemma 3.3, we have:

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0^+} \sup_{0 \leq s \leq C_1 \varepsilon} \mathbf{H}_{N-1}\left(\sum_i^s S_j^{C_1 \varepsilon}\right) \\ \leq \limsup_{\varepsilon \rightarrow 0^+} \sup_{0 \leq s \leq C_1 \varepsilon} \mathbf{H}_{N-1}\left(\sum_i^s S_j^\rho\right) = \mathbf{H}_{N-1}((\partial S_i \cap (\Omega \setminus S_j)) \setminus S_j^\rho). \end{aligned}$$

Passing to the infimum for  $\rho > 0$ , we finally have  $\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, \mathbf{K}_{ij}^\varepsilon) = 0$ ,

and then

$$\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, \Omega_0^\varepsilon) = 0.$$

It remains to estimate the terms of (3.3) of the form  $\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon, \Omega_{ij}^\varepsilon)$ .

We have:

$$\begin{aligned} F_\varepsilon(u_\varepsilon; \Omega_{ij}^\varepsilon) &= \int_{\Omega_{ij}^\varepsilon} \left[ \varepsilon |Du_\varepsilon|^2 + \frac{1}{\varepsilon} W(u_\varepsilon) \right] dx \\ &= \int_{\Omega_{ij}^\varepsilon} \left[ \varepsilon |D(\gamma_{ij} \circ y_\varepsilon^{ij} \circ h_i)(x)|^2 + \frac{1}{\varepsilon} W(\gamma_{ij} \circ y_\varepsilon^{ij} \circ h_i)(x) \right] |Dh_i(x)| dx \end{aligned}$$

(by using the coarea formula)

$$\int_0^{C_1 \varepsilon} \left[ \varepsilon \gamma_{ij}'^2(y_\varepsilon^{ij}(t)) (y_\varepsilon^{ij})'(t)^2 + \frac{1}{\varepsilon} W(\gamma_{ij} \circ y_\varepsilon^{ij})(t) \right] \mathbf{H}_{N-1}\left(\sum_i^t \cap S_j\right) dt$$

(by remarking that  $\sigma_\varepsilon = \sup_{0 \leq t \leq C_1 \varepsilon} \mathbf{H}_{N-1}\left(\sum_i^t \cap S_j\right) \rightarrow \mathbf{H}_{N-1}(\partial S_i \cap \partial S_j)$  as  $\varepsilon \rightarrow 0$ )

$$\leq \sigma_\varepsilon \int_0^{C_1 \varepsilon} \left[ \varepsilon \gamma_{ij}'^2(y_\varepsilon^{ij}(t)) ((y_\varepsilon^{ij})'(t))^2 + \frac{1}{\varepsilon} (\delta + W(\gamma_{ij} \circ y_\varepsilon^{ij})(t)) \right] dt$$



[by using eq. (3.1)]

$$\begin{aligned} \sigma_\varepsilon \int_0^{C_1 \varepsilon} \{ |\gamma'_{ij}(y_\varepsilon^{ij}(t))| |y_\varepsilon^{ij'}(t)| [\delta + W(\gamma_{ij} \circ y_\varepsilon^{ij})(t)]^{1/2} \} dt \\ \leq 2 \sigma_\varepsilon \int_0^1 [\delta + W(\gamma_{ij}(s))]^{1/2} |\gamma'_{ij}(s)| ds. \end{aligned}$$

By collecting all previous inequalities, we conclude that

$$\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) \leq \sum_{i,j=1}^k H_{N-1}(\partial S_i \cap \partial S_j) \int_0^1 [\delta + W(\gamma_{ij}(s))]^{1/2} |\gamma'_{ij}(s)| ds.$$

Recall that the functions  $u_\varepsilon = u_\varepsilon^\delta$  depend on  $\delta$ . Passing now to the infimum for  $\delta > 0$  we obtain:

$$\inf_{\delta > 0} \limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon^\delta) \leq F_0(u_0) \quad (3.4)$$

and, by a simple diagonal argument, (2.9) is proved. The proof is now almost completed, modulo the existence of geodesics connecting any two zeros of  $W$ . If such geodesics do not exist, we could choose approximate geodesics  $\gamma_{ij}^h$  such that

$$\int_0^1 W^{1/2}(\gamma_{ij}^h(s)) |\gamma_{ij}^{h'}(s)| ds \leq d(\alpha_i, \alpha_j) + \frac{1}{h};$$

reasoning as above, we construct a sequence  $u_\varepsilon^h$  such that

$$\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon^h) \leq \sum_{ij} H_{N-1}(\partial S_i \cap \partial S_j) (d(\alpha_i, \alpha_j) + 1/h)$$

and again a diagonal argument completes the proof. ■

#### 4. FINAL REMARKS

In this Section we prove a compactness result in  $L^1(\Omega)$  for the family of minima.

**PROPOSITION 4.1.** — *Let  $\{u_\varepsilon\}_{\varepsilon > 0}$  be a sequence such that, for each  $\varepsilon > 0$ ,  $u_\varepsilon$  is a minimum point of  $F_\varepsilon$ . Let us suppose that there exists a constant  $M > 0$  such that  $u_\varepsilon(x) \leq M$  for almost all  $x \in \Omega$  and for all  $\varepsilon > 0$ . Then there exists a sequence  $\varepsilon_h \downarrow 0$  such that  $\{u_{\varepsilon_h}\}_{h \in \mathbb{N}}$  converges in  $L^1(\Omega)$ .*

*Proof.* — Since the sequences  $\{\varphi_i \circ u_\varepsilon\}_{\varepsilon > 0}$  are bounded in  $L^1(\Omega)$  for every  $i = 1, \dots, k$ , using the results obtained in Section 3 to prove the

inequality (2.8), we have:

$$\int_{\Omega} |D(\varphi_i \circ u_\varepsilon)| \leq \frac{1}{2} F_\varepsilon(u_\varepsilon), \quad \forall \varepsilon > 0, \quad i = 1, \dots, k \quad (4.1)$$

Furthermore, reasoning as for the proof of (2.9), we obtain that  $\{F_\varepsilon(u_\varepsilon)\}$  is equibounded. Thus, by using Rellich's Theorem in  $BV(\Omega)$ , there exist  $k$  functions  $f_1, \dots, f_k$  and a sequence  $\varepsilon_h \downarrow 0$  such that  $\varphi_i \circ u_{\varepsilon_h} \rightarrow f_i$  in  $L^1(\Omega)$  and pointwise a.e., for every  $i = 1, \dots, k$ . We now define:

$$S_i = \{x \in \Omega : f_i(x) = 0\},$$

$$u_0(x) = \sum_{i=1}^k \alpha_i \mathbf{1}_{S_i}(x).$$

Let us turn to the convergence of  $\{u_{\varepsilon_h}\}$ : it is of course sufficient to prove pointwise a.e. convergence. Let  $A = \{x \in \Omega : \limsup_{\varepsilon \rightarrow 0^+} W(u_\varepsilon(x)) > 0\}$ ; the

equiboundedness of  $F_\varepsilon(u_\varepsilon)$  implies that  $|A| = 0$ ; hence, for a.a.  $x \in \Omega$ , the limit points of  $u_{\varepsilon_h}(x)$  are in the set  $\{\alpha_1, \dots, \alpha_k\}$ . On the other hand, for a.a.  $x \in S_i$ , if  $u_{\varepsilon_{h_k}}(x) \rightarrow \alpha_j$ , we have that  $\varphi_i(u_{\varepsilon_{h_k}}(x)) \rightarrow \varphi_i(\alpha_j)$ ,  $\varphi_i(u_{\varepsilon_{h_k}}(x)) \rightarrow f_i(x) = 0$ ; therefore  $\varphi_i(\alpha_j) = 0$  and  $i = j$ . It follows that  $u_{\varepsilon_h}$  converges pointwise to  $u_0$  and Proposition (4.1) is proved. ■

In order to check the equiboundedness-hypothesis of the previous proposition, we recall the following result, which is a straightforward generalization of a theorem by Gurtin and Matano (see [GM] and [LM]).

PROPOSITION 4.2. — Suppose  $W \in C^1(\mathbf{R}_+^n)$ ,

$$W(t) = W_1(t_1) + \dots + W_n(t_n)$$

(see the introduction). Denote  $W'_i(t_i) = \frac{\partial W(t)}{\partial t_i}$ , and suppose that for an index

$i \in \{1, \dots, k\}$  the following properties hold:

- (i)  $W'_i(0) < W'_i(\tau), \forall \tau \in \mathbf{R}^+$ ,
- (ii)  $\lim_{\tau \rightarrow +\infty} W'_i(\tau) = +\infty$ .

Then the family of minima  $\{u_\varepsilon\}$  is bounded in  $L^\infty(\Omega)$ .

### APPENDIX: PROOF OF LEMMA 3.1

For the moment, we ignore part (iii) of the Lemma. Let us recall the following result by T. Quentin de Gromard [QG]:

THEOREM A.1. — Let  $\Omega \subset \mathbf{R}^N$  be a bounded and open set, and let  $E \subset \mathbf{R}^N$  be a set such that  $\mathbf{1}_E \in BV(\Omega)$ . Then there exists a sequence  $E_h$  of sets with

bounded perimeter such that:

- (i)  $|D(\mathbf{1}_{E_h} - \mathbf{1}_E)|(\Omega) \rightarrow 0$  as  $n \rightarrow +\infty$ .
- (ii)  $\Omega \cap \partial E_h$  is contained in a finite union of  $C^1$ -hypersurfaces, for every  $h \in \mathbb{N}$ .
- (iii)  $|(E_h \Delta E) \cap \Omega| \rightarrow 0$ ,  $E_h \subset E + B(0, 1/h)$ ,  $\Omega \setminus E_h \subset (\Omega \setminus E) + B(0, 1/h)$ .
- (iv) If  $E$  contains a ball  $B(x, r)$ , then there exists  $r_0 > 0$  such that  $E_h$  contains  $B(x, r_0)$  for every  $h \in \mathbb{N}$ . The same holds for  $\Omega \setminus E$ .

[(iv) follows from the proof given by T. Quentin de Gromard, even if it is not explicitly stated.]

The idea of the proof of Lemma 3.1 is as follows: we approximate the partition  $S_1, \dots, S_k$  with sets as in Theorem A.1, and then we approximate these sets with polygonal domains.

More precisely, we begin with the proof of the following result:

PROPOSITION A.2. — Let  $S_1, \dots, S_k$  be a partition as in Lemma 3.1. Then there exists a sequence of partitions  $\{S_1^h, \dots, S_k^h\}_{h \in \mathbb{N}}$  such that:

- (i)  $S_i^h$  is a closed set, and  $\partial S_i^h$  is contained in a finite union of  $C^1$ -hypersurfaces.
- (ii)  $\mathbf{1}_{S_i^h} \rightarrow \mathbf{1}_{S_i}$  in  $BV(\Omega)$  with the strong topology as  $h \rightarrow +\infty$ .
- (iii) If  $S_i$  contains  $B(x, r)$ , then there exists  $r_0 > 0$  such that  $S_i^h$  contains  $B(x, r_0)$  for every  $h \in \mathbb{N}$ .

Proof. — For any fixed  $i = 1, \dots, k$  we pick a sequence  $\tilde{S}_i^h$  as in Theorem A.1, and we define:

$$S_1^h = \tilde{S}_1^h$$

$$S_2^h = \tilde{S}_2^h \setminus S_1^h$$

$$\dots$$

$$S_k^h = \tilde{S}_k^h \setminus \left( \bigcup_{i=1}^{k-1} S_i^h \right).$$

This sequence of partitions verifies (i) and (ii) by virtue of the following proposition, while (iii) is trivial to prove. ■

PROPOSITION A.3. — Let  $A, B$  be disjoint sets with bounded perimeter in  $\Omega$ , let  $\{A_h\}$  and  $\{B_h\}$  be sequences such that  $\mathbf{1}_{B_k} \rightarrow \mathbf{1}_B$  and  $\mathbf{1}_{A_k} \rightarrow \mathbf{1}_A$  in  $BV(\Omega)$  with the strong topology. Then we have:

- (i)  $\mathbf{1}_{A_k \cup B_k} \rightarrow \mathbf{1}_{A \cup B}$  in  $BV(\Omega)$ .
- (ii)  $\mathbf{1}_{B_k \setminus A_k} \rightarrow \mathbf{1}_B$  in  $BV(\Omega)$ .

To prove this proposition, we need another lemma. We recall the definition of the  $(N - 1)$ -dimensional Gross measure. If  $S, B$  are borel sets, we define:

$$\delta(S) = \sup \{ L_{N-1}(\pi_a(S)) : a \in S^{N-1} \}$$

$$G_{N-1}(B) = \sup_{\varepsilon > 0} \inf \left\{ \sum_{i=1}^{\infty} \delta(S_i) : B \subset \bigcup_{i=1}^{\infty} S_i, S_i \text{ are borel sets, } \text{diam } S_i < \varepsilon \right\}$$

where  $L_{N-1}$  is the Lebesgue measure on the plane  $\Pi_a = \{ \langle x, a \rangle = 0 \}$  and  $\pi_a$  denotes the orthogonal projection on  $\Pi_a$ . Furthermore, define  $B_* = \{ x \in \Omega : B \text{ has density } 1 \text{ at } x \}$  and  $B^* = \{ x \in \Omega : \Omega \setminus B \text{ has density zero at } x \}$ .

LEMMA A.4. — Define  $A, B, \{A_h\}, \{B_h\}$  as above. Then we have:

$$\lim_{h \rightarrow +\infty} H_{N-1}(\partial^* A \cap B_{h^*}) = 0.$$

*Proof.* — This proof has been communicated to me by L. Ambrosio. The set  $\partial^* A \cap B_{h^*}$  is rectifiable and then, by Theorem 3.2.26 of Federer [F], we have:

$$H_{N-1}(\partial^* A \cap B_{h^*}) = G_{N-1}(\partial^* A \cap B_{h^*}).$$

Define:

$$S = \{ x \in B_* : x \notin B_{h^*} \text{ freq.} \} \cup \{ x \in B^* : x \notin B_h^* \text{ freq.} \} \\ \cup \{ x \in \partial^* B : x \notin \partial^* B_h \text{ freq.} \},$$

where freq. means for infinitely many  $h$ . Assume that  $\delta(S) = 0$ . Then  $G_{N-1}(S) = 0$  and

$$G_{N-1}(\partial^* A \cap B_{h^*}) = \int_{\Omega} \mathbf{1}_{\partial^* A \cap B_{h^*}}(x) dG_{N-1}(x) \\ \leq \int_{B^* \cup \partial^* B} \mathbf{1}_{B_{h^*}}(x) dG_{N-1}(x) \rightarrow 0:$$

the inequality comes from the fact that  $A \cap B = \emptyset$  and  $\partial^* A \subset B^* \cup \partial^* B$  up to a set of null Gross-measure.

We now prove that  $\delta(S) = 0$ . This is obvious if  $N = 1$  because the sets with bounded perimeter in  $\mathbf{R}$  are finite unions of pairwise disjoint intervals, and then the strong convergence in  $BV(\Omega)$  entails that the endpoints of the intervals definitively coincide. If  $N > 1$  we choose  $a \in S^{N-1}$  and we write  $\mathbf{R}^N = \Pi_a \times \mathbf{R}$ .

By the Vol'pert's decomposition of sets with bounded perimeter (see [Vol], th. 1.6) we obtain that, for  $L_{N-1}$  almost every  $x \in \Pi_a$ , the set

$$S_h(x) = \{ t \in \mathbf{R} : (x, t) \in B_* \Delta B_{h^*} \text{ or } (x, t) \in B^* \Delta B_h^* \text{ or } (x, t) \in \partial^* B \Delta \partial^* B_h \}$$

is definitively empty. Thus:

$$L_{N-1}(\pi_a(S)) = L_{N-1}(\{ x \in \Pi_a(B) : \exists t \in \mathbf{R} : t \in S_h(x) \text{ definitively} \}) \\ \leq L_{N-1}(\{ x \in \Pi_a \text{ such that } S_h(x) \neq \emptyset \text{ freq.} \}) = 0$$

As this relation holds for every  $a \in S^{N-1}$ , we conclude that  $\delta(S) = 0$ . ■

The following result is well known (see, for example, Giusti [Gi]).

PROPOSITION A.5. — *Let A, B sets with bounded perimeter. Then we have:*

$$H_{N-1}(\{x \in \partial^* A \cap \partial^* B : v_A(x) \neq \pm v_B(x)\}) = 0.$$

*Proof of Proposition A.3.* — (i) We first prove that  $\mathbf{1}_{A_h \cup B_h} \rightarrow \mathbf{1}_{A \cup B}$  in  $BV(\Omega)$ . It suffices to prove that  $P_\Omega(A_h \cap B_h) \rightarrow 0$ . Let  $G$  be a borel set; by using Lemma 2 of [QG] we obtain the following formula (see also [Vo]):

$$\begin{aligned} D \mathbf{1}_{E \cap F}(C) &= D \mathbf{1}_F(C \cap E_*) + D \mathbf{1}_E(C \cap F_*) \\ &\quad + 1/2 \int_{\partial^* A_h \cap \partial^* B_h \cap C} |v_{A_h}(x) + v_{B_h}(x)| dH_{N-1}(x). \end{aligned}$$

Then:

$$\begin{aligned} (A.1) \quad P_\Omega(A_h \cap B_h) &\leq H_{N-1}(\partial^* A_h \cap B_{h^*}) + H_{N-1}(\partial^* B_h \cap A_{h^*}) \\ &\quad + 1/2 \int_{\partial^* A_h \cap \partial^* B_h} |v_{A_h}(x) + v_{B_h}(x)| dH_{N-1}(x). \end{aligned}$$

By using again Lemma 2 of [QG], we obtain:

$$\begin{aligned} H_{N-1}(\partial^* A_h \Delta \partial^* A) &= H_{N-1}(\partial^* A \cap (\Omega \setminus \partial^* A_h)) \\ &\quad + H_{N-1}(\partial^* A_h \cap (\Omega \setminus \partial^* A)) \\ &\leq |D \mathbf{1}_A|(\Omega \setminus \partial^* A_h) + |D \mathbf{1}_A|(\Omega \setminus \partial^* A) \\ &= |D \mathbf{1}_{A_h \Delta A}|(\Omega) \leq \int_\Omega |D(\mathbf{1}_{A_h} - \mathbf{1}_A)|; \end{aligned}$$

hence

$$\lim_{h \rightarrow +\infty} H_{N-1}(\partial^* A_h \Delta \partial^* A) = 0.$$

Observing that

$$\partial^* A_h \cap B_{h^*} = ((\partial^* A_h \setminus \partial^* A) \cap B_{h^*}) \cup ((\partial^* A_h \cap \partial^* A) \cap B_{h^*})$$

we infer by Lemma A.4 that

$$\lim_{h \rightarrow +\infty} H_{N-1}(\partial^* A_h \cap B_{h^*}) = 0$$

and analogously:

$$\lim_{h \rightarrow +\infty} H_{N-1}(\partial^* B_h \cap A_{h^*}) = 0.$$

Finally, we have by Proposition A.5

$$\lim_{h \rightarrow +\infty} \int_{\partial^* A_h \cap \partial^* B_h \cap \partial^* A \cap \partial^* B} |v_{A_h}(x) - v_{B_h}(x)| dH_{N-1}(x) = 0$$

and also

$$\lim_{h \rightarrow +\infty} H_{N-1}((\partial^* A_h \cap \partial^* B_h) \Delta (\partial^* A \cap \partial^* B)) = 0$$

because of

$$\begin{aligned} (\partial^* A_h \cap \partial^* B_h) \setminus (\partial^* A \cap \partial^* B) &= ((\partial^* A_h \setminus \partial^* A) \cap \partial^* B_h) \cup ((\partial^* B_h \setminus \partial^* B) \cap \partial^* A_h) \\ (\partial^* A \cap \partial^* B) \setminus (\partial^* A_h \cap \partial^* B_h) &= ((\partial^* A \setminus \partial^* B_h) \cap \partial^* B) \cup ((\partial^* B \setminus \partial^* B_h) \cap \partial^* A) \end{aligned}$$

(ii) It follows from

$$\mathbf{1}_{B_k \setminus A_h} = \mathbf{1}_{B_h} - \mathbf{1}_{B_h \cap A_h}. \quad \blacksquare$$

We have now to construct an approximation of a  $C^1$  partition ( $C^1$ -regular up to a set with null  $H_{N-1}$  measure) by polygonal domains. We have a partition of  $\Omega$ , say  $S_1, \dots, S_k$ , made by sets with boundaries contained in a finite union of  $C^1$ -hypersurfaces, and we look for a sequence  $\{S_1^h, \dots, S_k^h\}_{h \in \mathbb{N}}$  of partitions such that the  $S_i^h$ 's are polygonal domains and such that the property (iv) of Lemma 3.1 holds. By the Fleming-Rishel formula, this will be proved if we verify the following property:

$$\forall J \subset \{1, \dots, k\}, \lim_{h \rightarrow +\infty} P_\Omega(\bigcup_{i \in J} S_i^h) = P_\Omega(\bigcup_{i \in J} S_i). \quad (A.2)$$

The first step consists in eliminating the singularities of the boundaries of the  $S_i$ 's by including them in a family of cubes with little volume and perimeter, and by adding all these cubes to the set  $S_1$ .

Let  $B$  be the singular set of the given partition. It is possible to construct, for every  $h \in \mathbb{N}$ , a finite family of cubes  $\{Q_1^h, \dots, Q_{v(h)}^h\}$  such that:

(i)  $B \subset \bigcup_{i=1}^{v(h)} Q_i^h$ .

(ii)  $P_\Omega\left(\bigcup_{i=1}^{v(h)} Q_i^h\right) < 2N/h, \left|\bigcup_{i=1}^{v(h)} Q_i^h\right| < 1/h$ .

(iii)  $\partial Q_i$  is transverse to  $\partial S_j, \forall i = 1, \dots, v(h); j = 1, \dots, k$ .

For the sequence of partitions given by the sets  $S_1 \cup \bigcup Q_j^h; S_2 \setminus \bigcup Q_j^h, \dots; S_k \setminus \bigcup Q_j^h$ , (A.2) holds. The boundaries of these sets are piecewise- $C^1$  hypersurfaces, whose singular sets are transverse intersections of regular surfaces with cubes. We now approximate the portions of the interfaces not belonging to the cubes by polygons, in such a way that (A.2) holds. Let  $\Sigma$  be one of these hypersurfaces, closed in the open set  $\Omega' = \Omega \setminus \bigcup_i Q_i$ .  $\Sigma$  is an oriented hypersurface transverse to each one of the

$Q_i$ 's. Putting

$$\Omega_p = ((\Sigma + B(0, \rho)) \setminus \bigcup_i Q_i) \cap \Omega$$

we have that, if  $\rho$  is small enough,  $\Sigma$  gives a partition of  $\Omega_\rho$  in two parts, each of them belonging to one of the sets  $S_j$ . Let  $E$  be one of these portions:  $E$  is an open set with  $C^1$  boundary in  $\Omega_\rho$ .

LEMMA A.7. — *There exists a sequence  $E_h$ , with  $\partial E_h$  a piecewise-linear hypersurface, such that:*

- (i)  $\lim_{h \rightarrow +\infty} H_{N-1}(\partial E_h \cap \Omega_{\rho/2}) = H_{N-1}(\Sigma)$ .
- (ii)  $\int_{\Omega_{\rho/2}} |\mathbf{1}_{E_h} - \mathbf{1}_E| dx \rightarrow 0$ .
- (iii) *The set  $\partial\Omega_{\rho/2} \setminus \partial\Omega'$  has a neighborhood in  $\Omega_{\rho/2}$  on which  $\mathbf{1}_E \equiv \mathbf{1}_{E_h}$ ,  $\forall h \in \mathbf{N}$ .*
- (iv)  $H_{N-1}(\partial E_h \cap \partial\Omega') = 0, \forall h \in \mathbf{N}$ .

*Proof.* — Let us call  $\tilde{f}$  an extension of  $\mathbf{1}_E|_{\Omega_{\rho/2}}$  to an open neighborhood  $\tilde{\Omega} \supset \bar{\Omega}_{\rho/2}$ , such that  $\int_{\partial\Omega_{\rho/2}} |D\tilde{f}| = 0$ . This extension is possible because  $\partial\Omega_{\rho/2} \cap \partial\Omega'$  is lipschitz continuous. Let  $\tilde{f}_\varepsilon$  be a sequence converging to  $\tilde{f}$  in  $L^1(\tilde{\Omega})$  such that  $\lim_{\varepsilon \rightarrow 0^+} \int_{\Omega_{\rho/2}} |D\tilde{f}_\varepsilon| = \int_{\Omega_{\rho/2}} |D\mathbf{1}_E| = H_{N-1}(\Sigma)$ . It is not restrictive to assume that  $f_\varepsilon \equiv \mathbf{1}_E$  on a neighborhood of  $\partial\Omega_{\rho/2} \setminus \partial\Omega'$  in  $\Omega_{\rho/2}$  and that  $0 \leq \tilde{f}_\varepsilon \leq 1$  for every  $x \in \tilde{\Omega}$ . Let  $f_h = \tilde{f}_{1/h}$ .

There exists a sequence of piecewise linear function  $g_h : \tilde{\Omega} \rightarrow \mathbf{R}$  such that:

$$\int_{\Omega_{\rho/2}} |g_h(x) - f_h(x)| < 1/h \tag{A.3}$$

$$\left| \int_{\Omega_{\rho/2}} |Dg_h(x)| dx - \int_{\Omega_{\rho/2}} |Df_h(x)| dx \right| < 1/h$$

and such that  $g_h \equiv \mathbf{1}_E$  on a neighborhood of  $\partial\Omega_{\rho/2} \setminus \partial\Omega'$ .

It follows that:

$$\left| \int_{\Omega_{\rho/2}} |Dg_h| - H_{N-1}(\Sigma) \right| < 2/h. \tag{A.4}$$

Let  $t \in [0, 1]$ , and put:  $E_{th} = \{x \in \tilde{\Omega} : g_h(x) > t\}$ . We remark that  $E_{th}$  is a polygonal domain for almost all  $t \in [0, 1]$ . We have:

$$g_h(x) - \mathbf{1}_E(x) > t \text{ on } E_{th} \setminus E$$

$$\mathbf{1}_E - g_h \geq 1 - t \text{ of } E \setminus E_{th}$$

Thus:

$$\int_{\Omega_{\rho/2}} |g_h - \mathbf{1}_E| \geq \int_{E_{th} \setminus E} |g_h - \mathbf{1}_E| + \int_{E \setminus E_{th}} |g_h - \mathbf{1}_E|$$

$$\geq t |E_{th} \setminus E| + (1-t) |E \setminus E_{th}| \geq \min(t, 1-t) \int_{\Omega_{\rho/2}} |\mathbf{1}_{E_{th}} - \mathbf{1}_E|$$

and so  $\mathbf{1}_{E_{th}} \rightarrow \mathbf{1}_E$  in  $L^1(\Omega_{\rho/2})$ . By the coarea formula:

$$\int_{\Omega_{\rho/2}} |Dg_h| = \int_0^1 P_{\Omega_{\rho/2}}(E_{th}) dt. \tag{A.5}$$

By the lower semicontinuity of the total variation:

$$\liminf_{h \rightarrow +\infty} \int_{\Omega_{\rho/2}} |D\mathbf{1}_{E_{th}}| \geq \int_{\Omega_{\rho/2}} |D\mathbf{1}_E|, \quad \forall t \in [0, 1] \tag{A.6}$$

By the Fatou's Lemma and (A.6) we have:

$$\int_{\Omega_{\rho/2}} |D\mathbf{1}_E| = \lim_{h \rightarrow +\infty} \int_{\Omega_{\rho/2}} |Dg_h| \geq \int_0^1 \liminf_{h \rightarrow +\infty} P_{\Omega_{\rho/2}}(E_{th}) dt \geq \int_{\Omega_{\rho/2}} |D\mathbf{1}_E|$$

Then, by using (A.5), we have that for almost all  $t \in [0, 1]$  the following formula holds:

$$\liminf_{h \rightarrow +\infty} \int_{\Omega_{\rho/2}} |D\mathbf{1}_{E_{th}}| = \int_{\Omega_{\rho/2}} |D\mathbf{1}_E| = H_{N-1}(\Sigma) \tag{A.7}$$

By using that  $H_{N-1}(\partial\Omega') < +\infty$ , we have that, for almost all  $t \in [0, 1]$ ,  $H_{N-1}(\partial E_{ht} \cap \partial\Omega') = 0$ . Thus, by choosing the appropriate  $t$  and passing to a subsequence in (A.7), we conclude the proof. ■

It remains to fulfil the condition (iii) of Lemma 3.1. By repeating the argument used by Modica [M1], and by remarking that the property of having non-empty interior is preserved by our construction of approximating partitions, we have only to indicate how to modify the original partition in such a way that each non-empty set  $S_i$  of the partition has non-empty interior. Take a point  $x \in \Omega \cap \partial^* S_1 \cap \partial^* S_2$ : the density of both  $S_1$  and  $S_2$  at  $x$  is exactly 1/2, and then, if  $\rho_h \downarrow 0$ , we have that  $0 < |B_{\rho_h}(x) \cap S_2| < |B_{\rho_h}(x)|$  for every  $h \in \mathbb{N}$ . Then, for each  $h \in \mathbb{N}$ , there exists  $\tilde{\rho}_h$  such that  $|B_{\tilde{\rho}_h}(x)| = |B_{\rho_h}(x) \setminus S_1|$ . We can now divide the ball  $B_{\rho_h}(x)$  into  $(k-1)$  sectors  $\tilde{B}_{hl}$  with area  $|B_{\tilde{\rho}_h} \cap S_l|$ ,  $l \neq 1$ . Since obviously  $H_{N-1}(\partial \tilde{B}_{hl}) \leq K \tilde{\rho}_h$ , where the constant  $K$  depends only on  $N$ , it follows that the perimeter of the (nonempty) sets  $\tilde{B}_{hl}$  tends to zero. For the following sequence of partitions:

$$S_1^h = S_1 \cup (B_{\rho_h}(x) \setminus B_{\tilde{\rho}_h}(x))$$

$$S_l^h = (S_l \setminus B_{\rho_h}(x)) \cup \tilde{B}_{hl}, \quad l \neq 1$$



one has  $|S_m^h| = |S_m|$  for each  $m = 1, \dots, k$ , and for each  $J \subset \{1, \dots, k\}$ ,

$$\lim_{h \rightarrow +\infty} P_\Omega(\cup_{l \in J} S_l^h) = P_\Omega(\cup_{l \in J} S_l).$$

Moreover the sets  $S_1^h$  and  $S_2^h$  have nonempty interior. Repeating the same construction for the other elements of the partition, the proof is complete. ■

## REFERENCES

- [A] L. AMBROSIO (to appear).
- [CH] J. W. CAHN and J. E. HILLIARD, Free Energy of a Nonuniform System: I. Interfacial Energy, *J. Chem. Ph.*, Vol. **28**, 1958, pp. 258-267.
- [DGF] E. DE GIORGI and T. FRANZONI, Su un tipo di convergenza variazionale, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Mat. Fis. Natur.*, Vol. **58**, 1975, pp. 842-850.
- [F] H. FEDERER, *Geometric Measure Theory*, Springer Verlag, Berlin 1968.
- [FT] I. FONSECA and L. TARTAR, *The Gradient Theory of Phase Transitions for Systems with two Potential Wells*, Preprint Carnegic-Mellon Univ., Pittsburgh, 1988.
- [Gi] E. GIUSTI, *Minimal Surfaces and Functions of Bounded Variation*, Birkhauser Verlag, 1984.
- [GM] M. E. GURTIN and H. MATANO, On the Structure of Equilibrium Phase Transitions within the Gradient Theory of Fluids, *Quarterly of Appl. Math.*, Vol. **46**, 1988.
- [LM] S. LUCKAUS and L. MODICA, *The Gibbs-Thompson Relation within the Gradient Theory of Phase Transitions* (to appear).
- [M1] L. MODICA, The Gradient Theory of Phase Transitions and the Minimal Interface Criterion, *Arch. Rat. Mech. & Analysis*, **98**, 1987, pp. 123-142.
- [M2] L. MODICA, Gradient Theory of Phase Transitions with Boundary Contact Energy, *Ann. Inst. H. Poincaré. Anal. nonlin.*, Vol. **4**, 1987, pp. 487-512.
- [QG] T. Q. DE GROMARD, Approximation forte dans  $BV(\Omega)$ , *C. R. Acad. Sci. Paris*, T. **301**, 1985, pp. 261-264.
- [S] P. STERNBERG, The Effect of a Singular Perturbation on Nonconvex Variational Problems, *Arch. Rat. Mech. & Analysis*, Vol. **101**, 1988, pp. 209-260.
- [Vo] A. I. VOLPERT, The Spaces  $BV$  and Quasilinear Equations, *Mat. Sbornik.*, Vol. **73**, 1967, pp. 225-254.
- [VW] J. D. VAN DER WAALS, The Thermodynamic Theory of Capillarity under the Hypothesis of a Continuous Variation of Density, *Verhaendel. Kronik. Akad. Wetten. Amsterdam*, Vol. **1**, 1893.

(Manuscript received August 1st, 1988.)