

ANNALES SCIENTIFIQUES
DE L'UNIVERSITÉ DE CLERMONT-FERRAND 2
Série Mathématiques

MIRCEA SOFONEA

**Some remarks concerning a class of nonlinear evolution
equations in Hilbert spaces**

*Annales scientifiques de l'Université de Clermont-Ferrand 2, tome 95, série Mathéma-
tiques, n° 26 (1990), p. 13-20*

<http://www.numdam.org/item?id=ASCFM_1990__95_26_13_0>

© Université de Clermont-Ferrand 2, 1990, tous droits réservés.

L'accès aux archives de la revue « Annales scientifiques de l'Université de Clermont-Ferrand 2 » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

SOME REMARKS CONCERNING A CLASS OF NONLINEAR
EVOLUTION EQUATIONS IN HILBERT SPACES

Mircea Sofonea

Department of Mathematics, INCREST, Bucharest, Romania

1. Introduction

Let H be a real Hilbert space and let X, Y be two orthogonal subspaces of H such that $H = X \oplus Y$. Let Λ be a real normed space and let $T > 0$. In this paper we consider evolution problems of the form

$$(1.1) \quad \dot{y}(t) = F(\lambda(t), x(t), y(t), \dot{x}(t)) \quad \text{for all } t \in [0, T],$$

$$(1.2) \quad x(0) = x_0, \quad y(0) = y_0$$

in which the unknowns are the functions $x : [0, T] \rightarrow X$ and $y : [0, T] \rightarrow Y$, $F : \Lambda \times X \times Y \times H \rightarrow H$ is a nonlinear operator and $\lambda : [0, T] \rightarrow \Lambda$ is a parameter function (in (1.1) and everywhere in this paper the dot above represents the derivative with respect to the time variable t). Such type of problems arise in the study of quasistatic processes for semilinear rate-type materials (see for example [1] - [3]). In this case the unknowns x and y are the small deformation tensor and the stress tensor and F is an operator involving the constitutive law of the material; the parameter λ may be interpreted as the absolute temperature or an internal state variable.

For particular forms of F existence and uniqueness of the solution and error estimates of a numerical method for problems of the form (1.1), (1.2) were already given in [3], [4].

In this paper we prove the existence and uniqueness of the solution for problem (1.1), (1.2) using a technique based on the equivalence between (1.1), (1.2) and a Cauchy problem for an ordinary differential equation in the product Hilbert space $X \times Y$ (section 2). We also study the dependence of the solution with respect to the parameter λ and the initial data (section 3). In some applications (see

for example [5]) the function λ in (1.1) is needed to be considered as an unknown function whose evolution is given by

$$(1.3) \quad \dot{\lambda}(t) = G(\lambda(t), x(t), y(t)) \quad \text{for all } t \in [0, T] ,$$

$$(1.4) \quad \lambda(0) = \lambda_0$$

where $G : \Lambda \times X \times Y \rightarrow \Lambda$ is a nonlinear operator. For this reason we also consider problem (1.1)-(1.4) for which we prove the existence and uniqueness of the solution (section 4). Let us finally notice that the results presented here complete and generalize some results of [2] and may be applied in the study of some evolution problems for rate-type materials (see [1] - [5]).

2. An existence and uniqueness result

Everywhere in this paper if V is a real normed space we utilise the following notations : $\|\cdot\|_V$ - the norm of V ; 0_V - the zero element of V ; $C^0([0, T], V)$ - the space of continuous functions on $[0, T]$ with values in V ; $C^1([0, T], V)$ - the space of derivable functions with continuous derivative on $[0, T]$ with values in V ; $\|\cdot\|_{0, T, V}$ - the norm on the space $C^0([0, T], V)$ i.e. $\|z\|_{0, T, V} = \max_{t \in [0, T]} \|z(t)\|_V$ for all $z \in C^0([0, T], V)$; $\|\cdot\|_{1, T, V}$ - the norm on the space $C^1([0, T], V)$ i.e. $\|z\|_{1, T, V} = \|z\|_{0, T, V} + \|\dot{z}\|_{0, T, V}$ for all $z \in C^1([0, T], V)$. If moreover V is a real Hilbert space we denote by $\langle \cdot, \cdot \rangle_V$ the inner product of V . Finally, if V_1 and V_2 are real Hilbert spaces we denote by $V_1 \times V_2$ the product space endowed with the canonical inner product and by $v = (v_1, v_2)$ the elements of $V_1 \times V_2$.

Let us consider the following assumptions :

$$(2.1) \quad \begin{aligned} & \text{there exists } m > 0 \text{ such that } \langle F(\lambda, x, y, z_1) - F(\lambda, x, y, z_2), z_1 - z_2 \rangle_H \geq \\ & \geq m \|z_1 - z_2\|_H^2 \quad \text{for all } \lambda \in \Lambda, x \in X, y \in Y, z_1, z_2 \in X \end{aligned}$$

$$(2.2) \quad \begin{aligned} & \text{there exists } M > 0 \text{ such that } \|F(\lambda, x_1, y_1, z_1) - F(\lambda, x_2, y_2, z_2)\|_H \leq \\ & \leq M (\|x_1 - x_2\|_H + \|y_1 - y_2\|_H + \|z_1 - z_2\|_H) \quad \text{for all } \lambda \in \Lambda, x_i \in X, y_i \in Y, \\ & z_i \in H, i=1, 2; \end{aligned}$$

(2.3) $\lambda \rightarrow F(\lambda, x, y, z) : \Lambda \rightarrow H$ is an continuous operator , for all $x \in X$, $y \in Y$ and $z \in H$.

(2.4) $\lambda \in C^0(0, T, \Lambda)$

(2.5) $x_0 \in X$, $y_0 \in Y$.

The main result of this section is the following :

Theorem 2.1. Let (2.1)-(2.5) hold. Then problem (1.1), (1.2) has a unique solution $x \in C^1(0, T, X)$, $y \in C^1(0, T, Y)$.

In order to prove theorem 2.1 let us denote by Z the product Hilbert space $Z = X \times Y$ (which in fact is isomorph with H). We have :

Lemma 2.1. Let $\lambda \in \Lambda$, $x \in X$ and $y \in Y$: then there exists a unique element $z = (u, v) \in Z$ such that $v = F(\lambda, x, y, u)$.

Proof. The uniqueness part is a consequence of (2.1) ; indeed, if the elements $z = (u, v)$, $\tilde{z} = (\tilde{u}, \tilde{v}) \in Z$ are such that $v = F(\lambda, x, y, u)$, $\tilde{v} = F(\lambda, x, y, \tilde{u})$, using (2.1) we have $\langle v - \tilde{v}, u - \tilde{u} \rangle_H \geq m \|u - \tilde{u}\|_H^2$ hence by the orthogonality in H of $v - \tilde{v}$ and $u - \tilde{u}$ we deduce $u = \tilde{u}$ which implies $v = \tilde{v}$.

For the existence part let us denote by $P_1 : H \rightarrow X$ the projector map on X . Using (2.1) and (2.2) we get that the operator $P_1 F(\lambda, x, y, \cdot) : X \rightarrow X$ is a strongly monotone and Lipschitz continuous operator hence by Browder's surjectivity theorem we get that there exists $u \in X$ such that $P_1 F(\lambda, x, y, u) = 0_X$. It results that the element $F(\lambda, x, y, u)$ belongs to Y and we finish the proof taking $z = (u, v)$ where $v = F(\lambda, x, y, u)$.

Lemma 2.1 allows us to consider the operator $B : \Lambda \times Z \rightarrow Z$ defined by

(2.6) $B(\lambda, w) = z \quad \text{iff} \quad w = (x, y) , z = (u, v) \quad \text{and} \quad v = F(\lambda, x, y, u)$.

Moreover, we have :

Lemma 2.2. B is a continuous operator and there exists $L > 0$ such that

(2.7) $\|B(\lambda, w_1) - B(\lambda, w_2)\|_Z \leq L \|w_1 - w_2\|_Z \quad \text{for all } \lambda \in \Lambda , w_1, w_2 \in Z$.

Proof. Let $\lambda_i \in \Lambda$, $w_i = (x_i, y_i) \in Z$ and $z_i = (u_i, v_i) = B(\lambda_i, w_i)$, $i = 1, 2$. Using (2.6) we get :

$$(2.8) \quad v_i = F(\lambda_i, x_i, y_i, u_i), \quad i = 1, 2$$

which implies

$$(2.9) \quad P_1 F(\lambda_i, x_i, y_i, u_i) = 0_X, \quad i = 1, 2$$

From (2.1) and (2.9) we get

$$\begin{aligned} m \|u_1 - u_2\|_H^2 &\leq \langle F(\lambda_1, x_1, y_1, u_1) - F(\lambda_1, x_1, y_1, u_2), u_1 - u_2 \rangle_H = \\ &= \langle P_1 F(\lambda_2, x_2, y_2, u_2) - P_1 F(\lambda_1, x_1, y_1, u_2), u_1 - u_2 \rangle_H \leq \|F(\lambda_2, x_2, y_2, u_2) - F(\lambda_1, x_1, y_1, u_2)\|_H \|u_1 - u_2\|_H \end{aligned}$$

which implies

$$(2.10) \quad \|u_1 - u_2\|_H \leq \frac{1}{m} \|F(\lambda_1, x_1, y_1, u_2) - F(\lambda_2, x_2, y_2, u_2)\|_H.$$

Using now (2.8) and (2.2) we get

$$\|v_1 - v_2\|_H \leq M \|u_1 - u_2\|_H + \|F(\lambda_1, x_1, y_1, u_2) - F(\lambda_2, x_2, y_2, u_2)\|_H$$

hence by (2.10) it results

$$(2.11) \quad \|v_1 - v_2\|_H \leq (\frac{M}{m} + 1) \|F(\lambda_1, x_1, y_1, u_2) - F(\lambda_2, x_2, y_2, u_2)\|_H.$$

Using again (2.2) we get

$$(2.12) \quad \begin{aligned} \|F(\lambda_1, x_1, y_1, u_2) - F(\lambda_2, x_2, y_2, u_2)\|_H &\leq M (\|x_1 - x_2\|_H + \|y_1 - y_2\|_H) + \\ &+ \|F(\lambda_1, x_2, y_2, u_2) - F(\lambda_2, x_2, y_2, u_2)\|_H \end{aligned}$$

hence by (2.3) we obtain $\|F(\lambda_1, x_1, y_1, u_2) - F(\lambda_2, x_2, y_2, u_2)\|_H \rightarrow 0$ when $\lambda_1 \rightarrow \lambda_2$ in Λ , $x_1 \rightarrow x_2$ in X and $y_1 \rightarrow y_2$ in Y . Using now (2.10) and (2.11) we get the continuity of B and taking $\lambda_1 = \lambda_2$ from (2.10)-(2.12) we get (2.7).

Some remarks concerning in Hilbert spaces

Proof of theorem 2.1. Let $A : [0, T] \times Z \rightarrow Z$ and z_0 be defined by

$$(2.13) \quad A(t, z) = B(\lambda(t), z) \quad \text{for all } t \in [0, T] \text{ and } z \in Z$$

$$(2.14) \quad z_0 = (x_0, y_0) .$$

Using (2.6) we get that $x \in C^1(0, T, X)$ and $y \in C^1(0, T, Y)$ is a solution of (1.1), (1.2) iff $z = (x, y) \in C^1(0, T, Z)$ is a solution of the problem

$$(2.15) \quad \dot{z}(t) = A(t, z(t)) \quad \text{for all } t \in [0, T]$$

$$(2.16) \quad z(0) = z_0 .$$

In order to study (2.15), (2.16) let us remark that by lemma 2.2 and (2.4) we get that A is a continuous operator and

$$\|A(t, z_1) - A(t, z_2)\|_Z \leq L \|z_1 - z_2\|_Z \quad \text{for all } t \in [0, T] \text{ and } z_1, z_2 \in Z .$$

Moreover, by (2.5), (2.14) we get $z_0 \in Z$. Theorem 2.1 follows now from the classical Cauchy-Lipschitz theorem applied to (2.15), (2.16).

3. The continuous dependence of the solution with respect to the data

Let us now replace (2.2), (2.3) by a stronger assumption namely

$$(3.1) \quad \begin{aligned} & \text{there exists } M > 0 \text{ such that } \|F(\lambda_1, x_1, y_1, z_1) - F(\lambda_2, x_2, y_2, z_2)\|_H \leq \\ & \leq M (\|\lambda_1 - \lambda_2\|_{\Lambda} + \|x_1 - x_2\|_H + \|y_1 - y_2\|_H + \|z_1 - z_2\|_H) \text{ for all } \lambda_i \in \Lambda, x_i \in X, \\ & y_i \in Y, z_i \in H, i = 1, 2 . \end{aligned}$$

We have the following result :

Theorem 3.1. Let (2.1), (3.1) hold and let $x_i \in C^1(0, T, X)$, $y_i \in C^1(0, T, Y)$ be the solution of (1.1), (1.2) for the data λ_i , x_{0i} , y_{0i} satisfying (2.4), (2.5), $i = 1, 2$. Then there exists $C > 0$ such that

$$(3.2) \quad \|x_1 - x_2\|_{1, T, H} + \|y_1 - y_2\|_{1, T, H} \leq C (\|\lambda_1 - \lambda_2\|_{0, T, \Lambda} + \|x_{01} - x_{02}\|_H + \|y_{01} - y_{02}\|_H).$$

Remark 3.1. In (3.2) and everywhere in this section C are strictly positive generic constants which depend only on F and T .

Proof of theorem 3.1. Let $z_i = (x_i, y_i)$ and $z_{0i} = (x_{0i}, y_{0i})$, $i = 1, 2$. As it results from the proof of theorem 2.1 we have

$$(3.3) \quad \dot{z}_i(t) = A_i(t, z_i(t)) \quad \text{for all } t \in [0, T]$$

$$(3.4) \quad z_i(0) = z_{0i}$$

where the operators A_i are defined by (2.13) replacing λ by λ_i , $i=1,2$. Since (3.1) implies that $B : \Lambda \times Z \rightarrow Z$ is a Lipschitz continuous operator (see the proof of lemma 2.2), from (2.13) we get that there exists $L > 0$ such that

$$(3.5) \quad \|A_1(t, z_1(t)) - A_2(t, z_2(t))\|_Z \leq L(\|\lambda_1(t) - \lambda_2(t)\|_\Lambda + \|z_1(t) - z_2(t)\|_Z) \quad \text{for all } t \in [0, T]$$

Using now (3.3) and (3.5) we get

$$\langle \dot{z}_1(t) - \dot{z}_2(t), z_1(t) - z_2(t) \rangle_Z \leq L(\|\lambda_1(t) - \lambda_2(t)\|_\Lambda + \|z_1(t) - z_2(t)\|_Z) \|z_1(t) - z_2(t)\|_Z$$

for all $t \in [0, T]$ hence by (3.4) and a Gronwall-type lemma we deduce

$$\|z_1(s) - z_2(s)\|_Z \leq C \left(\int_0^s \|\lambda_1(t) - \lambda_2(t)\|_\Lambda dt + \|z_{01} - z_{02}\|_Z \right) \quad \text{for all } s \in [0, T]$$

which implies

$$(3.6) \quad \|z_1 - z_2\|_{0,T,Z} \leq C (\|\lambda_1 - \lambda_2\|_{0,T,\Lambda} + \|z_{01} - z_{02}\|_Z) .$$

Using again (3.3) and (3.5) we have

$$\|\dot{z}_1(t) - \dot{z}_2(t)\|_Z \leq C(\|\lambda_1(t) - \lambda_2(t)\|_\Lambda + \|z_1(t) - z_2(t)\|_Z) \quad \text{for all } t \in [0, T]$$

and by (3.6) it results

$$(3.7) \quad \|\dot{z}_1 - \dot{z}_2\|_{0,T,Z} \leq C (\|\lambda_1 - \lambda_2\|_{0,T,\Lambda} + \|z_{01} - z_{02}\|_Z) .$$

From (3.6) and (3.7) we get

$$\|z_1 - z_2\|_{0,T,Z} \leq C (\|\lambda_1 - \lambda_2\|_{0,T,\Lambda} + \|z_{01} - z_{02}\|_Z)$$

which implies (3.2).

Remark 3.2. From (3.2) we deduce in particular the continuous dependence of the

solution with respect the initial data i.e. the finite-time stability of every solution of (1.1),(1.2) (for definitions in the field see for instance [6] chap.5).

4. A second existence and uniqueness result

In this section we suppose that Λ is a real Hilbert space. We consider the operator $G : \Lambda \times X \times Y \rightarrow \Lambda$ and the element λ_0 such that

$$(4.1) \quad ||G(\lambda_1, x_1, y_1) - G(\lambda_2, x_2, y_2)||_{\Lambda} \leq L(||\lambda_1 - \lambda_2||_{\Lambda} + ||x_1 - x_2||_H + ||y_1 - y_2||_H) \\ \text{for all } \lambda_i \in \Lambda, x_i \in X, y_i \in Y, i = 1, 2 \quad (L > 0)$$

$$(4.2) \quad \lambda_0 \in \Lambda.$$

We have the following existence and uniqueness result :

Theorem 4.1. Let (2.1), (2.5), (3.1), (4.1), (4.2) hold. Then problem (1.1)-(1.4) has a unique solution $x \in C^1(0, T, X)$, $y \in C^1(0, T, Y)$, $\lambda \in C^1(0, T, \Lambda)$.

Proof. Let us consider the product Hilbert spaces $H = H \times \Lambda$, $X = X \times \{\lambda_0\}$, $Y = Y \times \Lambda$ and let $F : X \times Y \times H \rightarrow H$ be the operator defined by

$$(4.3) \quad F(x, y, z) = (F(\lambda, x, y, z), G(\lambda, x, y)) \quad \text{for all } x = (x, \lambda_0) \in X, y = (y, \lambda) \in Y, \\ z = (z, \mu) \in H.$$

Let us also denote

$$(4.4) \quad x_0 = (x_0, \lambda_0), \quad y_0 = (y_0, \lambda_0).$$

From (2.1), (3.1) and (4.1) we deduce

$$(4.5) \quad \langle F(x, y, z_1) - F(x, y, z_2), z \rangle_H \geq m ||z_1 - z_2||_H^2 \quad \text{for all } x \in X, y \in Y, z_1, z_2 \in H$$

$$(4.6) \quad ||F(x_1, y_1, z_1) - F(x_2, y_2, z_2)||_H \leq L(||x_1 - x_2||_H + ||y_1 - y_2||_H + ||z_1 - z_2||_H) \\ \text{for all } x_1, x_2 \in X, y_1, y_2 \in Y, z_1, z_2 \in H \quad (L > 0)$$

and from (4.4), (2.5), (4.2) we obtain

$$(4.7) \quad x_0 \in X, \quad y_0 \in Y.$$

Since (4.5)-(4.7) are fulfilled we may apply theorem 2.1 and we obtain the existence and the uniqueness of $x = (x, \theta_\lambda) \in C^1(0, T, X)$, $y = (y, \lambda) \in C^1(0, T, Y)$ such that

$$(4.8) \quad \dot{y}(t) = F(x(t), y(t), \dot{x}(t)) \quad \text{for all } t \in [0, T]$$

$$(4.9) \quad x(0) = x_0, \quad y(0) = y_0.$$

Theorem 4.1 follows now from (4.3) and (4.4).

Remark 4.1. As in the case of the problem (1.1), (1.2), applying theorem 3.1 to (4.8), (4.9) we deduce the finite-time stability of every solution of (1.1)-(1.4).

References

- [1] I.R.Ionescu and M.Sofonea , Quasistatic processes for elastic-viscoplastic materials , Quart. Appl. Math., 2(1988) ,p. 229-243.
- [2] M.Sofonea , Functional methods in thermo-elasto-viscoplasticity , Ph.D.Thesis , Univ. of Bucharest , 1988 (in romanian).
- [3] I.R.Ionescu , Functional and numerical methods in viscoplasticity , Ph.D.Thesis, Univ. of Bucharest , 1990 (in romanian).
- [4] I.R.Ionescu , Error estimates of a numerical method for a nonlinear evolution equation , An. Univ. Bucharest , 2(1988) , p. 64-74.
- [5] M.Sofonea , Quasistatic processes for elastic-viscoplastic materials with internal state variables (to appear in An.Univ.Blaise Pascal(Clermont II)).
- [6] W.Hahn , Stability of motion , Springer-Verlag , Berlin , 1967 .