

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

A. L. MAYER

Generating curves on abelian varieties and Riemann's theta-function

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 3^e série, tome 19,
n° 1 (1965), p. 107-111

http://www.numdam.org/item?id=ASNSP_1965_3_19_1_107_0

© Scuola Normale Superiore, Pisa, 1965, tous droits réservés.

L'accès aux archives de la revue « *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

GENERATING CURVES ON ABELIAN VARIETIES AND RIEMANN'S THETA-FUNCTION

A. L. MAYER

I. Introduction.

We shall show how to prove (and possibly illuminate) two well known theorems of Riemann and Poincaré about theta-functions, using Weil's ([9]) « intrinsic » approach to classical abelian varieties. Some variations of Matsusaka's criterion for jacobians ([2], [5]) are incidentally obtained.

II. Generating curves.

Let Z be a positive 1-cycle on an n -dimensional abelian variety A . Since we work modulo numerical equivalence, we may suppose that Z is a (possibly reducible) curve, all of whose components pass through the identity 0 of A . Let $\{Z\}$ be the smallest abelian subvariety containing Z , and call Z a generating curve if $\{Z\} = A$.

Let $Z^{(n)}$ denote the n -fold Pontrjagin product of Z with itself (see [10]). There is a non-negative integer $k(Z)$ such that, as a cycle, $Z^{(n)} = k(Z) n! A$.

PROPOSITION 1. Z is a generating curve $\iff k(Z) \neq 0$.

PROOF. The implication \Leftarrow is obvious. For the converse, first consider the case of an irreducible Z , and let r be the smallest integer for which $Z^{(r+1)} = 0$. Then $|Z^{(r)}|$ (the support of $Z^{(r)}$) is invariant under translations by points of Z , and since it contains 0 , must be a subgroup, so clearly must be equal to $\{Z\}$, and the proposition follows immediately in this case. Now for any Z , let its irreducible components be the Z_i , and $n_i = \dim \{Z_i\}$, so that by the preceding, there are positive integers k_i for which $Z_i^{(n_i)} = n_i! k_i \{Z_i\}$. Furthermore, one sees that the $\{Z_i\}$ generate $\{Z\}$.

If the $\{Z_i\}$ are direct summands of $\{Z\}$ then $n = \sum_i n_i$, and $k(Z) = \sum_i k_i \neq 0$. If $n = \sum_i n_i$, but the sum is not direct, then the direct sum of the $\{Z_i\}$ form a cover of degree $d > 1$ of $\{Z\}$, and $k(Z) = d \sum_i k_i > 1$. If, finally, $n < \sum_i n_i$, for some $i \neq j$ $\dim \{Z_i\} \cap \{Z_j\} > 0$. Choose the smallest $r_i \leq n_i$ so that $Z_i^{(n_i)} \oplus Z_j^{(r_i+1)} = 0$ (where \oplus means Pontrjagin product). Then by an argument similar to that above, we see that $|Z_i^{(n_i)} \oplus Z_j^{(r_j)}|$ is an abelian variety, in fact, it equals $\{Z_i + Z_j\}$. By symmetry, we can also find an $r_i \leq n_i$ so that $\{Z_i + Z_j\} = |Z_i^{(r_i)} \oplus Z_j^{(n_j)}|$.

Hence, in the formula

$$Z^{(n)} = \sum_{g_1 + \dots + g_q = n} \frac{n!}{g_1! \dots g_q!} Z_1^{(g_1)} \oplus \dots \oplus Z_q^{(g_q)}$$

the right-hand side has at least two nonvanishing terms, so in fact, $k(Z) \geq 2$.

PROPOSITION 2. If $k(Z) = 1$, the Z_i are non-singular curves of genus n_i , and A is the direct sum of the $\{Z_i\}$, which are the jacobians of the Z_i .

PROOF. The proof of Proposition 1 shows that A is the direct sum of the $\{Z_i\}$, and that $Z_i^{(n_i)} = n_i! \{Z_i\}$, so we are reduced to the case of Z irreducible. In that case, let J be the jacobian of its normalization \bar{Z} . Since Z is generating, the map $\bar{Z} \rightarrow Z$ gives rise to a surjection $J \rightarrow A$, so \bar{Z} has genus $g \geq n$. But $k(Z) = 1$ means that the n -fold symmetric product of \bar{Z} is birationally equivalent to A , so by a remark of Weil ([11] p. 37) $n = g$, and $J \rightarrow A$ is an isomorphism. In particular, $Z = \bar{Z}$.

III. The associated Kähler metric

Assume we are in the classical case, and define an hermitian form on $H^0(A, \Omega^1)$ (the holomorphic differentials) by

$$H(\alpha, \alpha') = \frac{\sqrt{-1}}{2} \int_Z \alpha \wedge \bar{\alpha}'$$

PROPOSITION 3. Z is generating $\iff H$ is positive definite.

PROOF. The implication \Leftarrow is obvious. Conversely, let Z generate A , and J_j be the jacobians of the normalizations \bar{Z}_j of the components Z_j of Z . The maps $\varphi_j: \bar{Z}_j \rightarrow Z_j$ induce a surjection $\varphi: \prod_j J_j \rightarrow A$, whose dual map

$\widehat{\varphi}: \widehat{A} \rightarrow \Pi_j \widehat{J}_j$ on the Picard varieties thus has finite kernel. The lifting of $\widehat{\varphi}$ to the universal covering spaces is the map $\Pi_j \varphi_j^*: H^0(A, \Omega^1) \rightarrow \Pi_j H^0(J_j, \Omega^1)$ gotten by pulling back differentials, which, being linear and having a discrete kernel, is an injection. So if $\alpha \neq 0$ is a differential on A , some $\varphi_j^*(\alpha) \neq 0$, and

$$H(\alpha, \bar{\alpha}) \geq \frac{\sqrt{-1}}{2} \int_{\bar{z}_j} \varphi_j^*(\alpha) \wedge \overline{\varphi_j^*(\alpha)} = \int_{\bar{z}_j} \|\operatorname{Re}(\varphi_j^*(\alpha))\|^2 > 0$$

(where $\|\cdot\|$ is the Dirichlet norm) by a standard result about harmonic forms on Riemann surfaces.

Now let $\alpha_1, \dots, \alpha_n$ be an orthonormal basis for H . Then $(\sum_i \alpha_i)^2$ gives an hermitian form on the tangent space to A at 0, and hence an invariant Kähler metric. Associated with this is the closed 2-form $u = \frac{\sqrt{-1}}{2} \sum_i \alpha_i \wedge \bar{\alpha}_i$, and Hodge's adjoint operation $*$ on differentials forms of cohomology. An easy computation (see Weil [9] p. 20) shows that $*u = \left(\frac{\sqrt{-1}}{2}\right)^{n-1} \sum_i \Omega_i$, where $\Omega_i = \prod_{i \neq j} \alpha_j \wedge \bar{\alpha}_j$. Let $\gamma = \int_A u^n/n!$ be the volume of A for this metric.

Let $[X]$ denote the cohomology class of either a form or cycle, X , denote either intersection or cup products by juxtaposition, and preserve the notation $X \oplus Y$ and $X^{(r)}$ for Pontrjagin products of cohomology classes, as well as cycles. (It is not hard to see that the two are compatible). Then we have:

PROPOSITION 4. $*[u] = \gamma[Z]$

PROOF. Let η be a closed (1,1) form on A , hence cohomologous to $\frac{\sqrt{-1}}{2} \sum_{ij} a_{ij} \alpha_i \wedge \bar{\alpha}_j$, for constant a_{ij} . Now

$$\frac{\sqrt{-1}}{2} \alpha_i \wedge \bar{\alpha}_j \wedge \Omega_k = \begin{cases} u^n/n! & \text{if } i = j = k \\ 0 & \text{otherwise} \end{cases}$$

so $[\eta \wedge *u] = [\sum_j a_{jj} u^n/n!]$. But on the other hand,

$$\int_Z \eta = \sum_{ij} \frac{\sqrt{-1}}{2} a_{ij} \int_Z \alpha_i \wedge \bar{\alpha}_j = \sum_{ij} a_{ij} H(\alpha_i, \alpha_j) = \sum_j a_{jj}$$

and the proposition follows immediately.

PROPOSITION 5. $[Z^{(n-1)}] = \gamma^{1-n} (n-1)! [u]$.

PROOF. By a general result ([4]) one has $*(XY) = (*X) \oplus (*Y)$ for X and Y any cohomology classes. Hence

$$[Z]^{(n-1)} = (*[u]/\gamma)^{(n-1)} = \gamma^{1-n} *[u]^{(n-1)}.$$

But by [9] p. 25 $[u]^{n-1} = (n-1)! *[u]$, and our assertion follows.

COROLLARY 1. $k(Z) = \gamma^{-n}$, so if $\gamma \geq 1$ (in particular, if the metric is Hodge, $k(Z) = \gamma = 1$ and A is a product of jacobians).

PROOF. $[Z^{(n)}] = k(Z) n! [A]$, but also $[Z]^{(n)} = (*[u]/\gamma)^{(n)} = \gamma^{-n} *[u^n] = \gamma^{-n} n! [A]$. The last assertion follows from Proposition 2.

Assume we are dealing with a product of jacobians, and let, as usual, $r! W_r = Z^{(r)}$ and $\Theta = W_{n-1}$.

COROLLARY 2. (Poincaré [6]. For numerical equivalence, in the abstract case, see [5]). $(n-r)! [W_r] = [\Theta^{n-r}]$.

$$\text{PROOF. } (n-r)! [W_r] = \frac{(n-r)!}{r!} Z^{(r)} = \frac{(n-r)!}{r!} *([Z])^r = (n-r)!/r! *[u^r].$$

But by an elementary algebraic identity ([1] p. 170) $*[u^r] = r!/(n-r)! [u^{n-r}]$, from which the corollary follows.

COROLLARY 3. (Riemann [7]. See e. g. [3] for a rigorous presentation). Let ϑ be « Riemann's » theta-function. Then a translate of Θ is « cut out by » ϑ .

PROOF. Let ϑ cut out the positive divisor X . Examining its « factors of automorphy » we see that ϑ « belongs to » the hermitian form H , in the sense of Weil [9] ch. VI. Hence ([9] p. 112) $[X] = [u] = [\Theta]$, and so ([9] p. 115) X is linearly equivalent to a translate of Θ . But since $\gamma = 1$, $\lambda(X) = 1$ by the (Frobenius) Riemann-Roch theorem, and the corollary follows.

Note that ϑ is even, so $X = X^-$, the image of X under the endomorphism $x \rightarrow -x$. On the other hand, $\Theta = \Theta_c$, where c is the « canonical point » (see Weil [8] p. 73). So if $\Theta = X_r$, then $2r = c$.

The coordinates of the point r are the traditional « Riemannian constants ».

REFERENCES

- [1] W. V. D. HODGE, *Theory and Application of Harmonic Integrals*, Cambridge, 1959.
- [2] W. L. HOYT, *On Products and algebraic families of jacobian varieties*, *Annals of Math.* 77 (1963) 415-423.
- [3] J. LEWITTES, *Riemann Surfaces and Theta-functions*, to appear in *Acta Math.*
- [4] A. L. MAYER, *The cohomology ring of a compact Lie group with a bi-invariant metric*, to appear in *Proc. Amer. Math. Soc.*
- [5] T. MATSUSAKA, *On a characterization of a Jacobian variety*, *Mem. Coll. Sci. Kyoto*, ser. A. 39, No. 1, pp. 1-19.
- [6] H. POINCARÉ, *Remarques divers sur les fonctions abeliennes*, *œuvres*, t. IV.
- [7] B. RIEMANN, *Theorie der Abelschen Funktionen*, *Werke* (Dover ed.) 88-142.
- [8] A. WEIL, *Variétés abeliennes et courbes algébriques*, Hermann 1948.
- [9] A. WEIL, *Variétés Kähleriennes*, Hermann 1958.
- [10] S. LANG, *Abelian Varieties*, Interscience, 1959.
- [11] A. WEIL, *Zum Beweis des Torellischen Satzes*, *Nachr. Akad. Wiss. Gött. Math. Phys. Kl.* (1957) No. 2.