

# *Astérisque*

PIERRE MICHEL

**Minimal sets generated by a substitution of  
non constant length**

*Astérisque*, tome 40 (1976), p. 143-146

<[http://www.numdam.org/item?id=AST\\_1976\\_40\\_143\\_0](http://www.numdam.org/item?id=AST_1976_40_143_0)>

© Société mathématique de France, 1976, tous droits réservés.

L'accès aux archives de la collection « Astérisque » ([http://smf4.emath.fr/  
Publications/Asterisque/](http://smf4.emath.fr/Publications/Asterisque/)) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>*

MINIMAL SETS GENERATED BY A  
SUBSTITUTION OF NON CONSTANT LENGTH

Pierre MICHEL

(Rennes)

Let  $P$  denote the alphabet  $\{0,1\}$  and let

$$\Omega = P^{\mathbb{Z}} = \{\omega = \dots \omega_{-1} \omega_0 \omega_1 \omega_2 \dots \mid \omega_i \in P\}$$

A substitution  $\theta$  is a map from  $P$  to  $\bigcup_{n \geq 2} P^n$ . If  $\text{card } \theta(0) = \text{card } \theta(1)$

the substitution is said of constant length. If not, it is said of non constant length.

For suitable  $i$  and  $j$  we can generate (1) a bisequence by the following:

$$\omega[-\text{card } \theta^{2n}(i), \text{card } \theta^{2n}(i)+\text{card } \theta^{2n}(j)] = \theta^{2n}(i) \theta^{2n}(j), \forall n \in \mathbb{N}$$

where  $\omega(a,k)$  is the  $k$ -block of  $\omega$  which begin at  $\omega_a$ .

Under a condition of non-degeneracy, the dynamical system  $(O_\omega, T)$  where  $O_\omega$  is the orbit closure of  $\omega$ , and  $T$  the shift, is minimal (1) and strictly ergodic (2).

We can prove that the dynamical system associated with the substitution  $\theta_1$  defined by

$$\begin{cases} \theta_1(0) = 0^{n+1-p} 1 0^p \\ \theta_1(1) = 1 0^n \end{cases} \quad \text{for } 0 \leq p \leq n,$$

has a purely discrete spectrum.  $(i^k, i \in P, k \in \mathbb{N}$  means  $\underbrace{i \ i \dots i}_{n \text{ times}}$ ).

---

(1) GOTTSCHALK. Substitution minimal sets. Trans. Amer. Math. Soc. 109 (1963)

(2) MICHEL. Stricte ergodicité d'ensembles minimaux de substitution. CRAS Paris, 278, (1974).

For this, we introduce the notion of "coincidence-value of the substitution  $\theta$ " by the following :

$$\begin{aligned} \text{if } \theta^\infty(0) &= \lim_{n \rightarrow \infty} \theta^{2n}(0) = \omega_1^0 \omega_2^0 \dots \omega_n^0 \dots \\ \theta^\infty(1) &= \lim_{n \rightarrow \infty} \theta^{2n}(1) = \omega_1^1 \omega_2^1 \dots \omega_n^1 \dots , \end{aligned}$$

we say that  $n$  is a "coincidence value of  $\theta$ " if and only if

$$\omega_n^0 = \omega_n^1 .$$

If we denote the density of the coincidence values of  $\theta$  in  $\mathbb{N}$  by  $\delta_C(\theta)$ , we can prove that  $\delta_C(\theta_1)$  is equal to one.

Conversely if we consider the substitution  $\theta_2$  :

$$\begin{cases} \theta_2(0) = 0\ 1 \\ \theta_2(1) = 1\ 1\ 0\ 0 \end{cases}$$

we have  $\delta_C(\theta_2) = \frac{1}{3}$  and we can prove that there is a continuous part in the spectrum of  $\theta_2$ .

Moreover, if we consider the two closely related substitutions  $\theta_2'$  and  $\theta_2''$  defined by

$$\theta_2'(0) = \theta_2''(0) = 0\ 1$$

and

$$\begin{cases} \theta_2'(1) = 1\ 0\ 1\ 0 \\ \theta_2''(1) = 1\ 0\ 0\ 1 \end{cases}$$

we can prove that  $\theta_2'$  and  $\theta_2''$  have purely discrete spectra and that

$$\delta_C(\theta_2') = \delta_C(\theta_2'') = 1 .$$

More generally, we conjecture that the substitution  $\theta_n$  :

$$\theta_n(0) = 0^n 1^n = \underbrace{0\dots 0}_{n \text{ times}} \underbrace{11\dots 1}_{n \text{ times}}$$

$$\theta_n(1) = 1^{2n} 0^{2n} = \underbrace{11\dots 1}_{2n \text{ times}} \underbrace{00\dots 0}_{2n \text{ times}} ,$$

has :

- 1) a purely discrete spectrum if  $n$  is even (then  $\delta_c = 1$ )
- 2) a mixed spectrum (partly continuous and partly discrete)  
if  $n$  is odd (then  $\delta_c = \frac{1}{3}$ ).

At last, if we consider the constant-length case (3), we know that the so-called

- regular Toeplitz substitution generates a dynamical system with discrete spectrum. In this case  $\delta_c = 1$ .
- generalized Morse substitution generates a dynamical system with mixed spectrum and in this case  $\delta_c = 0$ .

Thus we can draw the following table, and we state the question

Is the following true :

$$\begin{aligned}\delta_c = 1 &\implies \text{discrete spectrum} \\ \delta_c < 1 &\implies \text{mixed spectrum ?}\end{aligned}$$

(see the following page)

---

(3) COVEN and KEANE. The structure of substitution minimal sets. (1971)

SUBSTITUTION		$\delta_c$	SPECTRUM
constant length	Regular Toeplitz substitution $(\theta(0) \neq \tilde{\theta}(1))$ (1)	1	Discrete
	Generalized Morse substitution $(\theta(0) = \tilde{\theta}(1))$ (1)	0	Mixed
non constant length	$\begin{cases} \theta(0) = 0^{n+1-p} 1^p 0^p \\ \theta(1) = 1^p 0^n \end{cases}$ $0 \leq p \leq n$	1	Discrete
	$\begin{cases} \theta(0) = 01 \\ \theta(1) = 1010 \end{cases}$ and $\begin{cases} \theta(0) = 01 \\ \theta(1) = 1001 \end{cases}$	1	Discrete
	$\begin{cases} \theta(0) = 01 \\ \theta(1) = 1100 \end{cases}$	$\frac{1}{3}$	Mixed
	$\begin{cases} \theta(0) = 0^n 1^n \\ \theta(1) = 1^{2n} 0^{2n} \end{cases}$	n even n odd	Discrete Mixed
	other cases	?	?

(1) with  $\tilde{\omega}_k = 1 - \omega_k$