

# *Astérisque*

MAREK GOLASINSKI

**Some remarks on the rational homotopy type of  
diagrams and reduced  $K_0$**

*Astérisque*, tome 113-114 (1984), p. 187-191

[http://www.numdam.org/item?id=AST\\_1984\\_\\_113-114\\_\\_187\\_0](http://www.numdam.org/item?id=AST_1984__113-114__187_0)

© Société mathématique de France, 1984, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

SOME REMARKS ON THE RATIONAL HOMOTOPY  
TYPE OF DIAGRAMS AND REDUCED  $K_0$ .

by

Marek GOLASINSKI

1 - THE RATIONAL HOMOTOPY TYPE OF DIAGRAMS

Let  $\mathcal{C}$  be a closed model category in the Quillen's sense (see 5). If  $\mathbb{I}$  is a small category  $\mathcal{C}^{\mathbb{I}}$  denotes the functors category. A map in  $\mathcal{C}^{\mathbb{I}}$   $f : C \rightarrow C'$  is a fibration, respectively a weak equivalence if  $f(i)$  is a fibration, respectively a weak equivalence for every  $i \in \text{ob } \mathbb{I}$ . A cofibration is a map that has the left lifting property with respect to all trivial fibrations. We have the following result of Quillen-Bousfield-Kan (see 1, p. 313).

THEOREM 1.1.-  $\mathcal{C}^{\mathbb{I}}$  equipped as above is a closed model category.

Let  $\mathcal{G}$  be a discrete group.  $\mathcal{G}\text{-Set}$  is the category of left  $\mathcal{G}$ -sets and  $\mathbb{H}$  is the full subcategory of  $\mathcal{G}\text{-Set}$  determined by  $\mathcal{G}/\mathbb{H}$  as  $\mathbb{H}$  varies over all subgroups of  $\mathcal{G}$ . Denote by  $\mathcal{G}\text{-SS}$  the category of left  $\mathcal{G}$ -simplicial sets and  $\mathcal{G}\text{-Top}$  the category of left  $\mathcal{G}$ -topological spaces. Define functors  $J : \mathcal{G}\text{-SS} \rightarrow \text{SS}^{\mathbb{I}}$  by  $J(X)(\mathcal{G}/\mathbb{H}) = X^{\mathbb{H}}$ , where  $X^{\mathbb{H}} = \{x \in X ; h x = x, \text{ for all } h \in \mathbb{H}\}$  and  $T : \text{SS}^{\mathbb{I}} \rightarrow \mathcal{G}\text{-SS}$  by  $T(F) = F(\mathcal{G})$  provided with its natural  $\mathcal{G}$ -action acquired from  $\mathcal{G}\text{-Set}(\mathcal{G}, \mathcal{G}) = \mathcal{G}$ .

Let  $f : T(F) \rightarrow X$  be a map in  $\mathcal{G}\text{-SS}$ . Define  $f' : F \rightarrow J(X)$  by  $f'(\sigma) = f F(q)(\sigma)$  for  $\sigma \in F(\mathcal{G}/\mathbb{H})$  and  $q : \mathcal{G} \rightarrow \mathcal{G}/\mathbb{H}$  the natural quotient map. It is routine to check that  $f'$  is natural. Furthermore if  $h : F \rightarrow J(X)$  then  $h(\sigma) = h^{\vee} F(q)(\sigma)$  where  $h^{\vee} : F(\mathcal{G}) \rightarrow X$  is the  $\mathcal{G}$ -component of  $h$ , i.e.  $h$  is determined by  $h^{\vee}$ . We have thus established :

PROPOSITION 1.2.- J is full and faithful and right adjoint to T. Furthermore T preserves limits and both T and J preserve tensor products over SS.

Using J we view G-SS as a subcategory of  $SS^{\mathbb{I}}$ . A map  $f : X \rightarrow X'$  of G-SS is said to be a fibration, respectively a weak equivalence if  $J(f)$  is a fibration, respectively a weak equivalence of  $SS^{\mathbb{I}}$ . A cofibration in G-SS is a map of G-SS that has the left lifting property with respect to all trivial fibrations in G-SS. We have

PROPOSITION 1.3.- G-SS equipped as above is a closed model category. Furthermore each monomorphism of G-SS is a cofibration and thus any object of G-SS is cofibrant.

Consider the adjoint pair  $S : \text{Top} \rightleftarrows SS : | |$ , where S is the singular functor and  $| |$  is the geometric realization. The functors yield by naturality an adjoint pair  $S_{\mathbb{C}} : \mathbb{C}\text{-Top} \rightleftarrows \mathbb{C}\text{-SS} : | |_{\mathbb{C}}$  with natural isomorphism  $\mathbb{C}\text{-Top}(|F|_{\mathbb{C}}, X) \simeq \mathbb{C}\text{-SS}(F, S_{\mathbb{C}}(X))$ .

Let Q-DGA be the category of differential graded Q-algebras and  $A^* : SS \rightleftarrows Q\text{-DGA} : F^*$  the pair of de Rham adjoint functors (see 6). These functors determine an adjoint pair  $A^{*\mathbb{I}} : SS^{\mathbb{I}} \rightleftarrows Q\text{-DGA}^{\mathbb{I}} : F^{*\mathbb{I}}$ . If  $fQ\text{-SS}_N^{\mathbb{I}} \subset SS^{\mathbb{I}}$  is the full subcategory given by functors  $X \in SS^{\mathbb{I}}$  such that  $X(G/H)$  is nilpotent, rational and of finite Q-type for every subgroup  $H \subset G$  and  $fQ\text{-DGA}^{\mathbb{I}} \subset Q\text{-DGA}^{\mathbb{I}}$  is the full subcategory given by those functors  $A \in Q\text{-DGA}^{\mathbb{I}}$  that  $A(G/H)$  is equivalent to a minimal algebra with finitely many multiplicative generators in each dimension for every subgroup  $H \subset G$ . Then we obtain a generalization of the Sullivan-de Rham result (cf. 6) :

THEOREM 1.4.- Let G be a finite group. The adjoint pair  $A^{*\mathbb{I}} : SS^{\mathbb{I}} \rightleftarrows Q\text{-DGA}^{\mathbb{I}} : F^{*\mathbb{I}}$  induces an equivalence of homotopy categories

$$\text{Ho}(fQ\text{-SS}_N^{\mathbb{I}}) \xrightarrow[\simeq]{\simeq} \text{Ho}(fQ\text{-DGA}^{\mathbb{I}}).$$

Let  $fQ\text{-SS}_N$  be the full subcategory of G-SS given by nilpotent, rational and of finite Q-type G-simplicial sets. The functor  $J : \mathbb{C}\text{-SS} \rightarrow SS^{\mathbb{I}}$  is full and faithful, then we have.

COROLLARY 1.5.- The above equivalence induces a bijection between equivariant rational homotopy types of  $fQ\text{-SS}_N$  on the one hand and isomorphism classes of minimal systems of DGA's in the Triantafillou sense (see 7) on the other.

2 - REDUCED  $K_0$  OF 0-FORMS ON A FINITE SIMPLICIAL COMPLEX

Sullivan has proved (see 6, cf. also 4) that for a finite simplicial complex  $X$  with vertices  $v_1, \dots, v_n$  and corresponding barycentric coordinates  $b_1, \dots, b_n$  the algebra of rational forms on  $X$

$$A_Q^0 X = Q[b_1, \dots, b_n] \otimes \Lambda(db_1, \dots, db_n) / I,$$

where  $Q[b_1, \dots, b_n]$  is the ring of rational polynomials in  $b_1, \dots, b_n$ ,  $\Lambda(db_1, \dots, db_n)$  is the exterior algebra on  $db_1, \dots, db_n$  and  $I$  is the ideal generated by  $b_1 + \dots + b_n - 1$ ,  $db_1 + \dots + db_n$ ,  $b_{i_1} \dots b_{i_p} db_{j_1} \dots db_{j_q}$  if there is no  $p+q$ -simplex of  $X$  with vertices  $v_{i_1}, \dots, v_{i_p}, v_{j_1}, \dots, v_{j_q}$ .

Kan and Miller have shown (see 3) that the weak homotopy type of a finite simplicial set  $X$  can be reconstructed from  $R$ -algebra  $A_R^0 X$  of 0-forms on  $X$ , when  $R$  is a unique factorization domain.

If  $\text{pro } R\text{-}\mathcal{A}$  denote the pro-category of  $R$ -algebras then Jardine has proved (see 2) that there are functors  $\hat{A} : \text{SS} \rightleftarrows \text{pro } R\text{-}\mathcal{A} : \hat{F}$  inducing an equivalence of suitable homotopy categories

$$\text{Ho}(\text{SS}) \xrightarrow[\cong]{\quad} \text{Ho}(\text{pro } R\text{-}\mathcal{A}).$$

Our purpose is to show that there exists a simplicial set  $G_\infty(\infty)$  (the simplicial Grassman variety) such that for a finite simplicial complex  $X$ ,  $\tilde{K}_0(A_k^0 X) = [X, G_\infty(\infty)]$  where  $\tilde{K}_0$  is the reduced Grothendieck's group of  $A_k^0 X$  and  $k$  is a field.

The Grassman variety  $G_m(n)$  is defined as a functor from the  $K$ -algebras category  $k\text{-}\mathcal{A}$  to the category of sets, for  $1 \leq n < m$  and  $R$  in  $k\text{-}\mathcal{A}$  by

$$G_m(n)(R) = \{Q \subset R^m; Q \text{ is } R\text{-split projective of rank } n\}.$$

The assignment  $Q \mapsto Q \otimes_R S$  associated to the  $k$ -algebra homomorphism  $\theta : R \rightarrow S$  defines the function  $\theta_* : G_m(n)(R) \rightarrow G_m(n)(S)$ .

Let  $P(R)$  ( $P_n(R)$ ) be the set of isomorphism classes of  $R$ -modules finitely generated and projective over  $R$  (of rank  $n$ ),  $\tilde{K}_0(R)$  the Grothendieck's group of  $P(R)$  and  $K_0(R)$  reduced  $K_0$ .

The natural embedding  $R^m \rightarrow R^{m+1}$  induces a map  $G_m(n) \rightarrow G_{m+1}(n)$ .

Put  $G_\infty(n) := \text{colim}_n G_m(n)$ .

Then there is a natural surjective function  $\tau_R : G_\infty(n)(R) \rightarrow P_n(R)$  which is induced by the assignment  $(P \rightarrow R^m) \mapsto P$ .

Let  $X$  be a finite simplicial complex, thought of as a member of the category  $SS$  of simplicial sets, and let  $k$  be an arbitrary field. Recall that there is a natural simplicial set map

$$\eta_X : X \longrightarrow \text{Spec}(A_k^0 X) (A_k^0 \Delta_*) = k\text{-}A(A_k^0 X, A_k^0 \Delta_*)$$

where  $\Delta_n$  is the standard  $n$ -simplex.

Let  $\text{Sch}_k$  denotes the category of schemes over  $k$ , thought of as a full subcategory of the functors category from  $k\text{-}A$  to  $\text{Set}$ .

$\eta_X$  may be used to define a function

$$\psi : \text{Sch}_k(\text{Spec } A_k^0 X, Y) \longrightarrow SS(X, Y(A_k^0 \Delta_*))$$

for arbitrary  $k$ -schemes  $Y$  in such a way that  $\psi$  associates to a  $k$ -scheme map  $f : \text{Spec } A_k^0 X \longrightarrow Y$  the composition

$$X \xrightarrow{\eta_X} \text{Spec}(A_k^0 X) (A_k^0 \Delta_*) \xrightarrow{f_*} Y(A_k^0 \Delta_*) .$$

PROPOSITION 2.1.-  $\psi$  induces a bijection

$$\psi_* : \text{Sch}_k(\text{Spec } A_k^0 X, Y) \xrightarrow{\cong} SS(X, Y(A_k^0 \Delta_*))$$

for all finite simplicial complexes  $X$  and all schemes  $Y$ .

Then the above map  $\tau$  gives rise to a natural surjective function

$$\tau_X : SS(X, G_\infty(n) (A_k^0 \Delta_*)) \longrightarrow P_n(A_k^0 X)$$

in view of above theorem and the Yoneda lemma.

THEOREM 2.2.- The map  $\tau_X : SS(X, G_\infty(n) (A_k^0 \Delta_*)) \longrightarrow P_n(A_k^0 X)$  factors through a bijection

$$(\tau_X)_* : [X, G_\infty(n) (A_k^0 \Delta_*)] \xrightarrow{\cong} P_n(A_k^0 X) .$$

The map  $P_n(A_k^0 X) \longrightarrow P_{n+1}(A_k^0 X)$  which is defined by  $P \longmapsto A_k^0 X \otimes P$  clearly fits into a commutative diagram

$$\begin{array}{ccc} [X, G_\infty(n) (A_k^0 \Delta_*)] & \xrightarrow{\quad\quad\quad} & [X, G_\infty(n+1) (A_k^0 \Delta_*)] \\ \downarrow (\tau_X)_* & & \downarrow (\tau_X)_* \\ P_n(A_k^0 X) & \xrightarrow{\quad\quad\quad} & P_{n+1}(A_k^0 X) \end{array}$$

DIAGRAMS AND REDUCED  $K_0$

Formal nonsense now shows that

$$[X, G_\infty^{(\infty)}(A_k^0 \Delta_*)] = \text{colim} (P_n(A_k^0 X) \longrightarrow P_{n+1}(A_k^0 X))$$

may be identified with  $\check{K}_0(A_k^0 X)$  via a map which is induced by the assignments

$$P \longmapsto P - (A_k^0 X)^{\text{rk} P}, \text{ where } G_\infty^{(\infty)} = \text{colim}_n G_\infty^{(n)}.$$

There is also a similar result for finite  $G$ -simplicial complexes.

Let  $G$  be a finite group such that  $\chi(k) \chi|G|$ ,  $\chi(k)$  is the characteristic of  $k$ ,  $V_1, \dots, V_\ell$  all irreducible  $G$ -modules over  $k$  and  $V := \bigoplus_{i=1}^{\ell} V_i$ . The  $\mathbb{C}$ -Grassman variety  $\mathbb{G}_m^{\mathbb{C}}(n)$  is defined as a functor from the  $k$ -algebras category  $k\text{-}\mathbb{A}$  to the category of sets, for  $1 \leq n < m$  and  $R$  in  $k\text{-}\mathbb{A}$  by

$$\mathbb{G}_m^{\mathbb{C}}(n)(R) := \{Q \subset R^m \otimes V; Q \text{ is } R\text{-split projective of rank } n\}.$$

Remark that for a  $k$ - $\mathbb{C}$ -algebra  $R$  the category of  $R$ - $\mathbb{C}$ -modules is equivalent to the category of  $R * \mathbb{C}$ -modules, where  $R * \mathbb{C}$  is the twisted product of  $R$  and  $\mathbb{C}$ . Let  $P^{\mathbb{C}}(R)$  denotes the set of isomorphism classes of  $R * \mathbb{C}$ -modules finitely generated and projective over  $R$ ,  $K_0^{\mathbb{C}}R$  the Grothendieck's group of  $P^{\mathbb{C}}(R)$  and  $\check{K}_0^{\mathbb{C}}(R)$  reduced  $K_0$ . Then we have.

THEOREM 2.3.- For a finite group  $G$  such that  $\chi(k) \chi|G|$  and a finite  $G$ -simplicial complex  $X$

$$K_0^{\mathbb{C}}(A_k^0 X) = [X, G_\infty^{(\infty)}(A_k^0 \Delta_*)]_{\mathbb{C}}.$$

R E F E R E N C E S.

- [1] A.K. BOUSFIELD, D.M. KAN - Lecture Notes in Math. 304 1972.
- [2] J.F. JARDINE - Algebraic homotopy theory, Can. J. Math. 33 1981, pp. 302-319.
- [3] D.M. KAN, E.Y. MILLER - Homotopy types and Sullivan's algebras of 0-forms, Topology vol. 16 1977, pp. 193-197.
- [4] M.A. PENNA - On a theorem of Sullivan, Can. Math. Bull. Vol. 21 1978 pp. 201-206.
- [5] D.G. QUILLEN - Lecture Notes in Math. 43 1967.
- [6] D. SULLIVAN - Differential forms and the topology of manifolds, Proc. of Congress on Manifolds, Tokyo, Japan 1973 pp. 37-49.
- [7] G. TRIANTAFILLOU - Equivariant minimal models, Trans. of A.M.S. Vol.274 2 1982, pp. 509-532.