

Astérisque

CLAS LÖFWALL

On the centre of graded Lie algebras

Astérisque, tome 113-114 (1984), p. 263-267

http://www.numdam.org/item?id=AST_1984__113-114__263_0

© Société mathématique de France, 1984, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE CENTRE OF GRADED LIE ALGEBRAS

by

Clas Löfwall

If \underline{a} is a graded Lie algebra over a field k in general, its centre may of course be any abelian graded Lie algebra. But if some restrictions are imposed on \underline{a} such as

- 1) $cd(\underline{a}) (= \text{gldim } U(\underline{a})) < \infty$
- 2) $U(\underline{a}) = \text{Ext}_R(k, k)$ R local noetherian ring
- 3) $\underline{a} = \pi_* S \otimes \mathbb{Q}$, $\text{cat}_0(S) < \infty$

what can be said about the centre?

Notation. For a graded Lie algebra \underline{a} , let $Z(\underline{a})$ denote its centre.

Felix, Halperin and Thomas have results in case 3) (cf [1]): Suppose $\dim_{\mathbb{Q}}(\underline{a}) = \infty$ then for each $k \geq 1$, $\sum_{n=k}^{2k-1} \dim_{\mathbb{Q}}(Z_{2n}(\underline{a})) < \text{cat}_0(S)$.

In case 1) we have the following result.

Theorem 1 Suppose $cd(\underline{a}) = n < \infty$. Then $\dim_k Z(\underline{a}) \leq n$ and $Z_{\text{odd}}(\underline{a}) = 0$. Moreover if $\dim_k Z(\underline{a}) = n$, then \underline{a} is an extension of an abelian Lie algebra on odd generators by its centre $Z(\underline{a})$.

Proof. We have that $U(Z(\underline{a}))$ is a sub Hopf algebra of $U(\underline{a})$ and hence $U(\underline{a})$ is free over $U(Z(\underline{a}))$ (cf [5] , th 4.4) and from this it follows that $cd(Z(\underline{a})) \leq cd(\underline{a}) = n$. But $U(Z(\underline{a}))$ is a tensor product of a polynomial algebra on the even generators of $Z(\underline{a})$ and an exterior algebra on the odd generators of $Z(\underline{a})$. Since the global dimension of an exterior algebra is infinite, we must have $Z_{\text{odd}}(\underline{a}) = 0$ and since the global dimension of a polynomial algebra is the number of variables, we also get $\dim_k Z(\underline{a}) \leq n$. Suppose now $\dim_k Z(\underline{a}) = n$. If $x \in \underline{a}$ and $\deg(x)$ is even and $x \notin Z(\underline{a})$, then $Z(\underline{a}) \otimes k \cdot x$ is a sub Lie algebra of \underline{a} of cohomological dimension $n+1$ which is impossible. Hence

$x \notin Z(\underline{a})$ implies $\deg(x)$ odd, and then it follows that $\underline{a}/Z(\underline{a})$ is abelian on odd generators.

Problem. Characterise those graded Lie algebras \underline{a} having $\text{cd}(\underline{a}) < \infty$ and which are an extension of an abelian finite dimensional Lie algebra on odd generators by its centre $Z(\underline{a})$.

In case 2) we have the following result.

Theorem 2. Suppose (R, m) is an equi-characteristic local noetherian ring with $m^3 = 0$. Let \underline{g} be the underlying Lie algebra of $\text{Ext}_R(k, k)$ ($k = R/m$). Then $Z(\underline{g}) = 0$ or $R = k[x]/(x^3)$ or $\text{cd}(\underline{g}) = 2$ (the last case is equivalent to saying that \underline{g} is generated by its one-dimensional elements).

This theorem is a consequence of the following one.

Theorem 3. Suppose \underline{a} is a graded Lie algebra and $V \neq 0$ is a syzygy in a (not necessarily minimal) resolution of k over $U(\underline{a})$. Let $\underline{g} = \underline{a} \ltimes F(V)$ be the semi-direct product of \underline{a} by $F(V) =$ the free Lie algebra on V . Then $Z(\underline{g}) = 0$ or \underline{a} is abelian on one single odd generator.

Notations. If A is an augmented ring, we will use $I(A)$ as a notation for the augmentation ideal. If I is an ideal in a ring A we denote by $\text{Ann}(I)$ the ideal $\{x \in A ; x \cdot I = 0\}$.

We will use the following lemma (cf e.g. [6]).

Lemma 1. Suppose \underline{a} is a graded Lie algebra and $\text{Ann}(IU(\underline{a})) \neq 0$, then \underline{a} is abelian and generated by finitely many odd elements.

We also have the following lemma.

Lemma 2. Suppose A is a graded ring, $A = \bigoplus_{n \geq 0} A_n$, $I(A) = \bigoplus_{n \geq 1} A_n$ and $a \in A$ is a homogeneous element satisfying $a^2 = 0$ and $\{x ; xa = 0\} = Aa$. Suppose further $F_n \xrightarrow{d} F_{n-1} \xrightarrow{d} \dots \rightarrow F_0 \xrightarrow{\epsilon} A_0 \rightarrow 0$ is the beginning of a graded free resolution of A_0 as a right A -module, and let $V = \ker(F_n \xrightarrow{d} F_{n-1})$.

Suppose also $V \cdot a = 0$. Then $V \cdot I(A) = 0$. In particular, if $V \neq 0$ then $\text{Ann}(I(A)) \neq 0$.

Proof. Take a homogeneous element v of V . Since v is also an element of the free A -module F_n and $v \cdot a = 0$, we can use the assumption $\{x \in A ; xa = 0\} = Aa$ to get an element x_n of F_n such that $v = x_n a$. But $0 = dv = (dx_n)a$ so in the same way we have an element x_{n-1} of F_{n-1} such that $dx_n = x_{n-1} a$. Finally we get $dx_1 = x_0 a$ where $x_0 \in F_0$. If x_0 has positive degree, there is $y_1 \in F_1$ with $dy_1 = x_0$. The equality $d(x_1 - y_1 a) = 0$ implies that there is $y_2 \in F_2$ such that $x_1 - y_1 a = dy_2$. From $dx_2 = x_1 a = (dy_2)a$ it follows that there is $y_3 \in F_3$ such that $x_2 - y_2 a = dy_3$ etc. At last $d(x_n - y_n a) = 0$, hence $x_n - y_n a \in V$ and since $V \cdot a = 0$ it follows that $v = x_n a = 0$. Suppose now x_0 is of degree zero. Then $\deg(x_1) = \deg(a)$, ..., $\deg(x_n) = n \cdot \deg(a)$ and $\deg(v) = (n+1) \cdot \deg(a)$. Hence V is concentrated in one single degree and therefore $V \cdot I(A) = 0$.

Remark. Lemma 2 is valid also for a local commutative ring A (with A_0 equal to the residue field).

Lemma 3. Suppose \underline{a} is a graded Lie algebra. Suppose $V \neq 0$ is a syzygy in a free (not necessarily minimal) resolution of k over $U(\underline{a})$ such that $V \cdot I(U(\underline{a})) = 0$. Then \underline{a} is abelian generated by one single odd element.

Proof. Since V is contained in a free module, the assumption $V \cdot I(U(\underline{a})) = 0$ implies that $\text{Ann } I(U(\underline{a})) \neq 0$. Hence by lemma 1 \underline{a} is abelian on finitely many odd generators. If V is a n^{th} syzygy it follows that

$$\text{Tor}_{n+1+i}^{U(\underline{a})}(k, k) = \text{Tor}_i^{U(\underline{a})}(k, k) \otimes_k V \text{ for } i \geq 1.$$

Hence $P_{U(\underline{a})}(z) = \text{Pol}(z)/(1 - \dim(V)z^{n+1})$ where $\text{Pol}(z)$ is a polynomial in z .

But we also know that there are numbers e_1, \dots, e_r such that

$$P_{U(\underline{a})}(z) = \prod_{i=1}^r (1 - z^{e_i})^{-1}.$$

The first expression shows that $z=1$ is a pole of order at most one, while the second expression shows that $z=1$ is a pole of order r . Hence $r=1$.

Proof of Theorem 3. Suppose $z \neq 0$ is an element of $Z(\underline{g})$ and $z = x + a$ where $x \in F(V)$ and $a \in \underline{a}$. 1) Assume first that $x \neq 0$. For each $y \in V$, $[x,y] + [a,y] = 0$. Now $F(V)$ is graded by the Lie degree "deg" defined by letting the elements of V have degree one. Since $\deg([a,y]) = \deg(y) = 1$ and $\deg([x,y]) = \deg(x) + 1$, it follows that $[x,y] = 0$. Since $F(V)$ is free, V must be one-dimensional and hence $V \cdot I(U(\underline{a})) = 0$. Since V is contained in a free $U(\underline{a})$ -module, it follows that $\text{Ann } I(U(\underline{a})) \neq 0$ and by lemma 1 and 3 \underline{a} is abelian on one odd generator. 2) Assume now that $x=0$, i.e. $z \in \underline{a}$. Then $y \cdot z = [y,z] = 0$ for all $y \in V$. Since V is non-zero and contained in a free $U(\underline{a})$ -module, z must be a zero-divisor on $U(\underline{a})$. But then z must be of odd degree and $z^2 = 0$. This follows from the Poincaré-Birkhoff-Witt theorem. Also from this theorem we get that $\{b \in U(\underline{a}) ; bz = 0\} = U(\underline{a}) \cdot z$. Since also $V \cdot z = 0$, lemma 2 may be applied to get $\text{Ann } I(U(\underline{a})) \neq 0$ and $V \cdot I(U(\underline{a})) = 0$ and then also in this case lemma 1 and 3 may be applied to get the result. ■

Finally, Theorem 2 follows from Theorem 3 since we know the structure of $\text{Ext}_R(k,k)$ if (R,m) is an equi-characteristic local ring with $m^3 = 0$. This may be deduced (with some effort) from [4], and hopefully it will appear in a forth-coming paper by the author. The structure of the underlying Lie algebra \underline{g} of $\text{Ext}_R(k,k)$ is given as follows. Let \underline{a} be the underlying Lie algebra of $\text{Ext}_R^{(1)}(k,k)$ = the sub algebra of $\text{Ext}_R(k,k)$ generated by its one-dimensional elements. Let V be the third syzygy in a minimal resolution of k over $U(\underline{a})$. Then $\underline{g} = \underline{a} \ltimes F(V)$.

An application.

Notation. For a local ring R , let $e_i(R)$ denote $\dim(\underline{g}_i)$ where \underline{g} is the Lie algebra of R .

Theorem 4. If (R,m) is local with $m^3 = 0$, then $e_i(R) > 0$ for all $i \geq 1$ or R is a complete intersection (which is possible only if $\dim(m/m^2) \leq 2$).

ON THE CENTRE OF GRADED LIE ALGEBRAS

Proof. We may assume that R is equi-characteristic since $P_R = P_{\text{gr}(R)}$
($\text{gr}(R) = \bigoplus_{i \geq 0} \mathfrak{m}^i / \mathfrak{m}^{i+1}$) (a result by Levin, cf [4]) and $e_i(R)$ may be computed
from P_R . During this conference I learned from Yves Felix that if $e_i(R) = 0$
for some i , then a "special variable" in the sense of André is defined (or, if
you prefer, there exists a "Gottlieb element"). But according to Jacobsson [3]
this defines an element in the centre of \mathfrak{g} = the Lie algebra of R . And by
Theorem 2 in this paper the centre of \mathfrak{g} is trivial, unless \mathfrak{g} is generated
by \mathfrak{g}_1 (or $R = k[x]/(x^3)$ which is a complete intersection). In this
case we have the following. If $e_i(R) = 0$ for some i then $e_j(R) = 0$ for all
 $j \geq i$. Hence by Gulliksen's theorem [2], R is a complete intersection.

References

- [1] Felix-Halperin-Thomas, The homotopy Lie algebra for finite complexes,
Publications mathématiques de l'IHES, n° 56 (1982), p. 387-410.
- [2] Gulliksen, A homological characterization of local complete intersection,
Compositio Math. 23, 1971, p. 251 - 255.
- [3] Jacobsson, On local flat homomorphisms and the Yoneda Ext-algebra of the fibre,
Reports, Dep. of Math., Univ. of Stockholm, no 8, 1982.
- [4] Löfwall, On the subalgebra generated by the one-dimensional elements of the
Yoneda Ext-algebra, Reports, Dep. of Math., Univ. of Stockholm,
no. 5, 1976.
- [5] Milnor-Moore, On the structure of Hopf algebras, Annals of Math., 81, 1965,
p. 210 - 264.
- [6] Sweedler, Hopf algebras, Benjamin, 1969, p. 108.

Acknowledgement. The author wants to thank Rikard Bøgvad and Calle Jacobsson
for many fruitful discussions.