

Astérisque

LÉONARD LIPSCHITZ

ZACHARY ROBINSON

**Rings of separated power series and quasi-affinoid
geometry - Pages préliminaires**

Astérisque, tome 264 (2000), p. I-2

http://www.numdam.org/item?id=AST_2000__264__R1_0

© Société mathématique de France, 2000, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ASTÉRIQUE 264

**RINGS OF SEPARATED POWER SERIES
AND QUASI-AFFINOID GEOMETRY**

**Leonard Lipshitz
Zachary Robinson**

L. Lipshitz

Department of Mathematics, Purdue University, West Lafayette,
IN 47907-1395 USA.

E-mail : lipshitz@math.purdue.edu

Z. Robinson

Department of Mathematics, East Carolina University, Greenville,
NC 27858-4353 USA.

E-mail : robinson@math.ecu.edu

1991 Mathematics Subject Classification. — 32P05, 32B05, 32B20, 32C35, 26E30,
12J25, 03C10, 13J05, 13B40.

Key words and phrases. — Rigid analytic geometry, Non-Archimedean analysis,
Subanalytic sets, Quantifier elimination, Model Completeness, Artin approximation,
Acyclicity.

Supported in part by the NSF.

RINGS OF SEPARATED POWER SERIES AND QUASI-AFFINOID GEOMETRY

Leonard Lipshitz, Zachary Robinson

Abstract. — The papers in this volume present a theory of rigid analytic geometry over an ultrametric field K that generalizes the classical, affinoid, theory to the setting of relative rigid analytic geometry over an “open” polydisc. The theory is based on the commutative algebra of power series rings $S_{m,n}$ that is developed in the first paper in this volume, *Rings of Separated Power Series*. Quasi-affinoid algebras (quotients $S_{m,n}/I$) share many properties with affinoid algebras (quotients T_m/I of a ring of strictly convergent power series.) Among the principal results are the Nullstellensatz for quasi-affinoid algebras A and the Universal Property for a broad class of open subdomains of $\text{Max } A$, the R -subdomains. The second paper, *Model Completeness and Subanalytic Sets*, obtains a structure theory for images of analytic maps based on any subcollection of $S = \cup S_{m,n}$ that satisfies certain closure properties; for example $T = \cup T_m$. The argument exploits the existential definability of the Weierstrass data as well as a difference between affinoid and quasi-affinoid rigid analytic geometry; namely, that a quasi-affinoid variety $\text{Max } A$ in general may be covered by finitely many disjoint quasi-affinoid subdomains, just as the valuation ring K° is the union of its maximal ideal $K^{\circ\circ}$ and its multiplicative units. A crucial role is played by the theory of generalized rings of fractions developed in the first paper. The third paper, *Quasi-Affinoid Varieties*, defines the category of $S_{m,n}$ -analytic varieties $X = \text{Max } A$ and establishes the acyclicity of quasi-affinoid covers. The proofs employ results from the first paper; in particular, the fact that the assignment $U \mapsto \mathcal{O}_X(U)$ is a presheaf of A -algebras for R -subdomains U of X . The quantifier elimination of the second paper is used to relate quasi-affinoid and affinoid covers, a key step in the proof of the Acyclicity Theorem. The fourth paper, *A Rigid Analytic Approximation Theorem*, gives a global Artin Approximation theorem between a “Henselization” $H_{m,n}$ of a ring T_{m+n} of strictly convergent power series and its “completion” $S_{m,n}$. This links the algebraic properties of affinoid and quasi-affinoid algebras.

Résumé (Anneaux de séries séparées et géométrie quasi-affinoïde)

Les articles de ce volume présentent une théorie de la géométrie analytique rigide sur un corps ultramétrique K qui généralise la théorie affinoïde classique au cas de la géométrie analytique rigide relative sur un polydisque « ouvert ». Cette théorie est basée sur l'étude algébrique des anneaux de séries convergentes $S_{m,n}$ développée dans le premier article, *Rings of Separated Power Series*. Les algèbres quasi-affinoïdes (les quotients $S_{m,n}/I$) partagent de nombreuses propriétés avec les algèbres affinoïdes (les quotients T_m/I d'un anneau de séries strictement convergentes). Parmi les résultats principaux signalons le Nullstellensatz pour les algèbres quasi-affinoïdes A ainsi que la Propriété Universelle pour une large classe de sous-domaines ouverts de $\text{Max } X$, les R -sous-domaines. Le second article, *Model Completeness and Subanalytic Sets*, contient des résultats sur la structure des images de familles de fonctions analytiques provenant par extension d'une famille quelconque de fonctions de $S = \cup S_{m,n}$ satisfaisant certaines propriétés de fermeture ; par exemple $T = \cup T_m$ est une telle famille. La preuve utilise le fait que les données de Weierstrass sont définissables ainsi que le fait, témoignant de la différence entre géométrie affinoïde et quasi-affinoïde, qu'une variété quasi-affinoïde $\text{Max } A$ peut généralement être recouverte par un nombre fini de sous-domaines quasi-affinoïdes disjoints, de la même façon que l'anneau de valuation K° est l'union de son idéal maximal $K^{\circ\circ}$ et de ses unités multiplicatives. La théorie des anneaux généralisés de fractions développée dans le premier article joue un rôle crucial. Dans le troisième article, *Quasi-Affinoid Varieties*, on définit la catégorie des variétés $S_{m,n}$ -analytiques $X = \text{Max } A$ et on établit l'acyclicité des recouvrements quasi-affinoïdes. Les démonstrations emploient des résultats du premier article, notamment le fait que le foncteur $U \mapsto \mathcal{O}_X(U)$ est un préfaisceau d' A -algèbres pour des R -sous-domaines U de X . On utilise également le résultat d'élimination des quantificateurs obtenu dans le second article pour établir un rapport entre les recouvrements quasi-affinoïdes et les recouvrements affinoïdes, ce qui est une étape cruciale dans la démonstration du théorème d'acyclicité. Le quatrième article, *A Rigid Analytic Approximation Theorem*, donne un théorème d'approximation globale d'Artin entre un « hensélisé » $H_{m,n}$ d'un anneau T_{m+n} de séries strictement convergentes et son complété $S_{m,n}$. Ce résultat permet de relier les propriétés algébriques des algèbres quasi-affinoïdes et affinoïdes.

CONTENTS

Introduction	1
Rings of Separated Power Series	3
1. Introduction	3
2. Rings of Separated Power Series	7
3. Restrictions to Polydiscs	24
4. The Commutative Algebra of $S_{m,n}$	60
5. The Supremum Semi-Norm and Open Domains	69
6. A Finiteness Theorem	101
References	107
Model Completeness and Subanalytic Sets	109
1. Introduction	109
2. Existentially Defined Analytic Functions	110
3. Existential Definability of Weierstrass Data	115
4. The Elimination Theorem	120
5. Subanalytic Sets	125
References	126
Quasi-Affinoid Varieties	127
1. Introduction	127
2. G -Topologies and the Structure Presheaf	130
3. Coverings and Acyclicity	139
References	149
A Rigid Analytic Approximation Theorem , by Zachary Robinson	151
1. Introduction	151
2. The Rings of Henselian Power Series	152
3. Flatness	156
4. Regularity	161
5. Approximation	164
References	168
Index	169

INTRODUCTION

Let K be a field, complete with respect to the non-trivial ultrametric absolute value $|\cdot| : K \rightarrow \mathbb{R}_+$. By K° denote the valuation ring, by $K^{\circ\circ}$ its maximal ideal, and by \tilde{K} the residue field $K^\circ/K^{\circ\circ}$. Let K' be an algebraically closed field containing K and consider the polydisc

$$\Delta_{m,n} := ((K')^\circ)^m \times ((K')^{\circ\circ})^n.$$

In 1961, Tate introduced rings T_m of analytic functions on the closed polydiscs $\Delta_{m,0}$. These rings lift the affine algebraic geometry of the field \tilde{K} . In particular, the Euclidean Division Theorem for $\tilde{K}[\xi]$ lifts to a global Weierstrass Division Theorem for T_m . The basic properties of T_m that follow from Weierstrass Division include Noetherianness, Noether Normalization, unique factorization, and a Nullstellensatz. These results pave the way for the development of rigid (affinoid) analytic geometry.

The representation

$$\Delta_{m,n} = \varinjlim_{\varepsilon} ((K')^\circ)^m \times (\varepsilon(K')^\circ)^n,$$

where $\varepsilon \in (K')^{\circ\circ}$, yields a ring of analytic functions on $\Delta_{m,n}$ by taking a corresponding inverse limit of Tate rings. This gives the polydisc $\Delta_{m,n}$ the structure of a rigid analytic variety. But its global functions are, in general, unbounded. Even if one restricts attention to those functions with finite supremum norm, the geometric behavior can be pathological.

In the first paper, *Rings of Separated Power Series*, we define rings $S_{m,n}$ of bounded analytic functions on $\Delta_{m,n}$ with a tractable algebraic and geometric behavior. Those rings share many of the nice properties of the Tate rings T_m , though the proofs are often rather more difficult. We show that the rings $S_{m,n}$ are Noetherian rings (often K -Banach algebras) of bounded analytic functions on $\Delta_{m,n}$, that satisfy a Nullstellensatz, are unique factorization domains, and are regular rings of dimension $m + n$.

We call quotients of the $S_{m,n}$ *quasi-affinoid algebras*. Quasi-affinoid algebras share most of the properties of affinoid algebras. For example, the residue norms arising from different presentations of a quasi-affinoid algebra are all equivalent; quasi-affinoid morphisms are continuous; in characteristic zero (and often in characteristic p) the residue norms and the supremum norm on a reduced quasi-affinoid algebra are equivalent; and quasi-affinoid rational domains satisfy the appropriate universal mapping property. These results pave the way for the development of a relative rigid analytic geometry over open polydiscs.

We give three applications of the general theory. In the second paper, *Model Completeness and Subanalytic Sets*, we present a quantifier elimination theorem which lays the foundation for the theory of rigid subanalytic sets based on the Tate Rings. The third paper, *Quasi-Affinoid Varieties*, applies the results of the first two papers to treat the basic sheaf theory of quasi-affinoid varieties and to prove the quasi-affinoid Acyclicity Theorem. In the fourth paper, *A Rigid Analytic Approximation Theorem*, a global Artin Approximation Theorem is presented for the pair of rings $H_{m,n} \hookrightarrow S_{m,n}$, where $H_{m,n}$ is the algebraic closure of T_{m+n} in $S_{m,n}$. In this context the rings $S_{m,n}$ play the role of a kind of completion of the Tate rings.

Acknowledgements. — Both authors thank MSRI and the ICMS, Edinburgh for support and hospitality, Jan Denef for his encouragement in this project, and Judy Mitchell for her patience in typing numerous versions of this manuscript. The second author also gratefully acknowledges the support of the CNR and the hospitality of the University of Pisa. Diagrams were created by using `diagram.sty` by Paul Taylor of Queen Mary and Westfield College, London.