

# COMPOSITIO MATHEMATICA

JAMES R. MOSHER

**Semirings with descending chain condition and without nilpotent elements**

*Compositio Mathematica*, tome 23, n° 1 (1971), p. 79-85

[http://www.numdam.org/item?id=CM\\_1971\\_\\_23\\_1\\_79\\_0](http://www.numdam.org/item?id=CM_1971__23_1_79_0)

© Foundation Compositio Mathematica, 1971, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## SEMIRINGS WITH DESCENDING CHAIN CONDITION AND WITHOUT NILPOTENT ELEMENTS

by

James R. Mosher

### 1. Introduction

Several decades ago, Artin, Nesbitt, and Thrall [1] published their classic work on rings with descending chain condition on left ideals. In recent years Herstein [5], Divinsky [3], Kertész [6], Szaász [8], and others have compiled these and other results in their books or articles. In this paper the author extends to a class of semirings some of these results in ring theory.

### 2. Definitions

A *semiring* is a non-empty set  $R$  on which two associative binary operations, addition and multiplication, are defined such that the multiplication distributes over the addition from both sides and such that there exists  $e \in R$  with  $x+e = e+x = x$  and  $ex = xe = e$  for all  $x \in R$ . We call  $e$  the *zero* of  $R$  and denote it by  $0$ .

A semiring  $R$  is *left semisubtractive* if for each  $x, y \in R$  there exists  $z \in R$  with  $z+x = y$  or  $x = z+y$ .

A *left semi-ideal* of a semiring  $R$  is a non-empty subset  $A$  of  $R$  such that for each  $x, y \in A$  and  $r \in R$  it is true  $x+y, rx \in A$ . A left semi-ideal  $A$  of  $R$  is a *left  $l$ -ideal* if  $x, y+x \in A$  imply  $y \in A$  and is a *left  $r$ -ideal* if  $x, x+y \in A$  imply  $y \in A$ . Similarly one defines these concepts using ‘right’ instead of ‘left’. A subset that is both a left and right semi-ideal is called a *semi-ideal*. Similarly one defines  *$l$ -ideal* and  *$r$ -ideal*. If a subset is a left [right]  *$l$ -ideal* and left [right]  *$r$ -ideal*, it is called a *left [right] ideal*. An *ideal* is a subset that is both a left and right ideal.

A semiring  $R$  satisfies the *descending chain condition of left  $l$ -ideals* (abbreviated DCC) if for each sequence  $R \supseteq L_1 \supseteq L_2 \supseteq \dots$  of left  *$l$ -ideals* there is a positive integer  $n$  such that  $L_n = L_{n+1} = L_{n+2} = \dots$ . This is clearly equivalent to the property that each non-empty set of left  *$l$ -ideals* of  $R$  contains a minimal member.

The definitions of *nilpotent* and *idempotent* elements of a semiring are

the same as in ring theory. The element 0 will be excluded when considering nilpotent or idempotent elements.

The *zeroid* of a semiring  $R$ , as introduced by Bourne and Zassenhaus [2], is  $\{x \in R \mid z+x = z \text{ or } x+z = z \text{ for some } z \in R\}$ . A semiring  $R$  has *right additive cancellation* if  $x+z = y+z$  for  $x, y, z \in R$  implies  $x = y$ . It follows that a left semisubtractive semiring with zero as its zeroid has right additive cancellation.

CONVENTION. We will let  $R$  denote a left semisubtractive semiring with DCC, with zero as its zeroid, and without nilpotent elements.

### 3. Preliminary results

PROPOSITION 1. *Each nonzero left  $l$ -ideal  $A$  of  $R$  contains an idempotent  $x$  with  $A = Rx$ .*

PROOF. By DCC,  $A$  contains a minimal nonzero left  $l$ -ideal  $B$ . For each  $c \neq 0$  in  $B$ ,  $Bc$  is a nonzero left semi-ideal of  $R$  in  $B$ . Let  $Bc^*$  be the left  $l$ -ideal of  $R$  generated by  $Bc$  (see [7]). Since  $Bc^* \subseteq B$ ,  $Bc^* = B$ . Hence  $c \in Bc^*$  which means  $xc = c+yc$  for some  $x, y \in B$ . By left semisubtractivity,  $b+x = y$  or  $x = b+y$  for some  $b \in B$ . If  $x = b+y$ , then  $bc = c$ . If  $b+x = y$ , then  $0 = c+bc$  and hence  $c = c+b(c+bc) = c+bc+b^2c = b^2c$ . In either case there exists  $e \in B$  such that  $c = ec$ .

For some  $d \in B$ ,  $d+e^2 = e$  or  $e^2 = d+e$ . If  $J = \{x \in B \mid xc = 0\}$ , then  $J$  is a left ideal of  $R$ , so that  $J = (0)$ . If  $d+e^2 = e$ , then  $ec+dc = e^2c+dc = ec$  and hence  $d \in J$ . If  $e^2 = d+e$ , again  $d = 0$ . Therefore  $A$  contains an idempotent  $e$ . We now show  $A$  has an idempotent  $x$  such that, if  $y \in A$  with  $yx = 0$ , then  $y = 0$ . For each idempotent  $e \in A$ , let  $M_e = \{x \in A \mid xe = 0\}$  which is a left ideal of  $R$ . Choose idempotent  $x$  of  $A$  such that  $M_x$  is minimal, and suppose  $M_x \neq (0)$ . Now  $M_x$  has an idempotent  $g$ ; note that  $gx = 0$ . For some  $h \in A$ ,  $h+xg = g+x$  or  $xg = h+g+x$ . In the first case, from  $hx+xgx = gx+x^2$  we get  $hx = x$  and similarly  $hg = gh = g$ . Thus  $h^2+xg = h^2+hxg = hg+hx = g+x = h+xg$ , so that  $h^2 = h$ . Clearly  $M_h \subseteq M_x$ ; since  $gx = 0$  and  $gh = g \neq 0$ ,  $M_h \neq M_x$ , a contradiction. For the other case, we have  $hg+g = gh+g = hx+x = ghx = 0$ . Hence for  $k = hxg+g+x$ , we have  $k^2 = k \in A$ . For  $z \in M_k$ ,  $zk = 0$  and hence  $zg+zx = zhxhg+zhxg+zg+zx = zhxhg$ , so that  $zx = zgx+zx^2 = zhxhg = 0$ , meaning  $z \in M_x$ . Thus  $M_k \subseteq M_x$  but  $M_x \neq M_k$ , a contradiction. Therefore  $M_x = (0)$ .

Finally we show  $y = yx$  for each  $y \in A$  and that  $A = Rx$ . If  $y \in A$ , then  $z+y = yx$  or  $y = z+yx$  for some  $z \in A$ . If  $z+y = yx$ , then  $yx = yx^2 = zx+yx$  and hence  $zx = 0$  meaning  $z = 0$ . The other case gives

the same result. Therefore  $y = yx$  for all  $y \in A$ . Since  $Rx \subseteq A = Ax \subseteq Rx$ ,  $A = Rx$ . This completes the proof.

Observe from Proposition 1 that each non-zero left  $l$ -ideal of  $R$  has a right identity. Also, if  $e$  is an idempotent of  $R$ , then  $Re$  [ $eR$ ] is a left [right] ideal. Clearly,  $Re$  is a left semi-ideal. If  $xe, y+xe = ze \in Re$ , then  $ye+xe = ye+xe^2 = ze^2 = ze = y+xe$ , so that  $y = ye \in Re$  and  $Re$  is a left  $l$ -ideal, and similarly  $Re$  is a left  $r$ -ideal. Consequently any left  $l$ -ideal is a left ideal by Proposition 1.

**THEOREM 2.** *If  $A$  is a non-zero ideal of  $R$ , then  $A$  contains an idempotent element  $e$  such that  $A = eR$  and such that  $e$  is the identity of  $A$ .*

**PROOF.** By Proposition 1,  $A$  contains an idempotent element  $e$  such that  $A = Re$ . Let  $B = \{x \in A | ex = 0\}$ . Now  $B$  is a right ideal of  $R$ . Since  $e$  is a right identity of  $A$ ,  $Be = B$ . Since  $B^2 = (Be)B = B(eB) = (0)$ , we have  $B = (0)$ . Letting  $y \in A$ , there exists  $z \in A$  such that  $z+y = ey$  or  $y = z+ey$ . If  $z+y = ey$ , then  $ey = e^2y = ez+ey$ , so that  $ez = 0$  and  $z = 0$ . By the other case  $z = 0$  also. Thus  $y = ey$  for each  $y \in A$ , so that  $e$  is the identity of  $A$ , and  $A = eR$ . This completes the proof.

**COROLLARY.** *The semiring  $R$  contains an identity 1.*

For  $a, b \in R$ ,  $(a+b)(1+1)$  is  $a+b+a+b$  and also  $a+a+b+b$ . Thus  $a+b+a = a+a+b$ . For some  $y \in R$ ,  $y+a+b = b+a$  or  $a+b = y+b+a$ . In the first case  $a+a+b = a+b+a = a+y+a+b$ , so that  $a = a+y$  and  $y = 0$ . Similarly  $y = 0$  in the other case. Consequently  $R$  is a hemiring, that is, a semiring with commutative addition.

The center of  $R$  is the set  $C = \{x \in R | yx = xy \text{ for every } y \in R\}$ . The following proposition is analogous to a theorem in ring theory [4].

**PROPOSITION 3.** *Each idempotent element  $e$  of  $R$  is in  $C$  if and only if  $e$  is the identity for some non-zero ideal of  $R$ .*

We now are able to prove that any left ideal of  $R$  has DCC.

**THEOREM 4.** *If  $A$  is a left ideal of  $R$ , then any left semi-ideal [ideal] of  $A$  is also a left semi-ideal [ideal] of  $R$ .*

**PROOF.** The proof is the same as the proof of the analogous ring theory theorem.

**COROLLARY.** *Any left ideal of  $R$  has DCC.*

It is to be observed from Theorem 4 that, if  $B$  is a right semi-ideal [ideal] of an ideal  $A$ , then  $B$  is a right semi-ideal [ideal] of  $R$ . This fact will be useful to us later in this paper.

#### 4. Central idempotent elements

An idempotent of a hemiring is *central* if it belongs to the center of the hemiring. Further, an idempotent is *semiprimitive* if it is central and if it cannot be expressed as  $u+v$  where  $u$  and  $v$  are central idempotents with  $uv = 0$ . The concepts of *orthogonal* and *pairwise orthogonal* idempotents in hemirings are defined analogously as in rings. At this point two characterizations of semiprimitives can be given.

**PROPOSITION 5.** *A central idempotent  $e$  of  $R$  is semiprimitive if and only if there does not exist a central idempotent  $u \neq e$  such that  $eu = u$  (that is,  $e$  is the only central idempotent of  $R$  in  $eR$ ).*

**PROOF.** Let  $e$  be semiprimitive and suppose there is a central idempotent  $u \neq e$  such that  $eu = u$ . For some  $v \in R$ ,  $v+u = e$  or  $u = v+e$ . If  $v+u = e$ , then  $vu + u = vu + u^2 = eu = u$  and hence  $uv = vu = 0$ . Thus  $v+u = v^2 + u$ , so that  $v^2 = v$ . Clearly  $v \neq 0$  and  $v \in C$ . Consequently,  $v$  is a central idempotent. Since  $v+u = e$  and  $uv = 0$  we have a contradiction to  $e$  being semiprimitive. If  $u = v+e$ , then  $ev + e = v+e$ , so that  $ev = v$ . Also  $uv = 0$ , so that  $0 = (v+e)v = v^2 + v$  and  $0 = u^3v = v^4 + 3v^3 + 3v^2 + v = v^4 + v$  which implies  $v^2 = v^4$ . Since  $v^2 \in C$ ,  $v^2$  is a central idempotent with  $e = v^2 + u$ , a contradiction. The converse follows easily from the contrapositive.

Before giving the second characterization, two definitions are necessary. A hemiring is *simple* if the only ideals it contains are (0) and itself. An ideal of a hemiring is *simple* if it is simple as a hemiring.

**PROPOSITION 6.** *A central idempotent  $e$  of  $R$  is semiprimitive if and only if  $Re$  is simple.*

**PROOF.** If  $e$  is semiprimitive, then it is the only central idempotent of  $Re$  by Proposition 5. Let  $J$  be a non-zero ideal of  $Re$ . By the observation before Theorem 2,  $Re$  is an ideal, so that  $J$  is an ideal of  $R$  by Theorem 4. Thus  $J = Ru$ , where  $u$  is a central idempotent. Since  $u \in J \subseteq Re$ ,  $u = e$ . Hence,  $J = Re$  and  $Re$  is simple. The converse is proved the same as in ring theory.

**THEOREM 7.** *Every central idempotent  $e$  of  $R$  which is not semiprimitive is a sum of a finite number of pairwise orthogonal semiprimitive idempotents.*

**PROOF.** The ideal  $Re$  contains semiprimitive idempotents. Suppose  $u$  and  $v$  are distinct semiprimitive idempotents of  $R$  in  $Re$ . By Proposition 6,  $Ru$  and  $Rv$  are simple ideals. If  $uv \neq 0$ , then  $Ru = Ru \cap Rv = Rv$ . By Proposition 5,  $u = v$  which is a contradiction. Hence  $uv = 0$ .

Let  $M$  be the set of all semiprimitive idempotents of  $R$  in  $Re$ . The elements of  $M$  are pairwise orthogonal. Consider any finite sum of elements of  $M$ , say  $\sum u_i = u$ . Clearly  $u^2 = u = ue = eu$ . For some  $x \in Re$ ,  $x+u = e$  or  $u = x+e$ . If  $x+u = e$ , then  $ux = 0$  and as well  $xu = 0$ ; with this  $x = x^2$ . Clearly  $x \in C$ , so that  $Rx$  is an ideal in  $Re$ . If  $u = x+e$ , then  $ex = x$ . Hence  $x+e = x^2+2x+e$  and  $x^2+x = 0$ . Also  $ux = xu = 0$ , so that  $x^2 = x^2+x^4+x^3 = x^4+x(x+x^2) = x^4$ . Since  $x^2 \in C$ ,  $Rx^2$  is an ideal in  $Re$ . Considering the set  $N$  of all these  $Rx$  or  $Rx^2$ , as the case might be, choose a minimal member of  $N$ . If it is not  $(0)$ , then it is equal to  $Rf$ , where  $f$  is a central idempotent of  $Re$  such that  $e = f + \sum v_i$ ,  $v_i \in M$ , or it is equal to  $Rf^2$ , where  $f^2$  is a central idempotent of  $Re$  such that  $f+e = \sum w_i$ ,  $w_i \in M$ . Considering the first case we observe that, by the corollary to Theorem 4,  $Rf$  contains a minimal non-zero ideal  $K$  which is also an ideal of  $R$ . By Theorem 2,  $K = Rv$  where  $v$  is a central idempotent of  $R$ . Since  $K$  is simple,  $v$  is semiprimitive, and hence  $v \in M$ .

Suppose  $v = v_j$  for some  $j$ . Hence  $v_j \in Rf$  which implies  $v_j = xf$  for some  $x \in R$ . Since  $e = f + \sum v_i$ ,  $xv^j = xfv_j + x(\sum v_i)v_j = v_j + xv_j$ , so that  $v_j = 0$  which is a contradiction. Therefore  $v \neq v_i$  for every  $i$ .

Take  $w = v + \sum v_i$ ; then  $w = y+e$  or  $y+w = e$  for some  $y \in Re$ . As before  $w^2 = w = we = ew$ , and  $w \in C$ . Suppose  $w = y+e$ ; then as before  $y^4 = y^2$ ,  $y^2 \in C$ , and hence  $Ry^2$  is an ideal of  $Re$ . Thus  $v + \sum v_i = y+f + \sum v_i$ , so that  $v = y+f$ . Since  $Rf$  is an ideal,  $y \in Rf$ . Therefore  $Ry^2 \subseteq Rf$ . Assume  $f \in Ry^2$ ; then  $f = ry^2$  for some  $r \in R$ . Since  $vy = 0$ ,  $v = vf = vry^2 = 0$ , a contradiction. Thus  $f \notin Ry^2$  and  $Ry^2 \neq Rf$ , a contradiction.

Suppose then that  $y+w = e$ ; then as before  $y^2 = y$ ,  $y \in C$ , and hence  $Ry$  is an ideal in  $Re$ , and as well  $y+v + \sum v_i = f + \sum v_i$  and  $y+v = f$ , so that  $Ry \subseteq Rf$ . As well  $Ry \neq Rf$ , a contradiction. Consequently, for this case the minimal member of  $N$  has to be  $(0)$ .

Consider now the second case; again  $Rf^2$  contains a non-zero ideal of the form  $Rv$  where  $v$  is a semiprimitive idempotent and hence in  $M$ . If  $v = w_j$  for some  $j$ , then  $w_j \in Rf^2$  and hence  $w_j = xf^2$  for some  $x \in R$ . Thence  $xw_j = x(\sum w_i)w_j = xfw_j + xew_j = xfw_j + xw_j$  and  $xfw_j = 0$ . Thus  $0 = f(xfw_j) = w_j^2 = w_j$ , a contradiction. Therefore  $v \neq w_i$  for every  $i$ .

Take  $w = v + \sum v_i$ ; then  $w = y+e$  or  $y+w = e$  for some  $y \in Re$ . As before  $w^2 = w = we = ew$ , and  $w \in C$ . Suppose  $w = y+e$ ; then as before  $y^4 = y^2$ ,  $y^2 \in C$ , and hence  $Ry^2$  is an ideal of  $Re$ . Thus  $v + \sum v_i = y+f + \sum v_i$ , so that  $v = y+f$ . Since  $Rf$  is an ideal,  $y \in Rf$ . Therefore  $Ry^2 \subseteq Rf$ . Assume  $f \in Ry^2$ ; then  $f = ry^2$  for some  $r \in R$ . Since  $vy = 0$ ,  $v = vf = vry^2 = 0$ , a contradiction. Thus  $f \notin Ry^2$  and  $Ry^2 \neq Rf$ , a contradiction.

Suppose then that  $y+w = e$ ; then as before  $y^2 = y$ ,  $y \in C$ , and hence  $Ry$  is an ideal in  $Re$ , and as well  $y+v+\Sigma v_i = f+\Sigma v_i$  and  $y+v = f$ , so that  $Ry \subseteq Rf$ . As well  $Ry \neq Rf$ , a contradiction. Consequently, for this case the minimal member of  $N$  has to be  $(0)$ .

Consider now the second case; again  $Rf^2$  contains a non-zero ideal of the form  $Rv$  where  $v$  is a semiprimitive idempotent and hence in  $M$ . If  $v = w_j$  for some  $j$ , then  $w_j \in Rf^2$  and hence  $w_j = xf^2$  for some  $x \in R$ . Thence  $xw_j = x(\Sigma w_i)w_j = xfw_j + xew_j = xfw_j + xw_j$  and  $xfw_j = 0$ . Thus  $0 = f(xfw_j) = w_j^2 = w_j$ , a contradiction. Therefore  $v \neq w_i$  for every  $i$ .

Take  $w = v + \Sigma w_i$ ; then  $w = y + e$  or  $y + w = e$  for some  $y \in Re$ . As before  $w^2 = w$ ,  $we = w$ , and  $w \in C$ . Suppose  $w = y + e$ ; then  $y^4 = y^2$ ,  $y^2 \in C$ , and  $Ry^2$  is an ideal of  $Re$ . Since  $f^2 + f = 0$ ,  $v + \Sigma w_i = e + y = f^2 + f + e + y = f^2 + \Sigma w_i + y$  and hence  $v = f^2 + y$ . Since  $Rf^2$  is an ideal,  $y \in Rf^2$ . Therefore  $Ry^2 \subseteq Rf^2$ . By assuming  $f^2 \in Ry^2$ ,  $f^2 = ry^2$ ,  $r \in R$ , and thus, since  $vy = 0$ ,  $v = vf^2 = vry^2 = 0$ , a contradiction. Therefore  $Ry^2 \neq Rf^2$ , a contradiction.

If  $y + w = e$ , then  $y^2 = y$ ,  $y \in C$ , and  $Ry$  is an ideal in  $Re$ . As well  $f + y + v + \Sigma w_i = f + y + w = f + e = \Sigma w_i$  and  $y + v = f^2 + f + y + v = f^2$ , so that  $y \in Rf^2$ . Thus  $Ry \subseteq Rf^2$  and  $Ry \neq Rf^2$ , a contradiction. Consequently, for this case the minimal member of  $N$  has to be  $(0)$ . Therefore  $e$  is a finite sum of semiprimitive idempotents, as we wanted to prove.

## 5. Direct sums and a structure theorem

The concept of *direct sum* in hemirings is the same as in ring theory. Hence we have the following theorem which is proved the same as in ring theory:

**PROPOSITION 8.** *If  $A_1, \dots, A_m$  are distinct simple ideals of  $R$  and if  $A = A_1 + \dots + A_m$ , then  $A$  is their direct sum.*

We conclude the section with the main theorem of the paper. It is a generalization to hemirings of a well-known structure theorem discussed by Artin, Nesbitt, and Thrall [1].

**THEOREM 9.** *The hemiring  $R$  has only a finite number of non-zero simple ideals and is their direct sum.*

**PROOF.** By the corollary to Theorem 2,  $R$  contains an identity 1 which is a central idempotent. If 1 is semiprimitive, then  $R$  is simple by Proposition 6 and the proof is complete. Assume 1 is not semiprimitive. By Theorem 7,  $1 = \Sigma e_i$  where the  $e_i$  are pairwise orthogonal semiprimitive idempotents. Since  $R = R \cdot 1 = R(\Sigma e_i) \subseteq \Sigma Re_i \subseteq R$ ,  $R = \Sigma Re_i$ . By Proposition 6, each  $Re_i$  is a simple ideal. By Proposition 8,  $R$  is the direct sum of  $Re_i$ .

Let  $I$  be a non-zero simple ideal of  $R$ . If  $RIR = (0)$ , then  $I^3 = (0)$ , a contradiction. Hence  $I = RIR$ . Thus  $I = RIR \subseteq I(\sum Re_i) \subseteq \sum IRe_i \subseteq I$ , so that  $I = \sum IRe_i$ . Some  $IRe_i \neq (0)$  since  $I \neq (0)$ ; hence  $I \cap Re_i \neq (0)$  which implies  $I = I \cap Re_i = Re_i$ . Therefore,  $R$  has only a finite number of simple ideals and the proof is complete.

## REFERENCES

E. ARTIN, C. J. NESBITT and R. M. THRALL

[1] Rings with minimum condition (University of Michigan Publication in Mathematics, no. 1, 1944.

S. BOURNE and H. ZASSENHAUS

[2] On the semiradical of a semiring, Proc. Nat. Acad. Sci. U.S.A., vol. 44 (1958), pp. 907-914.

N. J. DIVINSKY

[3] Rings and Radicals, Mathematical Expositions no. 14, Toronto: University of Toronto Press, 1965.

A. FORSYTHE and N. MCCOY

[4] On the commutativity of certain rings, Bull. Amer. Math. Soc., vol. 52 (1946), pp. 523-526.

I. N. HERSTEIN

[5] Noncommutative Rings, The Carus Mathematical Monographs no. 15, The Mathematical Association of America, 1968.

A. KERTÉSZ

[6] Vorlesungen über Artinsche Ringe, Akad. Kiado, 1968.

J. R. MOSHER

[7] Generalized quotients of hemirings, Compositio Mathematica, vol. 22 (1970), pp. 275-281.

F. SZÁSZ

[8] Über Ringe mit Minimalbedingung für Hauptideale, III, Acta Math., 14 (1963), pp. 447-461

(Oblatum: 9-III-70)

Prof. J. R. Mosher  
Texas A & M University  
Department of Mathematics  
COLLEGE STATION, Texas 77843  
U.S.A.