

# COMPOSITIO MATHEMATICA

ERNST-ULRICH GEKELER

## **Quasi-periodic functions and Drinfeld modular forms**

*Compositio Mathematica*, tome 69, n° 3 (1989), p. 277-293

<[http://www.numdam.org/item?id=CM\\_1989\\_\\_69\\_3\\_277\\_0](http://www.numdam.org/item?id=CM_1989__69_3_277_0)>

© Foundation Compositio Mathematica, 1989, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Quasi-periodic functions and Drinfeld modular forms

ERNST-ULRICH GEKELER

*The Institute for Advanced Study, Princeton, NJ 08540; present address:  
Max-Planck-Institut für Mathematik, Gottfried-Claren-Strasse, 26, D-5300 Bonn 3, FRG*

Received 7 March 1988; accepted 14 June 1988

### 1. Introduction

Let  $E = \mathbb{C}/\Lambda$  be a complex elliptic curve with period lattice  $\Lambda = \mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$ , and let  $p(z)$ ,  $\zeta(z)$  be the associated Weierstrass functions. Then  $p(z)$  and  $\zeta(z)$  are meromorphic on  $\mathbb{C}$  and  $p(z) = -\zeta'(z)$  is  $\Lambda$ -periodic, whereas  $\zeta(z)$  satisfies

$$\zeta(z + \omega_i) = \zeta(z) + \eta_i \quad (i = 1, 2) \quad (1.1)$$

with certain  $\eta_i \in \mathbb{C}$ . Between the periods of first kind  $\omega_i$  and that of second kind  $\eta_i$ , *Legendre's relation*

$$\omega_1\eta_2 - \omega_2\eta_1 = 2\pi i \quad (1.2)$$

holds (suppose  $\text{im}(\omega_1/\omega_2) > 0$ ). Further, the  $\eta_i$ , considered as functions on the complex upper half-plane, are closely related to the “false Eisenstein series of weight two,” i.e., the logarithmic derivative of the discriminant function  $\Delta(\omega)$  (See [8], App. 1 for a precise statement).

(1.3) Let now  $K$  be a function field in one variable having the field  $\mathbb{F}_q$  with  $q$  elements as exact constant field ( $q = \text{some power of the prime } p$ ). Fix a place  $\infty$  of  $K$  of degree  $\delta$ , say, and put

$$A = \{f \in K \mid f \text{ has poles at most at } \infty\}.$$

On  $A$ , we have the degree function  $\deg a = \delta$  times (pole order of  $a$  at  $\infty$ ). Further, let  $K_\infty$  be the completion of  $K$  at  $\infty$  with its normalized absolute value  $|\cdot|$ , and  $C$  the completion of an algebraic closure  $\bar{K}_\infty$  with respect to the unique extension, also named  $|\cdot|$ , of  $|\cdot|$  to  $\bar{K}_\infty$ . Note that  $C$  is again algebraically closed, i.e., is the minimal complete algebraically closed field containing  $K_\infty$ . The basic example (in fact, the only one considered later on)

is given by

$$\begin{aligned}
 K &= \mathbb{F}_q(T) \text{ the field of rational functions,} \\
 \text{“}\infty\text{”} &= \text{the place at infinity, so} \\
 A &= \mathbb{F}_q[T] \text{ is the polynomial ring, and} \\
 K_\infty &= \mathbb{F}_q((T^{-1})) \text{ the field of finite-tailed Laurent series in } T^{-1}.
 \end{aligned}
 \tag{1.4}$$

(1.5) It is well known that there is a deep analogy between the arithmetic of  $\mathbb{Q}$  and that of  $K$ , where  $A$  (resp.  $K_\infty$ ,  $C$ ) replaces the ring  $\mathbb{Z}$  of integers (resp.  $\mathbb{R}$ ,  $\mathbb{C}$ ). Roughly, the role of complex analysis in number theory is played by rigid analysis over the “complex” field  $C$ . (See [5] and the references given there.)

As explained below, the classical theory of elliptic modular forms corresponds to the theory of rank two Drinfeld modules. By following ideas of Pierre Deligne, as elaborated by Greg Anderson and Jing Yu, we will define quasi-periodic functions, periods of second kind, and de Rham cohomology for a Drinfeld module. Specializing to the case  $A = \mathbb{F}_q[T]$  and rank = two, where the theory of modular forms is sufficiently well developed, we obtain a non-vanishing result (Cor. 6.3) for the associated determinant which replaces (1.2). For weight reasons, the latter will not be a constant as in (1.2), but a meromorphic modular form of negative weight. Applying the results of [6], we get a very precise description of that form, in particular its expansion at  $\infty$ . As in the classical case, the maps  $\omega \mapsto \eta_i(\omega)$  ( $i = 1, 2$ ) which to each  $\omega$  associate its periods of second kind are related to a “false Eisenstein series” (Thm. 7.10). Finally, we calculate the Gauss-Manin connection  $\nabla$  on  $H_{\text{DR}}^*(\phi)$ , where the Drinfeld module  $\phi$  varies over the “upper half-plane”  $\Omega$  (Thm. 8.8). After a suitable normalization,  $\nabla$  is defined for the Tate–Drinfeld module, i.e., by a matrix whose entries are power series over  $A$ .

Certainly, some of our methods and results may be widely generalized. (For the “de Rham cohomology” introduced in section 3, see the forthcoming articles [2] and [9].)

For example, the local system over the moduli scheme defined by  $H_{\text{DR}}^*$ , which is rather simple in our case, will be of high interest if  $A$  is general. Nevertheless, the relation with modular forms presented here seems to be a feature special to the polynomial case  $A = \mathbb{F}_q[T]$  where one disposes of a “natural” basis for  $H_{\text{DR}}^*$ .

**2. Review of Drinfeld modules**

(See [3] or [5] for proofs.)

Let  $\Lambda$  be a rank  $r$  lattice in  $C$ , i.e., a projective, discrete  $A$ -submodule of  $C$  of  $A$ -rank  $r \in \mathbb{N}$ . Put

$$e_\Lambda(z) = z \prod_{\lambda \in \Lambda} (1 - z/\lambda), \tag{2.1}$$

where as usual,  $\prod$  will denote the product over the non-zero elements of a lattice. The product is easily seen to converge for  $z \in C$ ; it defines an entire surjective mapping  $e_\Lambda: C \rightarrow C$  which is additive and  $\mathbb{F}_q$ -linear. Further,  $e_\Lambda$  is  $\Lambda$ -periodic with kernel  $\Lambda$ .

Let  $C\{\tau\}$  be the non-commutative ring of polynomials over  $C$  of the form  $\sum l_i \tau^i$ , where  $\tau$  corresponds to  $X^q$ , and the multiplication  $f \circ g$  is defined by inserting  $g(X)$  into  $f(X)$ . Elements of  $C\{\tau\}$  will be referred to as  $q$ -additive polynomials. Note that  $\tau^0 = X$  is the identity of  $C\{\tau\}$ . We may consider

$$e_\Lambda(X) = \sum \alpha_i X^{q^i} = \sum \alpha_i \tau^i$$

as an element of the ring  $C\{\{\tau\}\}$  of “power series” in  $\tau$  with certain growth conditions on the coefficients  $\alpha_i$ .

Next, for  $a \in A$ , let  $\phi_a^\Lambda \in C\{\tau\}$  be defined by the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Lambda & \longrightarrow & C & \xrightarrow{e_\Lambda} & C \longrightarrow 0 \\ & & \downarrow a & & \downarrow a & & \downarrow \phi_a^\Lambda \\ 0 & \longrightarrow & \Lambda & \longrightarrow & C & \xrightarrow{e_\Lambda} & C \longrightarrow 0. \end{array} \tag{2.2}$$

A closer look shows  $\phi_a^\Lambda$  to be of the form

$$\phi_a^\Lambda = l_0 \tau^0 + \dots + l_{rd} \tau^{rd}, \tag{2.3}$$

where  $l_i \in C$ ,  $d = \deg a$ ,  $l_0 = a$ ,  $l_{rd} \neq 0$ . Further,  $a \mapsto \phi_a$  defines a ring homomorphism  $\phi^\Lambda: A \rightarrow C\{\tau\}$ .

(2.4) A ring homomorphism  $\phi: A \rightarrow C\{\tau\}$ ,  $a \mapsto \phi_a$  satisfying the conditions of (2.3) will be called a *Drinfeld module* over  $C$  of rank  $r$ . Given  $\phi$ , one may recover  $e_\Lambda$  and  $\Lambda$  (see, e.g., [5]). This sets up a bijection  $\Lambda \mapsto \phi^\Lambda$  between the set of rank  $r$  lattices in  $C$  and the set of rank  $r$  Drinfeld modules over  $C$ . By means of  $\phi$ , the additive group  $G_a|C$  gets another structure as an  $A$ -module which differs from the canonical one. Two lattices,  $\Lambda, \Lambda'$ , define

isomorphic Drinfeld modules if and only if they are similar, i.e., there exists  $c \in C^*$  such that  $\Lambda = c\Lambda$ .

(2.5) From now on, let us restrict to lattices  $\Lambda$  which are free over  $A$ . Let  $\Gamma$  be the group  $GL(r, A)$ , considered as a discrete subgroup of the Lie group  $GL(r, K_\infty)$  that operates on the projective space  $\mathbb{P}^{r-1}(C)$  by fractional linear transformations. We define

$$\Omega^r = \{(z_1 : \dots : z_r) \in \mathbb{P}^{r-1}(C) \mid z_1, \dots, z_r \text{ } K_\infty\text{-linearly independent}\}.$$

Then  $\Omega^r$  carries the structure of an  $r-1$ -dimensional rigid analytic manifold over  $C$ , and the induced action of  $\Gamma$  on  $\Omega^r$  is analytic. We have the bijection

$$\begin{aligned} \Gamma \backslash \Omega^r &\cong \{\text{Similarity classes of free lattices of rank } r\} \\ \mathbf{z} = (z_1 : \dots : z_r) &\mapsto \text{class of } \Lambda_{\mathbf{z}} = Az_1 + \dots + Az_r, \end{aligned}$$

which in fact gives an analytic description of the moduli scheme for Drinfeld modules of rank  $r$ .

2.6 EXAMPLE: Let  $A = \mathbb{F}_q[T]$ . Any rank  $r$  Drinfeld module is given by

$$\phi_T = T\tau^0 + \dots + l_r\tau^r = TX + \dots + l_rX^{q^r},$$

where  $l_r \neq 0$ . If further  $r = 2$ , then  $\Omega = \Omega^2 = \mathbb{P}^1(C) \backslash \mathbb{P}^1(K_\infty) = C \backslash K_\infty$ , and  $\Gamma = GL(2, A)$  acts by  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} (z) = (az + b)/(cz + d)$ . Any rank two  $\phi$  is given by  $\phi_T = T\tau^0 + g\tau + \Delta\tau^2$ , where  $g, \Delta \in C, \Delta \neq 0$ . Letting vary  $\phi$ ,  $g$  and  $\Delta$  are modular forms for  $\Gamma$ , see (4.9).

### 3. Quasi-periodic functions

(This section is developed entirely from the ideas of Greg Anderson, Pierre Deligne, and Jing Yu.)

Let  $\Lambda$  be a rank  $r$  lattice in  $C$  with exponential function  $e = e_\Lambda$  and Drinfeld module  $\phi = \phi^\Lambda$ . Let  $a \in A$  be non-constant and  $\partial \in \tau C\{\tau\}$  an element of  $C\{\tau\}$  without constant term. The difference equation

$$\partial(e(X)) = F(aX) - aF(X) \tag{3.1}$$

has a solution  $F \in C\{\{\tau\}\}$  which is well-defined up to adding a multiple of  $\tau^0 = X$ . Thus, assuming the  $\tau^0$ -coefficient to be zero (*which we will always assume in what follows*), there exists a unique solution  $F \in C\{\{\tau\}\}$ . Comparing

coefficients, one easily verifies  $F$  to define an entire function  $C \rightarrow C$ , also called  $F$ .

(3.2) Let  $C_\phi\{\tau\}$  be the  $A$ -bimodule  $C\{\tau\}$ , where the left action is as usual, and the right action of  $a \in A$  is right multiplication with  $\phi_a$ .

3.3. LEMMA: (i) Let  $\partial^{(0)}: A \rightarrow C\{\tau\}$   $a \mapsto \partial_a^{(0)}$  be defined by  $\partial_a^{(0)} = \phi_a - a\tau^0$ . Then  $\partial_{ab}^{(0)} = a\partial_b^{(0)} + \partial_a^{(0)} \circ \phi_b$ , i.e.,  $\partial^{(0)}$  is a derivation  $A \rightarrow C_\phi\{\tau\}$ .

(ii) Let  $\partial: a \mapsto \partial_a$  be any  $\mathbb{F}_q$ -linear derivation of  $A$  into  $C_\phi\{\tau\}$  with values in  $\tau C\{\tau\}$ . Then the unique solution  $F_\partial \in \tau C\{\{\tau\}\}$  of

$$\partial_a(e(X)) = F(aX) - aF(X) \tag{*}$$

does not depend on  $a$ , provided that  $a$  is non-constant. (For  $a = \text{constant}$ ,  $\partial_a = 0 = F$ .)

Proof: (i) is trivial:  $\partial_{ab}^{(0)} = \phi_{ab} - ab\tau^0 = a(\phi_b - b\tau^0) + (\phi_a - a\tau^0) \circ \phi_b$  since  $\phi_{ab} = \phi_a \circ \phi_b$ . (ii) Let  $F_a$  be the unique solution (without  $\tau^0$ -term) of (\*). From the derivation relations, it is straightforward to show

- (1)  $F_{ca} = F_a$  ( $c \in \mathbb{F}_q^*$ );
- (2)  $F_a = F_b \Rightarrow F_{a+b} = F_a = F_b$ ;
- (3)  $F_a = F_b \Rightarrow F_{ab} = F_a = F_b$ .

Clearly, (1), (2), (3) imply the assertion for  $A$  a polynomial ring  $A = \mathbb{F}_q[a]$ . Since any  $A$  is an integral extension of a polynomial ring, (1), (2), (3) give the result.

(3.4) In the following, derivations will always assumed to be  $\mathbb{F}_q$ -linear. Let  $D(\phi)$ ,  $D_i(\phi)$ ,  $D_{si}(\phi)$  respectively be the  $A$ -bimodule of derivations, inner derivations, strictly inner derivations respectively from  $A$  to  $C_\phi\{\tau\}$  with values in  $\tau C\{\tau\}$ . (Some  $\partial \in D(\phi)$  is called an *inner (strictly inner) derivation* if there exists  $m \in C\{\tau\}$  ( $m \in \tau C\{\tau\}$ ) such that for  $a \in A$ ,  $\partial_a = m \circ \phi_a - am$  holds.) The function  $F_\partial$  associated with some  $\partial \in D(\phi)$  will be called quasi-periodic for  $\Lambda$ .

3.5. Remarks: (i)  $F: C \rightarrow C$  to be quasi-periodic implies  $F(z + \lambda) = F(z) + F(\lambda)$  ( $\lambda \in \Lambda$ ), where  $F$  restricted to  $\Lambda$  is  $A$ -linear. This is analogous with (1.1).

(ii) If  $\partial = \partial^{(0)}$  is the inner derivation defined in (3.3i), then  $F_\partial(z) = e(z) - z$  which thus is quasi-periodic.

(iii) Let  $\partial$  be the strictly inner derivation associated with  $m \in \tau C\{\tau\}$ . Then  $F_\partial(z) = m(e(z))$  which in fact is periodic and vanishes on  $\Lambda$ .

3.6. EXAMPLE: Let  $A = \mathbb{F}_q[T]$  as in (1.4). Some  $\partial \in D(\phi)$  is given by  $\partial_T \in \tau C\{\tau\}$  which may be arbitrarily prescribed. The corresponding  $F_\partial$  is

the solution of

$$\partial_T(e(z)) = F(Tz) - TF(z).$$

Since  $\phi_T = T\tau^0 + \dots + l_r\tau^r$ ,  $l_r \neq 0$ , it is easy to see that  $H_{\text{DR}}^*(\phi) = D(\phi)/D_{\text{si}}(\phi)$ , considered as a left  $C$ -vector space, has dimension  $r$ . A basis is given by the  $r$  derivations  $\partial^{(i)}$  defined by  $\partial_T^{(i)} = \tau^i$ ,  $0 < i \leq r$ , or rather  $\{\partial^{(i)} \mid 0 \leq i < r\}$ , where  $\partial_T^{(0)} = \phi_T - T\tau^0$ . A generalization of this fact for arbitrary  $A$ , as well as an interpretation of  $H_{\text{DR}}^*(\phi)$  as *de Rham cohomology group* of the Drinfeld module  $\phi$ , will be given in [2], see also [1]. In that context, the map  $\Lambda \times H_{\text{DR}}^*(\phi) \rightarrow C: (\omega, \partial) \mapsto F_{\partial}(\omega)$  will appear as some sort of “path integration.”

#### 4. Relations with modular forms

From now on, we assume the situation of (1.4), (2.6), (3.6), i.e.,  $A = \mathbb{F}_q[T]$ . Let  $\omega = (\omega_1, \dots, \omega_r) \in C^r$ , the  $\omega_i$  linearly independent over  $K_\infty$ , and  $\Lambda_\omega$  the lattice  $A\omega_1 + \dots + A\omega_r$ . Let  $e = e_\Lambda$ , and  $\phi = \phi^\Lambda$  the associated Drinfeld module. We further put for  $0 \leq i < r$   $F^{(i)} = F_{\partial^{(i)}}$ . Thus

$$\begin{aligned} F^{(0)}(z) &= e(z) - z, \quad \text{and for } 0 < i < r, \\ e^{\partial^{(i)}}(z) &= F^{(i)}(Tz) - TF^{(i)}(z). \end{aligned} \tag{4.1}$$

We use a subscript  $\omega$  to indicate the dependence of all the data introduced from  $\omega$ , thus  $e = e_\omega$ ,  $\phi = \phi_\omega$ ,  $F^{(i)} = F_\omega^{(i)}$ . Now for any lattice  $\Lambda$ ,  $z$  and  $c \in C$ ,  $c \neq 0$ , we have

$$\begin{aligned} e_{c\Lambda}(cz) &= ce_\Lambda(z). \\ F_{c\omega}^{(i)}(cz) &= c^{\partial^{(i)}} F_\omega^{(i)}(z). \end{aligned} \tag{4.2}$$

Note that for  $i > 0$ ,  $F_\omega^{(i)}(\omega_j)$  is the analogue of a *period of second kind* on an elliptic curve.

We are interested in the *Legendre determinant*

$$L(\omega) = \det (F_\omega^{(i)}(\omega_j)) \quad (0 \leq i < r, 0 < j \leq r).$$

By the above,

$$L(c\omega) = c^k L(\omega) \quad \text{with } k = 1 + q + \dots + q^{r-1}. \tag{4.3}$$

4.4. LEMMA: For  $\gamma \in GL(r, A)$ , we have

$$L(\gamma\omega) = (\det \gamma)L(\omega).$$

*Proof:*  $F_\omega^{(i)}$  depends only on the lattice  $\Lambda_\omega = \Lambda_{\gamma\omega}$ . Thus,  $L(\gamma\omega) = \det(F_\omega^{(i)}((\gamma\omega)_j)) = (\det \gamma) \det(F_\omega^{(i)}(\omega_j)) = (\det \gamma)L(\omega)$  since  $F^{(i)}|_\Lambda$  is  $A$ -linear.

We now finally restrict to the case  $r = 2$  where we will give a description of the map  $\omega \mapsto L(\omega)$ . By (4.3), it suffices to consider  $\omega$  of the form  $\omega = (\omega, 1)$  with  $\omega \in \Omega = C \setminus K_\infty$ . The function  $L(\omega) = L((\omega, 1))$  now satisfies

(4.5) (i)  $L(\gamma\omega) = (\det \gamma)(c\omega + d)^{-q-1}L(\omega)$  ( $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma = GL(2, A)$  acts by  $\gamma\omega = (a\omega + b)/(c\omega + d)$ ), and (ii)  $L$  is holomorphic on  $\Omega$  in the rigid analytic sense.

Here, (i) is just a repetition of (4.3) and (4.4), whereas (ii) had to be shown. Since we did not give a description of the analytic structure on  $\Omega$  (which may be found e.g., in [7]), we omit the simple proof. But compare (5.3).

(4.6) In order to make a meromorphic modular form out of  $L(\omega)$ , we have to introduce cusp conditions. Let  $\varrho$  the rank one Drinfeld module (sometimes called the *Carlitz module*) defined by  $\varrho_T = T\tau^0 + \tau = TX + X^q$ . Via (2.4), it corresponds to the rank one lattice  $L = \bar{\pi}A$  with some number  $\bar{\pi} \in C$ . Hopefully, the double occurrence of the symbol  $L$  will cause no confusion. *Note:*  $\bar{\pi}$  is only defined up to a  $(q - 1)$ -st root of unity. We choose one  $\bar{\pi}$  and fix it for what follows. It is similar to the period  $2\pi i$  of the exponential function. Several additive and multiplicative expressions are known for  $\bar{\pi}$  ([6], (4.9)–(4.11)). In particular, its absolute value equals  $q^{q/(q-1)}$ . Let  $e_L$  be the lattice function associated with  $L$ , and  $t(z) = 1/e_L(\bar{\pi}z)$ . Obviously,  $t(z)$  is  $A$ -periodic. In our framework, it plays the part of  $q(z) = e^{2\pi iz}$  as a uniformizer at  $\infty$ .

(4.7) For  $z \in C$ , we define the “imaginary part”  $|z|_i = \inf |z - x|$ , where  $x$  runs through  $K_\infty$ . Also, let  $\Omega_s = \{z \in \Omega \mid |z|_i \geq s\}$ , where  $s \in \mathbb{R}$ . Then we have:

4.8. LEMMA ([6], LEMMA 5.5): For  $|z|_i > 1$ ,  $|e_L(\bar{\pi}z)|$  is a monotonically increasing function of  $|z|_i$ . Namely, if  $|z| = |z|_i = q^{d-\varepsilon}$  with some  $d \in \mathbb{N}$ ,  $0 \leq \varepsilon < 1$ , then  $\log_q |e_L(\bar{\pi}z)| = q^d(q/(q - 1) - \varepsilon)$ . There exists a constant



$c_0 > 1$  such that  $|z|_i \leq \log_q |e_L(\bar{\pi}z)| \leq c_0 |z|_i$ . Further,  $t(z)$  defines an analytic isomorphism of  $A \setminus \Omega_s$  ( $s > 1$ ) with some pointed ball  $\{z \in C \mid |z| \leq r\} - \{0\}$ .

4.9. DEFINITION: A function  $f: \Omega \rightarrow C$  is a holomorphic *modular form* of weight  $k \in \mathbb{Z}$  and type  $m \in \mathbb{Z}/(q - 1)$  if the following conditions hold:

- (i)  $f(\gamma z) = (\det \gamma)^{-m} (cz + d)^k f(z) \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ ;
- (ii)  $f$  is holomorphic;
- (iii)  $f$  is holomorphic at  $\infty$ .

The last point signifies that for  $|z|_i \gg 0$ ,  $f(z)$  has a series expansion  $f(z) = \sum_{i \geq 0} a_i t^i(z)$ , which, due to (4.8), corresponds to the cusp condition in the elliptic modular case. Replacing “holomorphic” by “meromorphic” and admitting Laurent series around  $\infty$  with a finite number of terms of negative order, we may define meromorphic modular forms. For a further discussion, see [5] or [6]. In the next sections, we will calculate the expansion of  $L(\omega)$  around  $\infty$ , thereby verifying it as meromorphic modular of weight  $-q - 1$  and type  $-1$ . For that purpose, we will need the following series of polynomials: for  $0 \neq a \in A$  of degree  $d$ , let  $\varrho_a(X) = aX + \dots \in A[X]$  be the polynomial derived from the Carlitz module. We have  $\deg(\varrho_a(X)) = q^d = |a|$ , and its leading coefficient agrees with that of  $a$  as a polynomial in  $T$ . Put  $f_a(X) = \varrho_a(X^{-1})X^{|a|}$ . Then for example,  $f_1(X) = 1, f_T(X) = 1 + TX^{q-1}, f_{T^2}(X) = 1 + (T + T^q)X^{q^2-q} + T^2X^{q^2-1}$ .

### 5. Behavior of $L$ at infinity

In all that follows,  $A = \mathbb{F}_q[T]$ , and we are considering the rank two case only.

For  $\omega \in \Omega$  and  $i = 0, 1$  let  $F_\omega^{(i)}(z) = F_{(\omega, 1)}^{(i)}(z)$  as defined in (4.1). Also, let  $e_\omega, \Lambda_\omega, \dots$  be the objects associated with  $\omega = (\omega, 1)$ . We are going to investigate the function

(5.1)  $\omega \mapsto L(\omega) = F_\omega^{(1)}(\omega) - \omega F_\omega^{(1)}(1)$  on  $\Omega$  which, in view of  $F^{(0)}(z) = e(z) - z$ , is the inhomogenous expression of  $L(\omega) = L((\omega, 1))$ . Actually, we will show

5.2. THEOREM:  $\lim_{|\omega|_i \rightarrow \infty} t(\omega)L(\omega) = \bar{\pi}^{-q}$ ,

the proof of which will occupy this section. Let us first treat the function  $F_\omega(z) = F_\omega^{(1)}(z)$  (there is no further need for the superscript (1)). From (4.1)

we get

$$\begin{aligned}
 F_\omega(z) &= e_\omega^q(z/T) + TF_\omega(z/T) \\
 &= e_\omega^q(z/T) + Te_\omega^q(z/T^2) + T^2F_\omega(z/T^2) \\
 &\vdots
 \end{aligned}$$

Looking at the definition of  $e_\omega$ , we see  $|e_\omega(z)| = |z|$  whenever  $|z| < \inf |a\omega + b| = \inf (|\omega|_i, 1)$ . Thus  $(0, 0) \neq (a, b) \in A^2$

$$|T^i e_\omega^q(z/T^{i+1})| = |z|^q q^{i-q(i+1)} \text{ for } i \geq 0$$

which shows the convergence (pointwise, in fact locally uniformly in  $z$  and in  $\omega$ ) of the series

$$F_\omega(z) = \sum_{i \geq 0} T^i e_\omega^q(z/T^{i+1}). \tag{5.3}$$

We are interested in the behavior of  $F_\omega(\omega)$  and  $F_\omega(1)$  for  $|\omega|_i \geq 0$ . The above shows:

$$F_\omega(1) \text{ is bounded on } \Omega_1 = \{\omega \in \Omega \mid |\omega|_i \geq 1\}. \tag{5.4}$$

In order to handle  $F_\omega(\omega)$ , we have to control the different terms in (5.3).

(5.5) Let  $u = (u_1, u_2) \in K^2 \setminus A^2$ , where for  $i = 1, 2, u_i = s_i/n$ , the  $s_i$  and  $n$  in  $A$ ,  $\deg s_i < \deg n$ , and  $n$  monic. Let further  $\Gamma(n)$  be the congruence subgroup  $\{\gamma \in \Gamma \mid \gamma \equiv 1 \pmod n\}$  of  $\Gamma$ . In [4], we studied the functions

$$e_u(\omega) = e_\omega(u_1\omega + u_2)$$

on  $\Omega$  which are meromorphic modular forms of weight  $-1$  for  $\Gamma(n)$ . Their behavior at  $\infty$  is described using the uniformizer  $t_{1/n}(\omega) = t(\omega/n) = 1/e_L(\bar{\pi}\omega/n)$ . It is related with  $t(\omega)$  by

$$t = t_{1/n}^{|n|} / f_n(t_{1/n}). \tag{5.6}$$

(In loc. cit.,  $t_{1/n}$  had been labelled by  $t_n$  which, unfortunately, is inconsistent with notations in [6] and in Sections 6–8.) Now,  $e_u$  has a convergent product

expansion given by (*loc. cit.* (2.1)):

$$e_u(\omega) = \bar{\pi}^{-1} \frac{f_{s_1}(t_{1/n}) + \zeta}{t_{1/n}^{s_1}} \prod_{c \in A} \frac{f_{nc-s_1}(t_{1/n}) - \zeta t_{1/n}^{nc}}{f_{nc}(t_{1/n})}. \tag{5.7}$$

Here  $\zeta = \zeta_{u_2} = e_L(\bar{\pi}u_2)$  is an  $n$ -division point (= root of  $\varrho_n(X)$ ) of the Carlitz module  $\varrho$ . The factor corresponding to  $0 \neq c \in A$  in (5.7) equals  $e_L(\bar{\pi}((c - u_1)\omega - u_2))/e_L(\bar{\pi}c\omega)$ . Since  $|(c - u_1)\omega - u_2|_i = |c||\omega|_i$ , (4.8) implies its absolute value to be 1, provided that  $|\omega|_i > 1$ . Due to (5.3), we may write

$$F_\omega(\omega) = \sum_{i \geq 0} T^i e_{u(i)}^q(\omega) \quad \text{with } u(i) = (T^{-i-1}, 0). \tag{5.8}$$

5.9. LEMMA: For  $i > 0$ ,  $\lim t(\omega)T^i e_{u(i)}^q(\omega) = 0$  uniformly in  $i$ , where the lim is over those  $\omega \in \Omega$  that satisfy  $|\omega| = |\omega|_i$ , and  $|\omega| \rightarrow \infty$ .

*Proof:* Put ad hoc  $t_i = t_{1/n}$  where  $n = T^i$ , and let  $|\omega|_i = |\omega| > 1$ . Then by (5.7),  $e_{u(i)}(\omega) = \bar{\pi}^{-1} t_{i+1}^{-1}(\omega) \times \text{unit}$  since  $u(i) = (T^{-i-1}, 0)$ ,  $s_1 = 1$ , and  $u_2 = 0 = \zeta$ . Therefore,

$$\begin{aligned} t(\omega)T^i e_{u(i)}^q(\omega) &= t(\omega)T^i \bar{\pi}^{-q} t_{i+1}^{-q}(\omega) \times \text{unit} \\ &= \bar{\pi}^{-q} T^i e_L^q(\bar{\pi}\omega/T^{i+1})/e_L(\bar{\pi}\omega) \times \text{unit}. \end{aligned}$$

Using (4.8), one easily sees this tends to zero with  $|\omega| \rightarrow \infty$ . In order to get uniformity in  $i$ , we have to use (4.8) in full detail. Let  $|\omega| = |\omega|_i = q^{d-\varepsilon}$  where  $d \in \mathbb{N}$  and  $0 \leq \varepsilon < 1$ . We compute the  $q$ -logarithm of

$$(*) = |T^i e_L^q(\bar{\pi}\omega/T^{i+1})/e_L(\bar{\pi}\omega)|.$$

By (4.8)  $\log_q |e_L(\bar{\pi}\omega)| = q^d(q/(q-1) - \varepsilon)$ .

For the numerator, we distinguish the cases

- (a)  $|\omega/T^{i+1}| > 1$ , i.e.,  $i + 1 < d$
- (b)  $|\omega/T^{i+1}| \leq 1$ , i.e.,  $i + 1 \geq d$ .

In case (a),  $\omega' = \omega/T^{i+1}$  satisfies  $|\omega'| = |\omega'|_i = q^{d-i-1-\varepsilon}$ , thus

$$\log_q |e_L(\bar{\pi}\omega')| = q^{d-i-1}(q/(q-1) - \varepsilon).$$

We get the estimate

$$\begin{aligned} \log_q(*) &= q^{d-i}(q/q - 1) - \varepsilon + i - q^d(q/(q - 1) - \varepsilon) \\ &< d - 1 - q^{d-1} =: h_1(d) \end{aligned}$$

since  $i < d - 1$  and  $\varepsilon < 1$ .

In case (b),  $|e_L(\bar{\pi}\omega/T^{i+1})| = |\bar{\pi}\omega/T^{i+1}|$  as an immediate consequence of the definition of  $e_L$ . Thus

$$\log_q(*) = q \log_q |\bar{\pi}| + q(d - i - 1 - \varepsilon) + i - q^d(q/(q - 1) - \varepsilon).$$

Since  $\log_q |\bar{\pi}| = q/(q - 1)$ , this becomes

$$\begin{aligned} q(q/(q - 1) - 1 - \varepsilon) + qd + (1 - q)i - q^d(q/(q - 1) - \varepsilon) \\ \leq q/(q - 1) + qd - q^d/(q - 1) =: h_2(d). \end{aligned}$$

Clearly,  $\sup(h_1(d), h_2(d))$  tends to  $-\infty$  if  $d$  increases, which establishes the result.

Now we are able to prove Theorem 5.2: since  $L(\omega)$  is  $A$ -periodic, we may restrict the limit  $\lim_{|\omega|_i \rightarrow \infty}$  (simply denoted by “lim”) to those  $\omega$  satisfying  $|\omega| = |\omega|_i$ . Then

$$\lim t(\omega)L(\omega) = \lim t(\omega) \sum_{i \geq 0} T^i e_{u(i)}^q(\omega) - \lim t(\omega)\omega F_\omega(1).$$

The second limit vanishes by virtue of (4.8), (5.4), and the assumption  $|\omega| = |\omega|_i$ . As to the first, since the limits in (5.9) are uniform, we may interchange lim and  $\Sigma$ . Hence

$$\begin{aligned} \lim t(\omega)L(\omega) &= \lim t(\omega)e_{u(0)}^q(\omega) \\ &= \bar{\pi}^{-q} \lim e_L^q(\bar{\pi}\omega/T)/e_L(\bar{\pi}\omega) \end{aligned}$$

since the limit of the factors of  $\Pi'_{c \in A}(\dots)$  in (5.7) gives 1. Finally,

$$e_L(\bar{\pi}\omega) = \varrho_T(e_L(\bar{\pi}\omega/T)) = Te_L(\bar{\pi}\omega/T) + e_L^q(\bar{\pi}\omega/T),$$

thus

$$\begin{aligned} \lim e_L^q(\bar{\pi}\omega/T)/e_L(\bar{\pi}\omega) &= 1 - \lim Te_L(\bar{\pi}\omega/T)/e_L(\bar{\pi}\omega) \\ &= 1 \text{ by (4.8).} \end{aligned}$$

QED.

**6.  $L$  as a modular form**

Consider the inverse  $1/L(\omega)$  as a function on  $\Omega$ . It is meromorphic on  $\Omega$ , of weight  $q + 1$  and type 1 for  $\Gamma = GL(2, A)$ , and has a simple zero at  $\infty$  (see (4.9)). More precisely, its  $t$ -expansion begins  $1/L(\omega) = \bar{\pi}^q t(\omega) + \dots$ , due to Theorem 5.2. If  $0 \neq f(\omega)$  is any meromorphic modular form for  $\Gamma$  of weight  $k$  and type  $m$ , we have (see [6] for explanation):

$$\sum v_x(f) + v_0(f)/(q + 1) + v_\infty(f)/(q - 1) = k/(q^2 - 1), \tag{6.1}$$

where  $v_x(f)$  is the order of zero of  $f$  at  $x$  (negative if  $f$  has a pole at  $x$ ), the sum on the left hand side is over the non-elliptic points of  $\Gamma \backslash \Omega$ , and  $v_0, v_\infty$  is the order at the elliptic point, at  $\infty$ , respectively.

Since  $L$  is holomorphic on  $\Omega$  (so  $1/L$  has no zeroes on  $\Omega$ ), (6.1) shows  $1/L$  to have neither zeroes nor poles on  $\Omega$ . Thus  $1/L$  is a holomorphic modular form of weight  $q + 1$  and type 1. But the space of those forms is one-dimensional, generated by the form  $h(\omega)$  discussed in [6], Section 9. Using theorem 9.1 of that paper, we have

6.2. THEOREM:  $L(\omega) = -\bar{\pi}^{-q} h^{-1}(\omega)$ , where the modular form  $h$  of weight  $q + 1$  and type 1 has the product expansion around  $\infty$

$$h(\omega) = -t \prod_{a \in A \text{ monic}} f_a^{q^2-1}(t).$$

Note that  $h$  has several a priori different characterizations, e.g., as a Poincaré series, or as the ‘‘Serre derivation’’ of the normalized Eisenstein series of weight  $q - 1$  (which explains the minus sign).

6.3. COROLLARY: In the notation of (4.3),  $L(\omega) = L(\omega_1, \omega_2)$  is always non-zero.

We may even compute that value. The Drinfeld module  $\phi$  associated with  $A\omega_1 + A\omega_2$  is of the form

$$\phi_T = T\tau^0 + g\tau + \Delta\tau^2,$$

where  $\Delta(\omega)$ , considered as a modular form, equals  $-\bar{\pi}^{q^2-1} h^{q-1}(\omega)$ . Replacing  $\omega = (\omega_1, \omega_2)$  by  $c\omega$  replaces  $(g, \Delta)$  by  $(c^{1-q}g, c^{1-q^2}\Delta)$ , so we may assume  $\Delta(\omega) = -1$ . From the above, we get

6.4. COROLLARY: If  $\omega$  is normalized such that  $\Delta(\omega) = -1$  then  $L^{q-1}(\omega) = \bar{\pi}^{q-1}$ .

Clearly, this is a nice analogue of Legendre’s relation (1.2). The occurrence of  $(q - 1)$ -st powers reflect the fact that  $\bar{\pi}$  is not intrinsically defined, but  $\bar{\pi}^{q-1}$  is. Similarly, the definition of  $h$  (but not that of  $\bar{\pi}h$  or  $\bar{\pi}^q h$ ) depends on the choice of  $\bar{\pi}$ .

(6.5) For  $v \in \Lambda_\omega = A\omega_1 + A\omega_2$  and  $\partial \in D(\phi)$  we formally write

$$\int_v \partial = F_\partial(v)$$

which by remark 3.5(i) vanishes if  $\partial \in D_{\text{st}}(\phi)$ . We thus obtain a pairing of the two-dimensional  $C$ -vector spaces  $\Lambda_\omega \otimes C$  and  $H_{\text{DR}}^*(\phi_\omega)$  which by (6.3) is non-singular. For a generalization of this fact, see [9].

### 7. The periods of second kind

For the last time, we change notation and set for  $\omega \in \Omega$

$$\eta_1(\omega) = F_\omega^{(1)}(\omega), \quad \eta_2(\omega) = F_\omega^{(1)}(1), \tag{7.1}$$

the *periods of second kind* of the Drinfeld module associated with  $A\omega + A$ . They are related by

$$\eta_1(\omega) = \omega^q \eta_2(\omega^{-1}), \tag{7.2}$$

which follows immediately from (4.2). Further,

$$\eta_2(\gamma\omega) = \eta_2(\omega) \tag{7.3}$$

if  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$  with  $c = 0$ . Therefore,  $\eta_2(\omega)$  has a  $t$ -expansion that we will calculate. By means of section 5, we may write down two kinds of series representations for  $\eta_2$ . First, (5.3) and (5.5) give

$$\eta_2(\omega) = \sum_{i \geq 0} T^i e_{v(i)}^q(\omega), \quad v(i) = (0, T^{-i-1}).$$

Applying (5.7) and noting  $f_{nc}(t_{1/n}) = f_c(t)$  yields

$$\eta_2(\omega) = \bar{\pi}^{-q} \sum_{i \geq 0} T^i \zeta^q(i) \prod_{c \in A} (f_c(t) - \zeta(i)t^{c|})/f_c(t) \tag{7.4}$$

with  $\zeta(i) = e_L(\bar{\pi}/T^{i+1})$ .

Next, putting  $e_\omega(z) = \sum_{i \geq 0} \alpha_i(\omega) z^i$ , the  $\alpha_i(\omega)$  are modular forms of weight  $q^i - 1$  and type 0. Their  $t$ -expansions may be calculated with the methods of [6]. Let  $[i]$  be short for the element  $T^{q^i} - T$  of  $A$ . Comparing coefficients in the difference equation that defines  $F_\omega^{(1)}$ , we see

$$F_\omega^{(1)}(z) = \sum_{i \geq 0} \alpha_i^q(\omega) z^i / [i + 1],$$

in particular

$$\eta_2(\omega) = \sum_{i \geq 0} \alpha_i^q(\omega) / [i + 1]. \tag{7.5}$$

Note that (7.4) is an infinite sum of meromorphic modular forms of weight  $-q$  of different levels, whereas (7.5) is a sum of holomorphic modular forms of level one, but with different weights. Both sums converge in a neighborhood of  $\infty$ , but fail to give us rationality properties of the  $t$ -coefficients since for each term, an infinite number of summands contribute. At least we may read off the constant term:

7.6. LEMMA:  $\eta_2(\infty) = -\bar{\pi}^{1-q}$ .

*Proof:* As in (5.3), we have

$$F(z) = \sum_{i \geq 0} T^i e_L^q(z/T^{i+1})$$

for the unique solution  $F(z)$  without  $z$ -term of the difference equation

$$e_L^q(z) = F(Tz) - TF(z).$$

But  $e_L(Tz) = Te_L(z) + e_L^q(z)$ , thus  $F(z) = e_L(z) - z$  and

$$-\bar{\pi} = e_L(\bar{\pi}) - \bar{\pi} = \sum_{i \geq 0} T^i e_L^q(\bar{\pi}/T^{i+1}) = \sum_{i \geq 0} T^i \zeta^q(i).$$

Since  $f_c(0)$  = leading coefficient of  $c$  as a polynomial in  $T$ , the right hand factors of (7.4) evaluate to 1 at  $t = 0$ .

Now (6.2) combined with (7.2) gives the transformation law for  $\eta_2$ :

$$\omega^q \eta_2(\omega^{-1}) = \omega \eta_2(\omega) - \bar{\pi}^{-q} h^{-1}(\omega). \tag{7.7}$$

But  $h(\omega^{-1}) = -\omega^{q+1} h(\omega)$ , thus  $\tilde{\eta}_2(\omega) = \bar{\pi}^{q-1} h(\omega) \eta_2(\omega)$  satisfies

$$\tilde{\eta}_2(\omega^{-1}) = -\omega^2 \tilde{\eta}_2(\omega) + \bar{\pi}^{-1} \omega. \tag{7.8}$$

Let  $\theta$  be the differential operator  $\bar{\pi}^{-1}d/d\omega$ , given in terms of  $t$  by  $-t^2d/dt$ . (For this and the following, see [6], Section 8.) Let  $E = \theta(\Delta)/\Delta$  be the logarithmic derivative of the discriminant function  $\Delta(\omega)$ . Then  $E$  satisfies the same transformation rule (7.8) as  $\tilde{\eta}_2$ .  $E(\omega)$  is an analogue of the “false Eisenstein series of weight 2” which up to a constant gives the periods of second kind of a complex elliptic curve ([8], p. 166). Let us note the  $t$ -expansion of  $E(\omega)$ :

$$E(\omega) = \sum'_{a \in A \text{ monic}} at(a\omega) = \sum' at^{|a|}(\omega)/f_a(t) \tag{7.9}$$

which in particular has coefficients in  $A$ .

The function  $\omega \mapsto \tilde{\eta}_2(\omega) - E(\omega)$  has the following properties (which result from corresponding properties of  $\eta_2$ ,  $h$ , and  $E$ ):

- (i) It is of weight 2 and type 1 for  $\Gamma = GL(2, A)$ ;
- (ii) it is holomorphic on  $\Omega$ ;
- (iii) it has at least a double zero at  $\infty$ .

The last fact comes from Lemma 7.6,  $h(\omega) = -t + \dots$ , and  $E(\omega) = t + \dots$ . Thus by (6.1), it has to vanish identically, and we have shown:

7.10. THEOREM:  $\eta_2(\omega) = \bar{\pi}^{1-q}E(\omega)/h(\omega)$ .

7.11. COROLLARY: Up to the factor  $\bar{\pi}^{1-q}$ ,  $\eta_2(\omega)$  has its  $t$ -coefficients in  $A$ .

7.12. COROLLARY: Up to the factor  $\bar{\pi}^{1-q}$ , the  $t$ -series given by (7.4) and (7.5) have coefficients in  $A$ .

### 8. The Gauss–Manin connection

Let  $\phi = \phi_\omega$  be the generic rank two Drinfeld module associated with  $\Lambda_\omega = A\omega + A$ ,  $\omega$  varying over the upper half-plane  $\Omega$ , and  $\omega_1 = \omega$ ,  $\omega_2 = 1$ . Following Katz (see [8], A 1.3), we define the Gauss–Manin connection  $\nabla = \nabla_\theta$  of the differential operator  $\theta = \bar{\pi}^{-1}d/d\omega$  as the unique endomorphism  $\nabla: H_{\text{DR}}^*(\phi) \xrightarrow{\sim} H_{\text{DR}}^*(\phi)$  satisfying

$$\int_{\omega_i} \nabla(\partial) = \bar{\pi}^{-1} \frac{d}{d\omega} \int_{\omega_i} \partial \quad (i = 1, 2, \partial \in H_{\text{DR}}^*(\phi)). \tag{8.1}$$

$\nabla$  is well-defined in view of (6.5). Clearly, for  $f$  holomorphic on  $\Omega$  and  $\partial \in H_{\text{DR}}^*(\phi)$  a “differential form,”

$$\nabla(f\partial) = \theta(f)\partial + f\nabla(\partial) \tag{8.2}$$



holds. Now  $H_{\text{DR}}^*(\phi)$  is spanned by  $\partial^{(i)}$  ( $i = 0, 1$ ), and  $F_{\partial^{(i)}}(\omega_j)$  (where  $i = 0, 1$  and  $j = 1, 2$ ) is given by the matrix

$$\begin{pmatrix} -\omega & -1 \\ \eta_1(\omega) & \eta_2(\omega) \end{pmatrix}.$$

Trivially,  $(d/d\omega)\eta_i(\omega) = 0$  as results from (5.8) and (7.5). Hence

$$\left(\frac{d}{d\omega} F_{\partial^{(i)}}(\omega_j)\right) = \begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix},$$

and solving (8.1) gives

$$\nabla \begin{pmatrix} \partial^{(0)} \\ \partial^{(1)} \end{pmatrix} = \bar{\pi}^{q-1} h(\omega) \begin{pmatrix} \eta_2(\omega) & 1 \\ 0 & 0 \end{pmatrix} \begin{bmatrix} \partial^{(0)} \\ \partial^{(1)} \end{bmatrix}. \tag{8.3}$$

From [6], (8.5) we know that  $\theta h + Eh$  is a modular form of weight  $q + 3$  and type 2. Comparing  $t$ -expansions, we see it has a zero of order  $\geq 3$  at  $\infty$ . By (6.1), it vanishes identically, hence

$$\theta(h) = -Eh. \tag{8.4}$$

Similarly ([6], Theorem 9.1, note the different normalization used there):

$$\theta(g) = Eg + \bar{\pi}^{q-1} h. \tag{8.5}$$

Let now TD be the Tate–Drinfeld module over the power series ring  $R = A[[t]]$ , defined by

$$(\text{TD})_T = T\tau^0 + \tilde{g}\tau + \tilde{\Delta}\tau^2, \tag{8.6}$$

where for  $\tilde{g}$  (resp.  $\tilde{\Delta}$ ), we insert the  $t$ -expansion of  $\bar{\pi}^{1-q}g$  (resp.  $\bar{\pi}^{1-q^2}\Delta$ ) which in fact has coefficients in  $A$ . In order to have everything defined over  $R$ , we replace the basis  $\{\partial^{(0)}, \partial^{(1)}\}$  of  $H_{\text{DR}}^*(\phi)$  by  $\{\tilde{\partial}^{(0)}, \tilde{\partial}^{(1)}\}$ , where

$$\begin{aligned} \tilde{\partial}^{(0)} &= \bar{\pi}^{1-q^2}\partial^{(0)} + (\bar{\pi}^{1-q} - \bar{\pi}^{1-q^2})g(\omega)\partial^{(1)} \\ \tilde{\partial}^{(1)} &= h(\omega)\partial^{(1)}. \end{aligned} \tag{8.7}$$

Note that  $\tilde{\partial}_T^{(0)} = \tilde{g}\tau + \tilde{\Delta}\tau^2$ , i.e.,  $\tilde{\partial}^{(0)}$  is the inner derivation canonically associated with TD. Using (8.2)–(8.7) and some calculation, the following matrix representation for  $\nabla$  is yielded:

$$8.8. \text{ THEOREM: } \nabla \begin{pmatrix} \tilde{\partial}^{(0)} \\ \tilde{\partial}^{(1)} \end{pmatrix} = \begin{pmatrix} E(\omega) & 1 \\ 0 & -E(\omega) \end{pmatrix} \begin{pmatrix} \tilde{\partial}^{(0)} \\ \tilde{\partial}^{(1)} \end{pmatrix}.$$

By (7.9), the entries may be considered as elements of  $R$ . Stressing further the analogy with the elliptic modular case, the nilpotency of the matrix for  $\nabla$  evaluated at  $\omega = \infty$ , i.e.,  $t = 0$  should indicate  $\{\tilde{\partial}^{(0)}, \tilde{\partial}^{(1)}\}$  to be a basis for the canonical extension of  $(H_{\text{DR}}^*(\text{TD}), \nabla)$  to  $\infty$ .

### Acknowledgements

I am grateful to Greg Anderson, Pierre Deligne, David Goss, Dinesh Thakur, and Jing Yu for many helpful conversations on various topics centered around Drinfeld modules.

### References

1. G. Anderson:  $t$ -motives. *Duke Math. J.* 53 (1986) 457–502.
2. G. Anderson: Drinfeld modules and tensor products. In preparation.
3. V.G. Drinfeld: Elliptic modules (russian). *Math. Sbornik* 94 (1974) 594–627; English translation: *Math. USSR-Sbornik* 23 (1976) 561–592.
4. E.U. Gekeler: Modulare Einheiten für Funktionenkörper. *J. reine angew. Math.* 348 (1984) 94–115.
5. E.U. Gekeler: Drinfeld modular curves. *Lecture Notes in Mathematics* 1231. Springer-Verlag (1986).
6. E.U. Gekeler: On the coefficients of Drinfeld modular forms. *Inv. Math.* 93 (1988) 667–700.
7. L. Gerritzen and M. van der Put: Schottky groups and Mumford curves. *Lecture Notes in Mathematics* 817. Springer-Verlag (1980).
8. N. Katz:  $P$ -adic properties of modular schemes and modular forms. *Lecture Notes in Mathematics* 350 69–190. Springer-Verlag (1973).
9. E.U. Gekeler: On the de Rham isomorphism for Drinfeld modules. IAS-Preprint Princeton 1988.