

COMPOSITIO MATHEMATICA

WOLFRAM DECKER

**Monads and cohomology modules of rank
2 vector bundles**

Compositio Mathematica, tome 76, n° 1-2 (1990), p. 7-17

http://www.numdam.org/item?id=CM_1990__76_1-2_7_0

© Foundation Compositio Mathematica, 1990, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Monads and cohomology modules of rank 2 vector bundles

WOLFRAM DECKER *

*Universität Kaiserslautern, Fachbereich Mathematik, Erwin-Schrödinger-Straße, 6750
Kaiserslautern, FRG*

Received 19 August 1988; accepted 20 July 1989

Introduction

Monads are a useful tool to construct and study rank 2 vector bundles on the complex projective space \mathbb{P}_n , $n \geq 2$ (compare [O-S-S]). Horrocks' technique of eliminating cohomology [Ho 2] represents a given rank 2 vector bundle \mathcal{E} as the cohomology of a monad

$$(M(\mathcal{E})) \quad \mathcal{A} \xrightarrow{\varphi} \mathcal{B} \xrightarrow{\psi} \mathcal{C}$$

as follows.

First eliminate the graded $S = \mathbb{C}[z_0, \dots, z_n]$ -module $H^1\mathcal{E}(\ast) = \bigoplus_{m \in \mathbb{Z}} H^1(\mathbb{P}_n, \mathcal{E}(m))$ by the universal extension

$$0 \rightarrow \mathcal{E} \rightarrow \mathcal{Q} \rightarrow \tilde{L}_0 \rightarrow 0,$$

where

$$L_0 \rightarrow H^1\mathcal{E}(\ast) \rightarrow 0.$$

is given by a minimal system of generators (\sim stands for sheafification).

If $n = 2$ take this extension as a monad with $\mathcal{A} = 0$.

If $n \geq 3$ eliminate dually $H^{n-1}\mathcal{E}(\ast)$ by the universal extension

$$0 \rightarrow \tilde{L}_0^\vee(c_1) \rightarrow \mathcal{K} \rightarrow \mathcal{E} \rightarrow 0$$

(where $c_1 = c_1(\mathcal{E})$ is the first Chern-class). Then notice, that the two extensions

* Partially supported by the DAAD.

can be completed to the display

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & \tilde{L}_0^\vee(c_1) & \longrightarrow & \mathcal{X} & \longrightarrow & \mathcal{E} \longrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \tilde{L}_0^\vee(c_1) & \xrightarrow{\varphi} & \mathcal{B} & \longrightarrow & \mathcal{L} \longrightarrow 0 \\
 & & & & \downarrow \psi & & \downarrow \\
 & & & & \tilde{L}_0 & = & \tilde{L}_0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

of a monad

$$\tilde{L}_0^\vee(c_1) \xrightarrow{\varphi} \mathcal{B} \xrightarrow{\psi} \tilde{L}_0$$

for \mathcal{E} .

To get a better understanding for \mathcal{B} , φ and ψ consider first the case $n = 2, 3$. Then \mathcal{B} is a direct sum of line bundles by Horrocks' splitting criterion [Ho 1]. Taking cohomology we obtain a free presentation

$$B \xrightarrow{\psi} L_0 \longrightarrow H^1 \mathcal{E}(\ast) \longrightarrow 0$$

with $B = H^0 \mathcal{B}(\ast)$. The crucial point is that this is minimal [Ra]. Moreover, if $n = 3$, then B is self-dual [Ra]: $B^\vee(c_1) \simeq B$. We will see below that up to isomorphism φ is the dual map of ψ .

Let us summarize and slightly generalize. Consider an arbitrary graded S -module N of finite length with minimal free resolution (m.f.r. for short)

$$0 \longrightarrow L_{n+1} \xrightarrow{\alpha_n} L_n \longrightarrow \dots \longrightarrow L_1 \xrightarrow{\alpha_0} L_0 \longrightarrow N \longrightarrow 0.$$

If $n = 2$ then $N \simeq H^1 \mathcal{E}(\ast)$ for some rank 2 vector bundle \mathcal{E} on \mathbb{P}_2 iff $\widehat{\text{rk}} L_1 = \text{rk } L_0 + 2$ (compare [Ra]). In this case \mathcal{E} is uniquely determined as $\ker \alpha_0$:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \tilde{L}_0^\vee(c_1) & \xrightarrow{\alpha_0^\vee(c_1)} & \tilde{L}_1^\vee(c_1) & \xrightarrow{\alpha_1} & \tilde{L}_1 & \xrightarrow{\alpha_0} & \tilde{L}_0 & \longrightarrow & 0. \\
 & & & & \searrow & & \nearrow & & & & \\
 & & & & & & \mathcal{E} & & & & \\
 & & & & \nearrow & & \searrow & & & & \\
 & & & & 0 & & 0 & & & &
 \end{array}$$

(This sequence is self-dual by Serre-duality [Ho 1, 5.2], since $\mathcal{E}^\vee(c_1) \simeq \mathcal{E}$).

For $n = 3$ there is an analogous result. Answering Problem 10 of Hartshorne's list [Ha] we prove:

PROPOSITION 1. *N is the first cohomology module of some rank 2 vector bundle on \mathbb{P}_3 iff*

- (1) $\text{rk } L_1 = 2 \text{ rk } L_0 + 2$ and
- (2) *there exists an isomorphism $\Phi: L_1^\vee(c_1) \xrightarrow{\cong} L_1$ for some $c_1 \in \mathbb{Z}$ such that $\alpha_0 \circ \Phi \circ \alpha_0^\vee(c_1) = 0$.*

In this case any Φ satisfying (2) defines a monad

$$(M_\Phi) \quad \tilde{L}_0^\vee(c_1) \xrightarrow{\Phi \circ \alpha_0^\vee(c_1)} \tilde{L}_1 \xrightarrow{\alpha_0} \tilde{L}_0$$

and \mathcal{E} is a 2-bundle on \mathbb{P}_3 with $H^1 \mathcal{E}(\ast) \simeq N$ (and $c_1 = c_1(\mathcal{E})$) iff $(M(\mathcal{E})) \simeq (M_\Phi)$ for some Φ .

To complete the picture let us mention a result of Hartshorne and Rao (not yet published). If $N \simeq H^1 \mathcal{E}(\ast)$ as above then $L_0^\vee(c_1) \xrightarrow{\varphi} L_1$ is part of a minimal system of generators for $\ker \alpha_0$. In other words: There exists a splitting

$$0 \rightarrow L_4 \rightarrow L_3 \rightarrow L'_2 \oplus L_0^\vee(c_1) \rightarrow L_1 \rightarrow L_0 \rightarrow H^1 \mathcal{E}(\ast) \rightarrow 0$$

inducing the monad

$$(M(\mathcal{E})) \quad \tilde{L}_0^\vee(c_1) \rightarrow \tilde{L}_1 \rightarrow \tilde{L}_0$$

and the m.f.r.

$$0 \rightarrow \tilde{L}_4 \rightarrow \tilde{L}_3 \rightarrow \tilde{L}'_2 \rightarrow \mathcal{E} \rightarrow 0$$

resp.

For $n \geq 4$ there is essentially only one indecomposable 2-bundle known on \mathbb{P}_n : The Horrocks-Mumford-bundle \mathcal{F} on \mathbb{P}_4 with Chern-classes $c_1 = -1$, $c_2 = 4$. We prove:

PROPOSITION 2. *The m.f.r. of $H^2 \mathcal{F}(\ast)$ decomposes as*

$$\begin{array}{ccccccc}
 0 \rightarrow H_2 & \xrightarrow{\beta_1} & H_1 & \xrightarrow{\begin{pmatrix} \beta_0 \\ \beta'_0 \end{pmatrix}} & L_0^\vee(c_1) \oplus L_1 & \xrightarrow{\begin{pmatrix} 0 & \alpha_0 \\ \alpha_0^\vee(c_1) & \ast \end{pmatrix}} & L_0 \oplus L_1^\vee(c_1) \rightarrow \\
 & & & & & \searrow & \nearrow \\
 & & & & & B & \\
 & & & & 0 & \nearrow & \searrow \\
 & & & & & & 0
 \end{array}$$

$$\rightarrow H_1^\vee(c_1) \rightarrow H_2^\vee(c_1) \rightarrow H^2 \mathcal{F}(\ast) \rightarrow 0$$

with $B = H^0 \mathcal{B}(\ast)$, inducing the monad

$$(M(\mathcal{F})) \quad \tilde{L}_0^\vee(c_1) \rightarrow \mathcal{B} \rightarrow \tilde{L}_0$$

and the minimal free presentation

$$L_1 \xrightarrow{\alpha_0} L_0 \longrightarrow H^1 \mathcal{F}(\ast) \longrightarrow 0.$$

The corresponding m.f.r. decomposes as

$$\begin{aligned} 0 \rightarrow L_5 \rightarrow L_4 \rightarrow L_3 \oplus H_2 \xrightarrow{\begin{pmatrix} \ast & 0 \\ \ast & \beta_1 \end{pmatrix}} L_2' \oplus H_1 \xrightarrow{\begin{pmatrix} \ast & \beta_0' \end{pmatrix}} L_1 \xrightarrow{\alpha_0} L_0 \rightarrow \\ \rightarrow H^1 \mathcal{F}(\ast) \rightarrow 0 \end{aligned}$$

inducing the m.f.r.

$$0 \rightarrow \tilde{L}_5 \rightarrow \tilde{L}_4 \rightarrow \tilde{L}_3 \rightarrow \tilde{L}_2 \rightarrow \mathcal{F} \rightarrow 0.$$

$(M(\mathcal{F}))$ is the monad given in [H-M]. Using its display we can compute the above m.f.r.'s explicitly. Especially we reobtain the equations of the abelian surfaces in \mathbb{P}_4 ([Ma 1], [Ma 2]).

Of course we may deduce from \mathcal{F} some more bundles by pulling it back under finite morphisms $\pi: \mathbb{P}_4 \rightarrow \mathbb{P}_4$. The above result also holds for the bundles $\pi^* \mathcal{F}$ with $(M(\pi^* \mathcal{F})) = \pi^*(M(\mathcal{F}))$.

There is some evidence (but so far no complete proof), that a splitting as in Proposition 2 occurs for every indecomposable 2-bundle on \mathbb{P}_4 . This suggests a new construction principle for such bundles by constructing their H^2 -module first.

Proof of Proposition 1

Let $n = 3$ and N be a graded S -module of finite length with m.f.r.

$$0 \rightarrow L_4 \rightarrow L_3 \rightarrow L_2 \rightarrow L_1 \xrightarrow{\alpha_0} L_0 \rightarrow N \rightarrow 0.$$

Suppose first that $N \simeq H^1 \mathcal{E}(\ast)$ for some 2-bundle \mathcal{E} on \mathbb{P}_3 (with first Chern-class c_1). As seen in the introduction, Horrocks' construction leads to a monad

$$(M(\mathcal{E})) \quad \tilde{L}_0^\vee(c_1) \xrightarrow{\varphi} \tilde{L}_1 \xrightarrow{\alpha_0} \tilde{L}_0$$

for \mathcal{E} . The dual sequence

$$\tilde{L}_0^\vee(c_1) \xrightarrow{\alpha_0^\vee(c_1)} \tilde{L}_1^\vee(c_1) \xrightarrow{\varphi^\vee(c_1)} \tilde{L}_0$$

is a monad for $\mathcal{E}^\vee(c_1) \simeq \mathcal{E}$. The induced presentation of N has to be isomorphic to that one given by the m.f.r.:

$$\begin{array}{ccccccc} L_1^\vee(c_1) & \xrightarrow{\varphi^\vee(c_1)} & L_0 & \rightarrow & N & \rightarrow & 0 \\ \Phi^\vee(c_1) \downarrow & \parallel & & & \parallel & & \parallel \\ L_1 & \xrightarrow{\alpha_0} & L_0 & \rightarrow & N & \rightarrow & 0. \end{array}$$

Dualizing gives (2) since $\alpha_0 \circ \varphi = 0$ and thus also a monad (M_Φ) for \mathcal{E} , isomorphic to $(M(\mathcal{E}))$ (replace φ by $\Phi \circ \alpha_0^\vee(c_1)$).

Conversely if N satisfies (2), we obtain a monad (M_Φ) by sheafification. (Since $\tilde{N} = 0$, α_0 is a bundle epimorphism. Dually $\alpha_0^\vee(c_1)$ is a bundle monomorphism.) Let \mathcal{E} be the cohomology bundle of (M_Φ) . Then $H^1\mathcal{E}(\ast) \simeq N$. \mathcal{E} has rank 2, if N satisfies (1). □

REMARK 1. (i) Let $N \simeq H^1\mathcal{E}(\ast)$ as above with induced splitting

$$0 \rightarrow L_4 \rightarrow L_3 \rightarrow L_2 \oplus L_0^\vee(c_1) \rightarrow L_1 \rightarrow L_0 \rightarrow H^1\mathcal{E}(\ast) \rightarrow 0$$

as in the introduction. Recall that \mathcal{E} is stable iff $H^0(\mathbb{P}_3, \mathcal{E}(m)) = 0$ for $m \leq -c_1/2$. Thus \mathcal{E} is stable iff L_2' has no direct summand $S(m)$ with $m \geq c_1/2$. Notice that this condition only depends on N .

(ii) If N satisfies (1) and has only one generator, then (2) is obviously equivalent to the symmetry condition $L_1^\vee(c_1) \simeq L_1$. Thus [Ra, 3.1] is a special case of Proposition 1.

EXAMPLES. (i) The well-known Null correlation bundles are by definition the bundles corresponding to the S -module \mathbb{C} . Consider the Koszul-presentation

$$4S \xrightarrow{\alpha_0} S(1) \rightarrow \mathbb{C} \rightarrow 0, \quad \alpha_0 = (z_0, z_1, z_2, z_3).$$

The isomorphisms $4S \xrightarrow{\Phi} 4S$ with $\alpha_0 \circ \Phi \circ \alpha_0^\vee(c_1) = 0$ are precisely the 4×4 skew symmetric matrices with nonzero determinant. Two such matrices give isomorphic bundles iff they differ by a scalar (use [O-S-S, II, Corollary 1 to 4.1.3]). The moduli space of Null correlation bundles is thus isomorphic to $\mathbb{P}_5 \setminus \mathbb{G}$, where \mathbb{G} is the Plucker embedded Grassmanian of lines in \mathbb{P}_3 .

Unlike the case $n = 2$ the bundle is not uniquely determined by the module.

(ii) The S -module

$$6S \xrightarrow{\alpha_0} S(1) \oplus S(2) \rightarrow N \rightarrow 0, \alpha_0 = \begin{pmatrix} z_0 & z_1 & z_2 & z_3 & 0 & 0 \\ 0 & 0 & z_0^2 & z_1^2 & z_2^2 & z_3^2 \end{pmatrix}$$

satisfies (1) and the symmetry condition $L_1^\vee \simeq L_1$, i.e. the necessary conditions of [Ra]. But N does not satisfy (2).

Cohomology modules of the Horrocks-Mumford-bundle \mathcal{F}

We first recall the construction of \mathcal{F} [H-M]. Let

$$V = \text{Map}(\mathbb{Z}_5, \mathbb{C})$$

be the vector space of complex valued functions on \mathbb{Z}_5 . Denote by

$$H \subset N \subset \text{SL}(5, \mathbb{C})$$

the Heisenberg group and its normalizer in $\text{SL}(5, \mathbb{C})$ resp.

Let $V_0 = V, V_1, V_2, V_3$ and

$$W = \text{Hom}_H(V_1, \Lambda^2 V)$$

be defined as in [H-M]. The V_i are irreducible representations of H and N of degree 5. W is an irreducible representation of N/H of degree 2. It is unimodular, so it comes up with an invariant skew symmetric pairing.

Let $\mathbb{P}_4 = \mathbb{P}(V)$ be the projective space of lines in V . The Koszul-complex on $\mathbb{P}(V)$ is the exact sequence

$$(K) \quad \dots \rightarrow \mathcal{O}(-1) \otimes \Lambda^2 V \rightarrow \mathcal{O} \otimes \Lambda^3 V \rightarrow \dots,$$

$$\begin{array}{ccc} & \searrow & \nearrow \\ & (\Lambda^2 \mathcal{F})(-3) & \\ & \nearrow & \searrow \\ 0 & & 0 \end{array}$$

obtained by exterior multiplication with the tautological subbundle

$$\mathcal{O}(-1) \rightarrow V \otimes \mathcal{O}.$$

The exterior product provides (K) with a self-duality (with values in $\mathcal{O}(-1) \otimes \Lambda^5 V \simeq \mathcal{O}(-1)$).

This induces the natural pairings

$$(\Lambda^i \mathcal{F})(-i-1) \otimes (\Lambda^{4-i} \mathcal{F})(i-5) \xrightarrow{\wedge} (\Lambda^4 \mathcal{F})(-6) \simeq \mathcal{O}(-1)$$

and is compatible with the action of $\mathrm{SL}(5, \mathbb{C})$.

It can be extended to $(K) \otimes W$ by tensoring with the invariant form, then being compatible with the action of N .

As in the proof of [H-M, Lemma 2.4] it follows, that $(K) \otimes W$ decomposes as

$$\begin{array}{ccccccc} \dots & \rightarrow & \mathcal{O}(-1) \otimes V_1 & \xrightarrow{\begin{pmatrix} 0 & \alpha_0 \\ \alpha_0^\vee(-1) & \star \end{pmatrix}} & \mathcal{O} \otimes V_3 & \rightarrow & \dots, \\ & & \oplus & & \oplus & & \\ & & \mathcal{O}(-1) \otimes V_1 \otimes U & & \mathcal{O} \otimes V_3 \otimes U & & \\ & & & \searrow \quad \nearrow & & & \\ & & & (\Lambda^2 T)(-3) \otimes W & & & \\ & & \nearrow & & \searrow & & \\ & & 0 & & 0 & & \end{array}$$

given by the splitting into irreducible N -modules. Moreover the induced

$$\mathcal{O}(-1) \otimes V_1 \rightarrow (\Lambda^2 \mathcal{F})(-3) \otimes W \rightarrow \mathcal{O} \otimes V_3$$

is the self-dual Horrocks-Mumford-monad, whose cohomology is \mathcal{F} (normalized such that $c_1 \mathcal{F} = -1$).

To proof Proposition 2 consider the display

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & \mathcal{O}(-1) \otimes V_1 & \longrightarrow & \mathcal{H} & \longrightarrow & \mathcal{F} \rightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \rightarrow & \mathcal{O}(-1) \otimes V_1 & \rightarrow & (\Lambda^2 \mathcal{F})(-3) \otimes W & \rightarrow & \mathcal{L} \rightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & \mathcal{O} \otimes V_3 & = & \mathcal{O} \otimes V_3 \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0 \end{array}$$

It first follows that $H^2 \mathcal{F}(\ast) = W$ is a vector space, sitting in degree -2 (compare [H-M]). Its m.f.r. is the Koszul-complex obtained from $(K) \otimes W$ by taking global

sections. So it decomposes, inducing the presentation

$$S(-1) \otimes V_1 \otimes U \xrightarrow{\alpha_0} S \otimes V_3 \longrightarrow H^1 \mathcal{F}(\ast) \longrightarrow 0$$

and the Horrocks-Mumford-monad. But this is just

$$(M(\mathcal{F})) \quad \mathcal{O}(-1) \otimes V_1 \rightarrow \mathcal{B} \rightarrow \mathcal{O} \otimes V_3:$$

apply e.g. [O-S-S, II, Corollary 1 to 4.1.3] (notice that $H^0 \mathcal{F} = 0$ implies $H^0 \mathcal{B} = H^0 \mathcal{B}^*(-1) = 0$ by construction of $(M(\mathcal{F}))$).

It remains to show that α_0 is minimal and that the corresponding m.f.r. of $H^1 \mathcal{F}(\ast)$ decomposes, inducing the m.f.r., say,

$$0 \longrightarrow F_3 \xrightarrow{\gamma_2} F_2 \xrightarrow{\gamma_1} F_1 \xrightarrow{\gamma_0} F_0 \longrightarrow F \longrightarrow 0$$

of $F = H^0 \mathcal{F}(\ast)$.

From the second row of the display we obtain the m.f.r. of $Q = H^0 \mathcal{Q}(\ast)$:

$$\begin{array}{ccccccc}
 & & 0 & & & & 0 \\
 & & \downarrow & & & & \downarrow \\
 & & S(-3) \otimes W & = & & & S(-3) \otimes W \\
 & & \downarrow \beta_1 & & & & \downarrow \beta_1 \\
 & & S(-2) \otimes V \otimes W & = & & & S(-2) \otimes V \otimes W \\
 & & \downarrow \beta_0 & & & & \downarrow \beta_0'' \\
 0 \rightarrow & S(-1) \otimes V_1 \rightarrow & S(-1) \otimes V_1 \oplus (S(-1) \otimes V_1 \otimes U) & \rightarrow & S(-1) \otimes V_1 \otimes U & \rightarrow & 0 \\
 & \parallel & \downarrow & & \downarrow & & \\
 0 \rightarrow & S(-1) \otimes V_1 \longrightarrow & H^0(\Lambda^2 \mathcal{F})(\ast - 3) \otimes W & \longrightarrow & Q & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & & & 0
 \end{array}$$

Define

$$A = (a_{ij})_{\substack{0 \leq i \leq 4 \\ 0 \leq j \leq 1}} \text{ by}$$

$$\left. \begin{aligned} a_{i0} &= e_{i+2} \wedge e_{i+3} \\ a_{i1} &= e_{i+1} \wedge e_{i+4} \end{aligned} \right\} i \bmod 5.$$

Then w_0, w_1 , given by $w_j(e_i) = a_{ij}$ is a basis of W . Identifying $W \simeq \mathbb{C}^2$, the invariant form on W becomes the standard symplectic form $Q = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ on \mathbb{C}^2 .

We thus may rewrite $(M(\mathcal{F}))$ as

$$(M(\mathcal{F})) \quad 5\mathcal{O}(-1) \xrightarrow{Q \cdot A} 2(\Lambda^2 \mathcal{F})(-3) \xrightarrow{A} 5\mathcal{O},$$

the matrices operating by exterior multiplication.

(ii) From the explicit form of $(M(\mathcal{F}))$ we can compute α_0 explicitly. Choose a convenient basis of $\Lambda^2 V \otimes W = (V_1 \oplus U)$. Then

$$15S(-1) \xrightarrow{\alpha_0} 5S \rightarrow H^1 \mathcal{F}(\ast) \rightarrow 0$$

is the matrix

$$\alpha_0 = \begin{pmatrix} 0 & z_3 & 0 & 0 & z_2 & 0 & 0 & z_1 & z_4 & 0 & z_0 & 0 & 0 & 0 & 0 \\ z_3 & 0 & z_4 & 0 & 0 & 0 & 0 & 0 & z_2 & z_0 & 0 & z_1 & 0 & 0 & 0 \\ 0 & z_4 & 0 & z_0 & 0 & z_1 & 0 & 0 & 0 & z_3 & 0 & 0 & z_2 & 0 & 0 \\ 0 & 0 & z_0 & 0 & z_1 & z_4 & z_2 & 0 & 0 & 0 & 0 & 0 & 0 & z_3 & 0 \\ z_2 & 0 & 0 & z_1 & 0 & 0 & z_0 & z_3 & 0 & 0 & 0 & 0 & 0 & 0 & z_4 \end{pmatrix}.$$

Resolving it (use e.g. [B-S]), we obtain the m.f.r. of $H^1 \mathcal{F}(\ast)$. Its shape is

$$0 \rightarrow 2S(-8) \rightarrow 20S(-6) \rightarrow 35S(-5) \oplus 2S(-3) \rightarrow (15S(-4) \oplus 4S(-3)) \oplus 10S(-2) \\ \rightarrow 15S(-1) \rightarrow 5S \rightarrow H^1 \mathcal{F}(\ast) \rightarrow 0.$$

(iii) Consider the induced m.f.r. of F and its dual

$$\dots \rightarrow 35S(-5) \xrightarrow{\gamma_0} 15S(-4) \oplus 4S(-3) \xrightarrow{\Gamma} 15S(3) \oplus 4S(2) \xrightarrow{\gamma_0} 35S(4) \rightarrow \dots$$

$$\begin{array}{ccc} & \searrow & \nearrow \\ & F \simeq F^\vee(-1) & \\ & \nearrow & \searrow \\ 0 & & 0. \end{array}$$

Γ can be computed by resolving γ_0 (use again [B-S]). We thus obtain explicit

bases for the spaces of sections $H^0 \mathcal{F}(m)$. Especially we get the equations of the zero-schemes of sections of $\mathcal{F}(3)$, including the abelian surfaces in \mathbb{P}_4 .

(iv) Let $\pi: \mathbb{P}_4 \rightarrow \mathbb{P}_4$ be a finite morphism and d^4 its degree. Then $\pi^* \mathcal{F}$ is a stable 2-bundle with Chern-classes $c_1 = -d$, $c_2 = 4d^2$. Proposition 2 and the above remarks also hold for $\pi^* \mathcal{F}$: Replace (K) , $(M(\mathcal{F}))$ by $\pi^*(K)$, $(M(\pi^* \mathcal{F})) = (M(\pi^* \mathcal{F}))$ and z_0, \dots, z_4 in α_0 by f_0, \dots, f_4 , where f_0, \dots, f_4 are the forms of degree d defining π .

Acknowledgement

I'd like to thank R. Hartshorne for helpful discussions.

References

- [B-S] D. Bayer and M. Stillman: Macaulay, a computer algebra system for algebraic geometry.
- [Ha] R. Hartshorne: Algebraic vector bundles on projective spaces: A problem list. *Topology* 18 (1979) 117–228.
- [H-M] G. Horrocks and D. Mumford: A rank 2 vector bundle on \mathbb{P}^4 with 15,000 symmetries. *Topology* 12 (1973) 63–81.
- [Ho 1] G. Horrocks: Vector bundles on the punctured spectrum of a local ring. *Proc. London Math. Soc.* 14 (1964) 689–713.
- [Ho 2] G. Horrocks: Construction of bundles on \mathbb{P}^n . In: *Les équations de Yang-Mills*, A. Douady, J.-L. Verdier, séminaire E.N.S. 1977–1978. *Astérisque* 71–72 (1980) 197–203.
- [Ma 1] N. Manolache: Syzygies of abelian surfaces embedded in \mathbb{P}^4 . *J. reine angew. Math.* 384 (1988) 180–191.
- [Ma 2] N. Manolache: The equations of the abelian surfaces embedded in $\mathbb{P}^4(\mathbb{C})$. Preprint, Bukarest, 1988.
- [O-S-S] C. Okonek, M. Schneider and H. Spindler: Vector bundles on complex projective spaces. Boston, Basel, Stuttgart: Birkhäuser 1980.
- [Ra] P. Rao: A note on cohomology modules of rank two bundles. *J. Algebra* 86, (1984) 23–34.