

J.-C. FIOROT

P. JEANNIN

S. TALEB

**B-rational curves and reparametrization  
: the quadratic case**

*M2AN - Modélisation mathématique et analyse numérique*, tome  
27, n° 3 (1993), p. 289-311

[http://www.numdam.org/item?id=M2AN\\_1993\\_\\_27\\_3\\_289\\_0](http://www.numdam.org/item?id=M2AN_1993__27_3_289_0)

© AFCET, 1993, tous droits réservés.

L'accès aux archives de la revue « M2AN - Modélisation mathématique et analyse numérique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>



## B-RATIONAL CURVES AND REPARAMETRIZATION : THE QUADRATIC CASE

by J.-C. FIOROT (†), P. JEANNIN (‡) and S. TALEB (††)

Communicated by P. J. LAURENT

*Abstract.* — *The notion of massic vector, previously introduced, allows us to represent any rational curve as a B-rational (BR) curve, such a BR-curve being completely determined by its massic polygon. For computational and stability purposes we are led to determine the massic polygon resulting from a one-to-one quadratic change of parameter which allows a description of the whole support with a parameter belonging to [0, 1].*

*Keywords :* Rational curves, BR-curves, quadratic change of parameter, homogeneous coordinates, massic polygons,  $\rho$ -reciprocity, Bézier curves.

*Résumé.* — *La notion de vecteur massique, récemment introduite, nous permet de représenter toute courbe rationnelle comme une courbe B-rationnelle (BR), une telle courbe (BR) étant complètement déterminée par son polygone massique. Dans l'objectif de calculer une courbe rationnelle de manière stable nous sommes amenés à déterminer le polygone massique résultant d'un changement quadratique biunivoque du paramètre. Ce nouveau polygone massique permet alors une description de tout le support avec un paramètre parcourant [0, 1].*

### 1. INTRODUCTION

In [5, 6, 7] a general framework is given to represent any rational curve by a massic polygon. Such a representation is called a B-rational (BR) curve, it includes the Bézier (BP) curves [1, 2], i.e., polynomial curves defined over a bounded interval and the rational Bézier curves [4, 3]. BR-curves were introduced and analysed to enable rational curves to be exploited in computer

---

(\*) Manuscript received March 1992.

(\*) Manuscript reçu le 30 mars 1992.

(†) Université de Valenciennes et du Hainaut-Cambrésis, ENSIMEV, Laboratoire MMI, B.P. 311, F-59304 Valenciennes Cedex.

(‡) Université des Sciences et Technologies de Lille, UFR de Calais, Laboratoire ANO, Bât. M, F-59655 Villeneuve d'Ascq Cedex.

(††) Université de Valenciennes et du Hainaut-Cambrésis, IUT Département Génie Mécanique Productique, Laboratoire MMI, 59326 Valenciennes Cedex.

science, CAGD and CAM. Example 1 following proposition 1 illustrates the process in order to obtain a massic polygon of a  $BR$ -curve.

The problem of change of parameter crops up in the study of parametrized curves in order to give equivalent representation of their supports, and more significantly for computational and stability purposes. In [8] we studied the homographic (and affine) change of parameter, for  $BR$ -curves. This paper is devoted to the quadratic case. Usually a rational curve is parametrized over  $\tilde{\mathbb{R}}$ . We want to parametrize the support of such a curve over a finite interval which will be  $[0, 1[$ . The simplest function allowing us to make this change is a special quadratic function introduced in [6, section 2.3] which provides a one-to-one correspondence between  $\tilde{\mathbb{R}}$  and  $[0, 1[$ . Hence the support of a  $BR$ -curve of length  $n$  with parameter  $t \in \tilde{\mathbb{R}}$  is identical to that of a  $BR$ -curve of length  $2n$  with parameter  $u \in [0, 1[$ . In [6, section 2.3] such a representation is given for the circle, the folium of Descartes, the skew cubic and the window of Viviani. In this paper we propose a systematic study of the problem.

We are led to treat this problem for computational and stability purposes. For instance the question of overflow arises when calculating points of a rational curve which are at finite distance, image of parameter values belonging to a neighbourhood of infinity. The proposed quadratic change of parameter bringing back the parameter in  $[0, 1[$  provides an answer to this question. Moreover with a parameter belonging to  $[0, 1[$  we can calculate the curve by extensions of the De Castel'jau's algorithm, called ALBR1 and ALBR2 [6, sections 3.3 and 3.4], by handling affine combinations that are convex, which would not be the case with a curve defined over  $\tilde{\mathbb{R}}$ . This property offers a guarantee of stability.

The paper is organized as follows. In the first section we recall the general framework and some results concerning the  $BR$ -curves. This material comes from [6]. In the second section we make, for a  $BR$ -curve of length  $n$ , a one-to-one quadratic change of parameter. The new parameter belongs to  $[0, 1[$ . The curve obtained is again a rational curve, of length  $2n$ . The new massic vectors are determined as linear combinations of the previous ones. The new massic polygon presents a kind of symmetry called  $\rho$ -reciprocity. These  $(2n + 1)$  massic vectors are completely determined by the first  $(n + 1)$ . The third section gives an algorithm to calculate the triangular matrix defining the new massic vectors from the old ones. In the fourth section we apply this process to the polynomial curves. Three examples show the simplicity of it. Another development of this work is done in [9, 10]. This paper is written so as to be self-contained.

## 2. GENERAL FRAMEWORK. $BR$ -FORM OF A RATIONAL CURVE.

According to [5, 6, 7] we recall some basic definitions and results concerning the representation of rational curves by  $BR$ -curves.

Let us define  $\mathcal{E}$  (resp.  $\mathcal{F}$ ) a real affine space,  $\vec{\mathcal{E}}$  (resp.  $\vec{\mathcal{F}}$ ) its associated linear vector space such that  $\mathcal{E}$  (resp.  $\vec{\mathcal{E}}$ ) is an hyperplane of  $\mathcal{F}$  (resp.  $\vec{\mathcal{F}}$ ) and  $\tilde{\mathcal{E}}$  the projective completion of  $\mathcal{E}$ ; In general,  $\mathcal{E}$  is a 2 or 3-dimensional affine space.

Let  $\Omega$  be a point of  $\mathcal{F}$  not belonging to  $\mathcal{E}$ .

We define the linear vector space  $\hat{\mathcal{E}} = (\mathcal{E} \times \mathbb{R}^*) \cup \vec{\mathcal{E}}$  called the *massic vector space* [6]:  $\theta \in \hat{\mathcal{E}}$  is called a *massic vector*, it is either a weighted point or a pure vector.

The isomorphism  $\hat{\Omega} : \hat{\mathcal{E}} \rightarrow \vec{\mathcal{F}}$  defined by  $\hat{\Omega}(P, \alpha) = \alpha \overrightarrow{\Omega P}$ ,  $\hat{\Omega}(\vec{u}) = \vec{u}$  induces an addition operator and an external multiplication in  $\hat{\mathcal{E}}$ , respectively denoted by  $\oplus$  and  $*$ , such that  $\hat{\mathcal{E}}$  is a linear space and  $\hat{\Omega}$  is an isomorphism.

For any  $\theta$  and  $\theta' \in \hat{\mathcal{E}}$  and  $\lambda \in \mathbb{R}$ :  $\theta \oplus \theta' = \hat{\Omega}^{-1}(\hat{\Omega}(\theta) + \hat{\Omega}(\theta'))$  and  $\lambda * \theta = \hat{\Omega}^{-1}(\lambda \cdot \hat{\Omega}(\theta))$ . Proposition 1.2.1.6 in [6] gives the rules of use of  $\oplus$  and  $*$  operators. We consider the linear form  $\chi : \hat{\mathcal{E}} \rightarrow \mathbb{R}$ ;  $\chi(P; \alpha) = \alpha$ ,  $\chi(\vec{u}) = 0$ ;  $\chi(\theta)$  is called the *mass* of  $\theta$ .

Let  $\Pi\Omega : \vec{\mathcal{F}} - \{\Omega\} \rightarrow \vec{\mathcal{E}}$  be the conic projection of apex  $\Omega$  and  $\Pi : \hat{\mathcal{E}} - \{\vec{0}\} \rightarrow \vec{\mathcal{E}}$  be the natural projection:  $\Pi(P; \alpha) = P$ ,  $\Pi(\vec{u}) = (\vec{u})_\infty$ . They are linked by the relation  $\Pi = \Pi\Omega \circ \hat{\Omega}$  [6, proposition 1.2.2.3]. We have:  $\forall \lambda \neq 0, \Pi(\lambda \theta) = \Pi(\theta)$ .

**DEFINITION 1 OF A B-RATIONAL CURVE:** Let  $\theta_0, \theta_1, \dots, \theta_n$  be  $(n + 1)$  massic vectors not simultaneously null; a B-rational (BR)-curve of  $\vec{\mathcal{E}}$ , of controlling massic polygon  $\theta = (\theta_0, \theta_1, \dots, \theta_n)$ , denoted by  $BR[\theta_0, \theta_1, \dots, \theta_n]$  or  $BR[\theta]$ , is described by the point:

$$BR[\theta_0, \theta_1, \dots, \theta_n](t) = \Pi(BP[\theta_0, \theta_1, \dots, \theta_n](t));$$

$BP[\theta_0, \theta_1, \dots, \theta_n] = \sum_{i=0}^n B_i^n(t) * \theta_i$  is the Bézier curve in  $\hat{\mathcal{E}}$ ,  $n$  is called the length of the (BR)-curve;  $B_i^n(t) = \binom{n}{i} (1-t)^{n-i} t^i, i = 0, 1, \dots, n$  are the Bernstein basis polynomials relatively to  $[0, 1]$ .

**EQUIVALENT DEFINITION 2:** Considering  $R_i \in \mathcal{F}$  defined by  $\overrightarrow{\Omega R_i} = \hat{\Omega}(\theta_i)$  we have:

$$BR[\theta_0, \theta_1, \dots, \theta_n](t) = \Pi\Omega(BP[R_0, R_1, \dots, R_n](t)).$$

EXPLICIT FORM : Define  $I = \{i : \theta_i \in \mathcal{E} \times \mathbb{R}^*, \theta_i = (P_i ; \beta_i)\}$  and  $\bar{I} = \{i : \theta_i \in \vec{\mathcal{E}}, \theta_i = \vec{U}_i\}$ . We have  $I \cup \bar{I} = \{0, 1, \dots, n\}$ ,  $I \cap \bar{I} = \emptyset$ . From Definition 1 above and preliminary results [[6] proposition 1.3] we obtain the explicit form of a (BR) curve :

$$BR[\theta](t) = \frac{\sum_{i \in I} \beta_i B_i^n(t) P_i}{\beta(t)} + \frac{\sum_{i \in \bar{I}} B_i^n(t) \vec{U}_i}{\beta(t)} \tag{1}$$

if  $\beta(t) \neq 0$  and where  $\beta(t) = \sum_{i \in I} \beta_i B_i^n(t)$ .

$$BR[\theta](t) = \left( \sum_{i \in I} \beta_i B_i^n(t) \vec{P}_i + \sum_{i \in \bar{I}} B_i^n(t) \vec{U}_i \right)_{\infty} \tag{2}$$

if  $\beta(t) = 0$  and  $\vec{V}(t) = \sum_{i \in I} \beta_i B_i^n(t) \vec{P}_i + \sum_{i \in \bar{I}} B_i^n(t) \vec{U}_i \neq \vec{0}$

$$BR[\theta](t_0) = \lim_{t \rightarrow t_0} BR[\theta](t) \text{ if } \beta(t_0) = 0 \text{ and } \vec{V}(t_0) = \vec{0}. \tag{3}$$

Remark 1 :

- (i) When  $\bar{I} = \emptyset$ , the BR-curve is reduced to a rational Bézier curve [4, 3].
- (ii) When  $\bar{I} = \emptyset$  and  $\theta_i = (P_i ; c)$ ,  $i = 0, 1, \dots, n$ ,  $c$  being a constant, then  $\beta(t) = c$  and the BR-curve is reduced to a polynomial Bézier (BP) curve [1, 2].

THEOREM 1 [6, proposition 2.2.2.2] : Any rational (or unicursal) curve is a BR-curve and conversely.

### 3. QUADRATIC CHANGE OF PARAMETER FOR A BR-CURVE. BASIC RESULTS

Let  $BR[\omega]$  be a BR-curve of length  $n$  with massic polygon  $\omega = (\omega_0, \omega_1, \dots, \omega_n)$ .

We consider the quadratic change of parameter introduced in [6, section 2.3] :  $t = \Phi(u)$

$$\Phi(u) = \frac{\alpha B_0^2(u) + \beta B_1^2(u) + \gamma B_2^2(u)}{B_1^2(u)},$$

$\alpha, \beta, \gamma$  being three arbitrary constants with  $\alpha \gamma < 0$ .

This last condition ensures that whatever  $\beta$ ,  $\Phi$  is a one-to-one correspondence between  $[0, 1[$  and  $\tilde{\mathbb{R}}$  ( $\Phi$  is increasing if  $\alpha < 0$ , decreasing if  $\alpha > 0$ ). We notice that  $\Phi(0) = \Phi(1) = \infty$ . Then the curve described by

$BR[\omega](t)$  with  $t \in \tilde{\mathbb{R}}$  is identical to that described by  $BR[\omega](\Phi(u))$  with  $u \in [0, 1[$ . We will see that  $BR[\omega](\Phi(u))$  is a  $BR$ -curve of length  $2n$  :

$$BR[\omega](\Phi(u)) = BR[\theta_0, \theta_1, \dots, \theta_{2n}](u) .$$

The aim of this paper is to calculate the new massic polygon  $\theta$  as a function of  $\omega$ . Let us recall that if  $\theta$  describes a  $BR$ -curve and  $\lambda \in \mathbb{R}^*$ , then  $\lambda\theta = (\lambda\theta_1, \lambda\theta_2, \dots, \lambda\theta_{2n})$  describes the same curve.

LEMMA 1 : Let  $F(T_1, T_2)$  be a homogeneous polynomial (vector-valued or not) of degree  $n$  in  $T_1, T_2$  and let  $P_1(U_1, U_2), P_2(U_1, U_2)$  be two homogeneous polynomials of degree 2 in  $U_1, U_2$  such that

$$\begin{aligned} P_1(\gamma U_1, \alpha U_2) &= \alpha \gamma P_1(U_2, U_1) \\ P_2(\gamma U_1, \alpha U_2) &= \alpha \gamma P_2(U_2, U_1) . \end{aligned}$$

Then  $G(U_1, U_2) = F(P_1(U_1, U_2), P_2(U_1, U_2))$  is a homogeneous polynomial of degree  $2n$  in  $U_1, U_2$ . It can be written :

$$G(U_1, U_2) = \sum_{i=0}^{2n} b_i U_1^{2n-i} U_2^i$$

with  $b_i$  satisfying  $b_{2n-i} = \rho^{n-i} b_i, i = 0, 1, \dots, n$ , where  $\rho = \frac{\gamma}{\alpha}$ .

Proof : Successively we have :

$$\begin{aligned} G(\gamma U_1, \alpha U_2) &= F(P_1(\gamma U_1, \alpha U_2), P_2(\gamma U_1, \alpha U_2)) \\ &= F(\alpha \gamma P_1(U_2, U_1), \alpha \gamma P_2(U_2, U_1)) \\ &= \alpha^n \gamma^n F(P_1(U_2, U_1), P_2(U_2, U_1)) \\ G(\gamma U_1, \alpha U_2) &= \alpha^n \gamma^n G(U_2, U_1) . \end{aligned}$$

Replacing  $G$  by  $\sum_{i=0}^{2n} b_i U_1^{2n-i} U_2^i$  on both sides of this last equality we obtain the result.

DEFINITION 3 : A sequence  $b = (b_0, b_1, \dots, b_{2n})$  satisfying :

$$b_{2n-i} = \rho^{n-i} b_i, \quad i = 0, 1, \dots, n, \quad \rho \neq 0 ,$$

is said to be  $\rho$ -reciprocal.

THEOREM 2 : Let  $BR[\omega]$  be a  $BR$ -curve of length  $n$  and let  $\Phi$  be the quadratic function defined as above. Then  $BR[\omega](\Phi(u))$  can be written in the following form :

$$BR[\omega](\Phi(u)) = BR[\theta_0, \theta_1, \dots, \theta_{2n}](u) .$$

The massic polygon  $\theta = (\theta_0, \theta_1, \dots, \theta_{2n})$  is  $\frac{\gamma}{\alpha}$ -reciprocal :

$$\theta_{2n-i} = \left( \frac{\gamma}{\alpha} \right)^{n-i} \theta_i, \quad i = 0, 1, \dots, n.$$

*Proof :* By  $(T_1, T_2)$  we denote the homogeneous cartesian coordinates of  $t$  relatively to the cartesian projective frame of reference  $(\infty, 0, 1)$  of the projective line  $\tilde{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$  :

$$t = \frac{T_1}{T_2} \quad \text{if } T_2 \neq 0, \quad t = \infty \quad \text{for } T_2 = 0.$$

By  $(U_1, U_2)$  we denote the homogeneous barycentric coordinates of  $u$  relatively to the barycentric projective frame of reference  $(0, 1, \frac{1}{2})$  of  $\tilde{\mathbb{R}}$  :

$$u \in \mathbb{R} : \begin{pmatrix} U_1 \\ U_2 \end{pmatrix} = \lambda \begin{pmatrix} 1-u \\ u \end{pmatrix}, \quad u = \infty : \begin{pmatrix} U_1 \\ U_2 \end{pmatrix} = \lambda \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad \lambda \neq 0.$$

The quadratic change of parameter  $t = \Phi(u)$  can be written again in the following form :

$$\begin{aligned} T_1 &= \alpha U_1^2 + 2\beta U_1 U_2 + \gamma U_2^2 \\ T_2 &= 2 U_1 U_2. \end{aligned}$$

By  $T_1(U_1, U_2)$  and  $T_2(U_1, U_2)$  we denote the two above-mentioned polynomials.  $\Delta$  being the symbol for forward difference operator, successively we have :

$$\begin{aligned} BR[\omega](t) &= \Pi \left( \sum_{i=0}^n B_i^n(t) \omega_i \right) \\ &= \Pi \left( \sum_{i=0}^n \binom{n}{i} t^i \Delta^i \omega_0 \right) \\ &= \Pi \left( \frac{1}{T_2^n} \sum_{i=0}^n \binom{n}{i} T_1^i T_2^{n-i} \Delta^i \omega_0 \right) \\ &= \Pi \left( \sum_{i=0}^n \binom{n}{i} T_1^i T_2^{n-i} \Delta^i \omega_0 \right). \end{aligned}$$

Let us define  $F(T_1, T_2) = \sum_{i=0}^n \binom{n}{i} T_1^i T_2^{n-i} \Delta^i \omega_0$ .  $F$  is a vector-valued homogeneous polynomial of degree  $n$  in  $T_1, T_2$ .

$T_1(U_1, U_2)$  and  $T_2(U_1, U_2)$  are homogeneous polynomials of degree 2 in  $U_1, U_2$  having the following property :

$$\begin{aligned} T_1(\gamma U_1, \alpha U_2) &= \alpha \gamma T_1(U_2, U_1) \\ T_2(\gamma U_1, \alpha U_2) &= \alpha \gamma T_2(U_2, U_1). \end{aligned}$$

From  $F$  we define  $G(U_1, U_2) = F(T_1(U_1, U_2), T_2(U_1, U_2))$ , so  $G$  is a homogeneous polynomial of degree  $2n$  in  $U_1, U_2$ . It can be written

$$G(U_1, U_2) = \sum_{j=0}^{2n} \binom{2n}{j} U_1^{2n-j} U_2^j \theta_j \quad \text{with } \theta_j \in \hat{\mathcal{E}}.$$

We have :

$$\begin{aligned} BR[\omega](\Phi(u)) &= \Pi(F(T_1(U_1, U_2), T_2(U_1, U_2))) \\ &= \Pi\left(\sum_{j=0}^{2n} \binom{2n}{j} U_1^{2n-j} U_2^j \theta_j\right) \\ &= \Pi\left(\lambda^{2n} \sum_{j=0}^{2n} B_j^{2n}(u) \theta_j\right) \\ &= \Pi\left(\sum_{j=0}^{2n} B_j^{2n}(u) \theta_j\right) \end{aligned}$$

$$BR[\omega](\Phi(u)) = BR[\theta_0, \theta_1, \dots, \theta_{2n}](u).$$

The polynomials  $F(T_1, T_2), T_1(U_1, U_2), T_2(U_1, U_2)$  satisfying the hypothesis of lemma 1, it follows that coefficients of  $G$  verify the relations

$$\begin{aligned} \binom{2n}{2n-i} \theta_{2n-i} &= \left(\frac{\gamma}{\alpha}\right)^{n-i} \binom{2n}{i} \theta_i, \quad i = 0, 1, \dots, n \\ \text{i.e., } \theta_{2n-i} &= \left(\frac{\gamma}{\alpha}\right)^{n-i} \theta_i, \quad i = 0, 1, \dots, n. \end{aligned}$$

COROLLARY 1 : *The support of  $BR[\theta](u)$  when  $u \in [0, 1[$  is identical to that of  $BR[\omega](t)$  when  $t \in \tilde{\mathbb{R}}$ .*

*Proof :* It comes from  $BR[\omega](u) = BR[\omega](\Phi(u))$  and the properties of  $\Phi$ .

Remark 2 : Theorem 2 states that it is sufficient to know  $\theta_0, \theta_1, \dots, \theta_n$  to determine entirely  $\theta = (\theta_0, \theta_1, \dots, \theta_{2n})$ .

PROPOSITION 1 : *With the notation above, for  $i = 0, 1, \dots, n$  we let*

$$T_1^i T_2^{n-i} = \sum_{j=0}^{2n} \alpha_j^i U_1^{2n-j} U_2^j.$$



By  $K$  we denote the  $(2n + 1) \times (n + 1)$  matrix with  $a'_j$  as entries at row  $j$  and column  $i$  and by  $\hat{K}$  the submatrix of  $K$  reduced to its  $n + 1$  first rows Then it follows :

- (i) The sequence of the  $2n + 1$  rows of  $K$  are  $\frac{\gamma}{\alpha}$ -reciprocal ( $\hat{K}$  determines  $K$  entirely),
- (ii) the entries of  $\hat{K}$  above the secondary diagonal are null ( $a'_i = 0$  for  $i + j = n - 1, n - 2, \dots, 0$ ),
- (iii) the rising secondary diagonal entries of  $\hat{K}$  are  $(2^n, \alpha 2^{n-1}, \alpha^2 2^{n-2}, \dots, \alpha^n)$ ,
- (iv) determinant of  $\hat{K} = (-2\alpha)^{\frac{n(n+1)}{2}}$ ,
- (v) the massic vectors  $\theta_0, \theta_1, \dots, \theta_n$  are related to  $\omega_0, \omega_1, \dots, \omega_n$  by the following matrix relation :

$$\begin{pmatrix} \theta_0 \\ \theta_1 \\ \vdots \\ \theta_n \end{pmatrix} = D_1^{-1} \hat{K} D_2 \begin{pmatrix} \omega_0 \\ \Delta \omega_0 \\ \vdots \\ \Delta^n \omega_0 \end{pmatrix}$$

with

$$D_1 = \text{diag} \left( \binom{2n}{0}, \binom{2n}{1}, \dots, \binom{2n}{n} \right)$$

and

$$D_2 = \text{diag} \left( \binom{n}{0}, \binom{n}{1}, \dots, \binom{n}{n} \right).$$

*Proof:* (i)  $T_1^i T_2^{n-i}$  is a homogeneous polynomial of degree  $n$  in  $T_1, T_2$ ;  $T_1(U_1, U_2)$  and  $T_2(U_1, U_2)$  are homogeneous polynomials of degree 2 in  $U_1, U_2$ . According to the lemma 1 we deduce that  $T_1^i T_2^{n-i} = \sum_{j=0}^{2n} a'_j U_1^{2n-j} U_2^j$  with  $a'_i = \left(\frac{\gamma}{\alpha}\right)^{n-j} a'_j, j = 0, 1, \dots, n$ .

The sequence  $(a'_0, a'_1, \dots, a'_i, \dots, a'_n)$  is the  $i$ -th column of  $K$  and is  $\frac{\gamma}{\alpha}$ -reciprocal for each  $i = 0, 1, \dots, n$ .

It follows that the sequence of rows of  $K = (K^0, K^1, \dots, K^n, \dots, K^{2n})$  is  $\frac{\gamma}{\alpha}$ -reciprocal :  $K^{2n-j} = \left(\frac{\gamma}{\alpha}\right)^{n-j} K^j$  for  $j = 0, 1, \dots, n$ .

- (ii) Expanding  $T_1^i T_2^{n-i}$  in decreasing powers of  $U_1$  :

$$\begin{aligned} T_1^i T_2^{n-i} &= (\alpha U_1^2 + 2\beta U_1 U_2 + \gamma U_2^2)^i 2^{n-i} U_1^{n-i} 2^{n-i} \\ &= 2^{n-i} \alpha^i U_1^{n+i} U_2^{n-i} + \dots, \end{aligned}$$

it gives :  $a_i^0 = a_i^1 = \dots = a_i^{n-i-1} = 0$  for  $i = 0, 1, \dots, n-1$  and  $a_i^{n-i} = 2^{n-i} \alpha^i$  for  $i = 0, 1, \dots, n$ .

(iii) From above the rising secondary diagonal is  $(2^n, 2^{n-1} \alpha, \dots, 2^{n-i} \alpha^i, \dots, \alpha^n)$ .

(iv) Determinant of  $\hat{K}$  is equal to  $(-1)^{\frac{n(n+1)}{2}}$  times the product of the rising secondary diagonal entries i.e., determinant of  $\hat{K} = (-2 \alpha)^{\frac{n(n+1)}{2}}$ .

(v) We know (previous proof) that :

$$BR[\omega](t) = \Pi \left( \sum_{i=0}^n \binom{n}{i} \Delta^i \omega_0 T_1^i T_2^{n-i} \right),$$

then

$$\begin{aligned} BR[\omega](\Phi(u)) &= \Pi \left( \sum_{i=0}^n \binom{n}{i} \Delta^i \omega_0 \left( \sum_{j=0}^{2n} a_j^i U_1^{2n-j} U_2^j \right) \right) \\ &= \Pi \left( \sum_{j=0}^{2n} \left( \sum_{i=0}^n \binom{n}{i} a_j^i \Delta^i \omega_0 \right) U_1^{2n-j} U_2^j \right) \\ &= \Pi \left( \sum_{j=0}^{2n} \binom{2n}{j} U_1^{2n-j} U_2^j \theta_j \right) \end{aligned}$$

with

$$\theta_j = \frac{1}{\binom{2n}{j}} \sum_{i=0}^n \binom{n}{i} a_j^i \Delta^i \omega_0, \quad j = 0, 1, \dots, 2n.$$

By the  $\frac{\gamma}{\alpha}$ -reciprocal property of the  $\theta_i, i = 0, 1, \dots, 2n$ , we only calculate the  $(n+1)$  first massic vectors  $\theta_i$ . They are related with  $\omega_0, \omega_1, \dots, \omega_n$  by the matrix relation given in the proposition with

$$\hat{K} = \begin{pmatrix} 0 & 0 & \dots & 0 & \dots & 0 & a_n^0 \\ 0 & 0 & \dots & 0 & \dots & a_{n-1}^1 & a_n^1 \\ \vdots & \vdots & & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & a_i^{n-i} & \dots & a_{n-1}^{n-i} & a_n^{n-i} \\ \vdots & \vdots & & \vdots & & \vdots & \vdots \\ 0 & a_1^{n-1} & \dots & a_i^{n-1} & \dots & a_{n-1}^{n-1} & a_n^{n-1} \\ a_0^n & a_1^n & \dots & a_i^n & \dots & a_{n-1}^n & a_n^n \end{pmatrix}$$

*Remark 3.* — Determination of matrix  $D_1^{-1} \hat{K} D_2$  : The entries of the  $i$ -th column of matrix  $\hat{K}$  are the  $(n+1)$  first coefficients of the expansion of

$T_1 T_2^{n-1} = (\alpha U_1^2 + 2 \beta U_1 U_2 + \gamma U_2^2)' (2 U_1 U_2)^{n-1}$  with respect to  $U_1^{2n}, U_1^{2n-1} U_2, \dots, U_1^n U_2^n, \dots, U_2^{2n}$ .

For example we obtains, as matrix  $D_1^{-1} \hat{K} D_2$ ,

$n=2$  :

$$\begin{pmatrix} 0 & 0 & \alpha^2 \\ 0 & \alpha & \alpha \beta \\ \frac{2}{3} & \frac{4}{3} \beta & \frac{1}{3} (2 \beta^2 + \alpha \gamma) \end{pmatrix}$$

$n = 3$  :

$$\begin{pmatrix} 0 & 0 & 0 & \alpha^3 \\ 0 & 0 & \alpha^2 & \alpha^2 \beta \\ 0 & \frac{4}{5} \alpha & \frac{8}{5} \alpha \beta & \frac{1}{5} (4 \alpha \beta^2 + \alpha^2 \gamma) \\ \frac{2}{5} & \frac{6}{5} \beta & \frac{3}{5} (2 \beta^2 + \alpha \gamma) & \frac{1}{5} (3 \alpha \beta \gamma + 2 \beta^3) \end{pmatrix}$$

$n = 4$  :

$$\begin{pmatrix} 0 & 0 & 0 & 0 & \alpha^4 \\ 0 & 0 & 0 & \alpha^3 & \alpha^3 \beta \\ 0 & 0 & \frac{6}{7} \alpha^2 & \frac{12}{7} \alpha^2 \beta & \frac{1}{7} (6 \alpha^2 \beta^2 + \alpha^3 \gamma) \\ 0 & \frac{4}{7} \alpha & \frac{12}{7} \alpha \beta & \frac{1}{7} (12 \alpha \beta^2 + 3 \alpha^2 \gamma) & \frac{1}{7} (4 \alpha \beta^3 + 3 \alpha^2 \beta \gamma) \\ \frac{8}{35} & \frac{32}{35} \beta & \frac{8}{35} (6 \beta^2 + 3 \alpha \gamma) & \frac{8}{35} (4 \beta^3 + 6 \alpha \beta \gamma) & \frac{8}{35} \left( \beta^4 + 3 \alpha \beta^2 \gamma + \frac{3}{8} \alpha^2 \gamma^2 \right) \end{pmatrix}.$$

For greater values of  $n$ , the expansion of  $T_1 T_2^{n-1}$  and matrix  $D_1^{-1} \hat{K} D_2$  can be obtained using computer algebra systems.

Propositions 2 and 3 below lead to an algorithm for another calculation of  $\hat{K}$ .

*Example 1 : Folium of Descartes defined over  $[0, 1]$*

Suppose the plane  $\mathcal{E}$  and the space  $\mathcal{F}$  have a cartesian frame of reference respectively  $(\Omega_1, \vec{i}, \vec{j})$  and  $(\Omega, \vec{i}, \vec{j}, \vec{k} = \overrightarrow{\Omega \Omega_1})$ .

The Folium of Descartes (1638) is the curve represented parametrically by the point  $M(t)$  with coordinates :

$$x(t) = \frac{3 t}{1 + t^3}, \quad t \in \tilde{\mathbb{R}}$$

$$y(t) = \frac{3 t^2}{1 + t^3}$$

or in homogeneous cartesian coordinates :

$$X(t) = 3 t$$

$$Y(t) = 3 t^2$$

$$Z(t) = 1 + t^3 .$$

We know [6, proposition 2.1.4.1] that

$$M(t) = \Pi\Omega (X(t) \vec{i} + Y(t) \vec{j} + Z(t) \vec{k})$$

$$= \Pi\Omega (\vec{k} + 3 t \vec{i} + 3 t^2 \vec{j} + t^3 \vec{k}) .$$

As  $\Pi\Omega = \Pi \circ \hat{\Omega}^{-1}$  and  $\hat{\Omega}^{-1}(\vec{k}) = (\Omega_1 ; 1)$ ,  $\hat{\Omega}^{-1}(\vec{i}) = \vec{i}$ ,  $\hat{\Omega}^{-1}(\vec{j}) = \vec{j}$ , it becomes

$$M(t) = \Pi((\Omega_1 ; 1) \oplus 3 t \vec{i} \oplus 3 t^2 \vec{j} \oplus t^3 (\Omega_1 ; 1)) .$$

Therefore there exists a massic polygon  $\omega = (\omega_0, \omega_1, \omega_2, \omega_3)$  such that :

$$\omega_0 = (\Omega_1 ; 1)$$

$$\Delta \omega_0 = \vec{i}$$

$$\Delta^2 \omega_0 = \vec{j}$$

$$\Delta^3 \omega_0 = (\Omega_1 ; 1) .$$

Then

$$M(t) = \Pi(\omega_0 \oplus 3 t \Delta \omega_0 \oplus 3 t^2 \Delta^2 \omega_0 \oplus t^3 \Delta^3 \omega_0)$$

$$= \Pi \left( \sum_{i=0}^3 B_i^3(t) \omega_i \right)$$

$$= BR [\omega_0, \omega_1, \omega_2, \omega_3] (t)$$

with :  $\omega_0 = (\Omega_1 ; 1)$ ,  $\omega_1 = (\Omega_1 + \vec{i} ; 1)$ ,  $\omega_2 = (\Omega_1 + 2 \vec{i} + \vec{j} ; 1)$ ,  
 $\omega_3 = \left( \Omega_1 + \frac{3}{2} (\vec{i} + \vec{j}) ; 2 \right)$ .

Defining  $P = \Omega_1 + \vec{i}$ ,  $Q = \Omega_1 + 2\vec{i} + \vec{j}$ ,  $R = \Omega_1 + \frac{3}{2}(\vec{i} + \vec{j})$ , we obtain explicitly :

$$M(t) = \frac{B_0^3(t) \Omega_1 + B_1^3(t) P + B_2^3(t) Q + 2 B_3^3(t) R}{B_0^3(t) + B_1^3(t) + B_2^3(t) + 2 B_3^3(t)}, \quad t \in \tilde{\mathbb{R}}.$$

Again we find the massic polygon given in [6, section 2.3.2]. The image of  $[0, 1]$  is the anticlockwise arc  $(\Omega_1, R)$ , (fig. 1).




Figure 1.

We make the quadratic change of parameter  $t = \Phi(u)$ . We obtain

$$M \circ \Phi = BR[\omega] \circ \Phi = BR[\theta_0, \theta_1, \theta_2, \theta_3, \theta_4, \theta_5, \theta_6]$$

with  $\theta = (\theta_0, \theta_1, \dots, \theta_6)$ , a  $\rho$ -reciprocal massic polygon such that

$$\begin{pmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \theta_3 \end{pmatrix} = D_1^{-1} \hat{K} D_2 \begin{pmatrix} \omega_0 \\ \Delta \omega_0 \\ \Delta^2 \omega_0 \\ \Delta^3 \omega_0 \end{pmatrix}.$$

From remark 3 ( $n = 3$ ), within the specific case  $\alpha = -1$ ,  $\beta = 0$ ,  $\gamma = 1$  ( $\rho = -1$ ) we obtain

$$\begin{pmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \theta_3 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -\frac{4}{5} & 0 & \frac{1}{5} \\ \frac{2}{5} & 0 & -\frac{3}{5} & 0 \end{pmatrix} \begin{pmatrix} (\Omega_1; 1) \\ \vec{i} \\ \vec{j} \\ (\Omega_1; 1) \end{pmatrix}$$

and finally :

$$\begin{aligned} \theta_0 &= (\Omega_1; -1) \\ \theta_1 &= \vec{j} \\ \theta_2 &= \left( \Omega_1 - 4\vec{i}; \frac{1}{5} \right) \\ \theta_3 &= \left( \Omega_1 - \frac{3}{2}\vec{j}; \frac{2}{5} \right) \\ \theta_4 &= -\theta_2 \\ \theta_5 &= \theta_1 \\ \theta_6 &= -\theta_0. \end{aligned}$$

Defining  $A = \Omega_1 - 4\vec{i}$ ,  $B = \Omega_1 - \frac{3}{2}\vec{j}$ , we obtain explicitly

$$\begin{aligned} M(u) &= \frac{-B_0^6(u)\Omega_1 + \frac{1}{5}B_2^6(u)A + \frac{2}{5}B_3^6(u)B - \frac{1}{5}B_4^6(u)A + B_6^6(u)\Omega_1}{-B_0^6(u) + \frac{1}{5}B_2^6(u) + \frac{2}{5}B_3^6(u) - \frac{1}{5}B_4^6(u) + B_6^6(u)} \\ &+ \frac{B_1^6(u)\vec{j} + B_5^6(u)\vec{j}}{-B_0^6(u) + \frac{1}{5}B_2^6(u) + \frac{2}{5}B_3^6(u) - \frac{1}{5}B_4^6(u) + B_6^6(u)}, \quad u \in [0, 1[. \end{aligned}$$

In [6, section 2.3.2] we give a slightly different massic polygon corresponding to  $\alpha = 1$ ,  $\beta = 0$ ,  $\gamma = -1$ .

The anticlockwise loop (fig. 1) corresponds to  $u \in \left[ \frac{1}{2}, 1 \right[$  (resp. to  $t \in [0, +\infty[$ ), segment on the second quadrant to  $u \in \left[ 1 - \frac{\sqrt{2}}{2}, \frac{1}{2} \right]$  (resp. to  $t \in [-1, 0)$ ) and the segment on the fourth quadrant to  $u \in \left] 0, 1 - \frac{\sqrt{2}}{2} \right]$  (resp. to  $t \in ]-\infty, -1)$ ).

**COROLLARY 2 :** Denoting the mass of a massic vector  $\theta$  by  $\chi(\theta)$  we have the relation

$$\begin{pmatrix} \chi(\theta_0) \\ \chi(\theta_1) \\ \vdots \\ \chi(\theta_n) \end{pmatrix} = D_1^{-1} \hat{K} D_2 \begin{pmatrix} \chi(\omega_0) \\ \Delta\chi(\omega_0) \\ \vdots \\ \Delta^n\chi(\omega_0) \end{pmatrix}.$$

The massic sequence  $(\chi(\theta_0), \chi(\theta_1), \dots, \chi(\theta_n), \dots, \chi(\theta_{2n}))$  is  $\frac{\gamma}{\alpha}$ -reciprocal.

*Proof :* We know [6, proposition 1.2.2.8] that  $\chi$  is a linear form :  $\hat{\mathcal{E}} \rightarrow \mathbb{R}$ , the relation  $(\nu)$  of proposition 1 giving the  $\theta_i$  as a function of  $\omega_i$  remains true by considering the masses, on both sides of the equality.

**COROLLARY 3 :** Let  $\theta = (\theta_0, \theta_1, \dots, \theta_{2n})$  be a  $\rho$ -reciprocal massic polygon with  $\rho < 0$ . Choosing  $\alpha$  and  $\gamma$  such that  $\rho = \frac{\gamma}{\alpha}$ , then there exists a massic polygon  $\omega = (\omega_0, \omega_1, \dots, \omega_n)$  such that

$$BR[\omega] \circ \Phi = BR[\theta]$$

$\Phi$  is the quadratic function previously given ( $\beta$  is arbitrary). It is sufficient to define the massic polygon  $\omega$  by the relation :

$$\begin{pmatrix} \omega_0 \\ \Delta\omega_0 \\ \vdots \\ \Delta^n\omega_0 \end{pmatrix} = D_2^{-1} \hat{K}^{-1} D_1 \begin{pmatrix} \theta_0 \\ \theta_1 \\ \vdots \\ \theta_n \end{pmatrix}.$$

*Proof :* The scalar  $\alpha$  is different from zero, so  $\hat{K}$  is invertible and we deduce  $(\omega_0, \omega_1, \dots, \omega_n)$  from equation  $(\nu)$  proposition 1. The condition  $\rho < 0$  guarantees that the quadratic correspondence  $t = \Phi(u)$  is a one-to-one correspondence between  $[0, 1[$  and  $\tilde{\mathbb{R}}$ .

*Example 2 : Curve of Agnesi*

We consider the curve of Agnesi (1748) also named versiera, was first considered by Fermat (1666), is given parametrically by the point  $M(u)$  with cartesian coordinates :

$$\begin{aligned} x(u) &= \frac{2u - 1}{2u(1 - u)} \\ y(u) &= \frac{4u^2(1 - u)^2}{4u^4 - 8u^3 + 8u^2 - 4u + 1}, \quad u \in [0, 1[. \end{aligned}$$

Its homogeneous cartesian coordinates are :

$$\begin{aligned} X(u) &= 8 u^5 - 20 u^4 + 24 u^3 - 16 u^2 + 6 u - 1 \\ Y(u) &= -8 u^6 + 24 u^5 - 24 u^4 + 8 u^3 \\ Z(u) &= -8 u^6 + 24 u^5 - 32 u^4 + 24 u^3 - 10 u^2 + 2 u . \end{aligned}$$

We obtain the following *BR*-form of this rational curve by the method illustrated in example 1 or by that of [6, section 2.2.2.3] :

$$M(u) = BR[\theta_0, \theta_1, \theta_2, \theta_3, \theta_4, \theta_5, \theta_6] (u)$$

with  $\theta_0 = \vec{V}_0$ ,  $\vec{V}_0 = (-1, 0)$ ;  $\theta_1 = \left(Q_1; \frac{1}{3}\right)$ ,  $Q_1 = (0, 0)$ ,  $\theta_2 = \vec{V}_2$ ,  $\vec{V}_2 = \left(-\frac{1}{15}, 0\right)$ ,  $\theta_3 = \left(Q_3; \frac{1}{5}\right)$ ,  $Q_3 = (0, 2)$ ;  $\theta_4 = -\theta_2$ ,  $\theta_5 = \theta_1$ ,  $\theta_6 = -\theta_0$ . In figure 2 the length of pure vectors  $\theta_0, \theta_2, \theta_4, \theta_6$  has been multiplied by 2.




Figure 2.

Explicitly we have :

$$\begin{aligned} M(u) &= \frac{\frac{1}{3} B_1^6(u) Q_1 + \frac{1}{5} B_3^6(u) Q_3 + \frac{1}{3} B_5^6(u) Q_1}{\frac{1}{3} B_1^6(u) + \frac{1}{5} B_3^6(u) + \frac{1}{3} B_5^6(u)} \\ &+ \frac{B_0^6(u) \vec{V}_0 + B_2^6(u) \vec{V}_2 - B_4^6(u) \vec{V}_2 - B_6^6(u) \vec{V}_0}{\frac{1}{3} B_1^6(u) + \frac{1}{5} B_3^6(u) + \frac{1}{3} B_5^6(u)}, \quad u \in [0, 1[ . \end{aligned}$$

This massic polygon being  $\rho$ -reciprocal with  $\rho = -1$  then by corollary 3 this *BR*-curve comes from a *BR*-curve of length 3 by a quadratic change of



parameter. More precisely there exists a massic polygon  $\omega = (\omega_0, \omega_1, \omega_2, \omega_3)$  such that

$$BR[\omega] \circ \Phi = BR[\theta].$$

Taking  $\alpha = -1, \beta = 0, \gamma = 1, (\rho = -1)$  we obtain :

$$D_2^{-1} \hat{K}^{-1} D_1 = \begin{pmatrix} 0 & \frac{3}{2} & 0 & \frac{5}{2} \\ -\frac{1}{4} & 0 & -\frac{5}{4} & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$$

which gives  $\omega_0, \Delta\omega_0, \Delta^2\omega_0, \Delta^3\omega_0$  and finally :

$$\omega_0 = (P_0; 1), \quad P_0 = (0, 1); \quad \omega_1 = (P_1; 1), \quad P_1 = \left(\frac{1}{3}, 1\right);$$

$$\omega_2 = \left(P_2; \frac{4}{3}\right), \quad P_2 = \left(\frac{1}{2}, \frac{3}{4}\right); \quad \omega_3 = (P_3; 2), \quad P_3 = \left(1, \frac{1}{2}\right), \quad (\text{fig. 3}).$$

Explicitly  $M(\Phi^{-1}(t)) = BR[\omega](t)$  is given by :

$$BR[\omega](t) = \frac{B_0^3(t)P_0 + B_1^3(t)P_1 + \frac{4}{3}B_2^3(t)P_2 + 2B_3^3(t)P_3}{B_0^3(t) + B_1^3(t) + \frac{4}{3}B_2^3(t) + 2B_3^3(t)}, \quad t \in \tilde{\mathbb{R}}.$$




Figure 3.

4. ALGORITHM FOR CALCULATING MATRIX  $\hat{K}$

PROPOSITION 2 :

(i)  $T_1^i$  must be written

$$T_1^i = \sum_{j=0}^{2i} \alpha_j^i U_1^{2i-j} U_2^j.$$

(ii)  $(\alpha_j^i), j = 0, 1, \dots, 2i$ , are  $\frac{\gamma}{\alpha}$ -reciprocal i.e.,  $\alpha_{2i-k}^i = \left(\frac{\gamma}{\alpha}\right)^{i-k} \alpha_k^i, k = 0, 1, \dots, i$ .

(iii) The following algorithm gives the coefficients  $(\alpha_j^i)$  of polynomials  $T_1^i$ ,  $i = 2, 3, \dots, n$  ( $T_1^1 = \alpha U_1^2 + 2 \beta U_1 U_2 + \gamma U_2^2$ ): initialization :  $\alpha_{-2}^1 = \alpha_{-1}^1 = 0$ ,  $\alpha_0^1 = \alpha$ ,  $\alpha_1^1 = 2 \beta$ ,  $\alpha_2^1 = \gamma$ ,  $\alpha_3^1 = 0$ ,  $\rho = \frac{\gamma}{\alpha}$  ;

for  $i = 2, 3, \dots, n$   
 for  $j = 0, 1, \dots, i$  do  
 $\alpha_j^i = \alpha \alpha_j^{i-1} + 2 \beta \alpha_{j-1}^{i-1} + \gamma \alpha_{j-2}^{i-1}$   
 $\alpha_{2i-j}^i = \rho^{i-j} \alpha_j^i$  (if  $j < i$ )  
 $\alpha_{-2}^i = \alpha_{-1}^i = 0$ .

*Proof :*

(i) Expanding  $T_1^i = (\alpha U_1^2 + 2 \beta U_1 U_2 + \gamma U_2^2)^i$ ,  $T_1^i$  can be written

$$T_1^i = \sum_{j=0}^{2i} \alpha_j^i U_1^{2i-j} U_2^j.$$

(ii)  $T_1^i$  is a homogeneous polynomial of degree  $i$  in  $T_1, T_2$ , according to lemma 1,  $T_1^i$  has  $\frac{\gamma}{\alpha}$ -reciprocal coefficients.

(iii)

$$\begin{aligned} T_1^{i-1} &= \sum_{j=0}^{2i-2} \alpha_j^{i-1} U_1^{2i-2-j} U_2^j \\ T_1^i &= T_1 T_1^{i-1} = (\alpha U_1^2 + 2 \beta U_1 U_2 + \gamma U_2^2) T_1^{i-1} \\ T_1^i &= \sum_{j=0}^{2i-2} \alpha \alpha_j^{i-1} U_1^{2i-j} U_2^j + \sum_{j=0}^{2i-2} 2 \beta \alpha_j^{i-1} U_1^{2i-1-j} U_2^{j+1} + \\ &+ \sum_{j=0}^{2i-2} \gamma \alpha_j^{i-1} U_1^{2i-2-j} U_2^{j+2}. \end{aligned}$$

We define  $\alpha_{-2}^{i-1} = \alpha_{-1}^{i-1} = \alpha_{2i-1}^{i-1} = \alpha_{2i}^{i-1} = 0$ . By some direct change of indices,  $T_1^i$  becomes :

$$T_1^i = \sum_{j=0}^{2i} (\alpha \alpha_j^{i-1} + 2 \beta \alpha_{j-1}^{i-1} + \gamma \alpha_{j-2}^{i-1}) U_1^{2i-j} U_2^j.$$

Then  $\alpha_j^i = \alpha \alpha_j^{i-1} + 2 \beta \alpha_{j-1}^{i-1} + \gamma \alpha_{j-2}^{i-1}$ ,  $j = 0, 1, \dots, 2i$ .

The  $(\alpha_j^i)$  being  $\frac{\gamma}{\alpha}$ -reciprocal, we calculate  $(\alpha_j^i)$ , for  $j = 0, 1, \dots, i$ , by this relation and we deduce  $\alpha_{2i-j}^i = \left(\frac{\gamma}{\alpha}\right)^{i-j} \alpha_j^i$ ,  $j = 0, 1, \dots, i-1$ .

PROPOSITION 3 : The entries  $a_i^j$  of the  $(n + 1) \times (n + 1)$  matrix  $\hat{K}$  are given by :

$$\begin{aligned} a_i^j &= 0 \quad \text{for } i + j = n - 1, n - 2, \dots, 0 \\ a^{n-i+j} &= 2^{n-i} \alpha_j^i, \quad i = 0, 1, \dots, n, \quad j = 0, 1, \dots, i \\ &(\alpha_0^0 = 1). \end{aligned}$$

*Proof* : The first result was proved proposition 1(ii).

From  $T_1^i = \sum_{j=0}^{2i} \alpha_j^i U_1^{2i-j} U_2^j$  we obtain

$$\begin{aligned} T_1^i T_2^{n-i} &= \left( \sum_{j=0}^{2i} \alpha_j^i U_1^{2i-j} U_2^j \right) (2 U_1 U_2)^{n-i} \\ &= \sum_{j=0}^{2i} 2^{n-i} \alpha_j^i U_1^{n-j+i} U_2^{n+j-i}. \end{aligned} \tag{I}$$

In proposition 1 we have defined  $T_1^i T_2^{n-i} = \sum_{j=0}^{2n} a_j^i U_1^{2n-j} U_2^j$ . From (i) and (ii) of this proposition we deduce :

$$T_1^i T_2^{n-i} = \sum_{k=n-i}^{n+i} a_k^i U_1^{2n-k} U_2^k.$$

By changing the indices :  $j = k - n + i$ , it becomes

$$T_1^i T_2^{n-i} = \sum_{j=0}^{2i} a_i^{n+j-i} U_1^{n-j+i} U_2^{n+j-i}.$$

By comparing with equality (I) we deduce :

$$a_i^{n+j-i} = 2^{n-i} \alpha_j^i, \quad i = 0, 1, \dots, n, \quad j = 0, 1, \dots, i.$$

In proposition 1 (iii) we have seen that  $a_0^n = 2^n$ , so we define  $\alpha_0^0 = 1$ . Then we deduce the following algorithm giving the non-zero entries of  $\hat{K}$ .

ALGORITHM : Initialization :  $\alpha_{-2}^1 = \alpha_{-1}^1 = 0, \quad \alpha_0^1 = \alpha, \quad \alpha_1^1 = 2\beta,$   
 $\alpha_2^1 = \gamma, \quad a_0^n = 2^n, \quad a_1^{n-1} = \alpha 2^{n-1}, \quad a_i^n = \beta 2^n;$

for  $i = 2, 3, \dots, n$

for  $j = 0, 1, \dots, i$  do

$$\alpha_j^i = \alpha \alpha_j^{i-1} + 2\beta \alpha_{j-1}^{i-1} + \gamma \alpha_{j-2}^{i-1}$$

$$a_i^{n-i+j} = 2^{n-i} \alpha_j^i$$

$$\alpha_{i+1}^i = \frac{\gamma}{\alpha} \alpha_{i-1}^i$$

$$\alpha_{-2}^i = \alpha_{-1}^i = 0.$$

5. REPRESENTATION OF A COMPLETE POLYNOMIAL CURVE BY A BR-CURVE

If we have to calculate a complete polynomial curve, i.e., for parameter  $t \in \tilde{\mathbb{R}}$ , calculating with its Bézier representation can be inaccurate. Actually for points corresponding to values of  $t$  not belonging to  $[0, 1]$ , we use, in the De Casteljaou's algorithm, affine combinations of points which are not convex and coefficients of these combinations tend to infinity. Proposition 4 below allows us to describe a Bézier curve as a BR-curve with parameter  $u \in [0, 1[$ . Hence calculating any point of this BR-curve by algorithm ALBR1 or ALBR2 [6, sections 3.3 and 3.4] (algorithms related to De Casteljaou's), this time we manipulate convex affine combinations of points or massic vectors which offers a guarantee of stability.

PROPOSITION 4 : *If  $\omega_i = (P_i ; c)$ ,  $i = 0, 1, \dots, n$ ,  $c$  being the common mass, then*

(i)  $BR[\omega] = BP[P]$ , i.e., a Bézier curve,

(ii)  $BP[P] \circ \Phi(u) = BR[\theta](u)$  where the  $\theta_i$  are all pure vectors ( $\theta_i \in \vec{\mathcal{E}}$ ) except  $\theta_n$  which is a weighted point of mass equal to  $\frac{2^n}{\binom{2n}{n}} c$ .

More precisely

$$\theta_j = \frac{1}{\binom{2n}{j}} \sum_{i=n-j}^n \binom{n}{i} a_i^j \Delta^i P_0, \quad j = 0, 1, \dots, n-1$$

$$\theta_n = \left( P_0 + \frac{1}{2^n c} \sum_{i=1}^n \binom{n}{i} a_i^n \Delta^i P_0 ; \frac{2^n}{\binom{2n}{n}} c \right).$$

The massic polygon  $\theta$  being  $\left(\frac{\gamma}{\alpha}\right)$ -reciprocal.

(iii) *The support of  $BR[\theta](u)$  when  $u \in [0, 1[$  is identical to that of  $BP[P](t)$  when  $t \in \tilde{\mathbb{R}}$ .*

(iv) *Conversely if the massic polygon  $\theta = (\theta_0, \theta_1, \dots, \theta_{2n})$  is  $\rho$ -reciprocal and if all  $\theta_i$  are pure vectors except  $\theta_n$  then there exist points of  $\mathcal{E} : P_0, P_1, \dots, P_n$ , such that :*

$$BP[P_0, P_1, \dots, P_n](\Phi(u)) = BR[\theta](u)$$

with  $\Phi$  a quadratic function defined as previously, the constants  $\alpha$  and  $\gamma$  satisfying  $\frac{\gamma}{\alpha} = \rho$ ,  $\beta$  arbitrary.

*Proof :*

(i) We have  $\omega_i = (P_i ; c)$ ,  $i = 0, 1, \dots, n$ . From remark 1 (ii) :  $BR[\omega](t) = BP[P](t)$ .

(ii) The  $\omega_i$  have same mass then  $\Delta\chi(\omega_0) = \dots = \Delta^n\chi(\omega_0) = 0$ . Corollary 2 implies  $\chi(\theta_0) = \dots = \chi(\theta_{n-1}) = 0$  i.e.,  $\theta_0, \theta_1, \dots, \theta_{n-1}$  are pure vectors and  $\chi(\theta_n) = \frac{2^n}{\binom{2n}{n}} \chi(\theta_0)$ .

From proposition 1 ( $\nu$ ) we deduce that

$$\theta_j = \frac{1}{\binom{2n}{j}} \sum_{i=n-j}^n \binom{n}{i} \alpha_i \Delta^i \omega_0, \quad j = 0, 1, \dots, n.$$

For  $i = 1, 2, \dots, n$  :  $\Delta^i \omega_0 = \Delta^i P_0$ . Distinguishing the case  $j = n$  and  $j = 0, 1, \dots, n - 1$  we obtain the announced result.

(iii) is identical to corollary 1.

(iv) Conversely let  $\theta = (\theta_0, \theta_1, \dots, \theta_{2n})$  be a  $\rho$ -reciprocal massic polygon with all  $\theta_i$  being pure vectors except  $\theta_n$ . Following Corollary 3 we define  $\omega = (\omega_0, \omega_1, \dots, \omega_n)$  by :

$$\begin{pmatrix} \omega_0 \\ \Delta\omega_0 \\ \vdots \\ \Delta^n\omega_0 \end{pmatrix} = D_2^{-1} \hat{K}^{-1} D_1 \begin{pmatrix} \theta_0 \\ \theta_1 \\ \vdots \\ \theta_n \end{pmatrix}$$

with  $D_1, D_2, \hat{K}$  defined in proposition 1 ( $\rho = \frac{\gamma}{\alpha}$ ,  $\beta$  arbitrary). We have  $BR[\omega](\Phi(u)) = BR[\theta](u)$  and

$$\begin{pmatrix} \chi(\omega_0) \\ \Delta\chi(\omega_0) \\ \vdots \\ \Delta^{n-1}\chi(\omega_0) \\ \Delta^n\chi(\omega_0) \end{pmatrix} = D_2^{-1} \hat{K}^{-1} D_1 \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ x(\theta_n) \end{pmatrix}.$$

But matrices  $\hat{K}^{-1}$  and  $D_2 \hat{K}^{-1} D_1$  have zero entries under the secondary diagonal i.e., if  $b'_i$  denote the entries of  $\hat{K}^{-1}$  we have  $b'_i = 0$  for  $i + j = n + 1, n + 2, \dots, 2n$ .

Consequently  $\chi(\omega_0) \neq 0$  and  $\Delta\chi(\omega_0) = \dots = \Delta^n\chi(\omega_0) = 0$  i.e.,  $\chi(\omega_i) = \chi(\omega_0) = c$ ;  $\omega_i$  can be written  $(P_i ; c)$ ,  $P_i \in \mathcal{E}$  and  $BR[\omega] = BP[P]$  i.e., a Bézier curve of length  $n$ .

*Example 3. — Cubic of Tschirnausen*

The cubic of Tschirnausen (1690) also named trisectrix of Catalan (1501-1576) or L'Hospital's cubic (1696) is the polynomial curve given parametrically by the point  $M(t)$  with cartesian coordinates :

$$\begin{aligned} x(t) &= 3(t^2 - 3) \\ y(t) &= t(t^2 - 3), \quad t \in \tilde{\mathbb{R}}. \end{aligned}$$

Its Bézier representation follows directly :

$$M(t) = BP [P_0, P_1, P_2, P_3] (t), \quad t \in \tilde{\mathbb{R}},$$

with  $P_0 = (-9, 0)$ ,  $P_1 = (-9, -1)$ ,  $P_2 = (-8, -2)$ ,  $P_3 = (-6, -2)$  and  $BP [P] = BR [\omega]$  with  $\omega_i = (P_i ; 1)$ ,  $i = 0, 1, 2, 3$  (fig. 4).




Figure 4.

By making the quadratic change of parameter  $t = \Phi(u)$  we obtain :

$$BP [P_0, P_1, P_2, P_3] \circ \Phi(u) = BR [\theta_0, \theta_1, \dots, \theta_6] (u).$$

From  $\omega_i = (P_i ; 1)$  we deduce

$$\begin{aligned} \omega_0 = (P_0 ; 1), \quad P_0 = (-9, 0); \quad \Delta\omega_0 = \Delta P_0 = -\vec{j}; \\ \Delta^2\omega_0 = \Delta^2 P_0 = \vec{i}; \quad \Delta^3\omega_0 = \Delta^3 P_0 = \vec{j}. \end{aligned}$$

We take again  $\alpha = -1$ ,  $\beta = 0$ ,  $\gamma = 1$  ( $\rho = -1$ ) in the definition of  
vol. 27, n° 3, 1993

$\Phi$ , then proposition 1 ( $\nu$ ) and matrix  $D_1^{-1} \hat{K}D_2$  of example 1 give :

$$\begin{pmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \\ \theta_3 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -\frac{4}{5} & 0 & \frac{1}{5} \\ \frac{2}{5} & 0 & -\frac{3}{5} & 0 \end{pmatrix} \begin{pmatrix} (P_0; 1) \\ -\vec{j} \\ \vec{i} \\ \vec{j} \end{pmatrix}$$

$$\theta_0 = -\vec{j}$$

$$\theta_1 = \vec{i}$$

$$\theta_2 = \vec{j}$$

$$\theta_3 = \left( P_0 - \frac{3}{2} \vec{i}; \frac{2}{5} \right)$$

$$\theta_4 = -\theta_2$$

$$\theta_5 = \theta_1$$

$$\theta_6 = -\theta_0.$$

Defining  $Q_3 = P_0 - \frac{3}{2} \vec{i}$ ,  $Q_3 = \left( -\frac{21}{2}, 0 \right)$ . For more clarity in figure 4, the length of vectors  $\vec{i}$  and  $\vec{j}$  has been multiplied by 3.

$$BR[\theta](u) = Q_3 + \frac{-B_0^6(u)\vec{j} + B_1^6(u)\vec{i} + B_2^6(u)\vec{j} - B_4^6(u)\vec{j} + B_5^6(u)\vec{i} + B_6^6(u)\vec{j}}{B_3^6(u)},$$

$$u \in [0, 1[.$$

### 6. CONCLUSION

For a complete description of a rational curve or *a fortiori* of a polynomial curve we use its *BR*-representation. To describe its whole support we have resorted to special quadratic change of parameter which is a one-to-one correspondence between  $\tilde{\mathbb{R}}$  and  $[0, 1[$ . We have determined a new massic polygon from the old one by linear relations. This new massic polygon presents a kind of symmetry called  $\rho$ -reciprocity which reduces the computation. It enables us to store the whole support as *BR*-curve and to compute it by convex affine combinations of points or massic vectors.

## REFERENCES

- [1] P. BÉZIER, Procédé de définition numérique des courbes et surfaces non mathématiques, système Unisurf, *Automatisme 13*, 1968.
- [2] P. DE CASTELJAU, *Outillage, méthode de calcul*, André Citroën Automobile S.A., Paris 1959.
- [3] G. FARIN, Algorithms for rational Bézier curves, *Computer Aided Design. 15* (1983) pp. 73-77.
- [4] I. E. FAUX and M. J. PRATT, *Computational geometry for design and manufacture*, Ellis Horwood, Chichester, 1979.
- [5] J. C., FIOROT and P. JEANNIN, Nouvelle description et calcul des courbes rationnelles à l'aide de points et vecteurs de contrôle, *C.R. Acad. Sci. Paris*, t. 305, Série I (1987) pp. 435-440.
- [6] J. C. FIOROT and P. JEANNIN, *Courbes et Surfaces Rationnelles. Applications à la CAO, RMA12*, Masson, Paris, 1989 ; English version : *Rational Curves and Surfaces. Applications to CAD*, J. Wiley and Sons, 1992.
- [7] J. C. FIOROT and P. JEANNIN, *B-Rational curves and surfaces, N-Rational splines*, in *Curves and Surfaces*, P. J. Laurent, A. Le Méhauté and L. L. Schumaker (eds.), Academic Press, New York (1991) pp. 181-184.
- [8] J. C. FIOROT, P. JEANNIN and S. TALEB, *Homographic change of parameter for B-Rational curves*, preprint, 1992.
- [9] J. C. FIOROT and S. TALEB, Reparametrizations of Polynomial and Rational curves, in *Curves and Surfaces*, P. J. Laurent, A. Le Méhauté and L. L. Schumaker (eds.), Academic Press, New York (1991) pp. 185-188.
- [10] S. TALEB, *Incidence du paramétrage sur le contrôle des courbes rationnelles*, Thèse Université des Sciences et Technologies de Lille, 1991.