

RECHERCHE COOPÉRATIVE SUR PROGRAMME N° 25

ELLIOTT H. LIEB

Convex Trace Functions and the Wigner-Yanase-Dyson Conjecture

Les rencontres physiciens-mathématiciens de Strasbourg - RCP25, 1973, tome 19
« Conférences de : P. Lelong, A. Lichnerowicz, E. Lieb et C. Stanojevic », , exp. n° 4,
p. 0-35

http://www.numdam.org/item?id=RCP25_1973__19__A4_0

© Université Louis Pasteur (Strasbourg), 1973, tous droits réservés.

L'accès aux archives de la série « Recherche Coopérative sur Programme n° 25 » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

I.- INTRODUCTION

This paper is concerned with certain convex or concave mappings of linear operators on a Hilbert space into the reals. [$f(A)$ is convex if $f(\lambda A + (1-\lambda)B) \leq \lambda f(A) + (1-\lambda) f(B)$ for $0 < \lambda < 1$ and $f(A)$ is concave if $-f(A)$ is convex]. These mappings involve the trace operation which plays a central role in quantum statistical mechanics, and it is not surprising, therefore, that the mappings discussed here were motivated by considerations of physics. In particular, Theorem 1 solves affirmatively a conjecture due to Wigner, Yanase and Dyson [1] about a certain definition of information. In section III we use Theorem 1 to prove other convexity theorems when the Hilbert space is finite dimensional. One of those, Theorem 6, we extend to infinite dimensional spaces in section IV. Theorem 6 has a physical application ; it is the basis for proving that quantum mechanical entropy is strongly subadditive (cf. refs [2], [3] and [4]) . The proof of that fact will be given in a subsequent paper [5] .

From the work of Krauss and Benda and Sherman ([6] and the references quoted therein) it is known that certain convex functions from \mathbb{R} to \mathbb{R} extend to operator-valued convex functions. If $f(x)$ is such a function then $A \mapsto \text{Tr} K f(A)$ (where Tr means trace) is certainly convex when $K > 0$ and fixed. Simple examples are $f(A) = A^{-p}$ and $f(A) = -A^p$ for $A > 0$ and $0 < p \leq 1$. However, $A \mapsto \text{Tr} f(A)$ may be convex even when $f(A)$ is not convex as an operator-valued function. Examples of this are $f(A) = e^A$ for A selfadjoint and $f(A) = A^{-p}$ for $1 < p \leq 2$ and for $A > 0$ (cf. Theorem 8).

In this paper we shall be concerned with mappings more complicated than those just mentioned. One example, Theorem 6, is $A \mapsto -\text{Tr} \exp[L + \ln A]$ for $A > 0$ and L selfadjoint.

Theorem 1 is our main theorem and Theorems 2,3,6 and 7 are derived from it. Theorems 8 and 9 are a side issue and are independent of and simpler than Theorem 1. In section V we remark briefly on the logical connection of Theorems 1,2,3,6 and 7, namely that they can all be derived simply from each other (at least for finite dimensional Hilbert spaces).

II.- THE MAIN THEOREM AND THE WIGNER-YANASE-DYSON PROBLEM.

We begin by proving our main Theorem 1 which constitutes the basis for Theorems 2,3, 6 and 7 of the next section. Theorem 1 is also the Wigner-Yanase-Dyson (WYD) conjecture [1] (actually, it is a bit stronger) and at the end of this section we shall explain the WYD problem. We also discuss another problem concerning the WYD definition of information [1] and give a partial solution of it.

Theorem 1 will be proved directly for infinite dimensional Hilbert spaces and our notation is the following :

(1) H is a separable Hilbert space with inner product (x,y) which is linear in y and conjugate linear in x .

(2) $\mathcal{B}(H)$ is the set of bounded linear operators from H to H ; $\mathcal{B}^s(H) \subset \mathcal{B}(H)$ are the bounded selfadjoint operators ; $\mathcal{B}^+(H) \subset \mathcal{B}^s(H)$ are the positive operators ($A \in \mathcal{B}^+(H) \Rightarrow (x,Ax) \geq 0, \forall x$) ; $\mathcal{B}^{++}(H) \subset \mathcal{B}^+(H)$ are the strictly positive operators ($A \in \mathcal{B}^{++}(H) \Rightarrow (x,Ax) > 0, \forall x$).

(3) If $A \in \mathcal{B}^+(H)$ and $z \in \mathbb{R}$, we can use the spectral representation of A to define A^z . $z \mapsto A^z$ is entire analytic and $A^z \in \mathcal{B}^+(H)$ for $z \geq 0$.

(4) The \mathcal{J}_q classes ; If $A \in \mathcal{B}(H)$ we form $|A| = (A^\dagger A)^{1/2} \in \mathcal{B}^+(H)$. $A \in \mathcal{J}_q(H) \subset \mathcal{B}(H)$ ($q \geq 1$) if $\|A\|_q \equiv (\text{Tr } |A|^q)^{1/q} < \infty$, where Tr means trace. $\mathcal{J}_1(H)$ is the trace class and $\mathcal{J}_2(H)$ is the Hilbert-Schmidt class. $A \in \mathcal{J}_q(H)$ implies that A is compact and that

$A \in \mathfrak{J}_1(H)$. $\|A\|_q = (\sum_{j=1}^{\infty} \lambda_j^q)^{1/q}$ where the λ_j are the eigenvalues of $|A|$ in decreasing order, including multiplicity. If $A \in \mathfrak{B}^+(H)$ but $A \notin \mathfrak{J}_1(H)$, it is convenient to define $\text{Tr } A = \infty$.

(5) We recall that if $A \in \mathfrak{B}(H)$ and if K is a linear operator (not necessarily bounded) on a dense domain, $D(K)$, in H then AK may have a bounded extension to all of H . If so, it is unique and its adjoint is a bounded extension of $K^{\dagger}A^{\dagger}$.

Theorem 1 : Let K be a linear operator (not necessarily bounded) on H , let $A, B \in \mathfrak{B}^+(H)$ and let $\lambda, 0 < \lambda < 1$, be given. Form the convex combination $C = \lambda A + (1-\lambda) B$. Let p and r be given positive real numbers with $p + r \equiv s \leq 1$. If $M \equiv C^{p/2} K C^{r/2}$ has an extension to $\mathfrak{J}_2(H)$ then

(1) $A^{p/2} K A^{r/2}$ and $B^{p/2} K B^{r/2}$ have extensions to $\mathfrak{J}_2(H)$ and

(2) $\lambda \text{Tr } A^{r/2} K^\dagger A^p K A^{r/2} + (1-\lambda) \text{Tr } B^{r/2} K^\dagger B^p K B^{r/2}$
 $\leq \text{Tr } C^{r/2} K^\dagger C^p K C^{r/2}$, i.e.

$A \in \mathfrak{B}^+(H) \mapsto \text{Tr } A^{r/2} K^\dagger A^p K A^{r/2}$ is concave.

Proof : (a) We recall the theorem [6] that the map $A \in \mathfrak{B}^+(H) \mapsto A^q$ is concave on $\mathfrak{B}^+(H)$ when $0 < q \leq 1$. Thus, $\lambda A^q \leq C^q$ and $\text{Ker } (A^q) = \text{Ker } (A) \supset \text{Ker } (C) = \text{Ker } (C^q)$, and similarly for B . As A, B and C are bounded, their kernels are closed subspaces and $H = \text{Ker } (C) \oplus \text{Ker } (C)^\perp$. The above inequalities show that for $0 < q \leq 1$, $\alpha(q) \equiv A^{q/2} C^{-q/2}$ and $\alpha(q)^\dagger = C^{-q/2} A^{q/2}$ can be extended to bounded operators on $\text{Ker}(C)^\perp$ because $\| A^{q/2} C^{-q/2} \Psi \| \leq \lambda^{-1/2}$ for Ψ in the dense set $D_C = \{\text{vectors with support away from zero in the spectral representation of } C\}$. Similarly, we define $\beta(q) \equiv B^{q/2} C^{-q/2}$. Also, $\alpha(q)$ and $\alpha(q)^\dagger$ can be defined to be zero on $\text{Ker}(C)$ and thus are defined on all of H . Clearly, $C^{q/2} \alpha(q)^\dagger = A^{q/2} = \alpha(q) C^{q/2}$. Consequently, $A^{p/2} K A^{r/2} = \alpha(p) [C^{p/2} K C^{r/2}] \alpha(r)^\dagger = \alpha(p) M \alpha(r)^\dagger \in \mathfrak{J}_2(H)$, since $M \in \mathfrak{J}_2(H)$. Not only is the first part of the theorem thus proved, but we also see that if $\{\psi_i\}$ and $\{\varphi_i\}$ are orthonormal bases for $\text{Ker}(C)$ and $\text{Ker}(C)^\perp$ respectively, we can compute traces in the basis $\{\psi_i\} + \{\varphi_i\}$ and all terms involving $\{\psi_i\}$ will vanish. Thus, $\text{Ker}(C)$

is an irrelevant subspace, and we shall henceforth assume that

$$H = \text{Ker}(C)^\perp, \text{ i.e. } C > 0 .$$

(b) With the above definitions, part(2) is equivalent to the following : $\lambda T^A(p,r) + (1-\lambda) T^B(p,r) \leq \text{Tr} M^\dagger M$ for every $M \in \mathcal{J}_2(H)$, where $T^A(p,r) = \text{Tr} \alpha(r) M^\dagger \alpha(p)^\dagger \alpha(p) M \alpha(r)^\dagger$ and similarly for $T^B(p,r)$.

(c) Let $z = x + iy \in \mathbb{C}$ and consider the operator valued function $\alpha(z) \equiv A^{z/2} C^{-z/2} = A^{iy/2} A^{x/2} C^{-x/2} C^{-iy/2} = A^{iy/2} \alpha(x) C^{-iy/2}$. Since $C^{iy/2}$ is unitary and $\|A^{iy/2}\| \leq 1$, $\alpha(z)$ is uniformly bounded in $S = \{z \mid 0 \leq \text{Re}(z) \leq 1\}$. If $\bar{z} = x - iy$, $\alpha(\bar{z})^\dagger = C^{-z/2} A^{z/2}$. For $\Psi \in D_C$, $\alpha(z) \Psi$ is an entire analytic function of z because $C^{-z/2} \Psi$ is entire and $A^{z/2}$ is entire. Hence, by the boundedness of $\alpha(z)$ and a standard density argument, $\alpha(z) \Psi$ is regular on S (continuous on S and analytic in the interior of S) for all $\Psi \in H$. Since weak analyticity implies strong analyticity, we also have that $\alpha(z)$ is strongly continuous on S and is norm analytic in the interior of S . Furthermore, if $A_n \rightarrow A$ strongly and if $B \in \mathcal{J}_2(H)$, then $A_n B \rightarrow AB$ in the $\mathcal{J}_2(H)$ norm. (This is trivial if B is finite rank, but the finite rank operators are dense in the $\mathcal{J}_2(H)$ norm). Hence, $\alpha(z_1) M \alpha(\bar{z}_2)^\dagger$ is $\mathcal{J}_2(H)$ regular on $S \times S$, which means that $T^A(z_1, z_2) \equiv \text{Tr} \alpha(z_2) M^\dagger \alpha(\bar{z}_1)^\dagger \alpha(z_1) M \alpha(\bar{z}_2)^\dagger$ is bounded and regular on $S \times S$.

(d) We now set $z_1 = z$, $z_2 = s - z$ and consider $T^A(z) \equiv T^A(z, s-z)$ as a regular function on $\{z \mid 0 \leq \text{Re}(z) \leq s\}$. By (b) we need to show that $f(p) = \lambda T^A(p) + (1-\lambda) T^B(p) \leq \text{Tr} M^\dagger M$. By

the maximum modulus principle for bounded regular functions on a strip, $|f(p)| \leq \max \left\{ \sup_{\theta} |f(i\theta)|, \sup_{\theta} |f(s+i\theta)| \right\}$. We shall consider only the first case, $p = i\theta$, in detail because the second case, $p = s+i\theta$, is parallel. $|f(i\theta)| \leq \lambda |T^A(i\theta)| + (1-\lambda) |T^B(i\theta)|$. Using the facts that for $A \in \mathfrak{B}(H)$ and $B \in \mathfrak{J}_2(H)$, AB and $BA \in \mathfrak{J}_2(H)$, and for $B, C \in \mathfrak{J}_2(H)$, $\text{Tr } BC = \text{Tr } CB$ and $|\text{Tr } BC| \leq \frac{1}{2} \text{Tr } B^\dagger B + \frac{1}{2} \text{Tr } C^\dagger C$, we have that $2 |T^A(i\theta)| \leq \text{Tr } \alpha(s-i\theta) M^\dagger \alpha(-i\theta)^\dagger \alpha(-i\theta) M \alpha(s-i\theta)^\dagger + \text{Tr } \alpha(s+i\theta) M^\dagger \alpha(i\theta)^\dagger \alpha(i\theta) M \alpha(s+i\theta)^\dagger$.

However, $\|\alpha(-i\theta)^\dagger \alpha(-i\theta)\| \leq 1$, so the first term is at most $\text{Tr } \alpha(s-i\theta) M^\dagger M \alpha(s-i\theta)^\dagger = \text{Tr } M \alpha(s-i\theta)^\dagger \alpha(s-i\theta) M^\dagger = \text{Tr } M C^{-i\theta/2} \alpha(s)^\dagger \alpha(s) C^{i\theta/2} M^\dagger = \text{Tr } \alpha(s)^\dagger \alpha(s) C^{i\theta/2} M^\dagger M C^{-i\theta/2}$. Likewise, the second term is at most $\text{Tr } \alpha(s)^\dagger \alpha(s) C^{-i\theta/2} M^\dagger M C^{i\theta/2}$. If we add to these the corresponding two terms for $|T^B(i\theta)|$ we obtain

$$\lambda |T^A(i\theta)| + (1-\lambda) |T^B(i\theta)| \leq \frac{1}{2} \text{Tr} [\lambda \alpha^\dagger(s) \alpha(s) + (1-\lambda) \beta^\dagger(s) \beta(s)] P, \quad (2.1)$$

where $P = C^{i\theta/2} M^\dagger M C^{-i\theta/2} + C^{-i\theta/2} M^\dagger M C^{i\theta/2} \in \mathfrak{B}^\dagger(H)$. As we remarked before, $\lambda A^s + (1-\lambda) B^s \leq C^s$, whence $\lambda \alpha(s)^\dagger \alpha(s) + (1-\lambda) \beta(s)^\dagger \beta(s) = C^{-s/2} [\lambda A^s + (1-\lambda) B^s] C^{-s/2} \leq \mathbf{1}$. Substituting this in (2.1) proves the theorem. Q.E.D.

REMARK : If $C^{P/2} K$ has an extension to $\mathfrak{J}_1(H)$ then so does $A^{P/2} K$ and $B^{P/2} K$ since $\|C^{P/2} K\| \geq \lambda^{1/2} \|A^{P/2} K\|$. In this case $\text{Tr } C^{r/2} K^\dagger C^P K C^{r/2} = \text{Tr } C^r K^\dagger C^P K = \text{Tr } K^\dagger C^P K C^r$ and similarly for A and B .

COROLLARY 1.1 : With p and r as in Theorem 1, the function from

$\mathfrak{B}^+(H) \times \mathfrak{B}^+(H) \times \mathfrak{B}(H)$ to the nonnegative reals defined by

$$(A,B,K) \mapsto F(A,B,K) = \text{Tr } A^{r/2} K^\dagger B^p K A^{r/2}$$

(1) is jointly concave in (A,B)

(2) is convex in K .

Proof : Consider the Hilbert space $H' \cong H \oplus H$ and define the following operators in $\mathfrak{B}(H')$:

$$k : (x,y) \mapsto (0, Kx)$$

$$k^\dagger : (x,y) \mapsto (K^\dagger y, 0)$$

$$a : (x,y) \mapsto (Ax, By) , a \in \mathfrak{B}^+(H') .$$

Applying Theorem 1 to $\text{Tr } a^{r/2} k^\dagger a^p k a^{r/2}$ proves the first part. The second part follows from a Schwartz inequality type of argument since $F(A,B,K)$ is nonnegative and quadratic in K . Q.E.D.

COROLLARY 1.2 : With p and r as in Theorem 1, $p + r \equiv s \leq 1$,

the functions from $\mathfrak{B}^+(H) \times \mathfrak{B}^+(H) \times \mathfrak{B}(H)$ to the nonnegative reals defined by

$$(A,B,K) \mapsto F_q(A,B,K) = \{\text{Tr } A^{r/2} K^\dagger B^p K A^{r/2}\}^q$$

(1) are jointly concave in (A,B) when $0 < q \leq 1/s$

(2) are jointly convex in (A,B) when $q < 0$

(3) are convex in K when $q \geq 1/2$.

Proof : The proof is a standard one for homogeneous concave (or convex functions [7]). Let $x = (x_1, x_2) \in \mathbb{R}_+^2 = \{(x_1, x_2) \mid x_1 \geq 0, x_2 \geq 0\}$ and define $f(x) = F_1(x_1 A + x_2 A', x_1 B + x_2 B', K)$ for an arbitrary, but henceforth fixed, choice of $A, A', B, B' \in \mathfrak{B}^+(H)$.

Parts (1) and (2) are equivalent to showing that for all such choices, $f(x)^q$ is concave (or convex). By Corollary 1.1 $f(x)$ is nonnegative, concave and homogeneous of order s , i.e. $f(\lambda x) = \lambda^s f(x)$ for $\lambda \geq 0$.

For each $\alpha \geq 0$, define $G_\alpha = \{x \mid f(x) \geq \alpha, x \in \mathbb{R}_+^2\}$. It is easily seen from the properties of $f(x)$ that G_α is a convex subset of \mathbb{R}_+^2 and $G_\alpha = \alpha^{1/s} G_1$ for $\alpha > 0$. Define $k(x) = \sup \{\mu \geq 0 \mid x \in G_{\mu^s}\}$ for $x \in \mathbb{R}_+^2$. As $x \in G_{f(x)}$, $k(x)$ is everywhere defined. In fact, since $f(x) = \sup \{\alpha \geq 0 \mid x \in G_\alpha\}$, $f(x) = k(x)^s$. Obviously, $k(x)$ is nonnegative and homogeneous of order one and, since $k(x) = \sup \{\mu > 0 \mid x \in \mu G_1\}$ when $f(x) \neq 0$, it is easy to check that $k(x)$ is a concave function. For a nonnegative concave function, $k(x)$, $k(x)^p$ is concave when $0 < p \leq 1$ and $k(x)^p$ is convex when $p < 0$. This proves parts (1) and (2). For part (3) we define $f(x) = F_1(A, B, x_1 K + x_2 K')$, with $K, K' \in \mathfrak{B}(H)$. $f(x)$ is nonnegative, convex and homogeneous of order 2. We define: $G_\alpha = \{x \mid f(x) \leq \alpha, x \in \mathbb{R}_+^2\}$ which is convex; $k(x) = \inf \{\mu > 0 \mid x \in G_{\mu^2}\}$. Then $k(x)$ is nonnegative, convex and homogeneous of order one and $f(x) = k(x)^2$. For any nonnegative convex function, $k(x)$, $k(x)^p$ is convex when $p \geq 1$. Q.E.D.

The setting for the next corollary is the following :

Let H^1 and H^2 be two separable Hilbert spaces and $H^{12} \equiv H^1 \otimes H^2$ their tensor product. If $A_{12} \in \mathfrak{B}^+(H^{12})$ and $A_{12} \in \mathfrak{A}_1(H^{12})$ we can define $A_1 \in \mathfrak{B}^+(H^1)$ by means of the partial trace, i.e.

$$A_1 = \text{Tr}^1 A_{12}$$

which means that for $x, y \in H^1$

$$(x, A_1 y) = \sum_i (x \otimes e_i, A_{12} [y \otimes e_i])$$

where $\{e_i\}$ is any orthonormal basis in H^2 .

COROLLARY 1.3 : Let H^2 be of dimension $d_2 < \infty$, let K be a linear operator (not necessarily bounded) on H^1 and let $A_{12} \in \mathfrak{B}^+(H^{12})$, $A_{12} \in \mathfrak{J}_1(H^{12})$ with $A_1 = \text{Tr}^1 A_{12}$. Let $p > 0, r > 0, p + r = s \leq 1$. Define $L = K \otimes \mathbb{1}_2$ on H^{12} . If $A_1^{p/2} K A_1^{r/2}$ has an extension to $\mathfrak{J}_2(H^1)$ then $A_{12}^{p/2} L A_{12}^{r/2}$ has an extension to $\mathfrak{J}_2(H^{12})$ and

$$d_2^{1-s} \text{Tr}^1 A_1^{r/2} K^\dagger A_1^p K A_1^{r/2} \geq \text{Tr}^{12} A_{12}^{r/2} L^\dagger A_{12}^p L A_{12}^{r/2}. \quad (2.2)$$

Proof : If G is the group of unitary transformations on H^2 and if

dU is the normalized Haar measure on G then it is easy to see [3]

that $B_{12} \equiv \int_G U^\dagger A_{12} U \, dU = d_2^{-1} A_1 \otimes \mathbb{1}_2$. Let $F(A_{12})$ be the right hand side of (2.2). By Theorem 1, $A_{12} \mapsto F(A_{12})$ is concave so

$F(B_{12}) \geq \int_G F(U^\dagger A_{12} U) \, dU$. But $F(U^\dagger A_{12} U)$ is independent of U since $(U^\dagger A_{12} U)^p = U^\dagger A_{12}^p U$, $U^\dagger L U = L$ and $\text{Tr}^{12} U^\dagger X U = \text{Tr}^{12} X$.

However, $F(B_{12})$ is the left hand side of (2.2). Q.E.D.

REMARKS : (i) Theorem 1 can be regarded as a special case of Corollary 1.3

as may be seen by taking $H^2 = \mathbb{C}^2$ and $A_{12} = \frac{1}{2} A \otimes P^a + \frac{1}{2} B \otimes P^b$,

where P^a and P^b are two orthogonal projections on H^2 and

$A, B \in \mathfrak{B}^+(H^1)$. Then $A_1 = \frac{1}{2} A + \frac{1}{2} B = C$.

(ii) Similarly to Corollary 1.1, we can extend Corollary 1.3 to the following : Let $A_{12}, B_{12} \in \mathfrak{B}^+(H^{12})$, $A_{12}, B_{12} \in \mathfrak{A}_1(H^{12})$, and $A_1 = \text{Tr}^1 A_{12}$, $B_1 = \text{Tr}^1 B_{12}$. Then

$$d_2^{1-s} \text{Tr}^1 A_1^{r/2} K^\dagger B_1^p K A_1^{r/2} \geq \text{Tr}^{12} A_{12}^{r/2} K^\dagger B_{12}^p K A_{12}^{r/2} . \quad (2.3)$$

(iii) When $d_2 = \infty$, Corollary 1.3 makes no sense except when $s = 1$. In that special case, the corollary is true when $d_2 = \infty$. The proof, which we shall not give here, can be constructed in imitation of the proof of Theorem 1 itself. The principal idea is to define

$$\alpha(q) = A_{12}^{q/2} [A_1^{-q/2} \otimes \mathbb{1}_2] \in \mathfrak{B}(H^{12}) \quad \text{and} \quad M = A_1^{p/2} K A_1^{r/2} \in \mathfrak{A}_2(H^1).$$

(iv) If we let $A_{12} = \mathbb{1}_{12}$ in (2.3) and let K be a projection onto an arbitrary one dimensional subspace of H^1 we obtain the operator inequality (since $A_1 = d_2 \mathbb{1}_1$)

$$d_2^{1-p} B_1^p \geq \text{Tr}^2 B_{12}^p$$

for all $B_{12} \in \mathfrak{B}^+(H^{12})$ and $0 < p \leq 1$.

THE WIGNER - YANASE - DYSON CONJECTURE.

In quantum mechanics, a density matrix, ρ , on H satisfies $\rho \in \mathfrak{B}^+(H)$, $\rho \in \mathfrak{A}_1(H)$ and $\text{Tr } \rho = 1$. The entropy of ρ , as usually defined, is

$$S(\rho) = -\text{Tr } \rho \ln \rho .$$

Wigner and Yanase [1] extended this to the concept of the entropy of ρ relative to a self adjoint "observable", K , and defined it as $S(\rho, K) \equiv \frac{1}{2} \text{Tr } [\rho^{1/2}, K]^2$, where $[A, B] = AB - BA$. Dyson (cf. ref. [1]) proposed a generalization of this to

$$S_p(\rho, K) \equiv \frac{1}{2} \text{Tr } [\rho^p, K] [\rho^{1-p}, K]$$

for $0 < p < 1$. [Actually, Wigner and Yanase defined $I_p(\rho, K) \equiv -S_p(\rho, K)$ which they termed skew information].

It is well known and easy to prove that $S(\rho)$ is concave in ρ , and the WYD conjecture is that $S_p(\rho, K)$ is concave in ρ for each fixed K . They were able to prove this only when $p = 1/2$. In physical applications K may be unbounded, but it is always correct to assume that $\rho^p K$ and $K \rho^p$ have unique extensions to $\mathfrak{A}_1(H)$ for all $p > 0$. Thus (cf. the remark after Theorem 1)

$$S_p(\rho, K) = -\text{Tr } \rho K^2 + \text{Tr } \rho^{1-p} K \rho^p K .$$

The first term is linear, and hence concave, and the second term is concave by Theorem 1. [Remark : Theorem 1 is stronger than necessary because it allows K to be non-selfadjoint, i.e. $\text{Tr}[\rho^p, K^\dagger][\rho^{1-p}, K]$ is concave in ρ when K is non-selfadjoint. This generalization can be derived from the selfadjoint case when $p = \frac{1}{2}$ by a simple polarization argument, but not when $p \neq 1/2$]. Baumann and Jost [9,10] proved the concavity for general p , but for a special class of ρ and H .

Wigner and Yanase properly regarded the concavity of $\rho \mapsto S_p(\rho, K)$ as a necessary requirement in order that $I_p(\rho, K)$ be a sensible definition of information. Another absolute requirement is the subadditivity of $S_p(\rho, K)$. Subadditivity of the ordinary entropy, $S(\rho)$, means (in the terminology preceding Corollary 1.3 and with $\rho_2 \equiv \text{Tr}^2 \rho_{12}$) that

$$S(\rho_{12}) \leq S(\rho_1) + S(\rho_2) .$$

This inequality is well known.

For $S_p(\rho, K)$, Wigner and Yanase [1] take the following definition : Let K_1 (resp. K_2) be a selfadjoint operator on H^1 (resp. H^2) and define $L = K_1 \otimes \mathbb{1}_2 + \mathbb{1}_1 \otimes K_2$ on H^{12} . The subadditivity condition is that

$$S_p(\rho_{12}, L) \leq S_p(\rho_1, K_1) + S_p(\rho_2, K_2) . \tag{2.4}$$

It is easy to see that (2.4) is true when $\rho_{12} = \rho_1 \otimes \rho_2$. Wigner and Yanase proved (2.4) when ρ_{12} is a projection onto a one dimensional subspace of H^{12} and $p = 1/2$. In the general case, (2.4) becomes (with $r = 1-p$)

$$\begin{aligned}
& \text{Tr}^1 \rho_1^p K_1 \rho_1^r K_1 + \text{Tr}^2 \rho_2^p K_2 \rho_2^r K_2 \\
& \geq - 2 \text{Tr}^{12} \rho_{12} [K_1 \otimes K_2] + \text{Tr}^{12} \rho_{12}^p L \rho_{12}^r L . \quad (2.5)
\end{aligned}$$

We do not have a proof of this, but when K_1 or K_2 is zero, (2.5) is simply Corollary 1.3. Because (2.5) is true in these three special cases, there is reason to believe it is true generally.

III.- SOME FINITE DIMENSIONAL THEOREMS.

In this section we confine our attention to finite dimensional Hilbert spaces over the complex numbers, i.e. $H = \mathbb{C}^n$. Some of the results of this section will be generalized to the infinite dimensional case in the next section by approximation arguments. If $C \in \mathfrak{B}^{++}(H)$ then $C > \epsilon \mathbb{1}$ for some $\epsilon > 0$. We remark that for $A, B \in \mathfrak{B}^{++}(H)$ and $K \in \mathfrak{B}(H)$, $\text{Tr } A K^\dagger B K = \text{Tr}(B^{1/2} K A^{1/2})^\dagger (B^{1/2} K A^{1/2}) > 0$ for $K \neq 0$.

Theorem 2 : The function from $\mathfrak{B}^{++}(H) \times \mathfrak{B}(H)$ to the nonnegative reals defined by

$$(A, K) \mapsto F(A, K) = \text{Tr } A^{-r} K^\dagger A^{-p} K \quad (3.1)$$

is jointly convex in (A, K) whenever $p \geq 0$, $r \geq 0$ and $p + r \leq 1$, i.e. for all λ , $0 < \lambda < 1$, $\lambda F(A, K) + (1-\lambda) F(B, L) \geq F(C, M)$ when $A, B \in \mathfrak{B}^{++}(H)$; $K, L \in \mathfrak{B}(H)$; $C = \lambda A + (1-\lambda)B$; $M = \lambda K + (1-\lambda) L$.

Proof : We can think of $\mathfrak{B}(H)$ as a complex Hilbert space, $V(H)$ of dimension n^2 with the inner product $K, K' \in \mathfrak{B}(H) \mapsto \langle K, K' \rangle = \text{Tr } K^\dagger K'$.

The linear transformation of $V(H) \rightarrow V(H)$ defined by $K \mapsto \alpha K \beta$, with $\alpha, \beta \in \mathfrak{B}^{++}(H)$ is Hermitian and positive definite because

$$\langle K, \alpha K \beta \rangle > 0 \quad \text{when } K \neq 0. \quad \text{Thus, } K \mapsto F(A, K), \quad K \mapsto F(B, K) \text{ and}$$

$K \mapsto F(C, K)$ are positive definite quadratic forms on $V(H)$. Furthermore,

we can define $V_2(H) = V(H) \oplus V(H)$ with inner product

$$\langle (K, L), (K', L') \rangle = \text{Tr } K^\dagger K' + \text{Tr } L^\dagger L'. \quad \text{Clearly } D \equiv \lambda F(A, K) + (1-\lambda)F(A, L)$$

and $N \equiv F(C, \lambda K + (1-\lambda)L)$ are both positive quadratic forms on $V_2(H)$ and

D is definite. With A and B fixed, we form the variational quotient N/D

and maximize it with respect to $(K,L) \in V_2(H)$. Using the fact that $\text{Tr } AB = 0$ for all $A \in \mathfrak{B}(H)$ implies $B = 0$, we find the eigenvalue equations

$$\begin{aligned} \gamma A^{-p} K A^{-r} &= C^{-p} [\lambda K + (1-\lambda)L] C^{-r} \equiv M \in \mathfrak{B}(H) \\ \gamma B^{-p} L B^{-r} &= C^{-p} [\lambda K + (1-\lambda)L] C^{-r} \equiv M \quad , \end{aligned} \quad (3.2)$$

and the problem is to show that the eigenvalue $\gamma \leq 1$. If $\gamma \neq 0$, the equation $\gamma A^{-p} K A^{-r} = M$ has the unique solution $K = \gamma^{-1} A^p M A^r$. Likewise, $L = \gamma^{-1} B^p M B^r$ and $\lambda K + (1-\lambda)L = C^p M C^r$. Thus, finding $\gamma \neq 0$ solutions to (3.2) is equivalent to finding an $M \in \mathfrak{B}(H)$ such that $M \neq 0$ and $\lambda A^p M A^r + (1-\lambda) B^p M B^r = \gamma C^p M C^r$. However, if we multiply this equation on the left by M^\dagger and take the trace, we see that $\gamma \leq 1$ by Theorem 1 . Q.E.D.

COROLLARY 2.1 : Let $p \geq 0, r \geq 0$, $p + r \equiv s \leq 1$ and $q \in \mathbb{R}$, $q \neq 0$ be fixed. Then the functions from $\mathfrak{B}^{++}(H) \times \mathfrak{B}^{++}(H) \times \mathfrak{B}(H)$ to the nonnegative reals defined by

$$(A,B,K) \rightarrow F_q(A,B,K) = (\text{Tr } A^{-p} K^\dagger B^{-r} K)^q$$

is jointly convex in (A,B,K) when $q \geq (2-s)^{-1}$.

Proof : The same as for Corollaries 1.1 and 1.2 , since $F_1(A,B,K)$ is homogeneous of order $2-s > 0$. Q.E.D.

We next turn our attention to a function similar to (3.1) from $\mathfrak{B}^{++}(H) \times \mathfrak{B}(H)$ to the nonnegative reals defined by

$$(A,K) \mapsto Q(A,K) = \text{Tr} \int_0^\infty dx (A + x\mathbb{1})^{-1} K^\dagger (A + x\mathbb{1})^{-1} K .$$

We note that the linear transformation T from $V(H)$ (defined in the proof of Theorem 2) to $V(H)$ given by

$$T_A : K \mapsto \int_0^\infty dx (A + x \mathbb{1})^{-1} K (A + x \mathbb{1})^{-1} \quad (3.3)$$

is Hermitean and positive definite . In a basis in which A is diagonal it is easy to compute T explicitly :

$$T_A : \{K_{ij}\} \mapsto \{K_{ij} f(A_i, A_j)\}$$

where the $\{A_i\}$ are the eigenvalues of A and

$$f(x,y) = (x-y)^{-1} \ln (x/y) \text{ if } x \neq y, f(x,x) = x^{-1} .$$

The inverse transformation is

$$T_A^{-1} : \{K_{ij}\} \mapsto \{K_{ij} / f(A_i, A_j)\}$$

but this is the same as

$$T_A^{-1} : K \mapsto \int_0^1 dx A^x K A^{1-x} \quad (3.4)$$

as one sees by calculating the integral.

Theorem 3 : The function $Q(A,K)$ from $\mathfrak{B}^{++}(H) \times \mathfrak{B}(E)$ to the nonnegative reals is jointly convex in (A,K) .

Proof : The proof is exactly the same as for Theorem 2 up to the eigenvalue equation (3.2) which now reads

$$\begin{aligned} \gamma T_A(K) &= T_C(\lambda K + (1-\lambda)L) \equiv M \in \mathfrak{B}(H) \\ \gamma T_B(K) &= T_C(\lambda K + (1-\lambda)L) = M \end{aligned} \quad (3.5)$$

If $\gamma \neq 0$ we apply the inversion formula (3.4), whence

$$\int_0^1 dx [\lambda A^x M A^{1-x} + (1-\lambda) B^x M B^{1-x} - \gamma C^x M C^{1-x}] = 0.$$

By multiplying this on the left by M^\dagger and using Theorem 1, we see that $\gamma \leq 1$. Q.E.D.

COROLLARY 3.1 : The functions

$$(A,B,K) \mapsto Q_q(A,B,K) = \left[\text{Tr} \int_0^\infty dx (A + x\Pi)^{-1} K^\dagger (B + x\Pi)^{-1} K \right]^q$$

from $\mathfrak{B}^{++}(H) \times \mathfrak{B}^{++}(H) \times \mathfrak{B}(H)$ to the nonnegative reals are

- (1) jointly convex in (A,B,K) when $q \geq 1$
- (2) convex in K when $q \geq 1/2$
- (3) jointly convex in (A,B) when $q > 0$
- (4) jointly concave in (A,B) when $-1 \leq q < 0$ and $K \neq 0$.

Proof : The construction given in Corollary 1.1 allows us to replace A by (A,B) . We note that $Q_1(A,B,K)$ is homogeneous of order -1 in (A,B) , as may be seen by changing the integration variable to λx , and hence $Q_1(A,B,K)$ is homogeneous of order one in (A,B,K) . The proof of parts (1) and (2) is the same as that given in the second half of Corollary 1.2 for nonnegative, homogeneous, convex functions of positive order. To prove parts (3) and (4) we have to modify the proof given in Corollary 1.2. For $x \in \mathbb{R}_+^2 \setminus \{(0,0)\}$, we construct $f(x) \equiv Q_1(x_1 A + x_2 A', x_1 B + x_2 B', K)$ with $A, A', B, B' \in \mathfrak{B}^{++}(H)$. Assuming that $K \neq 0$ (otherwise there is nothing to prove) we note that $f(x)$ is strictly positive (as remarked before), convex and homogeneous of order $-t = -1$. We shall here give the construction for general $t > 0$ as we shall need it in Corollary 8.1 and Theorem 9. Define the convex sets $G_\alpha = \{x | f(x) \leq \alpha, x \in \mathbb{R}_+^2 \setminus \{(0,0)\}\}$, whence $f(x) = \inf \{ \alpha > 0 | x \in G_\alpha \}$. Define $k(x) \equiv \sup \{ \mu > 0 | x \in G_{\mu^{-t}} \} = \sup \{ \mu > 0 | x \in \mu G_1 \} = f(x)^{-1/t}$. As before, $k(x)$ is positive, concave and homogeneous of order one. The rest follows from the remarks in Corollary 1.2. Q.E.D.

The mapping $T_A : \mathfrak{B}(H) \rightarrow \mathfrak{B}(H)$ defined in (3.3) has a special significance when restricted to the self adjoint operators, $\mathfrak{B}^s(H)$. Let $A \in \mathfrak{B}^{++}(H)$ and $K \in \mathfrak{B}^s(H)$. Then

$$\frac{d}{dx} \ln (A + xK) \Big|_{x=0} = T_A(K) \quad . \quad (3.6)$$

[To derive this, use the representation

$$\ln A = \int_0^\infty dx (1+x)^{-1} (A - x\mathbb{1}) (A + x\mathbb{1})^{-1} \quad (3.7)$$

which can be verified in a basis in which A is diagonal ; then use

$$\frac{d}{dx} (A + x K)^{-1} \Big|_{x=0} = A^{-1} K A^{-1} \quad (3.8)$$

for any $K \in \mathfrak{H}$, $A \in \mathfrak{H}^{++}(\mathfrak{H})$] . Using (3.6), (3.7) and (3.8) one has

$$\begin{aligned} \frac{d^2}{dx^2} \ln(A+xK) \Big|_{x=0} &= -2 \int_0^\infty dx (A+x\mathbb{1})^{-1} K (A+x\mathbb{1})^{-1} K (A+x\mathbb{1})^{-1} \\ &\equiv -R_A(K) \quad . \end{aligned} \quad (3.9)$$

We note that $R_A(K) \in \mathfrak{H}^{++}(\mathfrak{H})$ when $K \neq 0$.

Proposition 4 : For any real number, γ , and $A \in \mathfrak{H}^{++}(\mathfrak{H})$, $K \in \mathfrak{H}^s(\mathfrak{H})$,
 $\gamma^2 R_A(K) + 2\gamma T_A(K) + \mathbb{1} \geq 0$.

Proof : $0 \leq R_A(K + A) = - \frac{d^2}{dx^2} \ln (A + x K + x A) \Big|_{x=0}$

$$= - \frac{d^2}{dx^2} \{ \ln (1 + x) \mathbb{1} + \ln (A + x(1+x)^{-1} K) \} \Big|_{x=0}$$

$$= \mathbb{1} + 2 T_A(K) + R_A(K) \quad .$$

Now replace K by γK . Q.E.D.

Proposition 4 is not necessary for what follows, but we mention it because it almost, but not quite, implies the following proposition, P : $R_A(K) \geq T_A(K)^2$. It does imply that $(\Psi, [R_A(K) - T_A(K)^2] \Psi) \geq 0$ when Ψ is any eigenvector of $T_A(K)$. P is false, however, as may be seen by the two-dimensional example :

$A = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ with $a > b > 0$ and $K = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Theorem 6 below would be trivial to prove if P were true and, in some sense, it may be taken as a substitute for P .

In order to prove Theorem 6, we need some preliminary remarks and Lemma 5. The remarks are that

$$Q(A, K) = \text{Tr } K^{\dagger} T_A(K) \quad . \quad (3.10)$$

Also, $Q(A + x B, K)$ is differentiable near $x = 0$ for all $B \in \mathfrak{H}^S(H)$ and

$$\frac{d}{dx} Q(A + x B, K) \Big|_{x=0} = -\text{Tr } B R_A(K) \quad (3.11)$$

for $K \in \mathfrak{H}^S(H)$.

Lemma 5 : Let C be a convex cone in a vector space and let $F : C \rightarrow \mathbb{R}$ be a convex function which is also right differentiable in the sense that

$$\lim_{x \downarrow 0} x^{-1} \{F(A + xB) - F(A)\} \equiv G(A, B)$$

exists for all $A, B \in C$. Assume, also, that F is homogeneous of order 1, i.e. $F(\lambda A) = \lambda F(A)$ for $\lambda > 0$. Then $G(A, B) \leq F(B)$.

Conversely, if F is two-sided differentiable in the sense above (with equal left and right derivatives), if $G(A + xB, B)$ is measurable on $\{x | x \geq 0\}$, if $G(A, B) \leq F(B)$ and if F is homogeneous of order 1, then F is convex.

Proof : For all $x > 0$, $F(A + xB) = F((1+x)^{-1} [(1+x)A + x(1+x)^{-1}B])$
 $= (1+x) F((1+x)^{-1}A + x(1+x)^{-1}B) \leq (1+x) \{ (1+x)^{-1} F(A) + x(1+x)^{-1} F(B) \}$
 $= F(A) + xF(B)$. Subtract $F(A)$ from both sides, divide by x , and
take the limit $x \downarrow 0$. For the second part, let $C = \lambda A + (1-\lambda) B$ and
note that $F(C) - \lambda F(A) = F(C) - F(\lambda A)$
 $= \int_0^{1-\lambda} dx \frac{dF(\lambda A + xB)}{dx} = \int_0^{1-\lambda} dx G(A+xB, B) \leq \int_0^{1-\lambda} dx F(B)$
 $= (1-\lambda) F(B)$. Q.E.D.

We shall return again to this lemma, but for now we note that
since $Q(A, K)$ is homogeneous of order 1 on the convex cone $\mathfrak{B}^{++}(H) \times \mathfrak{B}^S(H)$
then, using (3.10) and (3.11) ,

$$-\text{Tr } B R_A(K) + 2 \text{Tr } M T_A(K) \leq \text{Tr } M T_B(M) \quad (3.12)$$

for $A, B \in \mathfrak{B}^{++}(H)$ and $K, M \in \mathfrak{B}^S(H)$.

Theorem 6 : Let $L \in \mathfrak{B}^S(H)$ be fixed . Then the function from $\mathfrak{B}^{++}(H)$
to the nonnegative reals defined by

$$A \mapsto F_L(A) = \text{Tr} \exp(L + \ln A)$$

is concave on $\mathfrak{B}^{++}(H)$ for all L .

Proof : Choose $A \in \mathfrak{B}^{++}(H)$ and $K \in \mathfrak{B}^S(H)$ and consider the func-
tion $f(x) = \text{Tr} \exp(L + \ln(A + xK))$ which is defined and differentiable
for the real variable x in some neighborhood of $\{0\}$. The theorem is
equivalent to the statement that $d^2 f / dx^2 \leq 0$ when $x = 0$ for all
choices of A, L and K . Using the fact (which can be proved by a power
series expansion, for example) that

$$\frac{d}{dx} e^{F+xG} \Big|_{x=0} = \int_0^1 dy e^{yF} G e^{(1-y)F} = T_{\exp(F)}^{-1}(G)$$

for $F, G \in \mathfrak{B}^S(H)$, we compute

$$\begin{aligned} \frac{d^2 f}{dx^2} \Big|_{x=0} &= - \operatorname{Tr} BR_A(K) + \operatorname{Tr} \int_0^1 dy T_A(K) B^y T_A(K) B^{(1-y)} \\ &= - \operatorname{Tr} BR_A(K) + \operatorname{Tr} T_A(K) T_B^{-1} [T_A(K)] , \end{aligned} \quad (3.13)$$

where $B = \exp(L + \ln A)$. Now use inequality (3.12) with $M = T_B^{-1} [T_A(K)]$. Q.E.D.

COROLLARY 6.1 : Let k be a positive integer and p_1, \dots, p_k positive real numbers with $p_1 + \dots + p_k \equiv s \leq 1$. Let $L \in \mathfrak{B}^s(H)$ and $q \in \mathbb{R}$ be fixed. Then the functions from $\mathfrak{B}^{++}(H)^k$ to the positive reals defined by

$$F_q(A_1, \dots, A_k) = \{\operatorname{Tr} \exp [L + \sum_{j=1}^k p_j \ln(A_j)]\}^q$$

(1) are jointly concave in (A_1, \dots, A_k) when $0 < q \leq s^{-1}$

(2) are jointly convex in (A_1, \dots, A_k) when $q < 0$.

Proof : F_1 is homogeneous of order s and we have already explained in Corollaries 1.2 and 2.1 how to treat the cases $q \neq 1$. That F_1 is jointly concave seems like a stronger result than Theorem 6 but, surprisingly, it is not. We have to show that for every choice of $A_1, \dots, A_k \in \mathfrak{B}^{++}(H)$ and $K_1, \dots, K_k \in \mathfrak{B}^s(H)$, the k -square second derivative matrix of $f(x_1, \dots, x_k) \equiv F(A_1 + x_1 K_1, \dots, A_k + x_k K_k)$ is positive semidefinite when $x_1 = \dots = x_k = 0$. If $\frac{\partial^2}{\partial x_i \partial x_j} f(x_1, \dots, x_k) \Big|_{x_1 = \dots = x_k = 0} \equiv m_{ij} \equiv g_{ij} + \delta_{ij} h_i$, we compute

$$g_{ij} = p_i p_j \operatorname{Tr} T_B^{-1} [T_{A_i}(K_i)] T_{A_j}(K_j) ,$$

$$h_i = -p_i \operatorname{Tr} B R_{A_i} (K_i) \leq 0 ,$$

where $B = \exp [L + \sum_{j=1}^k p_j \ln (A_j)]$. Clearly, for any $F, G \in \mathfrak{B}(H)$, $\operatorname{Tr} F T_B^{-1}(G) = \operatorname{Tr} G T_B^{-1}(F)$ and $\operatorname{Tr} F^\dagger T_B^{-1}(F) \geq 0$ so, by a Schwarz inequality argument, $|\operatorname{Tr} F^\dagger T_B^{-1}(G)|^2 \leq [\operatorname{Tr} F^\dagger T_B^{-1}(F)] [\operatorname{Tr} G^\dagger T_B^{-1}(G)]$. Hence $|g_{ij}|^2 \leq g_{ii} g_{jj}$, and $g_{ii} \leq -p_i h_i$ by Theorem 6 . Thus, for any $\psi \in \mathbb{C}^k$,

$$\sum_{i=1}^k \sum_{j=1}^k \bar{\psi}_i m_{ij} \psi_j \leq \left\{ \sum_{i=1}^k (-p_i h_i) \right\}^{1/2} \sum_{i=1}^k |\psi_i|^2 + \sum_{i=1}^k h_i |\psi_i|^2$$

$$\leq (1-s) \sum_{i=1}^k h_i |\psi_i|^2 \leq 0 \text{ since } h_i \leq 0, \text{ all } i. \text{ Q.E.D.}$$

COROLLARY 6.2 : Let p_1, \dots, p_k be as in Corollary 6.1 and let $\psi \in H$ with $\|\psi\| = 1$. Then

$$(A_1, \dots, A_k) \mapsto \exp \left[\sum_{j=1}^k p_j (\psi, \ln A_j \psi) \right]$$

is a concave function on $\mathfrak{B}^{++}(H)^k$.

Proof : Let P be the projection onto the one-dimensional subspace of H spanned by ψ . Then take the limit $e^L \rightarrow P$ in Corollary 6.1. Q.E.D.

Remark : If we write $B = e^L$ in Theorem 6, then, by the Trotter product formula, $F_L(A) = \lim_{n \rightarrow \infty} F_L^{(n)}(A)$ with $F_L^{(n)}(A) = \operatorname{Tr} (B^{1/n} A^{1/n})^n$. Now, $F_L^{(1)}(A)$ is concave in A since it is linear, $F_L^{(2)}(A)$ is concave in A by Theorem 1 and $F_L(A)$ is concave in A by Theorem 6 . Hence, we are led to the

Conjecture : For each fixed $B \in \mathfrak{B}^+(H)$ and $n \in \mathbb{Z}^+$, the positive function $\operatorname{Tr} (B^{1/n} A^{1/n})^n$ is concave in $A \in \mathfrak{B}^+(H)$.

We remind the reader of the Golden-Thompson inequality

(GT) [11,12] which is that

$$\text{Tr } e^{A+B} \leq \text{Tr } e^A e^B \quad (3.14)$$

for $A, B \in \mathfrak{B}^S(H)$. This theorem can be extended to the infinite dimensional case [13]. The obvious generalization of GT to three operators in the form $\text{Tr } e^{A+B+C} \leq \text{Tr } e^A e^B e^C$ is false. We shall now show that Theorem 6 does provide a correct generalization of GT to three operators (and hence provides an alternative proof of GT itself).

Theorem 7 : Let $A, B, C \in \mathfrak{B}^S(H)$.

$$\text{Tr } e^C T_{\exp(-A)}(e^B) \geq \text{Tr } e^{A+B+C}.$$

If A commutes with B then

$$\text{Tr } e^C e^A e^B \geq \text{Tr } e^{A+B+C}.$$

Proof : Define $\alpha, \beta \in \mathfrak{B}^{++}(H)$ by $\alpha = e^{-A}$ and $\beta = e^B$ and define $L \in \mathfrak{B}^S(H)$ by $L = A+C$. Since $\alpha \mapsto -\text{Tr } e^{L+\ln \alpha}$ is homogeneous of order 1 and convex on the cone $\mathfrak{B}^{++}(H)$, we can use Lemma 5 to deduce that for $\beta \in \mathfrak{B}^{++}(H)$

$$\begin{aligned} \text{Tr } e^{A+B+C} &= \text{Tr } e^{L+\ln \beta} \leq \frac{d}{dx} \text{Tr } e^{L+\ln(\alpha + x\beta)} \Big|_{x=0} \\ &= \text{Tr } e^C T_{\alpha}(\beta). \end{aligned}$$

For the last part, we note that

$$T_{\alpha}(\beta) = \int_0^{\infty} dx (e^{-A} + x \mathbf{1}) e^B (e^{-A} + x \mathbf{1})^{-1} = e^A e^B \text{ if } [A, B] = 0.$$

Q.E.D.

Remark : An alternative formulation of Theorem 7 is this : Let

$A, C, D \in \mathfrak{B}^S(H)$. Then

$$\text{Tr } e^C e^D \geq \text{Tr } \exp \left[C + A + \ln \int_0^1 ds e^{-As} e^D e^{-A(1-s)} \right]$$

which follows from the definition $T_{\exp(-A)}(e^B) \equiv e^D$ and the inversion formula (3.4).

Additional Theorems :

The Theorems proved thus far all rely on Theorem 1 in an essential way. We shall now prove some theorems which appear to be similar to Theorems 2,3,6 and 7 but which in reality are less complicated because they can be proved directly by elementary methods.

Theorem 8 : Let $K \in \mathfrak{B}(H)$ and $1 \geq p > 0, 1 \geq r > 0$ be fixed. Then the function from $\mathfrak{B}^{++}(H)$ to the nonnegative reals defined by

$$A \mapsto F(A) = \text{Tr } A^{-p} K^\dagger A^{-r} K$$

is convex in A .

Proof : It is sufficient to show that for all $L \in \mathfrak{B}^S(H)$, $g(x) \equiv F(A+xL)$ has a positive second derivative at $x = 0$. Let

$$A(x) = A + xL, B_p = \frac{d}{dx} A(x)^{-p} \Big|_{x=0}, C_p = \frac{d}{dx} A(x)^p \Big|_{x=0}, D_p = \frac{d^2}{dx^2} A(x)^{-p} \Big|_{x=0}$$

and $E_p = \frac{d^2}{dx^2} A(x)^p \Big|_{x=0}$. Using the fact that $A(x)^p A(x)^{-p} = \mathbb{1}$, we compute that $B_p = -A^{-p} C_p A^{-p}$ and $D_p = 2B_p A^p B_p - A^{-p} E_p A^{-p}$. Hence

$$\frac{d^2}{dx^2} g(x) \Big|_{x=0} = -\text{Tr } E_p A^{-p} K^\dagger A^{-r} K A^{-p} - \text{Tr } E_r A^{-r} K A^{-p} K^\dagger A^{-r}$$

$$+ \text{Tr}(\gamma \delta_p + \delta_r \gamma)(\gamma \delta_p + \delta_r \gamma)^\dagger + \text{Tr}(\gamma \delta_p)(\gamma \delta_p)^\dagger + \text{Tr}(\delta_r \gamma)(\delta_r \gamma)^\dagger ,$$

where $\gamma = A^{-r/2} K A^{-p/2}$ and $\delta_p = A^{-p/2} C_p A^{-p/2}$. The theorem will be proved if we show that $E_p \leq 0$, but this fact follows from the integral representation (valid for $1 > p > 0$)

$$A^p = \pi^{-1} \sin \pi p \int_0^\infty x^{p-1} dx A (A + x \mathbb{1})^{-1} \quad \text{and (3.8) . Q.E.D.}$$

COROLLARY 8.1 : Let $1 \geq p > 0$ and $1 \geq r > 0$, $p + r \equiv s$ and $q \in \mathbb{R}$, $q \neq 0$ be fixed. Then the functions from $\mathfrak{B}^{++}(H) \times \mathfrak{B}^{++}(H) \times \mathfrak{B}(H)$ to the nonnegative reals, defined by

$$(A, B, K) \mapsto F_q(A, B, K) = (\text{Tr } A^{-p} K^\dagger B^{-r} K)^q$$

- (1) are convex in K when $q \geq 1/2$
- (2) are jointly convex in (A, B) when $q > 0$
- (3) are jointly concave in (A, B) when $-1/s \leq q < 0$.

Proof : The same as for Corollary 3.1. We note that the degree of homogeneity in K is 2 while in (A, B) it is $-s$. Q.E.D.

Remark : The map $A \mapsto A^{-p}$ is not convex for $p > 1$, but $A \mapsto \text{Tr} A^{-p}$ is convex for $2 \geq p > 0$.

Theorem 9 : Let k be a positive integer and p_1, \dots, p_k positive real numbers with $p_1 + \dots + p_k \equiv s$. Let $L \in \mathfrak{B}^s(H)$ and $q \in \mathbb{R}$ be fixed. Then the functions from $\mathfrak{B}^{++}(H)^k$ to the positive reals defined by

$$F_q(A_1, \dots, A_k) = \{\text{Tr } \exp[L - \sum_{j=1}^k p_j \ln(A_j)]\}^q$$

(1) are jointly convex in (A_1, \dots, A_k) when $q \geq 0$

(2) are jointly concave in (A_1, \dots, A_k) when $-1/s \leq q \leq 0$.

Proof : We need only consider the case $k = q = 1$. The extension to the general case is the same as in Corollary 3.1 . As in Theorem 8 we define $q(x) = F(A + x K)$, $K \in \mathfrak{B}^s(H)$, and compute its second derivative at $x = 0$ to be $-p \text{Tr } B R_A(K) + p^2 \text{Tr } T_A(K) T_B^{-1}[T_A(K)]$ where $B = \exp(L - p \ln A) > 0$. $R_A(K) \geq 0$ and, as remarked in the proof of Corollary 6.1, $\text{Tr } F^\dagger T_B^{-1}(F) \geq 0$ for all $F \in \mathfrak{B}(H)$. Q.E.D.

IV. - EXTENSION OF THEOREM 6 TO INFINITE DIMENSIONS.

We fix L , which is assumed to be self adjoint, and $e^L \in \mathcal{D}_1(H)$, which implies that L has purely discrete spectrum. For $A, B \in \mathcal{B}^+(H)$ and $C \equiv \lambda A + (1-\lambda) B$, $0 < \lambda < 1$, we want to show that

$$\lambda \operatorname{Tr} e^{L + \ln A} + (1-\lambda) \operatorname{Tr} e^{L + \ln B} \leq \operatorname{Tr} e^{L + \ln C} \quad (4.1)$$

which requires, among other things, giving meaning to these quantities.

Case 1. : We first assume that there exist real numbers ϵ and ω such that $\epsilon \mathbb{1} < A < \omega \mathbb{1}$, $\epsilon \mathbb{1} < B < \omega \mathbb{1}$ so that $\ln A$, etc... can be defined as bounded, self adjoint operators by means of the spectral representation of A , etc... We define $\alpha = L + \ln A$. Since $\ln A$ is bounded, α is a self adjoint operator on the domain of L . If we label the eigenvalues of L by $\mu_1(L) \geq \mu_2(L) \geq \dots$ we have, by the mini-max principle, that $\mu_k(\alpha) \leq \mu_k(L) + \ln \|A\|$ since $\ln A \leq \ln \|A\| \mathbb{1}$. The convergence of $\sum_1^\infty \exp(\mu_k(L))$ implies that $\mu_k(L) \rightarrow -\infty$, which implies that $\mu_k(\alpha) \rightarrow -\infty$, which implies that $\exp(\alpha)$ is compact, and since $t(A) \equiv \operatorname{Tr} e^\alpha \leq \|A\| \operatorname{Tr} e^L < \infty$, the trace is finite. Now let P_n be the projection onto R_n , the subspace spanned by the first n eigenvectors of L , and define

$$A_n \equiv P_n A P_n + \epsilon (\mathbb{1} - P_n)$$

$$\alpha_n \equiv L + \ln A_n$$

and likewise for B_n and C_n . Clearly,

$$t(A_n) = \text{Tr} \exp(\alpha_n) = \text{Tr} P_n \exp(L + \ln(A \uparrow R_n)) + r_n, \text{ where}$$

$$r_n = (\ln \epsilon) \text{Tr}(\mathbb{1} - P_n) e^L. \text{ From this we see that not only is } t(A_n)$$

finite, but that the terms involving r_n cancel from both sides of

(4.1) leaving an inequality about traces on the finite dimensional space

R_n . Thus by Theorem 6, $\lambda t(A_n) + (1-\lambda) t(B_n) \leq t(C_n)$. All we have

to do is prove that $t(A_n) \rightarrow t(A)$, etc... Now $A_n < \omega \mathbb{1}$ and

$$\alpha_n < L + \mathbb{1} \ln \omega. \text{ Since } \mu_k(\alpha_n) \leq \mu_k(L) + \ln \omega, \quad t(A_n) \rightarrow t(A)$$

by the dominated convergence theorem if we can show that $\mu_k(\alpha_n) \rightarrow \mu_k(\alpha)$

for each k . As L is bounded above, we can find a constant, d , such

that $\alpha_n < (d-1)\mathbb{1}$ and $\alpha < (d-1)\mathbb{1}$. Define $G_n = (\alpha_n - d\mathbb{1})^{-1} - (\alpha - d\mathbb{1})^{-1}$

$$= (\alpha_n - d\mathbb{1})^{-1} (\alpha - \alpha_n) (\alpha - d\mathbb{1})^{-1} \text{ and note that } (\alpha_n - d\mathbb{1})^{-1} \text{ is uniformly bounded,}$$

$\alpha_n - \alpha \rightarrow 0$ strongly, and $(\alpha - d\mathbb{1})^{-1}$ is compact since e^L is compact.

Hence, $\|G_n\| \rightarrow 0$. In general, $\|\mu_k(A) - \mu_k(B)\| \leq \|A-B\|$ by the mini-max

principle. Thus, $\mu_k(\alpha_n) \rightarrow \mu_k(\alpha)$ since $\mu_k((\alpha - d\mathbb{1})^{-1}) = [\mu_k(\alpha) - d]^{-1}$,

and the theorem is proved for case 1.

Case 2.: $0 < A < \omega \mathbb{1}$, $0 < B < \omega \mathbb{1}$. If we replace A by

$A_\epsilon = A + \epsilon \mathbb{1}$ and B by $B_\epsilon = B + \epsilon \mathbb{1}$, then C is replaced by

$$\lambda(A + \epsilon \mathbb{1}) + (1-\lambda)(B + \epsilon \mathbb{1}) = C + \epsilon \mathbb{1}. \text{ Also, } A_\epsilon, \text{ etc...}$$

are decreasing in ϵ . Thus, we can define

$$\text{Tr} \exp(L + \ln A) \equiv \lim_{\epsilon \downarrow 0} \text{Tr} \exp(L + \ln A_\epsilon) \tag{4.2}$$

because the right hand trace decreases as ϵ decreases. Theorem 6 is true

with this definition because it is true for every $\epsilon > 0$. The usefulness

of this definition stems from the following fact (B. Simon, private com-

munication): $\ln A$ is defined as an unbounded selfadjoint operator in

a natural way by its spectral decomposition.

If $D(\ln A) \cap D(L)$ is dense in H (where $D(\cdot)$ means domain), and if we define $-Q_\epsilon$ to be the Friedrichs extension of $-L - \ln A_\epsilon$, then

$$\text{Tr } e^{Q_\epsilon} = \lim_{\epsilon \downarrow 0} \text{exp } (L + \ln A_\epsilon) .$$

Finally, another case to which Theorem 6 can be extended (B. Simon, private communication) is where A and B are positive, not necessarily bounded, selfadjoint operators and $\ln A$ and $\ln B$ are form bounded perturbations of $-L$.

V. - ON THE "EQUIVALENCE" of THEOREMS 1, 2, 3, 6 and 7

It is a truism of logic that two true theorems, however disparate are always equivalent ; therefore the remarks in this section are formally heuristic. Although Theorem 1 is our main Theorem and Theorems 2, 3, 6 and 7 appear to be corollaries of it, the fact is that if one could find an independent proof of any of Theorems 2, 3, 6 or 7 the others could be derived from it in a simple way when the Hilbert space is finite dimensional. Thus, in some sense, all five theorems have equal content. We shall indicate very briefly some of the links among these theorems and the reader can easily supply the missing details as well as establish additional connections.

We have previously established the implications

$$1 \Rightarrow 2 \quad \text{and} \quad 1 \Rightarrow 3 \Rightarrow 6 \Rightarrow 7 \quad .$$

(a.) $2 \Rightarrow 1$. This is easily seen simply by reading the proof of Theorem 2 backwards, i.e. one minimizes the variational quotient N/D with respect to K , where $N = \lambda \text{Tr } A^r K^\dagger A^p K + (1-\lambda) \text{Tr } B^r K^\dagger B^p K$ and $D = \text{Tr } C^r K^\dagger C^p K$.

(b.) $7 \Rightarrow 6$. Theorem 7 was derived from Theorem 6 by using the derivative inequality, Lemma 5 . However, we can retrace our steps by using the second part of Lemma 5 .

(c.) $6 \Rightarrow (3.12)$. If, in (3.12), we fix A, B, K and minimize $\text{Tr } M [T_B(M) - 2 T_A(K)]$ with respect to M , we find $M = T_B^{-1}(T_A(K))$. Hence (3.12) is equivalent to the fact that the right hand side of (3.13) is negative and this, in turn, is equivalent to Theorem 6 .

(d.) (3.12) \Rightarrow 3 . The same remark as in (b.) above suffices.

(e.) 3 \Rightarrow 2 . Take 3 in the form of Corollary 3.1 with $q = 1$ and assume that $p + r \equiv s < 1$. For A (resp. B) substitute $\lambda A + \mathbb{1}$ (resp. $\gamma A + \mathbb{1}$) , where $\lambda, \gamma \geq 0$. Then integrate $\int_0^\infty d\lambda \lambda^{p-1} \int_0^\infty d\gamma \gamma^{r-1} Q_1(\lambda A + \mathbb{1}, \gamma A + \mathbb{1}, K)$ to obtain a positive constant times $F(A, K)$ of (3.1). When $p + r = 1$ one can appeal to continuity.

ACKNOWLEDGEMENT : The author is most grateful to B. Simon for explaining to him how to extend Theorems 1 and 6 to infinite dimensions. He is also indebted to D. Ruelle for stimulating his interest in this problem and to R. Jost, O. Lanford III and D.W. Robinson for useful conversations. The hospitality of the Institut des Hautes Etudes Scientifiques and the Chemistry Laboratory III, University of Copenhagen where parts of this work were carried out, is gratefully acknowledged.

REFERENCES

- [1.] E.P. Wigner and M.M. Yanase, Information Content of Distributions, Proc. Nat. Acad. Sci. U.S.A. 49 (1963), 910-918 . See also E.P. Wigner and M.M. Yanase, On the Positive Semidefinite Nature of Certain Matrix Expressions, Canad. J. Math. 16 (1964), 397-406.
- [2.] D.W. Robinson and D. Ruelle, Mean Entropy of States in Classical Statistical Mechanics, Commun. Math. Phys. 5 (1967), 288-300.
- [3.] O.E. Lanford III and D.W. Robinson, Mean Entropy of States in Quantum Statistical Mechanics, J.Math. Phys. 9 (1968), 1120-1125.
- [4.] H. Araki and E.H. Lieb, Entropy Inequalities, Commun. Math. Phys. 18 (1970), 160-170.
- [5.] E.H. Lieb and M.B. Ruskai, Proof of the Strong Subadditivity of Quantum Mechanical Entropy, to be submitted to J. Math. Phys. See also ~~A Fundamental Property~~ of Quantum Mechanical Entropy, Phys. Rev. Lett. (1973) to appear.
- [6.] J. Bendat and S. Sherman, Monotone and Convex Operator Functions, Trans. Amer. Math. Soc. 79 (1955), 58-71. F. Krauss, Math. Zeit. 41 (1936), 18 .
- [7.] R.T. Rockafellar, "Convex Analysis", Princeton University, Princeton, N.J. , 1970. cf. Theorem 15.3.

- [8.] The idea of representing A_1 in this manner is due to A. Uhlmann, Endlich Dimensionale Dichtematrizen II, preprint, submitted to Wissen. Z. Karl-Marx-Univ., Leipzig (1972).
- [9.] F. Baumann, and R. Jost, Remarks on a Conjecture of Robinson and Ruelle Concerning the Quantum Mechanical Entropy, in "Problems of Theoretical Physics ; Essays Devoted to N.N. Bogoliubov", (1969), 285-293, Moscow, Nauka.
- [10.] F. Baumann, Bemerkungen Ueber Quantenmechanische Entropie Ungleichungen, Helv. Phys. Acta 44 (1971), 95-100.
- [11.] S. Golden, Lower Bounds for the Helmholtz Function , Phys. Rev. 137 (1965) B 1127-1128.
- [12.] C.J. Thompson, Inequality with Applications in Statistical Mechanics, J. Math. Phys. 6 (1965), 1812-1813.
- [13.] M.B. Ruskai, Inequalities for Traces on Von Neumann Algebras, Commun. Math. Phys. 26 (1972), 280-289.