

STATISTIQUE ET ANALYSE DES DONNÉES

R. MOEANNADIN

H. TONG

Is a bilinear model an illusion ?

Statistique et analyse des données, tome 15, n° 1 (1990), p. 57-60

http://www.numdam.org/item?id=SAD_1990__15_1_57_0

© Association pour la statistique et ses utilisations, 1990, tous droits réservés.

L'accès aux archives de la revue « Statistique et analyse des données » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

IS A BILINEAR MODEL AN ILLUSION ?

R. MOEANNADIN et H. TONG

Department of Mathematics
University of Kent
Canterbury
CT2 7NF
GRANDE-BRETAGNE

Abstract

We have shown that paradoxically the better is the fit as measured by the residual sum of squares the less plausible is a bilinear model. Similar remarks apply to non-linear moving average and ARCH models.

Keywords : ARCH models, bilinear models, conditional least squares, method of moments, non-linear moving average models.

Classification AMS : 62 M 10, 93 E 99

Résumé

Nous avons montré que paradoxalement meilleur est l'ajustement mesuré par la somme des carrés des résidus moins un modèle bilinéaire est plausible. Des remarques similaires sont considérées pour des modèles ARCH et moyenne mobile non-linéaire.

Mots-clefs : modèles ARCH, modèles bilinéaires, moindres carrés conditionnels, méthode des moments, modèles moyenne mobile non-linéaires.

Let X_1, X_2, \dots, X_n denote observations from the simple diagonal bilinear model

$$X_t = \beta_0 X_{t-1} \varepsilon_{t-1} + \varepsilon_t, \quad (1)$$

where $\{\varepsilon_t\}$ is a sequence of independent and identically distributed random variables with zero mean and variance σ^2 . First it is easy to obtain a method-of-moments estimate, $\bar{\beta}$, of β_0 namely

$$\bar{\beta} = \bar{X}_n / \sigma^2, \quad (2)$$

if σ^2 is known. Here \bar{X}_n denotes the sample mean. The variance of $\bar{\beta}$ is

$$\frac{1}{n \sigma^2} + \frac{3 \beta_0^2}{n (1 - \beta_0^2 \sigma^2)} + \left[\frac{n-2}{n^2} \right] \beta_0^2, \quad (3)$$

which tends to infinity as σ^2 tends to zero !

Next, it is easy to obtain the least squares estimate, $\hat{\beta}$, of β_0 , conditional on X_C and ε_0 , by minimizing w.r.t. β

$$Q(\beta) = \sum_{t=1}^n \left[X_t - \beta X_{t-1} \varepsilon_{t-1}(\beta) \right]^2, \quad (4)$$

where n denotes the sample size. For the asymptotics to hold, we assume that the model (1) is invertible by imposing, for example, the condition $\beta^2 \sigma^2 < \frac{1}{2}$.

To evaluate the asymptotic variance of $\hat{\beta}$, we need the existence of the second moment of $\partial \varepsilon_t(\beta) / \partial \beta$, which unfortunately cannot be guaranteed. We thus resort to simulation. Results are given in Table 1.

Table 1

($\beta_0 = 0,5$, sample size = 1000, 100 replications)

σ^2	Interquartile Range	
	$\bar{\beta}$	$\hat{\beta}$
1	(0.466 , 0.530)	(0.496 , 0.502)
0.1	(0.286 , 0.702)	(0.374 , 0.629)
0.01	(-1.688 , 2.793)	(-0.780 , 1.674)
0.001	(-23.193 , 21.084)	(-13.250 , 12.696)

Now, equation (3) and Table 1 warn us against our natural instinct to strive for the smallest possible residual sum of squares in bilinear modelling, other things considered.

Our remark applies to the closely related ARCH models and non-linear moving average models. To save space, consider only the non-linear moving average model

$$X_t = \alpha \varepsilon_{t-1}^2 + \varepsilon_t \quad (4)$$

where $\{\varepsilon_t\}$ is as defined previously. Robinson (1977) has suggested the method of moments since the maximum likelihood estimate of α is usually not practical for non-linear moving average models. Let α denote the method-of-moments estimate of α , namely

$$\alpha = \bar{X}_n / \sigma^2, \quad (5)$$

where \bar{X}_n denotes the sample mean and σ^2 is assumed known. It is then easily verified that the variance of α is $1/(n \sigma^2) + (4 \alpha^2 / n)$, which may be compared with (3). Simulation produced a similar table to Table 1 in respect of conditional least-squares estimate of α .

Paradoxically, the better is the fit as measured by the noise variance the less plausible is a bilinear model and its like. This observation leads us to the question raised at the title of this note. The observation is quite general. For example, our result for the method-of-moments estimate generalises to the pure diagonal bilinear model

$$X_t = \sum_{j=1}^k \beta_j X_{t-j} \varepsilon_{t-j} + \varepsilon_t, \quad (6)$$

where $\{\varepsilon_t\}$ is as defined previously. Simulation studies produced similar conclusion for the conditional least-squares estimates. For non-diagonal bilinear models, the situation is more complex. However, a simulation study reported in the unpublished doctoral thesis of one of us (R.M.) suggests that a similar paradox exists.

ACKNOWLEDGEMENTS

Our investigation was first prompted by an incisive but unrecorded remark made by Professor M.S. Bartlett, FRS, at the 1978 Institute of Statisticians Annual Conference on Time Series Analysis at Cambridge. We are grateful to Professor A.J. Lawrance and Dr. I.T. Jolliffe for encouraging us to write up our results and to Professor K.S. Chan for pointing out an error in our first draft.

REFERENCE

Robinson, P.M. (1977), The estimation of a nonlinear moving average model. *Stoch. Proc. & Appl.*, **5**, 81-90.