



Centre de  
Mathématiques  
Laurent Schwartz



ÉCOLE  
POLYTECHNIQUE

SEMINAIRE

**Equations aux  
Dérivées  
Partielles**

**2005-2006**

Armen Shirikyan

**Controllability of three-dimensional Navier–Stokes equations and applications**

*Séminaire É. D. P.* (2005-2006), Exposé n° VI, 7 p.

<[http://sedp.cedram.org/item?id=SEDP\\_2005-2006\\_\\_\\_\\_A6\\_0](http://sedp.cedram.org/item?id=SEDP_2005-2006____A6_0)>

U.M.R. 7640 du C.N.R.S.  
F-91128 PALAISEAU CEDEX

Fax : 33 (0)1 69 33 49 49

Tél : 33 (0)1 69 33 49 99

cedram

*Article mis en ligne dans le cadre du  
Centre de diffusion des revues académiques de mathématiques*  
<http://www.cedram.org/>

# Controllability of three-dimensional Navier–Stokes equations and applications

Armen Shirikyan

Laboratoire de Mathématiques  
Université de Paris-Sud XI, Bâtiment 425  
91405 Orsay Cedex, France  
E-mail: Armen.Shirikyan@math.u-psud.fr

## Abstract

We formulate two results on controllability properties of the 3D Navier–Stokes (NS) system. They concern the approximate controllability and exact controllability in finite-dimensional projections of the problem in question. As a consequence, we obtain the existence of a strong solution of the Cauchy problem for the 3D NS system with an arbitrary initial function and a large class of right-hand sides. We also discuss some qualitative properties of admissible weak solutions for randomly forced NS equations.

**AMS subject classifications:** 35Q30, 60H15, 76D05, 93B05, 93C20

**Keywords:** Approximate controllability, exact controllability in projections, 3D Navier–Stokes system, Agrachev–Sarychev method, stationary solutions, irreducibility.

## 1 Main results

Let  $D \subset \mathbb{R}^3$  be a bounded domain with  $C^2$ -smooth boundary  $\partial D$ . Consider 3D Navier–Stokes (NS) equations

$$\dot{u} + (u, \nabla)u - \nu \Delta u + \nabla p = f(t, x), \quad \operatorname{div} u = 0, \quad x \in D, \quad (1)$$

where  $u = (u_1, u_2, u_3)$  and  $p$  are unknown velocity and pressure fields,  $\nu > 0$  is the viscosity, and  $f(t, x)$  is an external force. We introduce the spaces

$$H = \{u \in L^2(D, \mathbb{R}^3) : \operatorname{div} u = 0 \text{ in } D, \langle u, \mathbf{n} \rangle|_{\partial D} = 0\}, \\ V = H_0^1(D, \mathbb{R}^3) \cap H, \quad U = H^2(D, \mathbb{R}^3) \cap V,$$

where  $\mathbf{n}$  stands for the outward unit normal to  $\partial D$  and  $\langle \cdot, \cdot \rangle$  denotes the scalar product in  $\mathbb{R}^3$ . It is well known (e.g., see [Tem79]) that  $H$  is a closed vector

space in  $L^2(D, \mathbb{R}^3)$ , and we denote by  $\Pi$  the orthogonal projection in  $L^2(D, \mathbb{R}^3)$  onto  $H$ . Equations (1) are equivalent to the following evolution equation in  $H$ :

$$\dot{u} + \nu Lu + B(u) = f. \quad (2)$$

Here  $L = -\Pi\Delta$ ,  $B(u) = B(u, u)$ ,  $B(u, v) = \Pi\{(u, \nabla)v\}$ , and we use the same notation for the right-hand side of (1) and its projection to  $H$ . Equation (2) is supplemented with the initial condition

$$u(0) = u_0, \quad (3)$$

where  $u_0 \in V$ . Let us assume that the right-hand side of (2) is represented in the form

$$f(t, x) = h(t, x) + \eta(t, x), \quad (4)$$

where  $h \in L^2_{\text{loc}}(\mathbb{R}_+, H)$  is a given function and  $\eta$  is a control taking on values in a finite-dimensional subspace. To formulate the main results, we introduce some notation.

Define the space  $\mathcal{X}_T = C(J_T, V) \cap L^2(J_T, U)$ , where  $J_T = [0, T]$ . For any  $T > 0$ ,  $h \in L^2(J_T, H)$ , and  $u_0 \in V$ , we denote by  $\Theta_T(h, u_0)$  the set of functions  $\eta \in L^2(J_T, H)$  for which problem (2) – (4) has a unique solution  $u \in \mathcal{X}_T$ . It follows from the implicit function theorem that

$$\mathcal{D}_T := \{(u_0, \eta) \in V \times L^2(J_T, H) : \eta \in \Theta_T(h, u_0)\} \quad (5)$$

is an open subset of  $V \times L^2(J_T, H)$ , and the operator  $\mathcal{R}$  taking  $(u_0, \eta) \in \mathcal{D}_T$  to the solution  $u \in \mathcal{X}_T$  of (2) – (4) is locally Lipschitz continuous. We denote by  $\mathcal{R}_t$  the restriction of  $\mathcal{R}$  to the time  $t \in J_T$ . Let  $E \subset U$  and  $F \subset H$  be finite-dimensional subspaces, let  $\mathbf{P}_F : H \rightarrow H$  be the orthogonal projection onto  $F$ , and let  $X \subset L^2(J_T, E)$  be a vector space, not necessarily closed. We denote by  $B_F(R)$  the closed ball in  $F$  of radius  $R$  centred at origin.

**Definition 1.** Equations (2), (4) with  $\eta \in X$  are said to be *approximately controllable in time  $T$*  if for any  $u_0, \hat{u} \in V$  and any  $\varepsilon > 0$  there is a control  $\eta \in \Theta_T(h, u_0) \cap X$  such that

$$\|\mathcal{R}_T(u_0, \eta) - \hat{u}\|_V < \varepsilon. \quad (6)$$

Equations (2), (4) with  $\eta \in X$  are said to be  *$F$ -controllable in time  $T$*  if for any  $u_0 \in V$  and  $\hat{u} \in F$  there is  $\eta \in \Theta_T(h, u_0) \cap X$  such that

$$\mathbf{P}_F \mathcal{R}_T(u_0, \eta) = \hat{u}. \quad (7)$$

Equations (2), (4) with  $\eta \in X$  are said to be *solidly  $F$ -controllable in time  $T$*  if for any  $u_0 \in V$  and any  $R > 0$  there is a constant  $\delta > 0$  and a compact set  $\mathcal{C}$  in a finite-dimensional subspace  $Y \subset X$  such that  $\mathcal{C} \subset \Theta_T(h, u_0)$ , and for any continuous mapping  $\Phi : \mathcal{C} \rightarrow F$  satisfying the inequality

$$\sup_{\eta \in \mathcal{C}} \|\Phi(\eta) - \mathbf{P}_F \mathcal{R}_T(u_0, \eta)\|_F \leq \delta, \quad (8)$$

we have  $\Phi(\mathcal{C}) \supset B_F(R)$ .

For any finite-dimensional subspace  $G \subset U$ , we denote by  $\mathcal{F}(G)$  the largest vector space  $G_1 \subset U$  such that any element  $\eta_1 \in G_1$  is representable in the form

$$\eta_1 = \eta - \sum_{j=1}^k \lambda_j B(\zeta^j),$$

where  $\eta, \zeta^1, \dots, \zeta^k \in G$  are some vectors and  $\lambda_1, \dots, \lambda_k$  are non-negative constants. Since  $B$  is a quadratic operator continuous from  $U$  to  $V$ , we see that  $\mathcal{F}(G) \subset U$  is a well-defined vector space of finite dimension. Also note that  $\mathcal{F}(G) \supset G$ .

We now define a sequence of subspaces  $E_k \subset U$  by the rule

$$E_0 = E, \quad E_k = \mathcal{F}(E_{k-1}) \quad \text{for } k \geq 1, \quad E_\infty = \bigcup_{k=1}^{\infty} E_k. \quad (9)$$

The following theorem established in [Shi06a, Shi06b].

**Theorem 2.** *Let  $E \subset U$  be a finite-dimensional subspace such that  $E_\infty$  is dense in  $H$ . Then the following assertions take place for any  $T > 0$ ,  $\nu > 0$ , and  $h \in L^2(J_T, H)$ .*

- (i) *Equations (2), (4) with  $\eta \in C^\infty(J_T, E)$  are approximately controllable in time  $T$ .*
- (ii) *Equations (2), (4) with  $\eta \in C^\infty(J_T, E)$  are solidly  $F$ -controllable in time  $T$  for any finite-dimensional subspace  $F \subset H$ .*

In the general case, it is difficult to verify whether a subspace  $E \subset U$  satisfies the conditions of Theorem 2. However, if  $D$  is a torus in  $\mathbb{R}^3$ , then one can obtain a sufficient condition under which  $E_\infty$  is dense in  $H$ .

## 2 Case of a torus

In this subsection, we study controlled Navier–Stokes equations with periodic boundary conditions. More precisely, let us fix a vector  $q = (q_1, q_2, q_3)$  with positive components and set  $\mathbb{T}_q^3 = \mathbb{R}^3 / 2\pi\mathbb{Z}_q^3$ , where

$$\mathbb{Z}_q^3 = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 : x_i/q_i \in \mathbb{Z} \text{ for } i = 1, 2, 3\}.$$

Consider the Navier–Stokes system on  $\mathbb{T}_q^3$ . In other words, we consider Eqs. (1) with  $D = \mathbb{R}^3$  and assume that all functions are periodic of period  $2\pi q_i$  with respect to  $x_i$ ,  $i = 1, 2, 3$ . To simplify notation, we shall assume, without loss of generality, that the mean values of  $u$ ,  $h$ , and  $\eta$  with respect to  $x \in \mathbb{T}_q^3$  are zero. As in the case of a bounded domain with Dirichlet boundary condition, one can reduce (1) to an evolution equation in an appropriate Hilbert space. Namely, we set

$$H = \left\{ u \in L^2(\mathbb{T}_q^3, \mathbb{R}^3) : \operatorname{div} u \equiv 0, \int_{\mathbb{T}_q^3} u(x) dx = 0 \right\}$$

and denote by  $\Pi : L^2(\mathbb{T}_q^3, \mathbb{R}^3) \rightarrow H$  the orthogonal projection in  $L^2(\mathbb{T}_q^3, \mathbb{R}^3)$  onto the closed subspace  $H$ . Define the spaces

$$V = H^1(\mathbb{T}_q^3, \mathbb{R}^3) \cap H, \quad U = H^2(\mathbb{T}_q^3, \mathbb{R}^3) \cap H.$$

Projecting (1) to the space  $H$ , we obtain Eq. (2) in which  $L = -\Delta$  is the Stokes operator with the domain  $D(L) = U$  and  $B(u) = \Pi\{(u, \nabla)u\}$ . Theorem 2, which was formulated for the Dirichlet boundary condition, remains valid in this case as well. Our aim is to describe explicitly a finite-dimensional subspace  $E \subset U$  for which the hypothesis of Theorem 2 is fulfilled.

To this end, we first construct an orthogonal basis in  $H$  formed of the eigenfunctions of  $L$ . For  $x, y \in \mathbb{R}^3$ , let

$$\langle x, y \rangle_q = \sum_{i=1}^3 q_i^{-1} x_i y_i, \quad \langle x, y \rangle = \sum_{i=1}^3 x_i y_i, \quad |x| = \sum_{i=1}^3 |x_i|.$$

We set  $\mathbb{Z}_*^3 = \mathbb{Z}^3 \setminus \{0\}$  and  $\mathbb{R}_*^3 = \mathbb{R}^3 \setminus \{0\}$ . For  $a \in \mathbb{R}_*^3$ , denote by  $a^\perp$  the two-dimensional subspace in  $\mathbb{R}_*^3$  defined by the equation  $\langle x, a \rangle_q = 0$ . Note that  $a^\perp = (-a)^\perp$ . For any  $m \in \mathbb{Z}_*^3$ , let us choose a vector  $\ell(m) \in m^\perp$  so that  $\{\ell(m), \ell(-m)\}$  is an orthonormal basis in  $m^\perp$  with respect to the scalar product  $\langle \cdot, \cdot \rangle$ . We now set

$$c_m(x) = \ell(m) \cos\langle m, x \rangle_q, \quad s_m(x) = \ell(m) \sin\langle m, x \rangle_q \quad \text{for } m \in \mathbb{Z}_*^3.$$

It is a matter of direct verification to show that  $c_m$  and  $s_m$  are eigenfunctions of  $L$  and that  $\{c_m, s_m, m \in \mathbb{Z}_*^3\}$  is an orthogonal basis in  $H$ . For a finite family of functions  $\mathcal{A}$ , we denote by  $\text{span } \mathcal{A}$  the vector space spanned by  $\mathcal{A}$ .

**Theorem 3.** *For any vector  $q = (q_1, q_2, q_3)$  with positive components there is an integer  $d \geq 4$  such that if*

$$E = \text{span}\{c_m, s_m, |m| \leq d\},$$

*then the vector space  $E_\infty$  defined in (9) is dense in  $H$ .*

Theorems 2 and 3 imply the following result on controllability of the NS system by a force of finite dimension.

**Corollary 4.** *Let  $E \subset U$  be the subspace defined in Theorem 3. Then for any finite-dimensional subspace  $F \subset H$  and arbitrary constants  $T > 0$  and  $\nu > 0$  the Navier–Stokes equations (2), (4) with  $\eta \in C^\infty(J_T, E)$  are approximately controllable and solidly  $F$ -controllable in time  $T$ .*

The proofs of the above results are based on a development of a general approach introduced by Agrachev and Sarychev in the case of 2D Navier–Stokes equations (see [AS05, AS06]).

### 3 Applications

Our first application concerns the Cauchy problem for (2). Let  $G \subset H$  be a closed vector space. For any  $u_0 \in V$ ,  $T > 0$ , and  $\nu > 0$ , let  $\Xi_{T,\nu}(G, u_0)$  be the set of functions  $f \in L^2(J_T, G)$  for which problem (2), (3) has a unique solution  $u \in \mathcal{X}_T$ . If  $E \subset G$  is a closed subspace, then we denote by  $G \ominus E$  the orthogonal complement of  $E$  in  $G$  and by  $Q(T, G, E)$  the orthogonal projection in  $L^2(J_T, G)$  onto the subspace  $L^2(J_T, G \ominus E)$ . The following result is established in [Shi06a].

**Theorem 5.** *Let  $E \subset U$  be a finite-dimensional subspace such that  $E_\infty$  is dense in  $H$  and let  $G \subset H$  be a closed subspace containing  $E$ . Then  $\Xi_{T,\nu}(G, u_0)$  is a non-empty open subset of  $L^2(J_T, G)$  such that*

$$Q(T, G, E)\Xi_{T,\nu}(G, u_0) = L^2(J_T, G \ominus E) \quad \text{for any } T > 0, \nu > 0, u_0 \in V.$$

Our second application concerns the case in which Navier–Stokes equations are perturbed by a random force. Namely, suppose that

$$f(t, x) = h(x) + \eta(t, x), \quad (10)$$

where  $h \in H$  is a deterministic function and  $\eta$  is an  $H$ -valued random process satisfying the following condition.

- (C) There is an orthonormal basis  $\{f_k\}$  in  $V$  and a sequence of standard independent Brownian motions  $\{\beta_j(t), t \geq 0\}$  defined on a filtered probability space  $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$  such that

$$\eta(t) = \frac{\partial}{\partial t} \zeta(t), \quad \zeta(t) = \sum_{j,k=1}^{\infty} b_{jk} \beta_j(t) f_k,$$

where  $\{b_{jk}\}$  is a family of real constants satisfying the condition

$$B := \sum_{j,k=1}^{\infty} b_{jk}^2 < \infty.$$

Let us recall the concepts of an admissible weak solution and of a stationary measure for (2), (10). Define an Ornstein–Uhlenbeck process by the formula

$$z(t) = \int_0^t e^{-\nu(t-s)L} d\zeta(s).$$

It is well known that if Condition (C) is fulfilled, then  $z$  is a Gaussian process whose almost every trajectory belongs to the space  $C(\mathbb{R}_+, V) \cap L_{\text{loc}}^2(\mathbb{R}_+, U)$  and satisfies the Stokes equation

$$\dot{u} + \nu Lu = \eta(t).$$

**Definition 6.** An  $H$ -valued random process  $u(t)$  is called an *admissible weak solution* for (2), (10) if it is representable in the form

$$u(t) = v(t) + z(t),$$

where  $v(t)$  is an  $H$ -valued  $\mathcal{F}_t$ -adapted random process whose almost every trajectory belongs to the space  $L_{\text{loc}}^2(\mathbb{R}_+, V) \cap L_{\text{loc}}^\infty(\mathbb{R}_+, H)$  and satisfies the equation

$$\dot{v} + \nu Lv + B(v + z) = h$$

in the sense of distributions and the energy inequality

$$\begin{aligned} \frac{1}{2} \|v(t)\|^2 + \nu \int_0^t \|v(s)\|_V^2 ds + \int_0^t (B(v + z, z), v) ds \\ \leq \frac{1}{2} \|v(0)\|^2 + \int_0^t (h, v) ds, \quad t \geq 0, \end{aligned}$$

where  $(\cdot, \cdot)$  denotes the scalar product in  $H$ .

**Definition 7.** An admissible weak solution  $u(t)$  for (2), (10) is said to be *stationary* if its distribution does not depend on  $t$ :

$$\mathcal{D}(u(t)) = \mu \quad \text{for all } t \geq 0.$$

In this case,  $\mu$  is called a *stationary measure* for (2), (10).

Existence of admissible weak stationary solutions for 3D Navier–Stokes equations was established in [VF88, FG95]. Moreover, the construction of these works implies that

$$\int_H \|v\|_V^2 \mu(dv) < \infty. \quad (11)$$

Let us denote by  $Q$  the vector space of functions  $v \in V$  that are representable in the form

$$v = \sum_{j,k=1}^{\infty} b_{jk} u_j f_k,$$

where  $\{u_j\}$  is a sequence of real numbers such that  $\sum_j u_j^2 < \infty$ . Recall that the vector space  $E_\infty$  is defined in (9). For a finite-dimensional space  $F$ , denote by  $\ell_F$  the Lebesgue measure on  $F$ . The following theorem established in [Shi06c] provides some qualitative properties of stationary measures for (2), (10) (see also [AKSS06]).

**Theorem 8.** *Let  $\eta$  be a stationary process satisfying Condition (C), let  $E \subset U$  be a finite-dimensional vector space for which  $E_\infty$  is dense in  $H$ , and let  $\mu$  be a stationary measure for (2), (10) such that (11) holds. Suppose that  $Q \supset E$ . Then the following assertions take place.*

- (i) *The support of  $\mu$  coincides with  $H$ .*
- (ii) *Let  $F \subset H$  be a finite-dimensional subspace and let  $\mu_F$  be the projection of  $\mu$  to  $F$ . Then there is a function  $\rho_F \in C(F)$  such that  $\mu_F \geq \rho_F \ell_F$  and  $\rho_F(x) > 0$  for  $\ell_F$ -almost every  $x \in F$ .*

## References

- [AKSS06] A. Agrachev, S. Kuksin, A. Sarychev, and A. Shirikyan, *On finite-dimensional projections of distributions for solutions of randomly forced PDE's*, Preprint (2006).
- [AS05] A. A. Agrachev and A. V. Sarychev, *Navier–Stokes equations: controllability by means of low modes forcing*, J. Math. Fluid Mech. **7** (2005), 108–152.
- [AS06] A. A. Agrachev and A. V. Sarychev, *Controllability of 2D Euler and Navier–Stokes equations by degenerate forcing*, Commun. Math. Phys. (2006), to appear.
- [FG95] F. Flandoli and D. Gatarek, *Martingale and stationary solutions for stochastic Navier–Stokes equations*, Probab. Theory Related Fields **102** (1995), no. 3, 367–391.
- [Shi06a] A. Shirikyan, *Approximate controllability of three-dimensional Navier–Stokes equations*, Commun. Math. Phys. (2006), to appear.
- [Shi06b] A. Shirikyan, *Exact controllability in projections for three-dimensional Navier–Stokes equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire (2006), to appear.
- [Shi06c] A. Shirikyan, *Qualitative properties of stationary measures for three-dimensional Navier–Stokes equations*, in preparation (2006).
- [Tem79] R. Temam, *Navier–Stokes Equations*, North-Holland, Amsterdam, 1979.
- [VF88] M. I. Vishik and A. V. Fursikov, *Mathematical Problems in Statistical Hydromechanics*, Kluwer, Dordrecht, 1988.