

SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

JOHN B. WALSH

A non-reversible semi-martingale

Séminaire de probabilités (Strasbourg), tome 16 (1982), p. 212

<http://www.numdam.org/item?id=SPS_1982__16__212_0>

© Springer-Verlag, Berlin Heidelberg New York, 1982, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

A NON REVERSIBLE SEMI-MARTINGALE

John B. Walsh
University of British Columbia

The time-reversal of a semi-martingale may fail to be a semi-martingale. Here is a simple example.

Let B_t be a standard Brownian motion and, inspired by Barlow's example in [1], let ϕ be a measurable function which maps $C[0,1]$ one-to-one into $[0,1]$. Let $T(\omega) = \phi(\{B_s(\omega), 0 \leq s \leq 1\})$, and define

$$X_t = \begin{cases} B_t & \text{if } 0 \leq t \leq 1 \\ B_1 & \text{if } 1 \leq t \leq T + 1 \\ B_{t-T} & \text{if } t \geq 1 + T. \end{cases}$$

Then X is just a Brownian motion with a flat spot of length $T \leq 1$ interpolated from $t = 1$ to $t = T + 1$. T is $\sigma\{X_s, s \leq 1\}$ -measurable, so that it is easy to see that X is a martingale.

Now reverse X from $t = 2$: let $\tilde{X}_t = X_{2-t}$ for $0 \leq t \leq 2$. Let $(\tilde{\mathcal{F}}_t)$ be the natural filtration of \tilde{X} . Note that T is $\tilde{\mathcal{F}}_1$ -measurable, hence so is $\{\tilde{X}_t, t \leq 1\}$, since it is just the time-reversal of $\phi^{-1}(T)$. Consequently, $\tilde{\mathcal{F}}_t = \tilde{\mathcal{F}}_1$ for $t > 1$. Any martingale on these fields will be constant on $(1,2)$ and any semi-martingale will have finite variation there. But \tilde{X}_t has infinite variation on $(1,2)$, so it is not a semi-martingale relative to the $(\tilde{\mathcal{F}}_t)$. By Stricker's theorem [2], it can't be a semi-martingale relative to any filtration whatsoever.

References

- [1] Barlow, M.T. On Brownian Local Time. Preprint.
- [2] Stricker, C. Quasimartingales, martingales locales, semi-martingales, et filtrations naturelles, ZW 39, (1977) p 55-64.