

# SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

PETER JAGERS

OLLE NERMAN

## **The asymptotic composition of supercritical, multi-type branching populations**

*Séminaire de probabilités (Strasbourg)*, tome 30 (1996), p. 40-54

[http://www.numdam.org/item?id=SPS\\_1996\\_\\_30\\_\\_40\\_0](http://www.numdam.org/item?id=SPS_1996__30__40_0)

© Springer-Verlag, Berlin Heidelberg New York, 1996, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

# The Asymptotic Composition of Supercritical, Multi-Type Branching Populations

Peter Jagers and Olle Nerman\*

Department of Mathematics

Chalmers University of Technology and Gothenburg University

S-412 96 Göteborg, Sweden

## Abstract

The life, past and future are described of a typical individual in an old, non-extinct branching population, where individuals may give birth as a point process and have types in an abstract type space. The type, age and birth-rank distributions of the typical individual are explicitly given, as well as the Markov renewal type process that describes her history. The convergence of expected and actual compositions towards stable, asymptotic compositions is proved.

## 1 Introduction

If a proper, branching population does not die out, then its size grows indefinitely (*cf.* Jagers 1992, *e.g.*), and by some sort of law of large numbers its composition will stabilize. One aspect of this, the *stable age distribution* of demography has been known for a long time. Indeed, its roots can be traced back more than two centuries, to Euler, 1760. In such a grand perspective, the complete picture of the asymptotic composition of one-type general, supercritical branching populations is certainly recent (Jagers and Nerman and vice versa, 1984). The multi-type case was then investigated and presented by Nerman at the 16th Conference on Stochastic Processes and their Applications (1984). His results were, however, never published. A first account, informal and incomplete, appears in Jagers (1991 and, somewhat more extensively, 1992). A quite pertinent recent paper, from a different, graph and computer algorithm oriented, tradition is Aldous (1991).

The purpose of the present exposition is to give the strict description, that is lacking up to now, of the asymptotic composition of general super-critical

---

\*This work has been supported by a grant from the Swedish Natural Science Research Council

branching populations in abstract type spaces. It uses Markov renewal theory in Shurenkov's (1989) comprehensive formulation. The branching process framework and real time dynamics are from Jagers (1989).

## 2 The Population Space

Consider a typical individual of an old population. Call her *Ego* or, for short but less suggestively, 0; think of her as sampled at random from among all those born into the population, since its inception, long ago. She will have children, grandchildren *etc.*, whom we shall refer to in the classical Ulam-Harris manner,  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  is the  $x_n$ th child of the ... of the  $x_1$ :th child of 0, the set of 0 and all her possible descendants being denoted by  $I$ ,

$$I := \bigcup_{n \geq 0} N^n, \quad N^0 := \{0\}, \quad N := \{1, 2, \dots\}.$$

But 0 also has a mother, to be called -1, a grandmother, -2, and so forth. We concatenate vectors by writing them together, so that  $\mathbf{xy}$ ,  $\mathbf{x}, \mathbf{y} \in I$ , has first  $\mathbf{x}$ 's components, then  $\mathbf{y}$ 's, and we make the convention  $0\mathbf{x} = \mathbf{x}0 = \mathbf{x}$ . Then, all the possible progeny of -1, *except* 0 and her descendants, constitutes the set  $(-1)I = \{(-1)\mathbf{x}; \mathbf{x} \in I\}$ , and with  $Z_- := \{0, -1, -2, \dots\}$ , all the possible individuals of the whole population can be written

$$J := Z_- \times I.$$

A new-born individual is allotted (or chooses, depending upon your philosophy) a *life path* or life (career) from the *life space*  $(\Omega, \mathcal{A})$ . This should be thought of as abstract, with a countably generated  $\sigma$ -algebra, and rich enough to carry those functions that are of interest for the particular study. On  $\Omega$  there is a sequence  $0 \leq \tau(1) \leq \tau(2) \dots \leq \infty$  of random variables giving the successive ages at child-bearing. If  $\tau(j)(\omega) = \infty$ , then the interpretation is that the life career  $\omega$  involves fewer than  $j$  children.

At birth the new-born child inherits a *type* from a space  $(S, \mathcal{S})$ , again with a countably generated  $\sigma$ -algebra. In other words, there are also measurable functions  $\sigma(j) : \Omega \rightarrow S$ , giving the type of the  $j$ :th child. The *reproduction point process*  $\xi$  on  $S \times R_+$  is defined by

$$\xi(A \times B) := \#\{i \in N; \sigma(i) \in A, \tau(i) \in B\}.$$

In order to define the basic probability space for the whole population, the *population space*, we need the life space for each possible individual but also something that ties the individuals in  $Z_-$  to their mothers, information about their *birth ranks*. These are natural numbers, and we make the interpretation that if  $-j + 1$  has rank  $r$  then  $-j$  is  $-j$ 's  $i$ :th child for  $i = 1, \dots, r - 1$  and it is the  $i + 1$ :th child for  $i \geq r$ . Finally, to anchor the population in real time,

we need information about Ego's age at sampling. Thus the population space is defined to be

$$\Omega := R_+ \times N^\infty \times \Omega^J$$

with the obvious product  $\sigma$ -algebra to be denoted by  $\mathcal{C}$ . This space can be restricted to a tree-type space of the kind advocated by Neveu (1986) and Chauvin (1986). For the interpretation in terms of random trees (Aldous, 1991) that might indeed be an advantage.

Recall that a branching process started from a newborn 0-individual is suitably defined on  $S \times \Omega^J$ , the first coordinate being the ancestor's starting type. When referring to such a process, we shall write  $\sigma_x$  for the type of  $x \in I$  and  $\tau_x$  for her birth-time, defined recursively from  $\tau_0 = 0$  and the successive ages at child bearing in the line leading to  $x$ , cf. Jagers, 1989, though notation is slightly different there.

### 3 The Life Kernel and its Stable Population Law

The probability structure of the process is determined by a *life kernel*  $P_s$  on  $(\Omega, \mathcal{A})$ , the probability measure according to which the choice of life-career of an  $s$ -type is performed. In Jagers (1989) it is shown how such a kernel determines a unique Markov branching probability measure over  $(\Omega^J, \mathcal{A}^J)$ , once Ego's type has been fixed. We denote this measure, as well, by  $P_s$ ,  $s \in S$  being this starting type. More generally, if Ego's type is chosen according to a probability measure  $\pi$  on  $(S, \mathcal{S})$ , then  $P_\pi$  is the corresponding measure,  $P_\pi = \int_S P_s \pi(ds)$ . The expectations are  $E_s$  and  $E_\pi$ , respectively.

The crucial rôle for the development of a population is played by the *reproduction kernel*  $\mu$ , defined as the expected number of births of children of various types and at various ages:

$$\mu(r, ds \times dt) := E_r[\xi(ds \times dt)].$$

The population is supposed to be *Malthusian* and *supercritical*, this meaning that there is a number  $\alpha > 0$ , the Malthusian parameter, such that the kernel  $\hat{\mu}(\alpha)$ ,

$$\hat{\mu}(r, ds; \alpha) := \int_0^\infty e^{-\alpha t} \mu(r, ds \times dt)$$

has Perron root one and is what Shurenkov (1989) calls *conservative*. (This corresponds to irreducibility and  $\alpha$ -recurrence in the terminology of Niemi and Nummelin (1986).) By the abstract Perron-Frobenius theorem (Shurenkov, 1989, p. 43, or Nummelin, 1984, p. 70), there is then a  $\sigma$ -finite measure  $\pi$  on the type space  $(S, \mathcal{S})$ , and strictly positive a.e.  $[\pi]$  finite measurable function  $h$  on the same space, such that

$$\int_S \hat{\mu}(r, ds; \alpha) \pi(dr) = \pi(ds),$$

$$\int_S h(s) \hat{\mu}(r, ds; \alpha) = h(r).$$

Further we require strong or positive  $\alpha$ -recurrence in the sense that  $h \in L^1[\pi]$  and

$$0 < \beta = \int_{S \times S \times R_+} t e^{-\alpha t} h(s) \mu(r, ds \times dt) \pi(dr) < \infty.$$

(In population dynamics this entity might be interpreted as the stable age at childbearing, though some care has to be exercised about this in the multi-type case, as we shall see.) Then we can (and shall) norm to

$$\int_S h d\pi = 1.$$

Throughout we also make the homogeneity assumption that  $\inf h > 0$ . Then  $\pi$  is finite and can (and will) also be normed to a probability measure. These are the conditions (on  $\mu$  alone) for the general Markov renewal theorem of Shurenkov (1989), p. 107. Finally, we assume that the reproduction kernel is non-lattice and satisfies the natural condition

$$\sup_s \mu(s, S \times [0, \epsilon]) < 1$$

for some  $\epsilon > 0$ . Note that we assume only non-latticeness, rather than spread-outness of the kernel. We shall summarize all these conditions by referring to the population as non-lattice *strictly Malthusian*. (Clearly there is a lattice analog of our results, relying upon the lattice Markov renewal theorem, cf. Shurenkov (1989), p. 122. There you will also find the concept of latticeness developed in a multitype context, with the meaning that there is a stepping time unit, independent of starting and ending position in the type space, but a phase which may depend on both: for some  $d > 0$  and  $c : S \rightarrow [0, d)$ , and  $L_{dc}(s) := \{(r, t); r \in S, t \in R_+, t = c(r) - c(s) + nd, \text{ for some } n = 0, 1, \dots\}$

$$\pi(\{s; \mu(s, S \times R_+) > \mu(s, L_{dc}(s))\}) = 0).$$

In order, finally, to give a presentation of the stable population measure on  $(\Omega, \mathcal{A})$  we need notation for some random elements on this space:  $T_0$  will denote Ego's age at sampling,  $S_0$  her type, and  $R_0$  her rank, *i.e.* ordinal number in her sibship.  $T_1$  is Ego's mother's age, when she gave birth to Ego,  $S_1$  her type, and  $R_1$  her rank. And so on backwards. Similarly we let  $U_0, U_1, \dots$  denote the whole lives of Ego, Ego's mother,  $\dots$ , and  $Z^0$  the population initiated by Ego.  $Z^1, Z^2, \dots$  can be used to denote Ego's mother's life and daughter process *except Ego*, grandmother's daughter process, *except mother and her progeny etc.* Thus,  $Z^j$  is the coordinate projection  $\Omega \rightarrow \Omega^{-jI}$ . Similarly,  $T_0$  is the projection of the population space onto its first coordinate  $R_+$  and the sequence of ranks is the projection onto  $N^\infty$ , *cf.* the figure. Also recall that  $\sigma_i, \tau_i$  are the type and




Figure. The doubly infinite population space.

birth times in a branching process of the ancestor's  $i$ :th child (the latter equalling infinity, if no  $i$ :th child is ever born). In the following definition we shall interpret assertions ' $Z^j \in A_j$ ' also as assertions about a traditional branching process ( $-j$  being its ancestor). Then we must supplement it by information about which of the ancestor's children, who has been withdrawn from the process in order to play the rôle of  $-j + 1$ . Indeed, the very careful reader should interpret  $A_j, j = 1, 2, \dots$  on the right hand side beneath as the set

$$\{(\omega_{kx}; k \in Z_+, x \in I) \in \Omega^I; (\omega_{(k+1)_{\{k \geq i_{j-1}\}}}; k \in Z_+, x \in I) \in A_j\}.$$

Such subtleties are due to the convention we made about birth ranks in Section 2, where the concatenated vector  $kx$  was also defined.

**Definition 1** *The stable population measure  $\mathbf{P}$  on  $(\Omega, \mathcal{A})$  is determined by*

$$\begin{aligned} \mathbf{P}(Z^0 \in A_0, T_0 \in dt_0, S_0 \in ds_0, R_0 = i_0; Z^1 \in A_1, T_1 \in dt_1, S_1 \in ds_1, R_1 = i_1; \\ \dots; Z^n \in A_n, T_n \in dt_n, S_n \in ds_n, R_n = i_n) = \\ E_\pi[e^{-\alpha\tau_{i_n}}; \sigma_{i_n} \in ds_n] E_{s_n}[e^{-\alpha t_n}; A_n \cap \{\sigma_{i_{n-1}} \in ds_{n-1}, \tau_{i_{n-1}} \in dt_n\}] \dots \\ \dots E_{s_1}[e^{-\alpha t_1}; A_1 \cap \{\sigma_{i_0} \in ds_0, \tau_{i_0} \in dt_1\}] P_{s_0}(A_0) \alpha e^{-\alpha t_0} dt_0 \end{aligned}$$

for all  $n \in N, A_j \in \mathcal{A}^I, t_j \in R_+, s_j \in S, i_j \in N, j = 0, 1 \dots n$ .

By the eigenmeasure and eigenfunction properties of  $\pi$  and  $h$  it can be checked that changes in  $n$  yield a projective system defining  $\mathbf{P}$ , and in the next section we shall see how the stable population appears as a limit of growing branching processes. Here we shall try to understand its substance, by formulating some consequences of the definition. In them we write  $\hat{\mu}(r, ds)$  as an abbreviation for  $\hat{\mu}(r, ds; \alpha) = \int_0^\infty e^{-\alpha t} \mu(r, ds \times dt)$ . Powers  $\hat{\mu}^n$  are iterated kernels and  $\mu_\alpha(r, ds \times dt) := e^{-\alpha t} \mu(r, ds \times dt)$ .

**Proposition 1** *The sequence of types backwards from Ego,  $\{S_n\}_0^\infty$ , is a Markov chain with transition probabilities*

$$\mathbf{P}(S_{n+1} \in ds \mid S_n = r) = \pi(ds) \frac{\hat{\mu}(s, dr)}{\pi(dr)}.$$

*The distribution of  $S_0$  is  $\pi$ , whereas  $S_n \sim \hat{\mu}^n(s, S) \pi(ds) \rightarrow h(s) \pi(ds)$ , as  $n \rightarrow \infty$ , the latter limit also being the stationary distribution of the chain.*

**Proof** Integrating and summing in the theorem yields

$$\begin{aligned} \mathbf{P}(S_0 \in ds_0, \dots, S_n \in ds_n) = \\ = E_\pi \left[ \sum_{i_n} e^{-\alpha\tau(i_n)}; \sigma(i_n) \in ds_n \right] E_{s_n} \left[ \sum_{i_{n-1}} e^{-\alpha\tau(i_{n-1})}; \sigma(i_{n-1}) \in ds_{n-1} \right] \dots \end{aligned}$$

$$\begin{aligned}
& \dots E_{s_1} \left[ \sum_{i_0} e^{-\alpha \tau(i_0)}; \sigma(i_0) \in ds_0 \right] = \\
& = E_\pi [\hat{\xi}(ds_n; \alpha)] E_{s_n} [\hat{\xi}(ds_{n-1}; \alpha)] \dots E_1 [\hat{\xi}(ds_0; \alpha)] \\
& = \\
& \pi(ds_n) \hat{\mu}(s_n, ds_{n-1}) \dots \hat{\mu}(s_1, ds_0),
\end{aligned}$$

where

$$\hat{\xi}(A; \alpha) := \int_0^\infty e^{-\alpha t} \xi(A \times dt),$$

we have used the eigenmeasure property  $\int \hat{\mu}(s, ds_n) \pi(ds) = \pi(ds_n)$ , and written  $\sigma(i), \tau(i)$  instead of  $\sigma_i, \tau_i$ , since the  $P_{s_i}$  reduce to measures over the life space here. The asserted form of transition and marginal probabilities follow from this joint distribution of types. The convergence  $\hat{\mu}^n(s, S) \rightarrow h(s)$  follows directly from the lattice Markov renewal theorem (Shurenkov, 1989, p. 122). Of course, it can also be brought back to a limit theorem for Markov chains by the trick of norming  $\hat{\mu}$  to a kernel with mass one,  $h(r) \hat{\mu}(s, dr) / h(s)$ . The stationarity can be checked directly.

□

Without spelling this out as another proposition, let us state that the sequence  $\{(R_n, S_n)\}$  of ranks and types also constitutes a Markov chain. Indeed, given the sequence of types, the ranks even become conditionally independent. The rank marginals are given by  $\mathbf{P}(R_0 = i) = E_\pi[e^{-\alpha \tau(i)}]$ , and

$$\mathbf{P}(R_n = i) = \int_S \hat{\mu}^n(s, S) E_\pi[e^{-\alpha \tau(i)}; \sigma(i) \in ds] \rightarrow \int_S h(s) E_\pi[e^{-\alpha \tau(i)}; \sigma(i) \in ds],$$

as  $n \rightarrow \infty$ . Though the distribution of, at least,  $R_0$  is important for birth rank studies, the joint behaviour of types and times between births seems of greater import both mathematically, and in tracing populations backwards, *e.g.* in evolutionary genetics.

**Proposition 2** *The sequence of types and interbirth times backwards from Ego,  $\{S_n, T_n\}_0^\infty$  define a Markov renewal process. They have the transition kernel*

$$\mathbf{P}(S_{n+1} \in ds, T_{n+1} \in dt \mid S_n = r) = \pi(ds) \frac{\mu_\alpha(s, dr \times dt)}{\pi(dr)}.$$

*The distribution of  $S_0$  is  $\pi$ ,  $T_0$  is exponentially distributed with the Malthusian parameter, and independent of the rest.*

$$(S_n, T_n) \sim \int_S \hat{\mu}^{n-1}(r, S) \mu_\alpha(s, dr \times dt) \pi(ds) \rightarrow \int_S h(r) \mu_\alpha(s, dr \times dt) \pi(ds),$$

as  $n \rightarrow \infty$ .



**Proof** The proof follows the pattern of the preceding one, and is left out.  $\square$

Among other things, this shows that the expected age of the mother at the birth of a random child is

$$\mathbf{E}[T_1] = \int_{S \times R_+} t e^{-\alpha t} \mu(\pi, ds \times dt),$$

whereas the expectation of the asymptotic distribution of  $T_n, n \rightarrow \infty$  is

$$\beta = \int_{S \times R_+} t e^{-\alpha t} h(s) \mu(\pi, ds \times dt).$$

Here, of course,  $\mu(\pi, ds \times dt) = \int_S \mu(r, ds \times dt) \pi(dr)$

In analogy with Proposition 2, the sequence  $\{R_n, S_n, T_n\}$  constitutes a Markov renewal process, with a transition kernel that is easily determined from Definition 1. Actually, more generally:

**Proposition 3** *The sequence of ranks, types, and lives backwards from Ego has the Markov property*

$$\begin{aligned} \mathbf{P}(R_{n+1} = j, S_{n+1} \in ds, U_{n+1} \in A \mid R_n = i, S_n = r, U_n, R_{n-1}, S_{n-1} \dots) = \\ = E_\pi[e^{-\alpha\tau(j)}; \sigma(j) \in ds] \frac{E_s[e^{-\alpha\tau(i)}; A \cap \{\sigma(i) \in dr\}]}{E_\pi[e^{-\alpha\tau(i)}; \sigma(i) \in dr]}. \end{aligned}$$

The distribution of  $(R_0, S_0, U_0)$  is  $E_\pi[e^{-\alpha\tau(i)}; \sigma(i) \in ds] P_s(A), i \in N, s \in S, A \in \mathcal{A}$ , whereas in the notation

$$\hat{\mu}_A(s, B) := E_s[\hat{\xi}(B; \alpha); A],$$

$$\begin{aligned} \mathbf{P}(R_n = i, S_n \in ds, U_n \in A) = \\ = E_\pi[e^{-\alpha\tau(i)}; \sigma(i) \in ds] \int_S \hat{\mu}^{n-2}(r, S) \hat{\mu}_A(s, dr) \rightarrow \\ \rightarrow E_\pi[e^{-\alpha\tau(i)}; \sigma(i) \in ds] \int_S h(r) \hat{\mu}_A(s, dr), \end{aligned}$$

as  $n \rightarrow \infty$ .

**Proof** The proof is again (rather complicated but) not hard by insertion.  $\square$

The stable population measure  $\mathbf{P}$  describes a typical individual, her background and future when *sampling from among all those born*, dead or alive. Though this is artificial from, say, a biological viewpoint, it is not only mathematically convenient but also conceptually the fundamental situation. Being alive or not is a property of your age and your life career. Therefore the stable measure when *sampling in the live population* is obtained by conditioning in  $\mathbf{P}$  on Ego being alive, and correspondingly for sampling from other subsets of individuals.

To express this more formally, assume a *life span*,  $\lambda : \Omega \rightarrow [0, \infty]$ , defined and let  $L_0$  denote Ego's life span. Then,  $L_0 > T_0$  means that Ego is alive.

**Proposition 4** *The probability law describing a typical individual, sampled from among those alive is  $\mathbf{P}(\cdot \mid L_0 > T_0)$ .*

**Corollary 1** *The probability that a typical, live individual is of rank  $i$ , has type in  $ds$  and a life career in a set  $A \in \mathcal{A}$  is*

$$E_\pi[e^{-\alpha\tau(i)}; \sigma(i) \in ds] \int_0^\infty P_s(A, \lambda > t) \alpha e^{-\alpha t} dt / \int_0^\infty P_s(\lambda > t) \alpha e^{-\alpha t} dt.$$

*The probability of having just the property  $A$  is*

$$\frac{\int_0^\infty P_\pi(A, \lambda > t) e^{-\alpha t} dt}{\int_0^\infty P_\pi(\lambda > t) e^{-\alpha t} dt}.$$

In the next section we shall see that this is, indeed, the limit of the probability measure describing the properties of an individual sampled from among all those alive.

## 4 Convergence towards Stable Population Composition

Let  $J_n$  denote the class of individuals stemming from  $-n$ , i.e.  $\{-n, \dots, -1, 0\} \times I$ . Recall that by convention  $0 \in I$ ,  $0x = x$ , and  $-j0 = (-j, 0) = -j$ . Therefore  $-jI = \{-j\} \times I$  denotes  $-j$  and all her possible descendants except  $-j + 1$  and her progeny. Observe that if  $-n$  is mapped to 0, all her progeny being mapped onto  $I$  so as to preserve all family relations, then  $(\Omega^{J_n}, \mathcal{A}^{J_n})$  is mapped to  $(\Omega^I, \mathcal{A}^I)$ , and the two spaces can be thus identified. Fix  $n$ ,  $\iota = (i_{n-1} \dots i_0) \in N^n$ ,  $A \in \mathcal{A}^{J_n}$  and  $a \in R_+$ , and consider the subset  $E \in \mathcal{C}$ ,

$$E := [0, a] \times (i_0 \dots i_{n-1}) \times N^\infty \times A \times \Omega^{J \setminus J_n}.$$

Define  $\Pi_x$  as the projection mapping  $(s, \{\omega_y; y \in I\})$  to  $(\sigma_x, \{\omega_{xy}; y \in I\})$ , the *daughter process* of  $x \in I$ , cf. Jagers, 1989. In an obvious sense  $y \in I$  has the property  $E$  at time  $t$  if and only if

- $y = x_i$  for some  $x_i \in I$ ,
- $0 \leq t - \tau_{x_i} \leq a$ , and
- $\Pi_x \in S \times A$ .

Now, note that  $\tau_{x_i} = \tau_x + \tau_i \circ \Pi_x$ , so that defining a *random characteristic* on  $(S \times \Omega^I, \mathcal{S} \times \mathcal{A}^I)$  (cf. Jagers, 1989)

$$\chi^E(t) := 1_A 1_{[0,a]}(t - \tau_i),$$

( $A$  thus viewed as a subset of  $\Omega^I$ ), the number of individuals having the property  $E$  at time  $t$  in a branching population started at time 0 from a newborn ancestor will be

$$z_t^{\chi^E} := \sum_{x \in I} \chi^E(t) \circ \Pi_x = \sum_{x \in I} 1_{\{\Pi_x \in S \times A\}} 1_{\{0 \leq t - \tau_{x_i} \leq a\}}.$$

But adapting the convergence theorem for means of supercritical general branching populations (cf. *op. cit.*) to the Markov renewal theorems of Shurenkov (1989, pp. 107, 127, 134) we have:

**Theorem 1** *Consider a non-lattice, strictly Malthusian, supercritical branching population, counted with a bounded characteristic  $\chi$  such that the function  $e^{-\alpha t} E_s[\chi(t)]$  is directly Riemann integrable ( $\pi$ ). Then, for  $\pi$ -almost all  $s$ ,*

$$\lim_{t \rightarrow \infty} e^{-\alpha t} E_s[z_t^{\chi}] = h(s) \int_{S \times R} e^{-\alpha t} E_r[\chi(t)] \pi(dr) dt / \beta := h(s) E_\pi[\hat{\chi}(\alpha)] / \alpha \beta,$$

in the obvious notation for Laplace transform.

If the population is as above and some convolution power of the reproduction kernel is, further, non-singular (cf. beneath), then for  $\pi$ -almost all  $s \in S$

$$\lim_{t \rightarrow \infty} e^{-\alpha t} E_s[z_t^{\chi}] = h(s) E_\pi[\hat{\chi}(\alpha)] / \alpha \beta,$$

uniformly in all  $\chi$  with  $E_s[\chi(t)] \leq 1$  (without any Riemann integrability requirement).

The notion of direct Riemann integrability used is that of Shurenkov (1989) pp. 80 ff.: A measurable function  $g : S \times R_+ \rightarrow R$  is directly Riemann integrable ( $\pi$ ) if for any  $\epsilon > 0$  we can find  $\delta > 0$  and functions  $g^-$  and  $g^+$  both in  $L^1[\pi \times dt]$  such that for  $\pi$ -almost all  $s$ ,  $g^-(s, \cdot) \leq g(s, \cdot) \leq g^+(s, \cdot)$ ,  $g^\pm(s, t) = g^\pm(s, n\delta)$  for  $n\delta \leq t < (n+1)\delta$ , and the  $L^1[\pi \times dt]$ -distance between  $g^+$  and  $g^-$  is less than  $\epsilon$ . Convolution means convolution in time combined with transition in type. Non-singularity is Shurenkov's term (1989, p. 127) for spread-outness: For fixed  $r \in S$  and a Borel set  $B$ , the reproduction kernel  $\mu(r, \cdot \times B)$  is absolutely continuous with respect to  $\hat{\mu}(r, \cdot)$ . It is possible to choose a regular version of the Radon-Nikodym derivative,  $F(r, s, dt)$ , which is a measure on  $R_+$  in its

last coordinate. Non-singularity means that for almost all  $r, s$  with respect to  $\pi(dr)\hat{\mu}(r, ds)$  this measure is non-singular with respect to Lebesgue measure.

Thanks to this strong Markov renewal theory, the proof is rather straightforward, cf. Jagers (1992), by use of the regularity condition  $\sup_s \mu(s, S \times [0, \epsilon]) < 1$  for some  $\epsilon > 0$ , in order to guarantee boundedness of  $e^{-\alpha t} E_s[y_t]$ , where

$$y_t = \#\{x \in I; \tau_x \leq t\} = \sum_{x \in I} 1_{R_+}(t - \tau_x) = z_t^{1_{R_+}}$$

is the *total population* at time  $t$  of a branching process started at time 0. Of course, there is also a lattice variant of this result, cf. Shurenkov (1989, p. 122 and 134).

For sets  $E$  as above, we call

$$P_{s,t}^e(E) := E_s[z_t^{X^E}] / E_s[y_t],$$

the *composition in expectation* of a branching population at time  $t$ , started at time 0 from an ancestor of type  $s \in S$ . By summation over various  $\mathfrak{z} \in N^n$  and replacing the interval  $[0, a]$  by Borel sets  $B$ , this can obviously be extended to a probability measure over the measurable subsets of  $R_+ \times N^\infty \times \Omega^J$  which depend only upon  $n \in N$  steps backwards, i.e. belong to the  $\sigma$ -algebra generated by sets of the form  $E = B \times M \times N^\infty \times A \times \Omega^{J \setminus J_n}$ ,  $B \in \mathcal{B}(R_+)$ ,  $M \subset N^n$ ,  $A \in \mathcal{A}^{J_n}$  for fixed, but arbitrary  $n \in N$ . We denote the latter by  $\mathcal{C}_n$  and write  $\mathcal{C}_n(B)$  for the sub- $\sigma$ -algebra, where the first coordinate is fixed to be  $B$ .

**Corollary 2** *Under the assumptions of Theorem 1, consider a  $B \in \mathcal{B}(R_+)$ ,  $n \in N$ , and any  $E \in \mathcal{C}_n(B)$  such that  $R_j = i_j$ ,  $j = 0, \dots, n-1$ . Write  $\mathfrak{z} = (i_{n-1}, \dots, i_0)$  and assume that  $e^{-\alpha t} P_s(t - \tau_{\mathfrak{z}} \in B)$  is directly Riemann integrable. Then, the composition in expectation of a non-lattice, strictly Malthusian, and supercritical branching population at time  $t$ , started at time 0 from an ancestor of  $\pi$ -almost any type  $s \in S$ , satisfies*

$$P_{s,t}^e(E) = E_s[z_t^{X^E}] / E_s[y_t] \rightarrow \int_B \alpha e^{-\alpha t} dt E_\pi[e^{-\alpha \tau_{\mathfrak{z}}}; A] = \mathbf{P}(E),$$

as  $t \rightarrow \infty$ .

**Proof** This is only checking the direct Riemann integrability. Note that  $\mathfrak{z}$  is fixed, and of course matters in the coordinate projection singling out  $A$  from  $E$ , cf. the discussion preceding Definition 1. □

**Corollary 3** *If reproduction (i.e. some convolution power of the reproduction kernel) is non-singular,, besides the conditions of Theorem 1, then for  $\pi$ -almost all  $s$   $P_{s,t}^e \rightarrow \mathbf{P}$  in total variation, as  $t \rightarrow \infty$ .*

**Proof** By Theorem 1, the convergence is uniform over sets  $E \in \mathcal{C}_n$  for  $n$  fixed, at least if they are of the form  $E = B \times M \times N^\infty \times A \times \Omega^J \setminus J_n$ ,  $B \in \mathcal{B}(R_+)$ ,  $M \subset N^n$ ,  $A \in \mathcal{A}^J$ . But these sets,  $n \in N$  constitute an algebra that generates  $\mathcal{C} = \bigvee_n^\infty \mathcal{C}_n$ . The rest follows by approximation. □

Leaving composition in expectation, we turn to the *actual* composition,

$$P_{\sigma_0, t}(E) := z_t^{\chi^E} / y_t.$$

As for classical cases, convergence here requires the famed  $x \log x$ -condition. It has the following general form: Write

$$\bar{\xi} := \int_{S \times R_+} e^{-\alpha t} h(s) \xi(ds \times dt).$$

Then the condition is

$$E_\pi[\bar{\xi} \log^+ \bar{\xi}] < \infty.$$

From Jagers (1989) we have:

**Theorem 2** *Add the  $x \log x$ -condition and finiteness of  $\xi(S \times R_+)$  to the assumptions of Theorem 1. Further assume that, for fixed  $t$ ,  $y_t$  is uniformly integrable over its starting type  $\sigma_0 = s \in S$ . Then, as  $t \rightarrow \infty$ ,*

$$e^{-\alpha t} z_t^\chi \rightarrow w E_\pi[\hat{\chi}(\alpha)] / \alpha \beta$$

in  $L^1[P_s]$ , for  $\pi$ -almost all  $s \in S$ . Here  $w$  is a non-negative random variable with  $E_s[w] = h(s)$ .

Note that we have  $w$  with  $E_s[w] = h(s)$ , rather than expectation one as asserted in *op. cit.*. It is the unnormed random variable  $w$  that is the limit of the intrinsic martingale

$$w_L = \sum_{x \in L} e^{-\alpha \tau_x} h(\sigma_x)$$

(a.s. if only sequences of lines  $L$  are considered, in  $L^1$  otherwise, *c.f. op. cit.*).

From its definition (and Theorem 2) it is clear that  $w > 0 \Rightarrow y_t \rightarrow \infty$ . The converse of this is needed to show that for bounded characteristics  $\chi$

$$z_t^\chi / y_t \rightarrow E_\pi[\hat{\chi}(\alpha)]$$

if only  $y_t \rightarrow \infty$ .

**Lemma 1** *For a strictly Malthusian process, assume that  $\inf_{s \in S} P_s(w > 0) > 0$ . Then,  $w > 0 \Leftrightarrow y_t \rightarrow \infty$  a. s.  $P_s, s \in S$ .*

**Proof** Enumerate individuals in the order they are born into the population:  $X_0 = 0 \in I, 0 = \tau_{X_0} \leq \tau_{X_1} \leq \tau_{X_2} \leq \dots$  by some rule that guarantees that mothers precede their daughters (if individuals happen to appear simultaneously). Then

$$y_t = \sup\{n; \tau_{X_n} \leq t\}.$$

If  $y_t \rightarrow \infty$ , then the sequence  $\{X_n\}$  is already well defined. Otherwise, just continue somehow, respecting the rule that mothers must precede daughters. The assumption of strict Malthusianness prevents explosion in finite time. Hence  $\tau_{X_n} \rightarrow \infty$  whether  $y_t \rightarrow \infty$  or not.

But for any  $n \in N$

$$w \circ S_{X_n} > 0 \Rightarrow w > 0 \text{ or } y_t \not\rightarrow \infty.$$

(As the reader has noted we are far from finical about spelling out a.s.-qualifications.) If  $\mathcal{A}_n$  denotes the  $\sigma$ -algebra generated by the ancestor's type  $\sigma_0$  and the lives  $\omega_{X_0}, \omega_{X_1}, \dots, \omega_{X_n}$ , then Lévy's theorem yields that

$$\begin{aligned} 0 &< \inf_r P_r(w > 0) \leq P_{\sigma_{X_n}}(w \circ S_{X_n} > 0) = \\ &= P_s(w \circ S_{X_n} > 0 \mid \mathcal{A}_{n-1}) \leq P_s(w > 0 \text{ or } y_t \not\rightarrow \infty \mid \mathcal{A}_{n-1}) \rightarrow 1_{\{w > 0 \text{ or } y_t \not\rightarrow \infty\}}, \end{aligned}$$

as  $n \rightarrow \infty$ . Hence, a.s.  $y_t \rightarrow \infty \Rightarrow w > 0$ , the converse implication being already noted.

□

Note that under the conditions of Theorem 2,  $E_s[w] = h(s) > 0$ . Hence, for all  $s \in S$ ,  $P_s(w > 0) > 0$  and suitable compactness assumptions yield the same for the infimum.

**Corollary 4** *Let the assumptions of Theorem 2 hold and add that  $\inf_s P_s(w > 0) > 0$ . Then, for any  $E$  as in Corollary 2, the actual composition converges to the stable composition on the set of non-extinction:*

$$P_{s,t}(E) \rightarrow \mathbf{P}(E)$$

in  $P_s$ -probability on  $\{y_t \rightarrow \infty\}$  for  $\pi$ -almost any  $s \in S$ , as  $t \rightarrow \infty$ .

By invoking Aldous's paper (1991) we could have made the argument marginally simpler, proving Corollary 2 just in the setting where  $E$  depends only upon Ego's and her progeny's lives. That yields the convergence of his fringe tree, which is Ego's daughter process in our terminology. Realizations of our stable population process  $(\Omega, \mathcal{A}, \mathbf{P})$  are "sin-trees" in Aldous's parlance: they have a single infinite path. Thus the convergence of the extended fringe follows from Aldous's Proposition 11. Unfortunately, his main theorem on extremality of invariant laws, and the ensuing convergence in probability of the fringe distribution, is of

little avail here, since we have this type of convergence from the beginning and it is not easier to prove convergence in distribution.

Finally, note that in order to obtain average  $\chi$ -values among those *alive* we should just consider ratios  $z_t^X/z_t$  rather than  $z_t^X/y_t$ ,  $z_t = z_t^{1[0,\lambda)}$  denoting the number of individuals alive at time  $t$ . (This interpretation presumes that  $\chi$  counts only living individuals, i. e. that it vanishes outside the interval  $[0, \lambda)$ .)

## 5 References

1. Aldous, D. (1991). Asymptotic fringe distributions for general families of random trees. *Ann. Appl. Prob.* 1 228-266.
2. Chauvin, B. (1986). Arbres et processus de Bellman-Harris. *Ann. Inst. H. Poincaré* 22, 199-207.
3. Cohn, H. and Jagers, P. (1994) General branching processes in varying environment. *Ann. Appl. Prob.* 4 184-193.
4. Euler, L. *Recherches générales sur la mortalité et la multiplication du genre humain*. In: Histoire de l'Académie Royale des Sciences et Belles-Lettres 1760 (Berlin 1767), 144-164.
5. Jagers, P. (1975) *Branching Processes with Biological Applications*. J. Wiley & Sons, Chichester etc..
6. Jagers, P. (1989) General branching processes as Markov fields. *Stoch. Proc. Appl.* 32 183-242.
7. Jagers, P. (1991) The growth and stabilization of populations. *Statist. Sci.* 6 269-283.
8. Jagers, P. (1992) Stabilities and instabilities in population dynamics. *J. Appl. Prob.* 29.
9. Jagers, P. and Nerman, O. (1984) The growth and composition of branching populations. *Adv. Appl. Prob.* 16 221-259.
10. Nerman, O. (1984) *The Growth and Composition of Supercritical Branching Populations on General Type Spaces*. Dep. Mathematics, Chalmers U. Tech. and Gothenburg U. 1984-4.
11. Nerman, O. and Jagers, P. (1984) The stable doubly infinite pedigree process of supercritical branching populations. *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 64 445-460.
12. Neveu, J. (1986) Arbres et processus de Galton-Watson. *Ann. Inst. H. Poincaré* 22 199-207.

13. Niemi, S. and Nummelin, E. (1986) On non-singular renewal kernels with an application to a semigroup of transition kernels. *Stoch. Proc. Appl.* **22** 177-202.
14. Nummelin, E. (1984) *General Irreducible Markov Chains and Non-negative Operators*. Cambridge University Press, Cambridge.
15. Shurenkov, V. M. (1984) On the theory of Markov renewal. *Theory Prob. Appl.* **29** 247-265.
16. Shurenkov, V. M. (1989) *Ergodicheskie protsessy Markova*. Nauka, Moscow.