

LEV BIRBRAIR

MARINA SOBOLEVSKY

Realization of Hölder complexes

Annales de la faculté des sciences de Toulouse 6^e série, tome 8, n^o 1
(1999), p. 35-44

http://www.numdam.org/item?id=AFST_1999_6_8_1_35_0

© Université Paul Sabatier, 1999, tous droits réservés.

L'accès aux archives de la revue « Annales de la faculté des sciences de Toulouse » (<http://picard.ups-tlse.fr/~annales/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Realization of Hölder Complexes^(*)

LEV BIRBRAIR and MARINA SOBOLEVSKY⁽¹⁾

RÉSUMÉ. — Un complexe de Hölder est un graphe fini tel qu'à chaque arête est associé un nombre rationnel positif et on sait que c'est un invariant bi-lipschitzien des ensembles semi-algébriques singuliers de dimension 2. On montre dans cet article que tout complexe de Hölder peut être réalisé comme un ensemble semi-algébrique de dimension 2. Pour ce faire on plonge le graphe dans un tore de dimension n qu'on fait contracter sur un point singulier de telle sorte que les générateurs s'évanouissent avec les vitesses rationnelles et différentes.

ABSTRACT. — Hölder Complex, a graph and a rationally-valued function on the set of the edges of the graph, is a bi-Lipschitz invariant of 2-dimensional semialgebraic singular sets. Here we prove that each Hölder Complex can be realized as a 2-dimensional semialgebraic set. For this purpose we embed the graph to an n -dimensional torus. The torus is vanishing in a singular point such that the generators are vanishing with different rational rates.

1. Introduction

The paper is devoted to the local geometry of 2-dimensional semialgebraic sets. The local bi-Lipschitz classification theorem is proved in [1]. The main notion of the classification is a so-called Geometric Hölder Complex. It is a local version of a simplicial complex with some additional geometric information (see the definition below). A Hölder Complex can be considered as a combinatorial object – a finite graph with a rational-valued function defined on the set of edges.

(*) Reçu le 7 avril 1997, accepté le 30 septembre 1997

(1) Departamento de Matemática, Universidade Federal do Ceara, CEP 60455-760
BR-Fortaleza CE (Brazil)
e-mail: lev@mat.ufc.br
e-mail: marina@mat.ufc.br

The following question is natural. Let us define a Hölder Complex in a combinatorial way. Does it correspond to some semialgebraic set?

The answer is positive. To prove the Realization theorem we define a semialgebraic set $T(\beta_1, \dots, \beta_k)$. It is a generalization of the real algebraic set which gives an example of the noncoincidence of L_p -cohomology and Intersection Homology [2]. The set $T(\beta_1, \dots, \beta_k)$ has a toric link at the singular point and all generators of the torus have different vanishing rates in this point. It gives us a possibility to separate vanishing rates of all edges of a Hölder Complex.

2. Definitions and notations

Let us recall some definitions from [1]. Let Γ be a connected graph without loops, $V_\Gamma = \{a_1, a_2, \dots, a_k\}$ be the set of vertices and $E_\Gamma = \{g_1, g_2, \dots, g_r\}$ be the set of edges of the graph.

DEFINITION 2.1. — *A Hölder Complex (Γ, β) is a graph Γ with an associated function $\beta: E_\Gamma \rightarrow [1, \infty[\cap Q$ (here Q is the ring of rational numbers).*

DEFINITION 2.2. — *A Curvilinear triangle T is a subset of \mathbb{R}^n homeomorphic to a 2-dimensional simplex satisfying the following properties.*

- 1) *Each internal (in the induced topology) point $t \in T$ has an open neighbourhood $U_t \subset T$ such that U_t is a smooth 2-dimensional submanifold of \mathbb{R}^n at each point $t' \in U_t$.*
- 2) *The boundary of T is a union of three analytic curves $\gamma_1, \gamma_2, \gamma_3$ such that γ_i (for $i = 1, 2, 3$) has a neighbourhood at each internal (in the induced from \mathbb{R} topology on γ_i) point which is a smooth 1-dimensional submanifold of \mathbb{R}^n .*
- 3) *Locally T is a smooth manifold with a boundary at each smooth point of the boundary.*

Boundary points of γ_i we call vertices of T .

DEFINITION 2.3. — *A standard β -Hölder triangle ST_β is a subset of the plane \mathbb{R}^2 bounded by the following curves:*

$$\{y = 0\}, \quad \{y = x^\beta\}, \quad \{x = 1\}.$$

Realization of Hölder Complexes

Let us consider a cone CT over Γ . Let A_0 be the vertex of CT . We can consider CT as a topological space with the standard topology of a simplicial complex.

DEFINITION 2.4. — *A subset $H(\Gamma, \beta) \subset \mathbb{R}^n$ is called a Geometric Hölder Complex corresponding to (Γ, β) if it satisfies the following conditions.*

- 1) $H(\Gamma, \beta)$ is a subanalytic subset of \mathbb{R}^n .
- 2) There exists a homeomorphism $F: CT \rightarrow H(\Gamma, \beta)$.
- 3) The set $H(\Gamma, \beta) \cap S_{F(A_0), r}$ is empty or homeomorphic to Γ , for every r . (We use the notation $S_{F(A_0), r}$ for the sphere centered at the point $F(A_0)$ with the radius r .)
- 4) The image of the triangle (A_0, a_i, a_j, g) (where a_i and a_j are vertices of Γ , g is the edge connecting a_i and a_j , (A_0, a_i, a_j, g) is the subcone of CT over g) has the following properties :
 - (a) $F(A_0, a_i, a_j, g)$ is a subanalytic subset of \mathbb{R}^n ;
 - (b) $F(A_0, a_i, a_j, g)$ is subanalytically bi-Lipschitz equivalent to the standard $\beta(g)$ -Hölder triangle $ST_{\beta(g)}$;
 - (c) let $L: ST_{\beta(g)} \rightarrow F(A_0, a_i, a_j, g)$ be this subanalytic bi-Lipschitz map; then

$$L(0, 0) = F(A_0), \quad L(1, 0) = F(a_i), \quad L(1, 1) = F(a_j).$$

DEFINITION 2.5. — *A β -Hölder triangle HT_β is a subset of \mathbb{R}^n satisfying the following conditions.*

- 1) HT_β is a curvilinear triangle.
- 2) HT_β is bi-Lipschitz equivalent to some standard β -Hölder triangle ST_β .
- 3) The bi-Lipschitz map $L: ST_\beta \rightarrow HT_\beta$ is subanalytic. (The image of the point $(0, 0)$ is called a Hölder vertex of HT_β .)

DEFINITION 2.6. — *A standard β -horn SH_β (here $\beta \in \mathbb{Q} \cap [1, +\infty[)$ is a semialgebraic set in \mathbb{R}^3 defined by the following conditions:*

$$(x_1^2 + x_2^2)^q = y^{2p}, \quad 0 \leq y \leq 1,$$

(x_1, x_2, y) are coordinates of a point in \mathbb{R}^3 and $\beta = p/q$ with $\text{GCD}(p, q) = 1$.

We proved in [1] that every 2-dimensional semialgebraic (as well as semianalytic and subanalytic) set X is a Geometric Hölder Complex in a neighbourhood of a given point $a_0 \in X$ corresponding to some Hölder Complex. Here we are going to prove the following result.

REALIZATION THEOREM. — *Let (Γ, β) be a Hölder Complex. Then there exist a semialgebraic 2-dimensional set $X \subset \mathbb{R}^n$, a point $a_0 \in X$ and $\varepsilon > 0$ such that $X \cap B_{a_0, \varepsilon}$ is a Geometric Hölder Complex corresponding to the Hölder Complex (Γ, β) (here $B_{a_0, \varepsilon}$ is a closed ball in \mathbb{R}^n centered at the point a_0 with the radius ε).*

3. The set $T(\beta_1, \dots, \beta_k)$. Polar maps

We consider the space \mathbb{R}^{2k+1} with coordinates $(x_1, y_1, x_2, y_2, \dots, x_k, y_k, z)$. Let $D(\beta_1, \dots, \beta_k)$ (here $\beta_i = p_i/q_i$ with $p_i, q_i \in \mathbb{Z}$ and $\text{GCD}(p_i, q_i) = 1$) be a subvariety of \mathbb{R}^{2k+1} given by the following equations:

$$\begin{aligned} z^{2p_1} &= (x_1^2 + y_1^2)^{q_1} \\ &\vdots \\ z^{2p_i} &= (x_i^2 + y_i^2)^{q_i} \\ &\vdots \\ z^{2p_k} &= (x_k^2 + y_k^2)^{q_k}. \end{aligned} \tag{1}$$

(The set described in the paper [2] is a special 3-dimensional example of $D(\beta_1, \beta_2)$.)

Let

$$T(\beta_1, \dots, \beta_k) = D(\beta_1, \dots, \beta_k) \cap \{z \geq 0\}. \tag{2}$$

LEMMA 3.1

1) $\dim T(\beta_1, \dots, \beta_k) = k + 1$.

2) *The link of $T(\beta_1, \dots, \beta_k)$ at the point $(0, \dots, 0)$ is homeomorphic to T^k (a k -dimensional torus).*

(Remind that the link of $T(\beta_1, \dots, \beta_k)$ is the intersection of $T(\beta_1, \dots, \beta_k)$ with a small sphere centered at $(0, \dots, 0)$.)

Proof

1) Consider a section of $T(\beta_1, \dots, \beta_k)$ by the plane $z = c$. We obtain the equations

$$x_i^2 + y_i^2 = c_i,$$

where $c_i = c^{2p_i/q_i}$. Clearly, these equations define a k -dimensional torus. The variety $T(\beta_1, \dots, \beta_k)$ we obtain as a suspension of it. So, (1) is proved.

2) Let $r(z)$ be a function defined in the following way:

$$r(z) = \sqrt{z^2 + \sum_{i=1}^k z^{\beta_i}}.$$

This function $r(z)$ is a one-to-one function, for small z . Thus, for sufficiently small $\varepsilon > 0$, the link $T(\beta_1, \dots, \beta_k) \cap S_{0,\varepsilon}$ is equal to the torus $T(\beta_1, \dots, \beta_k) \cap \{(x_1, y_1, \dots, x_k, y_k, z) \in \mathbb{R}^{2k+1} \mid z = r^{-1}(\varepsilon)\}$. \square

Each point of $T(\beta_1, \dots, \beta_k)$ has uniquely defined polar coordinates $(\psi_1, \psi_2, \dots, \psi_k, z)$: ψ_i is the angle coordinate of the corresponding point of the circle $x_i^2 + y_i^2 = c_i$ and z is a z -coordinate in \mathbb{R}^{2k+1} . Let $x^0 = (\psi^0, z^0) = (\psi_1^0, \dots, \psi_k^0, z^0)$ be a point of $T(\beta_1, \dots, \beta_k)$. Let L_{x^0} be a curve on $T(\beta_1, \dots, \beta_k)$ defined as follows:

$$L_{x^0} = \{(\psi_1, \psi_2, \dots, \psi_k, z) \mid \psi_1 = \psi_1^0, \dots, \psi_k = \psi_k^0\}.$$

We call L_{x^0} a *polar line generated by x^0* . Now we can define a polar map in the following way.

Denote, for $\varepsilon > 0$, the set

$$T(\beta_1, \dots, \beta_k) \cap \{(x_1, y_1, \dots, x_k, y_k, z) \in \mathbb{R}^{2k+1} \mid z \leq \varepsilon\}$$

by $T^\varepsilon(\beta_1, \dots, \beta_k)$. Let $P_{\varepsilon_1, \varepsilon_2}: T^{\varepsilon_1}(\beta_1, \dots, \beta_k) \rightarrow T^{\varepsilon_2}(\beta_1, \dots, \beta_k)$ be a map defined as follows:

$$P_{\varepsilon_1, \varepsilon_2}(\psi_1, \dots, \psi_k, z) = \left(\psi_1, \dots, \psi_k, \frac{\varepsilon_1}{\varepsilon_2} z \right).$$

We call $P_{\varepsilon_1, \varepsilon_2}$ a *polar map*. Observe that $P_{\varepsilon_1, \varepsilon_2}$ is a bi-Lipschitz map.

Remark 3.1. — $T(\beta_1)$ is an usual β_1 -horn.

Remark 3.2. — $T(\beta_1, \dots, \beta_k)$ is included to $T(\beta_1, \dots, \beta_k, \dots, \beta_n)$ (here $n \geq k+1$) as a semialgebraic subset defined by the following equations $\psi_{k+1} = b_1, \psi_{k+2} = b_2, \dots, \psi_n = b_{n-k}, b_1, \dots, b_{n-k} \in \mathbb{R}$.

4. Proof of the Realization theorem

We use the induction on the number of edges. Suppose that each Hölder Complex (Γ, β) whose graph Γ has less or equal than k edges is realized as a semialgebraic subset of $T(\beta_1, \dots, \beta_k)$ such that all vertices of Γ belong to the section by the plane $z = 1$ and, for each vertex a , we have $\psi_i(a) = 0$ or $\psi_i(a) = \pi$. (We can identify the graph Γ and its image by the map F ; see Definition 2.4.)

For $k = 1$, the assertion is trivial: Γ has two vertices a_1 and a_2 . Set $\psi(a_1) = 0, \psi(a_2) = \pi$ and the edge connecting a_1 and a_2 be a half-circle. So, (Γ, β) is realized as a half of the standard β -horn.

Now consider a Hölder Complex (Γ, β) such that Γ has $(k + 1)$ edges. Let g be an edge such that $\beta(g) = \min_{\tilde{g} \in E_\Gamma} \beta(\tilde{g})$. Let us consider a graph $\tilde{\Gamma} = \Gamma - g$. We have two possibilities: $\tilde{\Gamma}$ is a connected graph or $\tilde{\Gamma}$ is not connected.

Suppose that $\tilde{\Gamma}$ is not connected. Then it is a union of two connected components $\tilde{\Gamma} = \tilde{\Gamma}^1 \cup \tilde{\Gamma}^2$ (we include also a case when one of these components is just a vertex). We can suppose that $g_1, \dots, g_\ell \in E_{\tilde{\Gamma}^1}, g_{\ell+1}, \dots, g_k \in E_{\tilde{\Gamma}^2}, g_{k+1} = g$. Now consider a set $T(\beta_1, \dots, \beta_k, \beta(g))$ and a section of that by the plane $z = 1$. This section is a $(k + 1)$ -dimensional torus (see the proof of the Lemma 3.1). By the induction hypotheses, the subcomplex $(\tilde{\Gamma}^1, \tilde{\beta}^1)$, where $\tilde{\beta}^1 = \beta|_{\tilde{\Gamma}^1}$, can be realized as a semialgebraic subset of $T(\beta_1, \dots, \beta_k)$ which can be considered as a semialgebraic subset of $T(\beta_1, \dots, \beta_k, \beta(g))$ given by the equation $\psi_{k+1} = 0$ (see the Remark 3.2). By the same way, $(\tilde{\Gamma}^2, \tilde{\beta}^2)$, where $\tilde{\beta}^2 = \beta|_{\tilde{\Gamma}^2}$, can be realized as a semialgebraic subset of $T(\beta_1, \dots, \beta_k)$ which can be considered as a semialgebraic subset of $T(\beta_1, \dots, \beta_k, \beta(g))$ given by the equation $\psi_{k+1} = \pi$. Suppose that g connects vertices $a_1 \in \tilde{\Gamma}^1$ and $a_2 \in \tilde{\Gamma}^2$;

Realization of Hölder Complexes

let a_1 has polar coordinates $(\psi_1(a_1), \dots, \psi_k(a_1), 0)$ and let a_2 has polar coordinates $(\psi_1(a_2), \dots, \psi_k(a_2), \pi)$. We connect these two vertices by the following curve $\Psi(\theta) = \{\psi_1(\theta), \psi_2(\theta), \dots, \psi_{k+1}(\theta), 1\}$ where

$$\psi_{k+1}(\theta) = \theta, \quad \psi_i(\theta) = \begin{cases} \psi_i(a_1) & \text{if } \psi_i(a_1) = \psi_i(a_2) \\ \theta & \text{if } \psi_i(a_1) = 0 \text{ and } \psi_i(a_2) = \pi \\ \pi + \theta & \text{if } \psi_i(a_1) = \pi \text{ and } \psi_i(a_2) = 0, \end{cases} \quad (3)$$

$1 \leq i \leq k, \theta \in [0, \pi]$. Clearly, $\Psi(0) = a_1$ and $\Psi(\pi) = a_2$. Define

$$H_{\beta(g)} := \bigcup_{\theta} L_{\Psi(\theta)},$$

the union of the polar lines generated by $\Psi(\theta)$.

LEMMA 4.1. — *The set $H_{\beta(g)}$ is a $\beta(g)$ -Hölder triangle.*

Proof. — $H_{\beta(g)}$ is a semialgebraic set because it is defined by the system (3) which can be written as a system of algebraic equations and inequalities in terms of variables x_i, y_i , for $1 \leq i \leq k+1$, and by the inequalities $0 \leq z \leq 1$. Hence, $H_{\beta(g)} \cap B_{0,\varepsilon}$ (here $B_{0,\varepsilon}$ is a closed ball in \mathbb{R}^{2k+3} centered at 0 with the radius ε) is a Geometric Hölder Complex $H(\bar{\Gamma}, \alpha)$ corresponding to some graph $\bar{\Gamma}$ with some rational-valued function α defined on its edges [1]. Since $H_{\beta(g)}$ is a curvilinear triangle (by the construction), $H_{\beta(g)} \cap B_{0,\varepsilon_0}$, for sufficiently small $\varepsilon_0 \leq \varepsilon$, is bi-Lipschitz equivalent to the standard α_0 -Hölder triangle where $\alpha_0 = \min_{\bar{g} \in E_{\bar{\Gamma}}} \alpha(\bar{g})$ [1, Second Structural Lemma]. But $H_{\beta(g)} \cap B_{0,\varepsilon_0}$ is bi-Lipschitz equivalent to $H_{\beta(g)}$ (the bi-Lipschitz equivalence is given by the polar map $P_{\varepsilon_0,1}$).

To complete the proof of the lemma we must show that $\alpha_0 = \beta(g)$. Let γ_ε be the equidistant line in $H_{\beta(g)}$, namely $\gamma_\varepsilon = H_{\beta(g)} \cap S_{0,\varepsilon}$. By [1], there exists a subanalytic bi-Lipschitz map $\Upsilon: H_{\beta(g)} \rightarrow \text{ST}_{\alpha_0}$ such that $\Upsilon(\gamma_\varepsilon) = \text{ST}_{\alpha_0} \cap \{(x, y) \in \mathbb{R}^2 \mid x = \varepsilon\}$. Denote by $\ell(\gamma_\varepsilon)$ the length of γ_ε . Since Υ is a bi-Lipschitz map, we have

$$c_1 \varepsilon^{\alpha_0} \leq \ell(\gamma_\varepsilon) \leq c_2 \varepsilon^{\alpha_0}, \quad (4)$$

for some positive constants c_1 and c_2 . To prove that $\alpha_0 = \beta(g)$ we will compute the length of γ_ε from another side. Consider the function

$$r(z) = \sqrt{z^2 + \sum_{i=1}^{k+1} z^{p_i/q_i}}$$

which is a one-to-one function, for small z . So, $r^{-1}(\varepsilon)$ is a well-defined function, for small ε . By the Lemma 3.1,

$$\gamma_\varepsilon = H_{\beta(g)} \cap \{(x_1, y_1, \dots, x_{k+1}, y_{k+1}, z) \in \mathbb{R}^{2k+3} \mid z = r^{-1}(\varepsilon)\}.$$

Consider the following set

$$\begin{aligned} T^\varepsilon &= T(\beta_1, \dots, \beta_k, \beta(g)) \\ &\cap \{(x_1, y_1, \dots, x_{k+1}, y_{k+1}, z) \in \mathbb{R}^{2k+1} \mid z = r^{-1}(\varepsilon)\}. \end{aligned}$$

It is a smooth manifold homeomorphic to a $(k+1)$ -dimensional torus. The equidistant line γ_ε belongs to this set. There are $(k+1)$ differential 1-forms $d\psi_1^\varepsilon, \dots, d\psi_k^\varepsilon$ and $d\psi_{k+1}^\varepsilon$ on T^ε corresponding to the coordinate system $\{\psi_1, \dots, \psi_k, \psi_{k+1}\}$. By (3), we have

$$\ell(\gamma_\varepsilon) = \int_{\gamma_\varepsilon} \sum_{i=1}^{k+1} m_i d\psi_i^\varepsilon \quad \text{where } m_i = \begin{cases} 1 & \text{if } \psi_i(a_1) \neq \psi_i(a_2) \\ 0 & \text{if } \psi_i(a_1) = \psi_i(a_2), \end{cases}$$

$$\int_{\gamma_\varepsilon} \sum_{i=1}^{k+1} m_i d\psi_i^\varepsilon \leq \sum_{i=1}^{k+1} \int_{\gamma_\varepsilon} m_i d\psi_i^\varepsilon.$$

By the definition of the equidistant line γ_ε ,

$$\int_{\gamma_\varepsilon} m_i d\psi_i^\varepsilon = m_i \pi z^{\beta_i}.$$

Using the above formula we obtain

$$\ell(\gamma_\varepsilon) \leq \sum_{i=1}^{k+1} m_i \pi z^{\beta_i}.$$

If z sufficiently small ($z < 1$) there exists $\tilde{C}_2 > 0$ such that

$$\ell(\gamma_\varepsilon) \leq \sum_{i=1}^{k+1} m_i \pi z^{\beta_i} \leq \tilde{C}_2 z^{\beta(g)},$$

because $\beta(g) = \min_{1 \leq i \leq k+1} \beta_i$.

By the definition of the function $r(\varepsilon)$, we have $r(\varepsilon) = a\varepsilon + o(\varepsilon)$, with $a > 0$.

Realization of Hölder Complexes

Hence, $\ell(\gamma_\varepsilon) \leq C'_2 \varepsilon^{\beta(g)}$, where $C'_2 = a\tilde{C}_2$. To obtain an estimate of $\ell(\gamma_\varepsilon)$ from below let us go back to the formulas (3)

$$\ell(\gamma_\varepsilon) = \int_{\gamma_\varepsilon} \sum_{i=1}^{k+1} m_i d\psi_i^\varepsilon \geq \int_{\gamma_\varepsilon} m_{k+1} d\psi_{k+1}^\varepsilon.$$

By (3), $m_{k+1} = 1$. Thus,

$$\ell(\gamma_\varepsilon) \geq \int_{\gamma_\varepsilon} d\psi_{k+1}^\varepsilon = \pi z^{\beta(g)} \geq C'_1 \varepsilon^{\beta(g)},$$

for some positive constant C'_1 . So,

$$C'_1 \varepsilon^{\beta(g)} \leq \ell(\gamma_\varepsilon) \leq C'_2 \varepsilon^{\beta(g)}. \quad (5)$$

From (4) and (5) we obtain that $\beta(g) = \alpha_0$.

Lemma 4.1 is proved. \square

Thus, the realization of (Γ, β) is given by the union of the realizations of $(\tilde{\Gamma}^1, \tilde{\beta}^1)$, $(\tilde{\Gamma}^2, \tilde{\beta}^2)$ and $H_{\beta(g)}$. It is a semialgebraic set because it is a finite union of semialgebraic sets.

Now consider the second case: $\tilde{\Gamma}$ is a connected graph. In this case, by the induction hypotheses, $(\tilde{\Gamma}, \tilde{\beta})$ (where $\tilde{\beta} = \beta|_{\tilde{\Gamma}}$) can be realized as a semialgebraic subset of $T(\beta_1, \dots, \beta_k)$ which can be considered as a semialgebraic subset of $T(\beta_1, \dots, \beta_k, \beta(g))$ defined by the equation $\psi_{k+1} = 0$. The edge g connects two vertices a_1 and a_2 . Now we can glue the realization of $(\tilde{\Gamma}, \tilde{\beta})$ and the curvilinear triangle $H_{\beta(g)}$ generated by the curve $\Psi(\theta) = \{\psi_1(\theta), \psi_2(\theta), \dots, \psi_{k+1}(\theta)\}$:

$$\psi_{k+1}(\theta) = \theta \text{ and } \psi_i(\theta) = \begin{cases} \psi_i(a_1) & \text{if } \psi_i(a_1) = \psi_i(a_2) \\ \frac{\theta}{2} & \text{if } \psi_i(a_1) = 0 \text{ and } \psi_i(a_2) = \pi \\ \pi + \frac{\theta}{2} & \text{if } \psi_i(a_1) = \pi \text{ and } \psi_i(a_2) = 0, \end{cases} \quad (6)$$

for $1 \leq i \leq k$, $\theta \in [0, 2\pi]$, $a_1 = (\psi_1(a_1), \dots, \psi_k(a_1), 0)$ and $a_2 = (\psi_1(a_2), \dots, \psi_k(a_2), \pi)$.

Set $H_{\beta(g)} := \bigcup_{\theta} L_{\Psi(\theta)}$. By the same arguments as in the Lemma 4.1, we can prove that $H_{\beta(g)}$ is a $\beta(g)$ -Hölder triangle.

The union of the realization of $(\tilde{\Gamma}, \tilde{\beta})$ and $H_{\beta(g)}$ is a semialgebraic realization of (Γ, β) .

The Realization theorem is proved. \square

Acknowledgments

The authors were supported by CNPq grants N 300985/93-2(RN) and N 142385/95-6. We are grateful to IMPA and to Mathematical Institute of PUC-Rio, where this work was done.

References

- [1] BIRBRAIR (L.) .— *Local bi-Lipschitz classification of 2-dimensional semialgebraic sets*, Preprint I.M.P.A. (1996).
- [2] BIRBRAIR (L.) and GOLDSHTEIN (V.) .— *An Example of Noncoincidence of L_p -cohomology and Intersection Homology for Real Algebraic varieties*. I.M.R.N. 6 (1994), pp. 265-271.