

ANNALES DE L'INSTITUT FOURIER

B. L. GUPTA

**On the Hausdorff summability of series associated
with a Fourier and its allied series**

Annales de l'institut Fourier, tome 21, n° 3 (1971), p. 173-179

http://www.numdam.org/item?id=AIF_1971__21_3_173_0

© Annales de l'institut Fourier, 1971, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE HAUSDORFF SUMMABILITY OF SERIES ASSOCIATED WITH A FOURRIER AND ITS ALLIED SERIES

by **B. L. GUPTA**

1. Let S_n be the n th partial sum of an infinite series $\sum_1^{\infty} a_n$ and let

$$t_n = \sum_{\nu=0}^n \binom{n}{\nu} (\Delta^{n-\nu} \mu_{\nu}) S_{\nu} . \quad (1.1)$$

Then the sequence $\{t_n\}$ is known as the Hausdorff means of sequence $\{S_n\}$, where $\{\mu_{\nu}\}$ is a sequence of real or complex numbers and the sequence $\{\Delta^p \mu_{\nu}\}$ denotes the differences of order p .

The series $\sum_1^{\infty} a_n$ is said to be summable by Hausdorff mean to the sum S , if $\lim t_n \rightarrow S$, whenever $S_n \rightarrow S$. The necessary and sufficient condition for the Hausdorff summability to be conservative is that the sequence $\{\mu_n\}$ should be a sequence of moment constant, i.e. ;

$$\mu_n = \int_0^1 x^n d\chi(x), \quad n \geq 0 ;$$

where $\chi(x)$ is a real function of bounded variation in $0 \leq x \leq 1$. We may suppose without loss of generality that $\chi(0) = 0$, if also $\chi(1) = 1$ and $\chi(+0) = \chi(0) = 0$, so that $\chi(x)$ is continuous at the origin, then μ_n is a regular moment constant and the Hausdorff method i.e. (H, μ_n) is a regular method of summation [2].

If

$$\sum_{n=0}^{\infty} |(t_n - t_{n-1})| < \infty , \quad (1.2)$$

then the series $\sum_1^{\infty} a_n$ is said to be absolutely summable (H, μ_n) or

summable $|H, \mu_n|$. It is also known that the Cesàro, Holder and Euler methods of summation are the particular cases of the above method.

2. Let $f(t)$ be a periodic function with period 2π and integrable in the sense of Lebesgue in $(-\pi, \pi)$. Let its Fourier series be

$$\frac{1}{2} a_0 + \sum_1^{\infty} (a_n \cos nt + b_n \sin nt) = \sum_{n=0}^{\infty} A_n(t)$$

and its allied series is

$$\sum_1^{\infty} (b_n \cos nt - a_n \sin nt) = \sum_1^{\infty} B_n(t) .$$

We write

$$\varphi(t) = \frac{1}{2} \{f(\theta + t) + f(\theta - t)\} ,$$

$$\psi(t) = \frac{1}{2} \{f(\theta + t) - f(\theta - t)\} .$$

Let $g(x)$ be integrable L in $(0, 1)$, then for $\varepsilon > 0$

$$g_{\varepsilon}^{+}(x) = \frac{1}{\Gamma(\varepsilon)} \int_0^x (x-u)^{\varepsilon-1} g(u) du ,$$

$$g_{\varepsilon}^{-}(x) = \frac{1}{\Gamma(\varepsilon)} \int_x^1 (u-x)^{\varepsilon-1} g(u) du .$$

Again, let

$$U_n(t) = \sum_{\nu=1}^n e^{i\nu t} ,$$

$$H(n, x, t) = E(n, x, t) + iF(n, x, t)$$

$$= \sum_{\nu=0}^n \nu^{\beta} \binom{n}{\nu} x^{\nu} (1-x)^{n-\nu} e^{i\nu t} .$$

The object of this paper is to prove the following :

THEOREM 1. — *If*

i) $\int_0^t |\varphi(u)| du = O(t)$

ii) (H, μ_n) is conservative

and

iii) $\left\{ \begin{array}{l} \text{either (a) } \chi(x) = g_{1+\beta+\varepsilon}^-(x) + c, \varepsilon > 0 ; \\ \text{or (b) } \chi(x) = g_{1+\beta+\varepsilon}^+(x) + c, \varepsilon > 0 ; \end{array} \right.$
 for some $g(x) \in L(0, 1)$;

then the series $\sum_{n=1}^{\infty} \frac{A_n(t)}{n^{1-\beta}}$, for $|\beta| \geq 0$ is summable (H, μ_n) at $t = \theta$, where c is an absolute constant.

THEOREM 2. — *If*

i) $\int_0^t |\psi(u)| du = O(t)$

ii) (H, μ_n) is conservative

and

iii) $\left\{ \begin{array}{l} \text{either (a) } \chi(x) = g_{1+\beta+\varepsilon}^-(x) + c, \varepsilon > 0 ; \\ \text{or (b) } \chi(x) = g_{1+\beta+\varepsilon}^+(x) + c, \varepsilon > 0 ; \end{array} \right.$
 for some $g(x) \in L(0, 1)$,

then the series $\sum_{n=1}^{\infty} \frac{B_n(t)}{n^{1-\beta}}$, for $|\beta| \geq 0$ is summable (H, μ_n) at $t = \theta$, where c is an absolute constant.

It may also be remarked if

$$\chi(x) = 1 - (1 - x)^\delta, \quad \delta > 0 ;$$

the method (H, μ_n) reduces to the well known Cesàro method of summation of order δ .

Further if we choose β such that $\delta > \beta + \varepsilon$ then it can be proved that $\chi(x) - 1$ is the $(1 + \beta + \varepsilon)$ th backward integral of

$$- \frac{\Gamma(1 + \delta)}{\Gamma(\delta - \beta - \varepsilon)} (1 - x)^{\delta - \beta - \varepsilon - 1}$$

and $\chi(x)$ is also the $(\varepsilon + \beta + 1)$ th forward integral of

$$\frac{\delta}{\Gamma(1 - \beta - \varepsilon)} \left\{ x^{-(\beta + \varepsilon)} + (1 - \beta) \int_0^x (1 - \nu)^{\delta - 2} (x - \nu)^{-(\beta + \varepsilon)} d\nu \right\}.$$

Hence the method $|C, \delta|$ satisfies the hypothesis of our theorem 1 and 2 for $\varepsilon > 0$, $\delta > \beta \geq 0$ and the following theorems of Cheng [1] becomes the corollary of our theorems.

THEOREM. — *The series $\sum \frac{A_n(t)}{n^{1-\beta}}$ for $0 \leq \beta < 1$ is summable $|C, \delta|$ for $\delta > \beta$, at the point θ , whenever i) of theorem I holds and similarly the series $\sum_1^\infty \frac{B_n(t)}{n^{1-\beta}}$, for $0 \leq \beta < 1$, is summable $|C, \delta|$, for $\delta > \beta$, at the point θ , whenever i) of theorem 2 holds.*

3. For the proof of the theorems, we require the following lemmas.

LEMMA 1. — *Uniformly in $0 < t \leq \pi$*

$$|U_n(t)| \leq \frac{k}{t}. \quad (3.1)$$

This can be easily proved.

LEMMA 2. — *If $g(x)$ and $h(x)$ be Lebesgue integrable in $(0, 1)$, then for $\varepsilon > 0$*

$$\int_0^1 g_\varepsilon^+(x) h(x) dx = \int_0^1 g(x) h_\varepsilon^-(x) dx. \quad (3.2)$$

This is known [3].

LEMMA 3. — *Uniformly in $0 \leq x \leq 1$*

$$\int_0^x H(n, \nu, t) d\nu = O\left(\frac{n^{\beta-1}}{t}\right) \quad (3.3)$$

LEMMA 4. — Let $\beta \geq 0$ $\varepsilon > 0$ and fixed, then for $\beta + \varepsilon < 1$

$$\int_0^x (x - u)^{\beta + \varepsilon - 1} H(n, u, t) du = O\left(\frac{n^{-\varepsilon}}{t^{\beta + \varepsilon}}\right) \quad (3.4)$$

uniformly in $0 \leq x \leq 1$ and similarly

$$\int_x^1 (u - x)^{\beta + \varepsilon - 1} \times H(n, u, t) du = O\left(\frac{n^{-\varepsilon}}{t^{\beta + \varepsilon}}\right) . \quad (3.5)$$

The lemma 3 and 4 are due to Tripathy [4].

Proof of Theorem 1. — If t_n and u_n denote the Hausdorff means of $\sum \frac{A_n(\theta)}{n^{1-\beta}}$ and the sequence $\{n A_n(\theta)\}$ then for $n \geq 1$

$$u_n = n(t_n - t_{n-1}) .$$

Hence, from (1.2) the series $\sum_{n=1}^{\infty} \frac{A_n(\theta)}{n^{1-\beta}}$ is summable $|H, \mu_n|$, if

$$I = \sum_{n=1}^{\infty} \frac{1}{n} \left| \sum_{\nu=1}^n \binom{n}{\nu} (\Delta^{n-\nu} \mu_\nu) \nu^\beta A_\nu(\theta) \right| < \infty .$$

Since (H, μ_n) is conservative, we have

$$\begin{aligned} I &= \sum_{n=1}^{\infty} \frac{1}{n} \left| \int_0^1 d\chi(x) \sum_{\nu=1}^n \binom{n}{\nu} x^\nu (1-x)^{n-\nu} \nu^\beta A_\nu(\theta) \right| \\ &= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \left| \int_0^1 d\chi(x) \sum_{\nu=1}^n \binom{n}{\nu} x^\nu (1-x)^{n-\nu} \nu^\beta \int_0^\pi \varphi(t) \cos \nu t dt \right| \\ &\leq \sum_{n=1}^{\infty} \frac{1}{n} \int_0^{\frac{1}{n}} |\varphi(t)| \left| \int_0^1 d\chi(x) \sum_{\nu=1}^n \binom{n}{\nu} x^\nu (1-x)^{n-\nu} \nu^\beta \cos \nu t \right| dt \\ &+ \sum_{n=1}^{\infty} \frac{1}{n} \int_{\frac{1}{n}}^\pi |\varphi(t)| \left| \int_0^1 d\chi(x) \sum_{\nu=1}^n \binom{n}{\nu} x^\nu (1-x)^{n-\nu} \nu^\beta \cos \nu t \right| dt \\ &= I_1 + I_2, \text{ say.} \end{aligned}$$

Since

$$\begin{aligned} |H(n, x, t)| &\leq n^\beta \left| \sum_{\nu=0}^n \binom{n}{\nu} x^\nu (1-x)^{n-\nu} \right| \\ &= n^\beta \end{aligned}$$

We have

$$\begin{aligned} I_1 &= O(1) \sum_{n=1}^{\infty} \frac{1}{n} n^{\beta} \int_0^{\frac{1}{n}} |\varphi(t)| dt \int_0^1 |d\chi(x)| \\ &= O(1) \sum_{n=1}^{\infty} \frac{1}{n^{1-\beta}} \cdot \frac{1}{n} \\ &= O(1) . \end{aligned}$$

Without loss of generality, we can suppose that $\beta + \varepsilon < 1$, if
a) $\chi(x) = g_{1+\beta+\varepsilon}^-(x) + c$, then

$$\begin{aligned} I_2 &= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \int_{\frac{1}{n}}^{\pi} |\varphi(t)| \cdot \left| \int_0^1 g_{\varepsilon+\beta}^-(x) E(n, x, t) dx \right| dt \\ &= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{n} \int_{\frac{1}{n}}^{\pi} |\varphi(t)| \cdot \left| \int_0^1 g(x) E_{\beta+\varepsilon}^+(n, x, t) dx \right| dt . \end{aligned}$$

Since

$$\begin{aligned} E_{\beta+\varepsilon}^+(n, x, t) &= \frac{1}{\Gamma(\beta + \varepsilon)} \int_0^x (x - u)^{\beta + \varepsilon - 1} E(n, u, t) du \\ &= \frac{1}{\Gamma(\beta + \varepsilon)} \int_0^x (x - u)^{\beta + \varepsilon - 1} I_m H(n, u, t) du \\ &= O\left(\frac{1}{n^{\varepsilon} t^{\beta + \varepsilon}}\right) , \text{ by lemma-4.} \end{aligned}$$

Therefore

$$\begin{aligned} I_2 &\leq \int_0^1 |g(x)| dx \sum_{n=1}^{\infty} \frac{1}{n} \int_{\frac{1}{n}}^{\pi} |\varphi(t)| \cdot O\left(\frac{1}{n^{\varepsilon} t^{\beta + \varepsilon}}\right) dt \\ &= \int_0^1 |g(x)| dx \sum_{n=1}^{\infty} \frac{1}{n^{1+\varepsilon}} \int_{\frac{1}{n}}^{\pi} \frac{|\varphi(t)|}{t^{\beta + \varepsilon}} dt \\ &= \int_0^1 |g(x)| dx \sum_{n=1}^{\infty} \frac{1}{n^{1+\varepsilon}} \{O(1) + O(n^{\beta + \varepsilon - 1})\} \\ &= O(1) \int_0^1 |g(x)| dx \\ &= O(1) . \end{aligned}$$

If b) $\chi(x) = g_{1+\beta+\varepsilon}^+(x) + c$, then proceeding in a similar way as in case a) and using estimate (4.5) of lemma 4, it can be proved that

$$I_2 = O(1) .$$

This completes the proof of theorem-1.

If we use the condition i) of Theorem-2 instead of the condition i) of theorem-1, we can prove that the series $\sum \frac{B_n(\theta)}{n^{1-\beta}}$ is summable $|H, \mu_n|$.

In the last, the author would like to express his deep sense of gratitude to Dr. P.L. Sharma for his encouragement in the preparation of this paper.

BIBLIOGRAPHY

- [1] M.T. CHENG, Summability factors of Fourier series, *Duke Math. Jour.* 15 (1948), 17-27.
- [2] G.H. HARDY, Divergent series. Oxford 1949.
- [3] B. KUTTNER, On the "second theorem of consistency" for riesz summability (II), *Jour. Lond. Math. Soc.* 27 (1952), 207-217.
- [4] N. TRIPATHY, On the absolute Hausdorff summability of some series associated with Fourier series and its allied series, *Jour. Ind., Math. Soc.* 32 (1960) 141-154.

Manuscrit reçu le 3 août 1970

B.L. GUPTA

Department of Mathematics,
Govt. Engineering College,
Rewa, M.P. India.