

ANNALES DE L'INSTITUT FOURIER

HARUO KITAHARA
SHINSUKE YOROZU

On the Cech bicomplex associated with foliated structures

Annales de l'institut Fourier, tome 28, n° 3 (1978), p. 217-224

http://www.numdam.org/item?id=AIF_1978__28_3_217_0

© Annales de l'institut Fourier, 1978, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE ČECH BICOMPLEX ASSOCIATED WITH FOLIATED STRUCTURES

by H. KITAHARA and S. YOROZU

1. We shall be in C^∞ -category. Let M be a paracompact connected n -dimensional manifold with a foliation \mathcal{F} of codimension q , and let $\mathcal{U} = \{U_\alpha\}$ be a simple covering of M such that each U_α is a flat neighborhood with respect to \mathcal{F} . Then there exists a decomposable q -form $w = w^1 \wedge \dots \wedge w^q$ on each U_α and, by Frobenius' theorem, there exists a 1-form η on each U_α satisfying $dw = w \wedge \eta$, where d denotes the exterior differentiation and \wedge the exterior product. The 1-form η is an interesting object; it is well known that $\eta \wedge (d\eta)^q$ defines a de Rham class in $H^{2q+1}(M, \mathbb{R})$ ([1], [2], [3]). Our aim is to show that η itself defines a certain cohomology class, that is,

THEOREM A. — $((-1)^{q-1}/2\pi)\eta$ defines a D -cohomology class in $H^2(\check{C}^{(*)}(\mathcal{U}; A^*(M)), D)$ depending only on \mathcal{F} .

The above theorem was announced in [4], where it contained misstatements.

THEOREM B. — Supposing M admits foliations $\mathcal{F}, \mathcal{F}'$ complementally transversal to each other, η defines a D' -cohomology class in $H^1(\check{C}^{(*)}(\mathcal{U}; A^*(M)), D')$ (Cf. [5], [6], [7]).

2. Since M has a foliation \mathcal{F} of codimension q , the tangent bundle TM of M has an integrable subbundle E with fibre dimension $n-q$. Let $Q = TM/E$ be a quotient bundle with fibre dimension q . Choosing a suitable Riemannian metric on TM , we obtain an isomorphism $TM \cong E \oplus Q$ (Whitney sum).

Then

$$\begin{aligned} d\bar{w}^i &= \sum_j dt_j^i \wedge w^j + \sum_j t_j^i dw^j \\ &= \sum_j \left(dt_j^i - \sum_k t_k^i \varphi_j^k \right) \wedge w^j, \end{aligned}$$

and on the other hand

$$\begin{aligned} d\bar{w}^i &= \sum_k \bar{w}^k \wedge \bar{\varphi}_k^i \\ &= \sum_j \left(- \sum_k t_j^k \bar{\varphi}_k^i \right) \wedge w^j. \end{aligned}$$

Thus

$$- \sum_k t_j^k \bar{\varphi}_k^i = dt_j^i - \sum_k t_k^i \varphi_j^k + \sum_k f_{jk}^i w^k$$

where f_{jk}^i are functions on $U_{\alpha_0} \cap U_{\alpha_1}$.

Let $\begin{pmatrix} s_j^i & 0 \\ 0 & s_b^a \end{pmatrix}$ denote the inverse matrix of $\begin{pmatrix} t_j^i & 0 \\ 0 & t_b^a \end{pmatrix}$.

Then

$$\sum_{i,k} s_l^i t_j^k \bar{\varphi}_k^i = - \sum_j s_l^j dt_j^i + \sum_{j,k} s_l^j t_k^i \varphi_j^k - \sum_{i,k} s_l^j f_{jk}^i w^k$$

and we obtain

$$\sum_i \bar{\varphi}_i^i = - \sum_{i,j} s_l^j dt_j^i + \sum_i \varphi_i^i - \sum_{i,j,k} s_l^j f_{jk}^i w^k.$$

From (4), $dt_j^i = \sum_k t_{jk}^i w^k$. Thus, by (3), we obtain $\sum_i \bar{\varphi}_i^i = \sum_i \varphi_i^i$.

Therefore we obtain $\bar{\eta} = \eta$ on $U_{\alpha_0} \cap U_{\alpha_1}$ and $\eta \in \check{C}^{0,1}(\mathcal{U}; A^*(M))$.
Q.E.D.

THEOREM B. — *Supposing M admits foliations $\mathcal{F}, \mathcal{F}'$ complementally transversal to each other, η defines a D' -cohomology class in $H^1(\check{C}^*(\mathcal{U}; A^*(M)), D')$ where η is defined by \mathcal{F} .*

We may suppose that $\mathcal{U} = \{U_\alpha\}$ is a simple covering such that each U_α is a locally trivial neighborhood of the bundle $Q \rightarrow M$.

Let ∇^α denote a local connection on $Q|_{U_\alpha}$, and let ω_α (resp. Ω_α) denote a connection form (resp. a curvature form) of ∇^α on U_α . Let Δ^p be a canonical p -simplex in R^{p+1} (with coordinates (t_0, t_1, \dots, t_p)). We define a connection form $\omega_{\alpha_0 \dots \alpha_p}$ on $U_{\alpha_0 \dots \alpha_p} \times \Delta^p$ ($U_{\alpha_0 \dots \alpha_p} = U_{\alpha_0} \cap U_{\alpha_1} \cap \dots \cap U_{\alpha_p}$) by

$$\begin{aligned} \omega_{\alpha_0 \dots \alpha_p} &= t_0 \omega_{\alpha_0} + \dots + t_p \omega_{\alpha_p} \\ &= (1 - t_1 - \dots - t_p) \omega_{\alpha_0} + t_1 \omega_{\alpha_1} + \dots + t_p \omega_{\alpha_p} \end{aligned}$$

and let $\Omega_{\alpha_0 \dots \alpha_p}$ denote a corresponding curvature form on $U_{\alpha_0 \dots \alpha_p} \times \Delta^p$.

For the set $A^k(U_{\alpha_0 \dots \alpha_p} \times \Delta^p)$ of all k -forms on $U_{\alpha_0 \dots \alpha_p} \times \Delta^p$, $\int_{\Delta^p} : A^k(U_{\alpha_0 \dots \alpha_p} \times \Delta^p) \rightarrow A^{k-p}(U_{\alpha_0 \dots \alpha_p})$ denotes the integration along the fibre. Then we obtain Stokes' theorem

$$\int_{\Delta^p} d \circ = (-1)^p d \circ \int_{\Delta^p} + \int_{\partial \Delta^p} \circ j^*$$

where $j : U_{\alpha_0 \dots \alpha_p} \times \partial \Delta^p \rightarrow U_{\alpha_0 \dots \alpha_p} \times \Delta^p$ denotes the inclusion.

We consider Čech bicomplex $\check{C}^{(*)}(\mathcal{U}; A^*(M))$: Let $\check{C}^{p,q} = \prod_{\alpha_0 \dots \alpha_p} A^q(U_{\alpha_0 \dots \alpha_p})$, and let $D' : \check{C}^{p,q} \rightarrow \check{C}^{p+1,q}$ denote the ordinary simplicial differential and $D'' = (-1)^p d : \check{C}^{p,q} \rightarrow \check{C}^{p,q+1}$ the de Rham differential. A multiplication: $\check{C}^{p,q} \otimes \check{C}^{p',q'} \rightarrow \check{C}^{p+p',q+q'}$ is defined by

$$(\Phi \cdot \Phi')_{\alpha_0 \dots \alpha_{p+q}} = (-1)^{qp'} \Phi_{\alpha_0 \dots \alpha_p} |_{U_{\alpha_0 \dots \alpha_{p+q}}} \wedge \Phi'_{\alpha_{p+1} \dots \alpha_{p+q}} |_{U_{\alpha_0 \dots \alpha_{p+q}}}$$

For $\check{C}^{(k)}(\mathcal{U}; A^*(M)) = \sum_{p+q=k} \check{C}^{p,q}$ and $D = D' + D'' : \check{C}^{(k)} \rightarrow \check{C}^{(k+1)}$, we obtain a graded algebra $(\check{C}^{(*)}(\mathcal{U}; A^*(M)), D, \cdot)$.

Let $I^*(\mathfrak{g}I_q)$ denote a graded algebra of invariant polynomials on a Lie algebra $\mathfrak{g}I_q$. A characteristic homomorphism

$$\gamma : I^*(\mathfrak{g}I_q) \rightarrow C^{(*)}(\mathcal{U}; A^*(M))$$

is defined by

$$\gamma \varphi = \sum_p (\gamma \varphi)^{p, 2k-p} \quad \varphi \in I^k(\mathfrak{g}I_q)$$

where

$$(\gamma\varphi)_{\alpha_0 \dots \alpha_p}^{p, 2k-p} = \int_{\Delta^p} \varphi(\Omega_{\alpha_0 \dots \alpha_p}, \dots, \Omega_{\alpha_0 \dots \alpha_p}).$$

Then we obtain,

LEMMA 1 (Cf. [8]). — For $\varphi \in \Gamma^k(\mathfrak{gl}_q)$,

$$(-1)^{[p(p-1)]/2} \frac{(k-p)!}{k!} (\gamma\varphi)_{\alpha_0 \dots \alpha_p}^{p, 2k-p} = \begin{cases} \int_{\Delta^p} dt_1 \wedge \dots \wedge dt_p \wedge \varphi(\omega_{\alpha_1} - \omega_{\alpha_0}, \omega_{\alpha_2} - \omega_{\alpha_0}, \dots, \omega_{\alpha_p} - \omega_{\alpha_0}, \Gamma_{\alpha_0 \dots \alpha_p}^{k-p}) & p \leq k \\ 0 & p > k \end{cases}$$

where $\Gamma_{\alpha_0 \dots \alpha_p} = \Omega_{\alpha_0 \dots \alpha_p} - \sum_i dt_i \wedge (\omega_{\alpha_i} - \omega_{\alpha_0})$.

Remark. — γ induces the Chern-Weil homomorphism

$$\gamma^*: I^*(\mathfrak{gl}_q) \longrightarrow H^*(\check{C}^*(\cdot), D) \xleftarrow{\cong} H^*(M).$$

The following lemma is easily proved.

LEMMA 2. — Let w^1, \dots, w^q be 1-forms on U_α such that $w^1 \wedge \dots \wedge w^q \neq 0$ on U_α . Put $w = w^1 \wedge \dots \wedge w^q$. Then (i) and (ii) are equivalent:

(i) There exists a (q, q) -matrix (φ_j^i) of 1-forms on U_α such that $dw^i = \sum_j \varphi_j^i \wedge w^j$.

(ii) There exists a 1-form η on U_α such that $dw = w \wedge \eta$.

Remark. — The existence of the matrix (φ_j^i) doesn't depend on choice of q 1-forms w^1, \dots, w^q on U_α .

Remark. — In the proof of this lemma, we obtain

$$\eta = (-1)^{q-1} \sum_i \varphi_i^i. \tag{1}$$

Let $\Gamma(\cdot)$ denote the space of all sections of bundle. The Bott connection $\tilde{\nabla}: \Gamma(E) \times \Gamma(Q) \longrightarrow \Gamma(Q)$ is defined by

$$\tilde{\nabla}_X Z = \pi_*([X, \tilde{Z}]) \quad X \in \Gamma(E), Z \in \Gamma(Q)$$

where $\tilde{Z} \in \Gamma(TM)$ such that $\pi_*(\tilde{Z}) = Z$ and $\pi: TM \longrightarrow Q$. Let

$\{e_i, e_a\}$ ($1 \leq i \leq q, q + 1 \leq a \leq n$) be a local basis dual to $\{w^i, w^a\}$ on U_α satisfying $e_i \in \Gamma(Q|_{U_\alpha})$ and $e_a \in \Gamma(E|_{U_\alpha})$ with respect to the isomorphism $TM \cong E \oplus Q$. Hereafter, we suppose that the indices run the following ranges : $1 \leq i, j, k, \dots \leq q, q + 1 \leq a, b, \dots \leq n$. We define a connection ∇^α on U_α by

$$\nabla_X^\alpha Z = \tilde{\nabla}_{X_E} Z + \sum_i X_Q(Z^i) e_i + \sum_{i,k} Z^i \varphi_i^k(X_Q) e_k \tag{2}$$

where $X = X_E + X_Q \in \Gamma(E|_{U_\alpha}) \oplus \Gamma(Q|_{U_\alpha})$ and $Z = \sum_i Z^i e_i \in \Gamma(Q|_{U_\alpha})$.

We put $\nabla_X^\alpha e_j = \sum_i \omega_{\alpha j}^i(X) e_i$, that is, $\omega_{\alpha j}^i$ denotes the connection form of ∇^α on U_α .

LEMMA 3. — $\omega_{\alpha j}^i = \varphi_j^i$ on U_α .

Proof. — We put $\tilde{\nabla}_{X_E} e_j = \sum_i \tilde{\omega}_j^i(X_E) e_i$, then

$\omega_{\alpha j}^i(X) = \tilde{\omega}_j^i(X_E) + \varphi_j^i(X_Q)$. Now we obtain

$$\begin{aligned} dw^i(e_a, e_j) &= \frac{1}{2} \{e_a(w^i(e_j)) - e_j(w^i(e_a)) - w^i([e_a, e_j])\} \\ &= -\frac{1}{2} \tilde{\omega}_j^i(e_a). \end{aligned}$$

On the other hand,

$$\begin{aligned} dw^i(e_a, e_j) &= \left(\sum_k w^k \wedge \varphi_k^i \right) (e_a, e_j) \\ &= -\frac{1}{2} \varphi_j^i(e_a). \end{aligned}$$

Thus we obtain $\tilde{\omega}_j^i(X_E) = \varphi_j^i(X_E)$. Therefore, for any $X \in \Gamma(TM|_{U_\alpha})$, $\omega_{\alpha j}^i(X) = \varphi_j^i(X_E) + \varphi_j^i(X_Q) = \varphi_j^i(X)$. Q.E.D.

LEMMA 4. — *If we consider the connection ∇^{α_0} defined by (2) on U_{α_0} and a Riemannian connection ∇^{α_1} on U_{α_1} , then, for $\varphi_1 \in I^1(\mathfrak{g}^1_q), (\gamma\varphi_1)_{\alpha_0\alpha_1}^{1,1} = ((-)^{q-1}/2\pi) \eta$.*

Proof. — By Lemma 1,

$$\begin{aligned}
 (\gamma\varphi_1)_{\alpha_0\alpha_1}^{1,1} &= \int_{\Delta^1} dt_1 \wedge \varphi_1(\omega_{\alpha_1} - \omega_{\alpha_0}) \\
 &= \varphi_1(\omega_{\alpha_1} - \omega_{\alpha_0}).
 \end{aligned}$$

Since $\varphi_k \in I^k(\mathfrak{g}I_q)$ is defined by

$$\det(\lambda I_q - (1/2\pi)X) = \sum_k \varphi_k(X) \lambda^{q-k}, \quad X \in \mathfrak{g}I_q \quad \text{and} \quad \text{trace}(\omega_{\alpha_{ij}}^i) = 0,$$

we obtain $(\gamma\varphi_1)_{\alpha_0\alpha_1}^{1,1} = (1/2\pi) \text{trace}(\omega_{\alpha_0j})$. By (1) and Lemma 3,

$$(\gamma\varphi_1)_{\alpha_0\alpha_1}^{1,1} = ((-1)^{q-1}/2\pi) \eta. \quad \text{Q.E.D.}$$

From this lemma, we obtain

THEOREM A. — $((-1)^{q-1}/2\pi) \eta$ defines a D-cohomology class in $H^2(\check{C}^{(*)}(\mathfrak{u}; A^*(M)), D)$ depending only on \mathfrak{F} .

Proof. — If $\gamma\varphi_1 \in \check{C}^{(2)}(\mathfrak{u}; A^*(M))$ is D-closed, then particular object $(\gamma\varphi_1)_{\alpha_0\alpha_1}^{1,1}$ is D-closed and, by Lemma 4,

$((-1)^{q-1}/2\pi) \eta \in \check{C}^{1,1}(\mathfrak{u}; A^*(M))$ define a D-cohomology class in $H^2(\check{C}^{(*)}(\mathfrak{u}; A^*(M)), D)$. Now we prove that $\gamma\varphi_1$ is D-closed.

$$\begin{aligned}
 (D(\gamma\varphi_1))_{\alpha_0\alpha_1}^{1,2} &= (D'(\gamma\varphi_1) + D''(\gamma\varphi_1))_{\alpha_0\alpha_1}^{1,2} \\
 &= (D'(\gamma\varphi_1))_{\alpha_0\alpha_1}^{1,2} + (D''(\gamma\varphi_1))_{\alpha_0\alpha_1}^{1,2} \\
 &= \{(\gamma\varphi_1)_{\alpha_1}^{0,2} - (\gamma\varphi_1)_{\alpha_0}^{0,2}\} + (-1)(d(\gamma\varphi_1))_{\alpha_0\alpha_1}^{1,2}.
 \end{aligned}$$

From Stokes' theorem,

$$(-1)d \cdot \int_{\Delta^1} \varphi_1(\Omega_{\alpha_0\alpha_1}) = \int_{\Delta^1} d \cdot \varphi_1(\Omega_{\alpha_0\alpha_1}) - \int_{\partial\Delta^1} j^* \varphi_1(\Omega_{\alpha_0\alpha_1})$$

and the left side of this is equal to $(-1)d \cdot (\gamma\varphi_1)_{\alpha_0\alpha_1}^{1,1}$, the first term of the right side vanishes and the second term of the right side is equal to $\varphi_1(\Omega_{\alpha_1}) - \varphi_1(\Omega_{\alpha_0})$. Thus $d \cdot (\gamma\varphi_1)_{\alpha_0\alpha_1}^{1,1} = \varphi_1(\Omega_{\alpha_1}) - \varphi_1(\Omega_{\alpha_0})$. From this and $(\gamma\varphi_1)_{\alpha_0}^{0,2} = \varphi_1(\Omega_{\alpha_0})$, we obtain

$$\begin{aligned}
 (D(\gamma\varphi_1))_{\alpha_0\alpha_1}^{1,2} &= \{\varphi_1(\Omega_{\alpha_1}) - \varphi_1(\Omega_{\alpha_0})\} \\
 &\quad + (-1)\{\varphi_1(\Omega_{\alpha_1}) - \varphi_1(\Omega_{\alpha_0})\} \\
 &= 0.
 \end{aligned}$$

From $(\gamma\varphi_1)_{\alpha_0}^{0,2} = \varphi_1(\Omega_{\alpha_0})$ and $d \circ \varphi_1(\Omega_{\alpha_0}) = 0$, we obtain

$$\begin{aligned}
 (D(\gamma\varphi_1))_{\alpha_0}^{0,3} &= (D''(\gamma\varphi_1))_{\alpha_0}^{0,3} \\
 &= (-1)^0 (d(\gamma\varphi_1))_{\alpha_0}^{0,3} \\
 &= 0.
 \end{aligned}$$

Now, from lemma 1, $(\gamma\varphi_1) \in \check{C}^{0,2} + \check{C}^{1,1}$. Thus we obtain

$$(D(\gamma\varphi_1))_{\alpha_0\alpha_1\alpha_2\alpha_3}^{3,0} = 0$$

and

$$\begin{aligned} (D(\gamma\varphi_1))_{\alpha_0\alpha_1\alpha_2}^{2,1} &= (D'(\gamma\varphi_1))_{\alpha_0\alpha_1\alpha_2}^{2,1} \\ &= \varphi_1(\omega_{\alpha_2}) - \varphi_1(\omega_{\alpha_1}) - \varphi_1(\omega_{\alpha_2}) + \varphi_1(\omega_{\alpha_0}) \\ &\quad + \varphi_1(\omega_{\alpha_1}) - \varphi_1(\omega_{\alpha_0}) \\ &= 0. \end{aligned}$$

Therefore we obtain $D(\gamma\varphi_1) = 0$. Q.E.D.

3. For q 1-forms w^1, \dots, w^q on U_α ($w^1 \wedge \dots \wedge w^q \neq 0$) we may choose $n-q$ 1-forms w^{q+1}, \dots, w^n on U_α such that

$$w^1 \wedge \dots \wedge w^q \wedge w^{q+1} \wedge \dots \wedge w^n \neq 0 \quad \text{on } U_\alpha.$$

Thus we obtain expressions

$$\varphi_j^i = \sum_k \varphi_{jk}^i w^k + \sum_a \varphi_{ja}^i w^a,$$

$$\eta = \sum_k \eta_k w^k + \sum_a \eta_a w^a.$$

Using same letters φ_j^i, η to simplify, we put

$$\varphi_j^i = \sum_a \varphi_{ja}^i w^a \quad \text{and} \quad \eta = \sum_a \eta_a w^a \quad (3)$$

on U_α .

Hereafter, we suppose that the manifold M admits foliations $\mathfrak{F}, \mathfrak{F}'$ complementally transversal to each other and that \mathfrak{F} (resp. \mathfrak{F}') is of codimension q (resp. $n-q$). Then we may consider that w^1, \dots, w^q are defined by \mathfrak{F} and that w^{q+1}, \dots, w^n are defined by \mathfrak{F}' .

Let 1-forms $\bar{w}^i, \bar{w}^a, \bar{\varphi}_j^i, \bar{\eta}$ on U_{α_1} correspond to 1-forms $w^i, w^a, \varphi_j^i, \eta$ on U_{α_0} respectively. Then we obtain

LEMMA 5. — On $U_{\alpha_0} \cap U_{\alpha_1} (\neq \emptyset)$, $\bar{\eta} = \eta$.

Proof. — On $U_{\alpha_0} \cap U_{\alpha_1}$, we may put

$$\bar{w}^i = \sum_j t_j^i w^j, \quad \bar{w}^a = \sum_b t_b^a w^b. \quad (4)$$

Proof. — We take η_{α_0} (resp. η_{α_1}) for η on U_{α_0} (resp. $\bar{\eta}$ on U_{α_1}). Then we obtain

$$\eta_{\alpha_1} - \eta_{\alpha_0} = 0 \quad \text{on } U_{\alpha_0} \cap U_{\alpha_1},$$

and

$$(D'\eta)_{\alpha_0\alpha_1}^{1,1} = \eta_{\alpha_1} - \eta_{\alpha_0} = 0.$$

Thus η is D' -closed.

Q.E.D.

BIBLIOGRAPHY

- [1] R. BOTT, Lectures on characteristic classes and foliations, *Lecture Notes in Math.*, Springer, 279 (1972), 1-94.
- [2] C. GODBILLON and J. VEY, Un invariant des feuilletages de codimension 1, *C.R. Acad. Sci.*, Paris, 273 (1971), A92-95.
- [3] F.W. KAMBER and P. TONDEUR, Foliated bundles and characteristic classes, *Lecture Notes in Math.*, Springer, 493 (1975).
- [4] H. KITAHARA and S. YOROZU, Sur l'homomorphisme de Chern-Weil local et ses applications au feuilletage, *C.R. Acad. Sci.*, Paris, 281 (1975), A703-706.
- [5] G. REEB, Sur certaines propriétés topologiques des variétés feuilletées, Hermann (1952).
- [6] Y. SHIKATA, On the spectral sequences associated to foliated structures, *Nagoya Math. J.*, 38 (1970), 53-61.
- [7] Y. SHIKATA, On the cohomology of bigraded forms associated with foliated structures, *Bull. Soc. Math. Grèce*, 15 (1974), 68-76.
- [8] A. SO, J.C. THOMAS and C. WATKISS, Sur la multiplicativité de l'homomorphisme de Chern-Weil local, *C.R. Acad. Sci.*, Paris, 280 (1975), A369-371.

Manuscrit reçu le 12 avril 1977

Proposé par G. Reeb.

H. KITAHARA and S. YOROZU,
 Department of Mathematics
 College of Liberal Arts
 Kanazawa University
 Kanazawa 920, Japan.