

ANNALES DE L'INSTITUT FOURIER

THIERRY FACK

Finite sums of commutators in C^* -algebras

Annales de l'institut Fourier, tome 32, n° 1 (1982), p. 129-137

http://www.numdam.org/item?id=AIF_1982__32_1_129_0

© Annales de l'institut Fourier, 1982, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

FINITE SUMS OF COMMUTATORS IN C*-ALGEBRAS

by Thierry FACK

Introduction.

Let A be a C^* -algebra and put

$$A_0 = \left\{ x \in A \mid x = \sum_{n \geq 1} x_n x_n^* - x_n^* x_n ; \text{norm convergence} \right\}.$$

By [4] (theorem 2.6), A_0 is exactly the null space of all finite traces on the self-adjoint part of A .

For von Neumann algebras, A_0 is spanned by finite sums of the above type (see for example [6]). This is not always true for C^* -algebras, as it is shown by Pedersen and Petersen ([8], lemma 3.5) for a very natural algebra. A reasonable question is then : when can this happen for C^* -algebras ?

The aim of this paper is to show that A_0 is spanned by finite sums for stable algebras and C^* -algebras with "sufficiently many projections" like infinite simple C^* -algebras or simple A.F-algebras (with unit).

We use the usual terminology of C^* -algebras as in [7]. A commutator of the form $[x, x^*] = xx^* - x^*x$ is called a self-adjoint commutator.

* * *
* *
*

I'd like to thank G. Skandalis for fruitful discussions and G.K. Pedersen who originally asked this question.

1. Stable C*-algebras.

Recall that a C*-algebra A is stable if $A \approx A \otimes \mathcal{K}$, where \mathcal{K} is the C*-algebra of compact operators. We have

THEOREM 1.1. — *Let A be a stable C*-algebra. Then, every hermitian element of A is the sum of five self-adjoint commutators.*

Every simple A.F.-algebra A without non zero finite trace being stable, it follows that A_0 is spanned by finite sums of self-adjoint commutators.

The proof of theorem 1.1 is based on the following lemmas.

LEMMA 1.2. — *Let A be a C*-algebra and $x = x^* \in A$. Let p be a projection in $M(A)$. Then, there exists $v \in A$ such that*

$$x = pxp + (1 - p)x(1 - p) + [v, v^*].$$

Proof. — Put

$$v = 1/2 |(1 - p)xp|^{1/2} - |(1 - p)xp|^{1/2} u^* + u |(1 - p)xp|^{1/2}$$

where u is the phase of $(1 - p)xp$. As $p \in M(A)$, we have $v \in A$. By direct calculation, we have $px(1 - p) + (1 - p)xp = [v, v^*]$.

LEMMA 1.3. — *Let A be a C*-algebra with unit and $x = x^* \in A$. Let $(\lambda_1, \dots, \lambda_n)$ be a sequence of real numbers satisfying*

$$0 \leq \sum_{i=1}^k \lambda_i \leq 1 \quad (k = 1, \dots, n - 1)$$

and

$$\sum_{i=1}^n \lambda_i = 0.$$

Then, there exists $u \in M_n(A)$, $\|u\| \leq \|x\|^{1/2}$, such that

$$\begin{bmatrix} \lambda_1 x & & & \circ \\ & \cdot & & \\ \circ & & \cdot & \\ & & & \lambda_n x \end{bmatrix} = [u, u^*].$$

Proof. — Write $x = x_+ - x_-$ and put

$$\mu_k^+ = \left(\sum_{i=1}^k \lambda_i \right)^{1/2} x_+^{1/2}$$

$$\mu_k^- = \left(\sum_{i=1}^k \lambda_i \right)^{1/2} x_-^{1/2} \quad (k = 1, \dots, n - 1).$$

Take $u = \sum_{k=1}^{n-1} (\mu_k^+ \otimes e_{k,k+1} + \mu_k^- \otimes e_{k+1,k})$, where $(e_{ij})_{1 \leq i, j \leq n}$ is the canonical system of matrix units. As $x_+ x_- = 0$, we get the result by direct calculation. \square

Let e be a rank one projection in \mathcal{K} .

LEMMA 1.4. — *Let A be a C*-algebra and $x = x^* \in A$. Then, $x \otimes e$ is the sum of two self-adjoint commutators of $A \otimes \mathcal{K}$.*

Proof. — Write $x \otimes e = \begin{bmatrix} x & & & \circ \\ & \lambda_1 x & & \\ & & \lambda_2 x & \\ \circ & & & \ddots \end{bmatrix} - \begin{bmatrix} \circ & & & \circ \\ & \lambda_1 x & & \\ & & \lambda_2 x & \\ \circ & & & \ddots \end{bmatrix},$

where $(\lambda_n)_{n \geq 1}$ is the sequence

$$\left(-\frac{1}{2}, -\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \underbrace{-\frac{1}{8}, \dots, -\frac{1}{8}}_{8 \text{ terms}}, \dots \right).$$

The result follows from lemma 1.3.

Proof of theorem 1.1. — Let x be a hermitian element of $A \otimes \mathcal{K}$. Take a projection $p \in M(\mathcal{K})$ with $p \sim 1 - p \sim 1$.

By lemma 1.2, there exists $v \in A \otimes \mathcal{K}$ such that

$$x = p x p + (1 - p) x (1 - p) + [v, v^*].$$

By lemma 1.4, $p x p$ and $(1 - p) x (1 - p)$ are both sums of two self-adjoint commutators. \square

2. Infinite simple C*-algebras.

The main result of this section is the following

THEOREM 2.1. — *Let A be a C*-algebra with unit. Suppose that there exist two orthogonal projections e and f such that $e \sim f \sim 1$ in A . Then, each hermitian element of A is the sum of five self-adjoint commutators.*

Recall that a simple C^* -algebra with unit is said to be *infinite* if it contains an element x such that $x^*x = 1$ and $xx^* \neq 1$. From theorem 2.1, we deduce

COROLLARY 2.2. — *Let A be an infinite simple C^* -algebra with unit. Then each hermitian element of A is the sum of five self-adjoint commutators.*

Apply theorem 2.1 and proposition 2.2 of [1]. The proof of theorem 2.1 is based on the following lemma :

LEMMA 2.3. — *Let A , e and f be as in theorem 2.1. Let p be a rank one projection in \mathcal{K} . Then, there exists a homomorphism*

$$\varphi : A \otimes \mathcal{K} \longrightarrow A \text{ such that}$$

$$\varphi(x \otimes p) = x \text{ for each } x \in (1 - f)A(1 - f).$$

Proof. — Let u, v be partial isometries such that

$$u^*u = v^*v = 1 \quad ; \quad uu^* = e, \quad vv^* = f.$$

Put $w_1 = 1 - f + vf$ and $w_n = vu^{n-1}v (n \geq 2)$.

The w_n are isometries with pairwise orthogonal ranges. Let (e_{ij}) be a system of matrix units for \mathcal{K} , with $e_{11} = p$. Put then

$$\varphi(z \otimes e_{ij}) = w_i z w_j^* \quad (z \in A). \quad \square$$

Proof of the theorem 2.1. — Let $x = x^* \in A$. By lemma 1.2, there exists $y \in A$ such that $x = exe + (1 - e)x(1 - e) + [y, y^*]$. By lemmas 2.3 and 1.4, both exe and $(1 - e)x(1 - e)$ are sums of two self-adjoint commutators (note that $exe \in (1 - f)A(1 - f)$). \square

For non simple infinite C^* -algebras with unit, we may combine corollary 2.2 with the following obvious lemma :

LEMMA 2.4. — *Let $0 \longrightarrow J \longrightarrow A \longrightarrow B \longrightarrow 0$ be an exact sequence of C^* -algebras. Suppose that each hermitian element of J (resp. of B) is a sum of n (resp. k) self-adjoint commutators. Then, any hermitian element of A is the sum of $n + k$ self-adjoint commutators.*

Example. — Let $A = (A(i, j))_{i, j \in \Sigma}$ be a transition matrix on a finite set Σ . Assume that A has no zero columns or rows. For $i, j \in \Sigma$, write $i \leq j$ if the transition from j to i is possible

(cf. [2]). We call i and j equivalent if $i \leq j \leq i$. Let F be the set of maximal states : $F = \{i \in \Sigma \mid \forall j \in \Sigma \ i \leq j \implies j \leq i\}$. F is an union of equivalence classes and every element of Σ is majorized by an element of F .

Assume that the restriction A_γ of A to each equivalence classe γ of F is not a permutation matrix. Then \mathcal{O}_A is defined in [2], [3] as the C*-algebra generated by *any* system $(S_i)_{i \in \Sigma}$ of non zero partial isometries with pairwise orthogonal ranges satisfying

$$S_i^* S_i = \sum_{j \in \Sigma} A(i, j) S_j S_j^* \quad (i \in \Sigma).$$

We claim that each hermitian element of \mathcal{O}_A is the sum of ten self-adjoint commutators.

Put $A' = A_{\Sigma - F}$ and $A'' = A_F$.

As $\mathcal{O}_{A''}$ is a finite direct sum of \mathcal{O}_B with B irreducible, each hermitian element of $\mathcal{O}_{A''}$ is the sum of five self-adjoint commutators by corollary 2.2 and theorem 2.14 of [3]. But it is easy to see that there exists an exact sequence

$$0 \longrightarrow \mathcal{O}_{A'} \otimes \mathcal{K} \longrightarrow \mathcal{O}_A \longrightarrow \mathcal{O}_{A''} \longrightarrow 0$$

and the result follows from lemma 2.4 and theorem 1.1.

3. Simple A.F-algebras.

In this section, we shall prove the following result :

THEOREM 3.1. — *Let A be a simple approximately finite dimensional C*-algebra with unit. Then, each element of A_0 is the sum of seven self-adjoint commutators.*

The proof is based on the following technical lemmas :

LEMMA 3.2. — *Let A be a C*-algebra and $x = x^* \in A$. Let p, q, r be orthogonal projections in A with $p + q + r = 1$. Then, there exists $u \in A$, $\|u\| \leq 2\sqrt{2}\|x\|^{1/2}$, such that*

$$x - pxp - qxq - rxr = [u, u^*].$$

Proof. – Put

$$u = p - r - \frac{1}{2}(pxq - qxp) - \frac{1}{4}(pxr - rxp) - \frac{1}{2}(qxr - rxq).$$

We have $x - pxp - qxq - rxr = [u, u^*]$ by direct calculation. Moreover, $\|x\| \leq 2$ implies $\|u\| \leq 4$. The lemma follows. \square

LEMMA 3.3. – Let A be a C^* -algebra and $x = x^* \in A$. Let p, q, r be orthogonal projections in A with $p + q + r = 1$ and $p \lesssim q \lesssim r$. Then, there exists $u \in A$, $\|u\| \leq 3\|x\|^{1/2}$ and $y \in A$ such that

$$\begin{aligned} x &= [u, u^*] + y \\ pyp &= qyq = 0 \\ \|ryr\| &\leq 3\|x\|. \end{aligned}$$

Proof. – Let v and w be partial isometries such that $vv^* = p$, $v^*v \leq q$, $ww^* = q$, $w^*w \leq r$. Put

$$u = \sqrt{(pxp)_+}v + v^*\sqrt{(pxp)_-} + \sqrt{(qxq + v^*xv)_+}w + w^*\sqrt{(qxq + v^*xv)_-}$$

and $y = x - [u, u^*]$. We have $\|u\| \leq 3\|x\|^{1/2}$, $pyp = qyq = 0$ and $\|ryr\| \leq 3\|x\|$ by direct calculation. \square

LEMMA 3.4. – Let A be a C^* -algebra and $x = x^* \in A$. Let p, q, r be orthogonal projections in A with $p + q + r = 1$ and $p \lesssim q \lesssim r$. Then, there exist $u, v \in A$; $\|u\| \leq 3\|x\|^{1/2}$, $\|v\| \leq 13\|x\|^{1/2}$ such that $x - [u, u^*] - [v, v^*] \in rAr$ and $\|x - [u, u^*] - [v, v^*]\| \leq 3\|x\|$.

Proof. – By lemma 3.3, we have $x = [u, u^*] + y$ with $\|u\| \leq 3\|x\|^{1/2}$, $pyp = qyq = 0$ and $\|ryr\| \leq 3\|x\|$. We deduce $\|y\| \leq 19\|x\|$, and the result follows from lemma 3.2. \square

LEMMA 3.5. – Let B be a finite dimensional C^* -algebra and $x \in B_0$. Then, there exists $u \in B$, $\|u\| \leq \sqrt{2}\|x\|^{1/2}$, such that $x = [u, u^*]$.

Proof. – Using the decomposition of B into simple components, we can assume that $B = M_n(\mathbb{C})$. One may also suppose x is diagonal. The proper values of x are real numbers $\lambda_1, \dots, \lambda_n$

with $\sum_{i=1}^n \lambda_i = 0$. As there exists a permutation τ of $\{1, \dots, n\}$ such that $0 \leq \sum_{i=1}^k \lambda_{\tau(i)} \leq 2 \sup_{1 \leq i \leq n} |\lambda_i|$ for $k = 1, \dots, n$, we can assume that $x = \sum_{i=1}^n \lambda_i e_{ii}$ and $0 \leq \sum_{i=1}^k \lambda_i \leq 2 \|x\|$ ($k = 1, \dots, n$), where $(e_{ij})_{1 \leq i, j \leq n}$ is some system of matrix units. Apply then lemma 1.3. \square

LEMMA 3.6. — *Let A be a simple A.F-algebra with unit. Suppose that A is non isomorphic to $M_n(\mathbf{C})$. Then, there exist sequences $(p_n)_{n \geq 1}$, $(q_n)_{n \geq 1}$ and $(r_n)_{n \geq 1}$ of projections such that*

- i) $p_1 + q_1 + r_1 = 1$
- ii) $p_n \lesssim q_n \lesssim r_n \quad (n \geq 1)$
- iii) *the r_n are mutually orthogonal,*
- iv) $r_{n-1} = p_n + q_n \quad (n \geq 2)$.

Proof. — It suffices to show that there exists, for each projection $p \neq 0$, an element $q \in K_0(A)_+$ such that $2q \leq p \leq 3q$. Passing to pAp , we may assume that $p = 1$. By [5] (lemma A.4.3), $K_0(A)$ is the limit of a system $Z^{r(1)} \xrightarrow{\varphi_1} Z^{r(2)} \xrightarrow{\varphi_2} \dots$ having the following properties :

- i) the φ_n are strictly positive, i.e. $\varphi_n = (\alpha_{ij}^n)$ with $\alpha_{ij}^n > 0$,
- ii) there exist order units $u_n \in Z^{r(n)}$ such that

$$u_1 \longrightarrow u_2 \longrightarrow \dots \longrightarrow 1.$$

One then may choose $q \in K_0(A)_+$ such that $2q \leq 1 \leq 3q$. \square

Proof of theorem 3.1. — The case $A = M_n(\mathbf{C})$ is trivial, so that we can assume $A \not\cong M_n(\mathbf{C})$. Let x be in A_0 . Let $(p_n)_{n \geq 1}$, $(q_n)_{n \geq 1}$ and $(r_n)_{n \geq 1}$ be sequences of projections as in lemma 3.6.

Apply first lemma 3.4 to get $x_1 \in r_1 A r_1$, $\|x_1\| \leq 3 \|x\|$, and $u, v \in A$ such that $x = [u, u^*] + [v, v^*] + x_1$. As r_1 is an order unit in $K_0(A)_+$, any finite trace on $r_1 A r_1$ extends uniquely to a finite trace on A , so that $x_1 \in (r_1 A r_1)_0$.

Starting from x_1 , we are going to construct sequences $(x_n)_{n \geq 1}$, $(u_n)_{n \geq 1}$, $(v_n)_{n \geq 1}$ and $(w_n)_{n \geq 1}$ satisfying

$$\alpha) x_n = [u_n, u_n^*] + [v_n, v_n^*] + [w_n, w_n^*] + x_{n+1},$$

$$\beta) u_n \in r_n A r_n; \quad v_n, w_n \in (r_n + r_{n+1}) A (r_n + r_{n+1}),$$

$$\gamma) x_n \in (r_n A r_n)_0,$$

$$\delta) \|x_n\| \leq \frac{3 \|x\|}{n}$$

$$\epsilon) \|u_n\| \leq 2 \|x_n\|^{1/2} \quad \text{and} \quad v_n, w_n \rightarrow 0 \quad (n \rightarrow \infty).$$

Suppose $(x_1, \dots, x_{n-1}, x_n)$, (u_1, \dots, u_{n-1}) , (v_1, \dots, v_{n-1}) and (w_1, \dots, w_{n-1}) constructed.

Put $\alpha = \frac{\|x\|}{n+1}$. As $x_n \in (r_n A r_n)_0$, we have

$$x_n = \sum_{p \geq 1} [c_p, c_p^*]$$

where $c_p \in r_n A r_n$ and the sum being norm convergent. By approximation, we can find a finite dimensional subalgebra B of $r_n A r_n$ and $y \in B_0$ such that $\|y\| \leq 2 \|x_n\|$ and $\|x_n - y\| \leq \alpha$.

By lemma 3.5, there exists $u_n \in r_n A r_n$,

$$\|u_n\| \leq \sqrt{2} \|y\|^{1/2} \leq 2 \|x_n\|^{1/2}$$

such that $x_n = [u_n, u_n^*] + z$, where $z = x_n - y$.

Note that $z \in ((r_n + r_{n+1}) A (r_n + r_{n+1}))_0$.

By lemma 3.4, there exist $v_n, w_n \in (r_n + r_{n+1}) A (r_n + r_{n+1})$ such that $z = [v_n, v_n^*] + [w_n, w_n^*] + x_{n+1}$ where $x_{n+1} \in r_{n+1} A r_{n+1}$ and

$$\|v_n\| \leq 3 \|z\|^{1/2} \leq 3\alpha^{1/2}$$

$$\|w_n\| \leq 13 \|z\|^{1/2} \leq 13\alpha^{1/2}.$$

We have

$$x_n = [u_n, u_n^*] + [v_n, v_n^*] + [w_n, w_n^*] + x_{n+1}$$

and hence $x_{n+1} \in (r_{n+1} A r_{n+1})_0$. Moreover,

$$\|x_{n+1}\| \leq 3 \|z\| \leq 3\alpha \leq \frac{3 \|x\|}{n+1}.$$

By induction, the existence of four sequences satisfying $\alpha)$, $\beta)$, $\gamma)$, $\delta)$ and $\epsilon)$ is then proved.

Put

$$U = \sum_{n > 1} u_n$$

$$V_{ev} = \sum_{n \geq 1} v_{2n} ; \quad V_{od} = \sum_{n \geq 0} v_{2n+1} ,$$

$$W_{ev} = \sum_{n \geq 1} w_{2n} ; \quad W_{od} = \sum_{n \geq 0} w_{2n+1} .$$

These sums make sense because they involve elements with disjoint support and norm converging to zero. Moreover, we have

$$x = [u, u^*] + [v, v^*] + [U, U^*] + [V_{ev}, V_{ev}^*] + [V_{od}, V_{od}^*] \\ + [W_{ev}, W_{ev}^*] + [W_{od}, W_{od}^*] .$$

The proof of theorem 3.1 is complete. \square

BIBLIOGRAPHY

- [1] J. CUNTZ, The structure of multiplication and addition in simple C*-algebras, *Math. Scand.*, 40 (1977).
- [2] J. CUNTZ, A class of C*-algebras and topological Markov chains II : Reducible Markov chains and the Ext-functor for C*-algebras, *Preprint Univ. Heidelberg*, n° 57 (March 1980).
- [3] J. CUNTZ and W. KRIEGER, A class of C*-algebras and topological Markov chains, *Inventiones Math.*, 56 (1980), 251-268.
- [4] J. CUNTZ and G.K. PEDERSEN, Equivalence and traces on C*-algebras, *J. Functional Analysis*, to appear.
- [5] E.G. EFFROS, Dimensions and C*-algebras, *Preprint UCLA* (1980).
- [6] T. FACK et P. DE LA HARPE, Sommes de commutateurs dans les algèbres de von Neumann finies continues, *Ann. Inst. Fourier*, Grenoble, 30,3 (1980), 49-73.
- [7] G.K. PEDERSEN, *C*-algebras and their Automorphism Groups*, Academic Press, New-York (1979).
- [8] G.K. PEDERSEN and N.H. PETERSEN, Ideals in a C*-algebra, *Math. Scand.*, 27 (1970), 193-204.

Manuscrit reçu le 25 mars 1981.

Thierry FACK,
Laboratoire de Mathématiques Fondamentales
Université Pierre et Marie Curie
4, Place Jussieu
Tour 45-46, 3^{ème} étage
75230 Paris Cedex 05.