ANNALES DE L'INSTITUT FOURIER

MANUEL VALDIVIA

A class of locally convex spaces without $\mathcal{C}$-webs

Annales de l'institut Fourier, tome 32, n° 2 (1982), p. 261-269

<http://www.numdam.org/item?id=AIF_1982__32_2_261_0>

© Annales de l'institut Fourier, 1982, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (http://annalif.ujf-grenoble.fr/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Ann. Inst. Fourier, Grenoble
32, 2 (1982), 261-269.

A CLASS OF LOCALLY CONVEX SPACES WITHOUT $\mathscr{C}$-WEBS

by Manuel VALDIVIA

The linear spaces we use are defined over the field K of the real or complex numbers. Space means non-zero separated locally convex topological linear space. Given a space E we denote by E' its topological dual. The dimension of E is denoted by $\dim E$. $\hat{E}$ is the completion of E. If E_n is a one-dimensional space, $n = 1, 2, \ldots$, we put $\varphi = \bigoplus_{n=1}^{\infty} E_n$.

A space E is unordered Baire-like if given a sequence (A_n) of closed absolutely convex sets of E, with $E = \bigcup_{n=1}^{\infty} A_n$, there is a positive integer n_0 such that A_{n_0} is a neighbourhood of the origin in E.

A space E is suprabarrelled if given an increasing sequence (E_n) of subspaces of E covering E there is a positive integer n_0 such that E_{n_0} is barrelled and dense in E, [7].

A space E is Baire-like if given an increasing sequence (A_n) of closed absolutely convex sets of E covering E there is a positive integer n_0 such that A_{n_0} is a neighbourhood of the origin in E.

If A is a bounded absolutely convex subsets of a space E we denote by E_A the linear hull of A endowed with the topology derived from the gauge of A. A is completing if E_A is a Banach space.

In what follows E and F are spaces and n is any non negative integer. We denote by $E \otimes_\pi F$ and $E \otimes_\varepsilon F$ the tensor product $E \otimes F$ endowed with the projective topology and the topology of the by-equicontinuous convergence respectively. $E \hat{\otimes}_\varepsilon F$ is the completion of $E \otimes_\varepsilon F$. We set

$$R_n(E, F) = \{z \in E \otimes F : \text{rank } z \leqslant n\}.$$

We write $\hat{R}_n(E, F)$ to denote the closure of $R_n(E, F)$ in $E \hat{\otimes}_\varepsilon F$.

MAIN LEMMA. — $\hat{R}_n(E, F) = R_n(\hat{E}, \hat{F})$.

Proof. — We proceed by complete induction. The result is obvious for $n = 0$. We suppose that the property is true for a positive integer p. Let

$$\left\{\sum_{j=1}^{p+1} x_j^i \otimes y_j^i : i \in I, \geqslant\right\} \tag{1}$$

be a net in $R_{p+1}(E,F)$ converging in $E \hat{\otimes}_\varepsilon F$ to an element $z \neq 0$. Since $E' \otimes F'$ separate points in $E \hat{\otimes}_\varepsilon F$, we can select $u \in E'$, $v \in F'$ such that $u \otimes v(z) \neq 0$. Obviously

$$\lim \left\{\sum_{j=1}^{p+1} u(x_j^i)v(y_j^i) : i \in I, \geqslant\right\} = u \otimes v(z) \neq 0. \tag{2}$$

Since (1) is a Cauchy net in $E \otimes_\varepsilon F$ given $\delta > 0$ and an equicontinuous subset V of F' there is an $i_0 \in I$ such that

$$\sup \left\{\left|\sum_{j=1}^{p+1} (u(x_j^h)w(y_j^h) - u(x_j^k)w(y_j^k))\right| : w \in V\right\} < \delta, \quad \text{for} \quad h, k \geqslant i_0$$

and therefore, according to (2),

$$\left\{\sum_{j=1}^{p+1} u(x_j^i)y_j^i : i \in I, \geqslant\right\} \tag{3}$$

converges in $\hat{F}$ to $y \neq 0$.

From (2) we deduce the existence of a subnet of (1), for which we use the same notation, such that at least one of the nets

$$\{u(x_j^i) : i \in I, \geqslant\}, \qquad j = 1, 2, \ldots, p + 1,$$

has all its elements different from zero. Without loss of generality we suppose that $u(x_1^i) \neq 0$, $\forall i \in I$.

Setting $\sum_{j=1}^{p+1} u(x_j^i)y_j = v_1^i$, $i \in I$, we have that

$$\sum_{j=1}^{p+1} x_j^i \otimes y_j^i = \frac{1}{u(x_1^i)} x_1^i \otimes \left(v_1^i - \sum_{j=2}^{p+1} u(x_j^i)y_j^i\right) + \sum_{j=2}^{p+1} x_j^i \otimes y_j^i,$$

and setting

$$z_1^i = \frac{1}{u(x_1^i)} x_1^i, \quad z_j^i = x_j^i - \frac{u(x_j^i)}{u(x_1^i)} x_1^i, \quad v_j^i = y_1^i, \quad j = 2, 3, \ldots, p + 1,$$

we have that the net (1) can be expressed as

$$\left\{\sum_{j=1}^{p+1} z_j^i \otimes v_j^i : i \in I, \geqslant\right\}.$$

Reasoning in the same way as we have done for the net (3) it follows that $\left\{\sum_{j=1}^{p+1} v(v_j^i)z_j^i : i \in I, \geqslant\right\}$ converges in $\hat{E}$ to an element $x \neq 0$. According to (2) again we suppose that $v(v_1^i) \neq 0$, $\forall i \in I$, and setting $\sum_{j=1}^{p+1} v(v_j^i)z_j^i = u_1^i$ it follows that

$$\begin{aligned}\sum_{j=1}^{p+1} x_j^i \otimes y_j^i &= \sum_{j=1}^{p+1} z_j^i \otimes v_j^i \\ &= \frac{1}{v(v_1^j)}\left(u_1^i - \sum_{j=2}^{p+1} v(v_j^i)z_j^i\right) \otimes v_1^i + \sum_{j=2}^{p+1} z_j^i \otimes v_j^i \\ &= \frac{1}{v(v_1^i)} u_1^i \otimes v_1^i + \sum_{j=2}^{p+1} z_j^i \otimes \left(v_j^i - \frac{v(v_j^i)}{v(v_1^i)} v_1^i\right).\end{aligned}$$

The net $\left\{\frac{1}{v(v_1^i)} u_1^i \otimes v_1^i : i \in I, \geqslant\right\}$ converges to $\frac{1}{u \otimes v(z)} x \otimes y$ in $E \hat{\otimes}_\varepsilon F$ and according to the induction hypothesis

$$\left\{\sum_{j=2}^{p+1} z_j^i \otimes \left(v_j^i - \frac{v(v_j^i)}{v(v_1^i)} v_1^i\right) : i \in I, \geqslant\right\}$$

converges to an element of $R_p(\hat{E},\hat{F})$ and therefore

$$\hat{R}_{p+1}(E,F) \subset R_{p+1}(\hat{E},\hat{F}).$$

Finally, since $R_{p+1}(E,F)$ is obviously dense in $R_{p+1}(\hat{E},\hat{F})$ it follows that $\hat{R}_{p+1}(E,F) = R_{p+1}(\hat{E},\hat{F})$. q.e.d.

COROLLARY 1. — *If* E *and* F *are complete then* $R_n(E,F)$ *is complete in* $E \otimes_\varepsilon F$.

COROLLARY 2. — $R_n(E,F)$ *is closed in* $E \otimes_\varepsilon F$.

Proof. — $R_n(E,F)$ coincides with $R_n(\hat{E},\hat{F}) \cap (E \otimes F)$ and according to Main Lemma $R_n(\hat{E},\hat{F})$ is complete in $E \hat{\otimes}_\varepsilon F$, hence $R_n(E,F)$ is closed in $E \otimes_\varepsilon F$. q.e.d.

THEOREM 1. – $E \otimes_\varepsilon F$ *has a* $\mathscr{C}$*-web if and only if one of the two following conditions is satisfied :*

1. E *has a* $\mathscr{C}$*-web and* $\dim F \leqslant \chi_0$;
2. F *has a* $\mathscr{C}$*-web and* $\dim E \leqslant \chi_0$.

Proof. – If F is finite dimensional and if E has a $\mathscr{C}$-web then $E \otimes_\varepsilon F$ is isomorphic to $E^{\dim F}$ which has a $\mathscr{C}$-web. If F has countable infinite dimension and if E has a $\mathscr{C}$-web then $E \otimes_\varepsilon F$ has a $\mathscr{C}$-web since the topology of $E \otimes_\varepsilon F$ is coarser than the topology ot $E \otimes_\varepsilon \varphi$ and this space has a topology coarser than the topology of $E \oplus E \oplus E \oplus \ldots$. We reach the same conclusion changing the roles of E and F.

We suppose now that $\dim E > \chi_0$, $\dim F > \chi_0$ and that $E \otimes_\varepsilon F$ has a $\mathscr{C}$-web $(C_{n_1, n_2, \ldots, n_p})$. Let $\{x_i : i \in I\}$ and $\{y_i : i \in I\}$ be families of linearly independent vectors in E and F respectively such that $\operatorname{card} I > \chi_0$. Since $\bigcup_{n=1}^{\infty} C_n = E \otimes F$ and $\bigcup_{n_{p+1}=1}^{\infty} C_{n_1, n_2, \ldots, n_{p+1}} = C_{n_1, n_2, \ldots, n_p}$ there is a sequence (n_p) of positive integers and there is a decreasing sequence (J_p) of subsets of I such that $x_i \otimes y_i \in C_{n_1, n_2, \ldots, n_p}$, $i \in J_p$, $\operatorname{card} J_p > \chi_0$, $p = 1, 2, \ldots$. It follows that we can take linearly independent elements $u_p \in E$, $v_p \in F$, $p = 1, 2, \ldots$, such that

$$u_p \otimes v_p \in C_{n_1, n_2, \ldots, n_p}.$$

We select a sequence (λ_p) of strictly positive real numbers such that $\{\lambda_p u_p \otimes v_p : p = 1, 2, \ldots\}$ is contained in a bounded completing absolutely convex subset A of $E \otimes_\varepsilon F$ [3, p. 75]. Since $(E \otimes F)_A$ is a Banach space and since $R_n(E,F)$ is closed in $E \otimes_\varepsilon F$ there is a positive integer n_0 such that $R_{n_0}(E,F) \cap (E \otimes F)_A$ has interior point in $(E \otimes F)_A$ and therefore

$$R_{2n_0}(E,F) \cap (E \otimes F)_A = (R_{n_0}(E,F) + R_{n_0}(E,F)) \cap (E \otimes F)_A$$

is a neighbourhood of the origin in $(E \otimes F)_A$ and since $R_{2n_0}(E,F)$ contains any scalar multiple of every vector $z \in R_{2n_0}(E,F)$ it follows that

$$A \subset R_{2n_0}(E,F).$$

But $\sum_{p=1}^{2n_0+1} \frac{1}{2^p} \lambda_p u_p \otimes v_p$ belongs to A and its rank is $2n_0 + 1$. This is a contradiction.

We suppose now that $E \otimes_\varepsilon F$ has a $\mathscr{C}$-web and $\dim F \leqslant \chi_0$. We

take $y_0 \in F$, $y_0 \neq 0$. Let T be the mapping from E into $E \otimes_\varepsilon F$ such that $T(x) = x \otimes y_0$, $x \in E$. Then T(E) is isomorphic to E (cf. [3], § 44, 1.(4)). Since T(E) is contained in $R_1(E,F)$ it is easy to prove that T(E) is closed in $E \otimes_\varepsilon F$ and thus has a $\mathscr{C}$-web. Analogously, if $E \otimes_\varepsilon F$ has a $\mathscr{C}$-web and $\dim E \leqslant \chi_0$, F has a $\mathscr{C}$-web. q.e.d.

THEOREM 2. — $E \otimes_\pi F$ *has a* $\mathscr{C}$*-web if and only if one of the two following conditions is satisfied :*

1. E *has a* $\mathscr{C}$*-web and* $\dim F \leqslant \chi_0$;
2. F *has a* $\mathscr{C}$*-web and* $\dim E \leqslant \chi_0$.

Proof. — On $E \otimes F$ the π-topology is finer than the ε-topology. Hence, if $E \otimes_\pi F$ has a $\mathscr{C}$-web then $E \otimes_\varepsilon F$ has a $\mathscr{C}$-web. Thus we apply the former result to obtain that 1 or 2 is satisfied.

If F is finite dimensional and if E has a $\mathscr{C}$-web then $E \otimes_\pi F = E \otimes_\varepsilon F$ has a $\mathscr{C}$-web. If F has countable infinite dimension and if E has a $\mathscr{C}$-web then $E \otimes_\pi F$ has a $\mathscr{C}$-web since the topology of $E \otimes_\pi F$ is coarser than the topology of $E \otimes_\pi \varphi = E \otimes_\varepsilon \varphi$. We reach the same conclusion changing the roles of E and F. q.e.d.

THEOREM 3. — *If* E *and* F *are Suslin spaces then* $E \otimes_\pi F$ *is a Suslin space.*

Proof. — Let $\psi : E \times F \to E \otimes_\pi F$ be the canonical bilinear mapping which is continuous and thus $\psi(E,F) = R_1(E,F)$ is a Suslin topological subspace of $E \otimes_\pi F$. On the other hand the mapping

$$\theta : R_1(E,F) \times R_1(E,F) \times \ldots \times R_1(E,F) \to E \otimes_\pi F$$

so that

$$\theta(x_1 \otimes y_1, x_2 \otimes y_2, \ldots, x_n \otimes y_n) = \sum_{j=1}^{n} x_j \otimes y_j$$

is π-continuous and therefore

$$\theta(R_1(E,F) \times R_1(E,F) \times \ldots \times R_1(E,F)) = R_n(E,F)$$

is a Suslin topological subspace of $E \otimes_\pi F$. Finally, $E \otimes_\pi F$ is a Suslin space since coincides with $\bigcup_{n=1}^{\infty} R_n(E,F)$, (cf. [2], § 6, N° 2, Prop. 8).

q.e.d.

COROLLARY 1.3. — *If* E *and* F *are Suslin spaces then* $E \otimes_\varepsilon F$ *is a Suslin space.*

Note 1. — If we consider two Suslin spaces E and F of dimension larger than χ_0 we can apply the former results to obtain that $E \otimes_\pi F$ is a Suslin space without a $\mathscr{C}$-web. Thus the closed graph theory of De Wilde [4] does not contain the closed graph theorem of L. Schwartz, [7] and [8].

Bourbaki [1, p. 43] proves that if E and F are metrizable and F is barrelled then every separately equicontinuous set of bilinear forms on $E \times F$ is equicontinuous. Since every barrelled metrizable space is Baire-like [1], Theorem 4 generalizes this result.

THEOREM 4. — *If* E *is metrizable and* F *is a Baire-like space then every separately equicontinuous set* $\mathscr{B}$ *of bilinear forms on* $E \times F$ *is equicontinuous.*

Proof. — Let (U_n) be a decreasing basis of closed absolutely convex neighbourhoods of the origin in E. We set

$$V_n = \{y \in F : |B(x,y)| \leqslant 1, x \in U_n, B \in \mathscr{B}\}, \quad n = 1, 2, \ldots,$$

which is a closed absolutely convex subset of F and the sequence (V_n) is increasing. On the other hand if $z \in F$ there is a positive integer p such that $|B(x,z)| \leqslant 1$, $x \in U_p$, $B \in \mathscr{B}$, according to the separate equicontinuity of $\mathscr{B}$. Thus $z \in V_p$ and since F is Baire-like there is a positive integer q such that V_q is a neighbourhood of the origin in F. Therefore $|B(x,y)| \leqslant 1$, $(x,y) \in U_q \times V_q$, $B \in \mathscr{B}$. q.e.d.

COROLLARY 1.4. — *If* E *is metrizable and if* F *is a barrelled space whose completion is Baire, then every set separately equicontinuous of bilinear forms on* $E \times F$ *is equicontinuous.*

Proof. — It is an immediate consequence of the former theorem and from the fact that every barrelled space whose completion is Baire is a Baire-like, [6]. q.e.d.

Note 2. — It is immediate that our Theorem 4 can be easily generalized to bilinear mapping with range in a third locally convex space. Then this theorem contains another result due to Bourbaki [3, Ex. 1, Chap. III, § 4, p. 44] for locally convex spaces.

PROPOSITION 1. — *If* E *is a barrelled metrizable space and if* F *is a Baire-like space then* $E \otimes_\pi F$ *is Baire-like.*

Proof. — Let (W_n) be an increasing sequence of closed absolutely convex subsets of $E \otimes_\pi F$ covering $E \otimes F$ and let (U_n) be a decreasing basis of closed absolutely convex neighbourhoods of the origin in E. We set

$$V_n = \{y \in F : x \otimes y \in W_n, x \in U_n\}, \qquad n = 1, 2, \ldots,$$

which is a closed absolutely convex subsets of F and (V_n) is an increasing sequence.

$$Z_n = \{x \in E : x \otimes z \in W_n\}, \qquad n = 1, 2, \ldots,$$

which is a closed absolutely convex subset of E and (Z_n) is an increasing sequence covering E. Then there is a positive integer n_0 such that Z_{n_0} is a neighbourhood of the origin in E. We can find a positive integer $p \geqslant n_0$ such that $U_p \subset Z_{n_0}$ and then $z \in V_p$ and so (V_n) covers F. Since F is a Baire-like there is a positive integer q such that V_q is a neighbourhood of the origin in E and thus W_q is a neighbourhood of the origin in $E \otimes_\pi F$.

q.e.d.

PROPOSITION 2. — *If* E *and* F *are unordered Baire-like spaces and if* E *is metrizable then* $E \otimes_\pi F$ *is unordered Baire-like.*

Proof. — Let (W_n) be a sequence of closed absolutely convex subsets of $E \otimes_\pi F$ covering $E \otimes F$ and let (U_n) be a decreasing basis of closed absolutely convex neighbourhood of the origin in E. We set

$$V_{n,m} = \{y \in F : x \otimes y \in W_n, \forall x \in U_m\}, \qquad , m = 1, 2, \ldots.$$

Obviously $V_{n,m}$ is a closed absolutely convex subset of F. We shall see that $\bigcup_{n,m=1}^{\infty} V_{n,m} = F$.

Indeed, let $z \in F$ and set

$$Z_n = \{x \in E : x \otimes z \in W_n\}, \quad n = 1, 2, \ldots,$$

which is a closed absolutely convex subset of E such that $\bigcup_{n=1}^{\infty} Z_n = E$. There is a positive integer p such that Z_p is a neighbourhood of the origin in E. We find a positive integer q such that $U_q \subset Z_p$. Then

$z \in V_{p,q}$. We can find positive integers r, s such that $V_{r,s}$ is a neighbourhood of the origin in F. If $\psi : E \times F \to E \otimes F$ is the canonical bilinear mapping then $\psi(U_s \times V_{r,s}) \subset W_r$ and thus W_r is a neighbourhood of the origin in $E \otimes_\pi F$. q.e.d.

PROPOSITION 3. — *If* E *is a metrizable suprabarrelled space and if* F *is an unordered Baire-like space, then* $E \otimes_\pi F$ *is suprabarrelled.*

Proof. — Let (G_n) be an increasing sequence of subspaces of $E \otimes_\pi F$ covering $E \otimes F$ and let us suppose G_n is not barrelled, $n = 1, 2, \ldots$. We can find in each G_n a barrel T_n which is not a neighbourhood of the origin in G_n. Let W_n be the closure of T_n in $E \otimes_\pi F$, $n = 1, 2, \ldots$. We use $V_{n,m}$ and Z_n with the same meaning as before. We set

$$S_n = \{x \in E : x \otimes z \in G_n\}, \quad n = 1, 2, \ldots.$$

(S_n) is an increasing sequence of subspace of E covering E and therefore there is a positive integer p such that S_p is barrelled and dense in E. If $x \in S_p$ there is a real number $h > 0$ such that $h(x \otimes z) \in T_p \subset W_p$ and thus $hz \in Z_p$, i.e., Z_p is absorbing and thus a barrel in E. It follows that $Z_p \cap S_p$ is a barrel in S_p and therefore a neighbourhood of 0. But S_p is dense in E, so $Z_p = \overline{Z_p \cap S_p}$ is a neighbourhood of the origin in E. Reasoning as we did in the proposition above, we obtain a positive number r such that W_r is a neighbourhood of the origin in $E \otimes_\pi F$, which is a contradiction. According to Proposition 1 there is a positive integer n_0 such that G_n is dense in $E \otimes_\pi F$, $n \geqslant n_0$. q.e.d.

PROPOSITION 4. — $E \otimes_\pi F$ *is a Baire space if and only if one of the following two conditions is satisfied :*

1. $\dim E < \infty$ *and* $F^{\dim E}$ *is a Baire Space*;
2. $\dim F < \infty$ *and* $E^{\dim F}$ *is a Baire space.*

Proof. — If the dimension of E is finite then $E \otimes_\pi F$ is isomorphic to $F^{\dim E}$. Analogously if F is finite dimensional then $E \otimes_\pi F$ is isomorphic to $E^{\dim F}$. Suppose that E and F are infinite dimensional and $E \otimes_\pi F$ is a Baire space. We can find a positive integer n_0 such that $R_{n_0}(E,F)$ is of second category in $E \otimes_\pi F$. According to Corollary 2.1, $R_{n_0}(E,F)$ is closed in $E \otimes_\varepsilon F$ and thus it is closed in $E \otimes_\pi F$ hence $R_{2n_0}(E,F)$ is a neighbourhood of the origin in $E \otimes_\pi F$ which is a contradiction since $R_{2n_0}(E,F)$ is not absorbing in $E \otimes F$.

The following Theorem is an immediate consequence of the preceding results, having in mind that every infinite dimensional Fréchet spaces has dimension larger or equal than $2^{\aleph_0}$, [6].

THEOREM 5. — *If* E *and* F *are infinite dimensional Fréchet spaces the following is true for* $E \otimes_\pi F$:

1. *is unordered Baire-like;*
2. *is not a Baire space;*
3. *has not a* $\mathscr{C}$*-web.*

If E *and* F *are separable then*

4. $E \otimes_\pi F$ *is a Suslin space.*

We gratefully thank Prof. H. Horváth for his valuable suggestions.

BIBLIOGRAPHY

[1] I. AMEMIYA and Y. KOMURA, Über nich-vollständinge Montelräume, *Math. Ann.*, 177 (1968), 273-277.

[2] N. BOURBAKI, Éléments de Mathématique, Topologie Générale Chap. IX, Hermann, Paris, 1974.

[3] N. BOURBAKI, Éléments de Mathématique, Espaces vectoriels topologiques, *Act. Sci. et Ind.*, Paris, vol. 1929 (1955).

[4] M. DE WILDE, Réseaux dans les espaces linéaires à semi-normes, *Mém. Soc. Royale Sci. Liège*, 18 (5), 2 (1969), 1-144.

[5] G. KÖTHE, Topological Vector Spaces II, Springer-Verlag, New York-Heidelberg-Berlin, 1979.

[6] H. LÖWIG, Über Die dimension linearer Räume, *Studia Math.*, (1934), 18-23.

[7] A. MARTINEAU, Sur le théorème du graphe fermé, *C. R. Acad. Sci.*, Paris, 263 (1966), 870-871.

[8] L. SCHWARTZ, Sur le théorème de graphe fermé. *C.R. Acad. Sci.*, Paris 263 (1966), 602-605.

[9] M. VALDIVIA, Absolutely convex sets in barrelled spaces, *Ann. Inst. Fourier*, Grenoble, 21, 2 (1971), 3-13.

[10] M. VALDIVIA, On suprabarrelled spaces. Functional Analysis, Holomorphy and Approximation Theory. Proocedings, Rio de Janeiro 1978, *Lecture Notes in Math.*, 843 (1981), 572-580, Springer-Verlag, New York-Heidelberg-Berlin.

Manuscrit reçu le 20 août 1981.

Manuel VALDIVIA,
Facultad de Matemáticas
Dr. Moliner s.n.
Burjasot (Valencia) Espagne.