

ANNALES DE L'INSTITUT FOURIER

SALAH BAOUENDI

LINDA P. ROTHSCHILD

Embeddability of abstract CR structures and integrability of related systems

Annales de l'institut Fourier, tome 37, n° 3 (1987), p. 131-141

http://www.numdam.org/item?id=AIF_1987__37_3_131_0

© Annales de l'institut Fourier, 1987, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

EMBEDDABILITY OF ABSTRACT CR STRUCTURES AND INTEGRABILITY OF RELATED SYSTEMS

by

M.S. BAOUENDI and L.P. ROTHCHILD

1. Introduction.

Let M be a smooth real manifold of dimension N , and \mathcal{V} a subbundle of the complex tangent bundle, \mathbf{CTM} , with $\dim \mathcal{V} = n$. We shall say that \mathcal{V} is *integrable* at a point $p_0 \in M$ if there exists a neighborhood Ω_0 of p_0 and smooth functions $\zeta_1, \dots, \zeta_{N-n}$ defined on Ω_0 with linearly independent differentials and satisfying

$$(1.1) \quad L \zeta_k = 0 \quad \text{in } \Omega_0, \quad k = 1, \dots, N - n,$$

for all $L \in \mathbf{L}_0$, where $\mathbf{L}_0 = C^\infty(\Omega_0, \mathcal{V})$, the space of smooth sections of \mathcal{V} over Ω_0 . In this paper we shall give a criterion for local integrability.

We call \mathcal{V} *formally integrable* if

$$(1.2) \quad [\mathcal{V}, \mathcal{V}] \subset \mathcal{V},$$

i.e. if for any sections $L, L' \in \mathbf{L}$, we have $[L, L'] \in \mathbf{L}$, where $\mathbf{L} = C^\infty(M, \mathcal{V})$. The Frobenius theorem then says that formal integrability implies integrability if \mathcal{V} is real (resp. real analytic), i.e. if \mathbf{L} has a basis of real (resp. real analytic) sections. In the general case it is easy to check by dimension that formal integrability is a necessary condition for integrability.

If, in addition, \mathcal{V} satisfies

$$(1.3) \quad \mathcal{V} \cap \overline{\mathcal{V}} = (0)$$

then \mathcal{V} is called an *abstract CR bundle*, and M an *abstract CR manifold*. In this case we have $N = 2n + \ell$ with $\ell \geq 0$. We say that \mathcal{V} is of *codimension* ℓ .

Key-words: Embeddability – CR structures – Complex Lie algebra.

A submanifold of $\mathbf{C}^{n+\ell}$ is a *generic CR manifold* if it is locally given by $\rho_j = 0, j = 1, \dots, \ell$, with ρ_j real valued, smooth, and satisfying $\partial\rho_1, \dots, \partial\rho_\ell$ linearly independent. It can be easily shown that an abstract CR manifold is integrable at p_0 if and only if near p_0 , M can be embedded as a generic CR manifold in $\mathbf{C}^{n+\ell}$, with the image of \mathcal{V} equal to the induced CR bundle i.e. the bundle whose sections are tangential, antiholomorphic vector fields.

For this reason an integrable CR structure is also called *embeddable* or *realizable*. The first example of a nonembeddable strictly pseudoconvex abstract hypersurface was given by Nirenberg [8]. (See also Jacobowitz-Treves [5]).

Our main result is the following :

THEOREM. — *Let M be a smooth manifold and $\mathcal{V} \subset \mathbf{CTM}$ a subbundle satisfying*

$$[\mathbf{L}, \mathbf{L}] \subset \mathbf{L},$$

where $\mathbf{L} = C^\infty(M, \mathcal{V})$. Then \mathcal{V} is locally integrable at $p_0 \in M$ if and only if there exist $\Omega_0 \subset M$, an open neighborhood of p_0 in M , and smooth complex vector fields R_1, \dots, R_ℓ defined in Ω_0 spanning a complex Lie algebra i.e.

$$(1.4) \quad [R_i, R_j] = \sum_{k=1}^{\ell} a_{ijk} R_k, \quad a_{ijk} \in \mathbf{C},$$

and satisfying

$$(1.5) \quad [\mathbf{L}_0, R_j] \subset \mathbf{L}_0, \quad j = 1, \dots, n,$$

with $\mathbf{L}_0 = C^\infty(\Omega_0, \mathcal{V})$, and for every $p \in \Omega_0$

$$(1.6) \quad \mathcal{V}_p + \overline{\mathcal{V}}_p + \mathcal{R}_p + \overline{\mathcal{R}}_p = \mathbf{CT}_p \Omega_0,$$

where \mathcal{V}_p is the fiber of \mathcal{V} at p , and \mathcal{R}_p is the span of the R_j at p . More precisely, if \mathcal{V} is integrable, we may find R_j so that $a_{ijk} = 0$ for all i, j, k and replace (1.6) by

$$(1.7) \quad \mathcal{V}_p + \overline{\mathcal{V}}_p \oplus \mathcal{R}_p = \mathbf{CT}_p \Omega_0.$$

For an integrable structure, the existence of vector fields R_j satisfying conditions similar to (1.4) with $a_{ijk} = 0$, (1.5) and (1.6) was proved and used in Baouendi-Treves [2]. However, the proof we

give here is more natural to the embedding and is used to establish the result for the general case.

For the case where \mathcal{V} is an abstract CR structure, the integrability result generalizes a theorem of Jacobowitz [4] where \mathcal{V} is of codimension one, and a theorem of the authors and Treves [1] for the case where the R_j are real independent vector fields. As in [1], the proof of integrability depends, in the CR case, on the Newlander-Nirenberg theorem [6], and in the general case on a corollary of Nirenberg [7], (see also Hörmander [3] and Treves [9]) which states that \mathcal{V} is integrable if $\mathcal{V} + \overline{\mathcal{V}} = \mathbf{CTM}$; we reprove this result by methods in the spirit of this paper.

Remark. – Note that we do not require the vector fields R_j satisfying (1.4), (1.5) and (1.6) to be linearly independent at every point of Ω_0 . However, when \mathcal{V} is integrable, we may choose them linearly independent, and such that the subbundle \mathcal{R} whose sections are spanned by them is totally real i.e.

$$\overline{\mathcal{R}} = \mathcal{R}.$$

2. Proof of the existence of the R_j .

We assume first that \mathcal{V} is CR i.e. $\mathcal{V} \cap \overline{\mathcal{V}} = \{0\}$. Assume M is integrable at p_0 , so that M may be regarded as a submanifold of $\mathbf{C}^{n+\ell}$ given by

$$(2.1) \quad \rho_j = 0, \quad j = 1, \dots, \ell$$

and $\partial\rho_1, \dots, \partial\rho_\ell$ linearly independent.

By relabeling the coordinates in $\mathbf{C}^{n+\ell}$ we may take $(z, w) \in \mathbf{C}^{n+\ell}$, $w \in \mathbf{C}^\ell$, and assume that

$$(2.2) \quad \rho_w = \begin{pmatrix} \frac{\partial\rho_1}{\partial w_1} & \dots & \frac{\partial\rho_1}{\partial w_\ell} \\ \cdot & & \cdot \\ \cdot & & \cdot \\ \frac{\partial\rho_\ell}{\partial w_1} & \dots & \frac{\partial\rho_\ell}{\partial w_\ell} \end{pmatrix}$$

is invertible near the origin. Similarly, we let

$$(2.3) \quad \rho_z = \begin{pmatrix} \frac{\partial \rho_1}{\partial z_1} & \cdots & \frac{\partial \rho_1}{\partial z_n} \\ \vdots & & \vdots \\ \frac{\partial \rho_\ell}{\partial z_1} & \cdots & \frac{\partial \rho_\ell}{\partial z_n} \end{pmatrix}$$

be an $\ell \times n$ matrix. Then a local basis for $C^\infty(M, \mathcal{V})$ is obtained

$$\text{as } (L) = \begin{pmatrix} L_1 \\ \vdots \\ L_n \end{pmatrix}, \text{ with}$$

$$(L) = \left(\frac{\partial}{\partial \bar{z}} \right) - {}^t \rho_z {}^t \rho_w^{-1} \left(\frac{\partial}{\partial \bar{w}} \right),$$

where we have written $\left(\frac{\partial}{\partial \bar{z}} \right)$ for $\begin{pmatrix} \frac{\partial}{\partial \bar{z}_1} \\ \vdots \\ \frac{\partial}{\partial \bar{z}_n} \end{pmatrix}$ and similarly for $\frac{\partial}{\partial \bar{w}}$.

We have

$$(2.4) \quad \text{PROPOSITION.} - \text{Set } (R) = \begin{pmatrix} R_1 \\ \vdots \\ R_\ell \end{pmatrix} \text{ where}$$

$$(R) = \left(\frac{\partial}{\partial w} \right) - {}^t \rho_w {}^t \rho_w^{-1} \left(\frac{\partial}{\partial w} \right).$$

Then the R_j are tangent to M , commute, and satisfy (1.5), and (1.7).

Proof. - Since $R_j \rho_k = 0$ by construction, the R_j are tangent to M . To prove (1.7) we observe that since $N = 2n + \ell$, and the L_j, \bar{L}_j and R_k are all linearly independent, the result holds by dimension.

For (1.4) and (1.5) we calculate $[L_j, R_k]$ and $[R_j, R_k]$. Each is again tangent to M , and from the form of the L 's and R 's, they contain only $\frac{\partial}{\partial \bar{w}_k}$, and hence are antiholomorphic. Since the L_j form a basis for the tangential antiholomorphic vector fields to M , each $[L_j, R_k]$ and $[R_j, R_k]$ is a linear combination of the L_j 's with smooth coefficients. These coefficients must be zero, since neither commutator contains a term of the form $\frac{\partial}{\partial \bar{z}_p}$. This proves (1.4) (with $a_{ijk} = 0$) and (1.5), and hence Proposition (2.4). □

We now assume that \mathcal{V} is integrable but not necessarily CR. We shall construct the R_j by adding variables in order to reduce to the case of a CR bundle. Let Ω be a small neighborhood of p_0 in M . First choose a basis L_j of $C^\infty(\Omega, \mathcal{V})$ and coordinates (x, y, t, s) in Ω vanishing at p_0 ,

$$x, y \in \mathbf{R}^r, t \in \mathbf{R}^{n-r}, s \in \mathbf{R}^{\varrho}$$

with $\varrho = N - n - r$, such that

$$(2.5) \quad L_j|_{p_0} = \frac{\partial}{\partial z_j} = \frac{1}{2} \left(\frac{\partial}{\partial x_j} + i \frac{\partial}{\partial y_j} \right), \quad 1 \leq j \leq r,$$

and

$$(2.6) \quad L_{j+r}|_{p_0} = \frac{\partial}{\partial t_j}, \quad r+1 \leq j \leq n.$$

We introduce $n-r$ new variables t'_1, \dots, t'_{n-r} and define new vector fields \tilde{L}_j in $\Omega' = \Omega \times \mathbf{R}^{n-r}$ by

$$\tilde{L}_j = L_j, \quad 1 \leq j \leq r,$$

and for $r+1 \leq j \leq n$, \tilde{L}_j is obtained from L_j by replacing $\frac{\partial}{\partial t_j}$ by $\frac{\partial}{\partial t_j} + i \frac{\partial}{\partial t'_j}$. Let \mathcal{V}' be the bundle with sections spanned by the \tilde{L}_j on Ω' . If $\xi_1, \dots, \xi_{r+\varrho}$ is a set of independent solutions for \mathcal{V} , then $\xi_1, \dots, \xi_{r+\varrho}, t_1 + it'_1, \dots, t_{n-r} + it'_{n-r}$ is a set of independent solutions for \mathcal{V}' . Since $\mathcal{V}' \cap \bar{\mathcal{V}}' = \{0\}$, we have proved

(2.7) LEMMA. — \mathcal{V}' is an integrable CR bundle on Ω' .

Let $\tau_j = t_j + it'_j$, $\tau = (\tau_1, \dots, \tau_{n-r})$ and $\zeta = (\zeta_1, \dots, \zeta_{r+\ell})$. The mapping

$$(x, y, t, t', s) \mapsto (\zeta(x, y, t, s), \tau)$$

is an embedding of Ω' onto a CR generic submanifold of $\mathbf{C}^{n+\ell}$. Therefore there exist real smooth functions $\rho_j(Z, \bar{Z})$ in $\mathbf{C}^{n+\ell}$ so that locally the image of Ω' is given by $\rho_j = 0, j = 1, \dots, \ell$, with $\partial\rho_1, \dots, \partial\rho_\ell$ linearly independent. Hence we have for $j = 1, \dots, \ell$

$$(2.8) \quad \rho_j(\zeta(x, y, t, s), \tau, \overline{\zeta(x, y, t, s)}, \bar{\tau}) \equiv 0$$

in Ω' .

We may assume that $\zeta(0) = 0$. If $Z_1, \dots, Z_{n+\ell}$ are the variables in $\mathbf{C}^{n+\ell}$, we write τ_k for $Z_{k+r+\ell}, k = 1, \dots, n-r$.

(2.9) LEMMA. — *We may assume that the ρ_j are independent of t'_k . Also we have for $j = 1, \dots, \ell$ and $k = 1, \dots, n-r$*

$$\frac{\partial \rho_j}{\partial \tau_k}(0) = 0.$$

Proof. — It suffices to differentiate (2.8) with respect to t_k and t'_k , and to use (2.6) and the fact that the ζ_j satisfy the equations

$$L_p \zeta_k = 0 \quad 1 \leq p \leq n, \quad 1 \leq k \leq r + \ell.$$

This proves the lemma. □

Since the ρ_j have independent complex differentials, the matrix

$$\begin{bmatrix} \rho_{1z_1} & \dots & \rho_{1z_{\ell+r}} & \rho_{1\tau_1} & \dots & \rho_{1\tau_{n-r}} \\ \rho_{2z_1} & \dots & \rho_{2z_{\ell+r}} & \rho_{2\tau_1} & \dots & \rho_{2\tau_{n-r}} \end{bmatrix}$$

has rank ℓ , therefore by Lemma (2.9) the submatrix

$$\left[\begin{array}{c} \partial \rho_j \\ \partial \bar{z}_k \end{array} \right]_{1 \leq j \leq \ell, 1 \leq k \leq \ell+r}$$

must have rank ℓ at 0 . Hence we may find new coordinates $(z, w) \in \mathbf{C}^r \times \mathbf{C}^{\ell}$ such that the matrix $\begin{bmatrix} \frac{\partial \rho}{\partial w} \end{bmatrix}$ is invertible at 0 . In these coordinates we may find a basis for \mathfrak{V}' in the form $(\tilde{L}) = \begin{pmatrix} \tilde{L}' \\ \tilde{L}'' \end{pmatrix}$, where

$$(2.11) \quad (\tilde{L}') = \left(\frac{\partial}{\partial \bar{z}} \right) - {}^t \rho_{\bar{z}} {}^t \rho_{\bar{w}}^{-1} \left(\frac{\partial}{\partial \bar{w}} \right),$$

and

$$(2.12) \quad (\tilde{L}'') = \left(\frac{\partial}{\partial \bar{\tau}} \right) - {}^t \rho_{\bar{\tau}} {}^t \rho_{\bar{w}}^{-1} \left(\frac{\partial}{\partial \bar{w}} \right),$$

where we use the notation conventions of § 2. Restricting to $t' = 0$ we find a basis (L) for \mathfrak{V} given by $(L) = \begin{pmatrix} L' \\ L'' \end{pmatrix}$:

$$(2.13) \quad (L') = \left(\frac{\partial}{\partial \bar{z}} \right) - {}^t \rho_{\bar{z}} {}^t \rho_{\bar{w}}^{-1} \left(\frac{\partial}{\partial \bar{w}} \right),$$

and

$$(2.14) \quad (L'') = \left(\frac{\partial}{\partial t} \right) - {}^t \rho_t {}^t \rho_{\bar{w}}^{-1} \left(\frac{\partial}{\partial \bar{w}} \right).$$

Now put

$$(R) = \left(\frac{\partial}{\partial w} \right) - {}^t \rho_w {}^t \rho_{\bar{w}}^{-1} \left(\frac{\partial}{\partial \bar{w}} \right)$$

as before. □

3. Proof of Integrability.

We now assume $\{R_j\}$ exist satisfying (1.4), (1.5), and (1.6) and prove \mathfrak{V} is integrable. First we give a new proof of the following result of Nirenberg [7].

(3.1) PROPOSITION. — *If \mathfrak{V} is a formally integrable subbundle of CTM for which*

$$(3.2) \quad \mathfrak{R}^{\mathfrak{Q}} + \overline{\mathfrak{R}^{\mathfrak{Q}}} = \mathbf{CTM},$$

then $\mathfrak{R}^{\mathfrak{Q}}$ is locally integrable.

Proof. — Let Ω be a small neighborhood of $p_0 \in \mathbf{M}$, and V_1, V_2, \dots, V_n be a commuting basis for $C^\infty(\Omega, \mathfrak{R}^{\mathfrak{Q}})$. After renumbering and multiplication by complex numbers we may assume V_1, \dots, V_r is a maximal set for which $V_1, \dots, V_r, \bar{V}_1, \dots, \bar{V}_r$ is linearly independent at p_0 , and that these, together with $\operatorname{Re} V_j, j > r$, span the section of $\mathbf{CT}\Omega$. Now let $\tilde{\mathfrak{R}}^{\mathfrak{Q}}$ be the bundle over $\Omega \times \mathbf{R}^{n-r}$ whose sections are spanned by $\tilde{V}_j = V_j, 1 \leq j \leq r$, and $\tilde{V}_j = V_j + i \frac{\partial}{\partial t_{r-j}}, j = r + 1, \dots, n$. Then $\tilde{\mathfrak{R}}^{\mathfrak{Q}}$ satisfies the conditions of the Newlander-Nirenberg theorem [6] since

$$\tilde{\mathfrak{R}}^{\mathfrak{Q}} \cap \overline{\tilde{\mathfrak{R}}^{\mathfrak{Q}}} = (0).$$

Hence there exist n solutions $f_1(u, t), \dots, f_n(u, t)$ for $\tilde{\mathfrak{R}}^{\mathfrak{Q}}$, where (u) is a coordinate system near p_0 in Ω vanishing at p_0 , and t is in a neighborhood of 0 in \mathbf{R}^{n-r} . We may assume $f_j(0) = 0, j = 1, \dots, n$.

We shall obtain solutions for $\mathfrak{R}^{\mathfrak{Q}}$ in the form

$$\zeta_k = F_k(f_1, \dots, f_n),$$

where each $F_k(Z)$ is holomorphic and satisfies

$$(3.3) \quad \frac{\partial}{\partial t_j} [F_k(f_1(u, t), \dots, f_n(u, t))] \equiv 0, \quad j = 1, \dots, n - r.$$

We shall prove that there exist F_1, \dots, F_r holomorphic satisfying (3.3) with linearly independent differentials. Indeed, for F holomorphic

$$(3.4) \quad \frac{\partial}{\partial t_j} F(f_1, \dots, f_n) = \sum_{p=1}^n \frac{\partial f_p}{\partial t_j} \frac{\partial F}{\partial Z_p}(f_1, \dots, f_n).$$

Since we may choose a basis for $\tilde{\mathfrak{R}}^{\mathfrak{Q}}$ taking vector fields with coefficients independent of the $t_j, \frac{\partial f_p}{\partial t_j}$ is again a solution for $\tilde{\mathfrak{R}}^{\mathfrak{Q}}$.

Hence there exists a holomorphic function H_{pj} such that

$$(3.5) \quad \frac{\partial f_p}{\partial t_j} = H_{pj}(f_1, \dots, f_n), \quad 1 \leq p \leq n, \quad 1 \leq j \leq n - r.$$

Substituting (3.4) and (3.5) into (3.3) we obtain the system

$$(3.6) \quad \sum_{p=1}^n H_{pj}(Z) \frac{\partial F}{\partial Z_p}(Z) = 0, \quad j = 1, \dots, n - r.$$

Since $df_1, \dots, df_n, d\bar{f}_1, \dots, d\bar{f}_n$ are linearly independent we conclude that the matrix

$$\left(\frac{\partial f_p}{\partial t_j} \right), \quad 1 \leq p \leq n, \quad 1 \leq j \leq n - r,$$

is of rank $n - r$. Therefore by (3.5) the same is true for the matrix (H_{pj}) at the origin. It follows by the Cauchy-Kovalevsky Theorem that there are $n - (n - r) = r$ linearly independent solution F_k of (3.6) near 0. Hence the functions

$$\zeta_k(u) = F_k(f_1(u, t), \dots, f_n(u, t)), \quad 1 \leq k \leq r,$$

provide a system of solutions for $\mathfrak{R}^{\mathfrak{Q}}$, proving integrability. □

We may now complete the proof of the theorem. We assume we are given the R_j satisfying (1.4), (1.5) and (1.6). We let S_1, \dots, S_ϱ be a basis for an abstract complex Lie algebra satisfying the same commutation relations as the R_j i.e.

$$(3.7) \quad [S_i, S_j] = \sum_{k=1}^{\varrho} a_{ijk} S_k.$$

By introducing local exponential coordinates on any corresponding connected complex Lie group we may find coordinates in an open neighborhood \mathfrak{O} of 0 in \mathbf{C}^ϱ near 0 in which we may represent the S_j as holomorphic vector fields with holomorphic coefficients i.e.

$$(3.8) \quad S_j = \sum_{k=1}^{\varrho} a_{jk}(t) \frac{\partial}{\partial t_k}$$

with $t_k = t'_k + it''_k \in \mathbf{C}$ and the matrix (a_{jk}) is invertible. Now we let $R'_j = R_j + S_j$. We claim that the bundle $\tilde{\mathfrak{V}}$ over $\Omega \times \mathfrak{O}$

spanned by \mathfrak{V} , $\{R'_j\}_{1 \leq j \leq \ell}$ and $\left\{ \frac{\partial}{\partial \bar{t}_k} \right\}_{1 \leq k \leq \ell}$ satisfies the condition of Proposition (3.1) for integrability.

Indeed, note that the S_j commute with $\frac{\partial}{\partial \bar{t}_j}$, as well as the R_j and L_0 . Hence

$$(3.9) \quad [R_i + S_i, R_j + S_j] = \sum a_{ijk} (R_k + S_k),$$

which proves that $\tilde{\mathfrak{V}}$ is formally integrable. Also, the span of the $\tilde{R}_j, \bar{\tilde{R}}_j, \frac{\partial}{\partial t_j}$ and $\frac{\partial}{\partial \bar{t}_j}$ is the same as that of the $R_j, \bar{R}_j, \frac{\partial}{\partial t_j}$ and $\frac{\partial}{\partial \bar{t}_j}$. Hence $\tilde{\mathfrak{V}}$ satisfies condition (3.2). By Proposition (3.1) there exist $N - n = N + 2\ell - (n + 2\ell)$ solutions $f_k(u, t', t'')$ which have linearly independent differentials.

Now let $\zeta_k(u) = f_k(u, 0, 0), k = 1, \dots, N - n$. Since the coefficients of elements of L are independent of (t', t'') , it is clear that the ζ_k are solutions of (1.1). It suffices to check that the ζ_k have linearly independent differentials. This will follow if the matrix $\left(\frac{\partial f_i}{\partial u_k} \right)_{\substack{1 \leq j \leq N-n \\ 1 \leq k \leq N}}$ has rank $N - n$. By the linear independence of the

f_k in the (u, t', t'') variables, it suffices to show that $\frac{\partial f_k}{\partial t'_j}$ and $\frac{\partial f_k}{\partial t''_j}$ are linear combinations of $\frac{\partial f_k}{\partial u}$. Since $\frac{\partial f_k}{\partial \bar{t}_i} = 0$ and $(R_j + S_j) f_k = 0, 1 \leq j \leq \ell$, this follows, and hence the proof of the theorem is complete.

□

BIBLIOGRAPHIE

- [1] M.S. BAOUENDI, L.P. ROTHSCHILD and F. TREVES, CR structures with group action and extendability of CR functions, *Invent. Math.*, 83 (1985), 359-396.

- [2] M.S. BAOUENDI and F. TREVES, A property of the functions and distributions annihilated by a locally integrable system of complex vector fields, *Ann. of Math.*, 113 (1981), 387-421.
- [3] L. HÖRMANDER, The Frobenius-Nirenberg theorem, *Arkiv För Mat.*, 5-29 (1964), 425-432.
- [4] H. JACOBOWITZ, The canonical bundle and realizable CR hypersurfaces, preprint.
- [5] H. JACOBOWITZ and F. TREVES, Non-realizable CR structures, *Invent. Math.*, 66 (1982), 231-149.
- [6] A. NEWLANDER and L. NIRENBERG, Complex coordinates in almost complex manifolds, *Ann. of Math.*, (2) 65 (1957), 391-404.
- [7] L. NIRENBERG, A Complex Frobenius Theorem, *Seminar on analytic functions I*, Princeton, (1957), 172-189.
- [8] L. NIRENBERG, On a question of Hans Lewy, *Russian Math. Surveys*, 29 (1974), 251-262.
- [9] F. TREVES, Approximation and Representation of Functions and Distributions Annihilated by a System of Complex Vector Fields, Ecole Polytechnique, Palaiseau, France, (1981).

Manuscrit reçu le 23 septembre 1986
révisé le 12 novembre 1986.

M.S. BAOUENDI,
Purdue University
Dept. of Mathematics
Mathematical Sciences Building
West Lafayette, IN 47907 (USA)

&

L.P. ROTHSCHILD,
University of California, San Diego
Dept. of Mathematics
La Jolla, CA 92093 (USA).