

ANNALES DE L'INSTITUT FOURIER

TEIMURAZ PIRASHVILI

On Leibniz homology

Annales de l'institut Fourier, tome 44, n° 2 (1994), p. 401-411

http://www.numdam.org/item?id=AIF_1994__44_2_401_0

© Annales de l'institut Fourier, 1994, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON LEIBNIZ HOMOLOGY

by Teimuraz PIRASHVILI

Introduction.

Let \mathfrak{g} be a Lie algebra over a field k and let M be a right \mathfrak{g} -module. Leibniz homology $HL_*(\mathfrak{g}, M)$ was defined by Jean-Louis Loday (see 10.6 in [CH]). Moreover, HL_* is well-defined on a much bigger category – the so called Leibniz algebras and their corepresentations (see [LP]). The purpose of this paper is to construct a few spectral sequences. These give a relation between Leibniz and ordinary homologies for Lie algebras and relate Leibniz homology of a given Leibniz algebra \mathfrak{h} with its Liezation $\mathfrak{h}_{\text{Lie}}$. As a consequence we compute $HL_*(\mathfrak{g}, k)$, where \mathfrak{g} is a semi-simple Lie algebra or has the dimension 2 or 3 ($\text{char } k = 0$), or is free Lie algebra. We also prove that in some cases (see Proposition 4.3) one has an isomorphism $HL_*(\mathfrak{h}, k) \cong HL_*(\mathfrak{h}_{\text{Lie}}, k)$.

The author would like to thank Friedhelm Waldhausen for hospitality at the University of Bielefeld while this paper was being written. The author is also indebted to Jean-Louis Loday for many useful discussions.

1. Relation between Chevalley-Eilenberg homology and Leibniz homology for Lie algebras.

Let M be a right module over the Lie algebra \mathfrak{g} . We denote

$$\begin{aligned} C_*(\mathfrak{g}, M) &= (M \otimes \Lambda^* \mathfrak{g}, d') \\ CL_*(\mathfrak{g}, M) &= (M \otimes T^* \mathfrak{g}, d), \end{aligned}$$

where T^*g is the tensor algebra over \mathfrak{g} , d' is the Chevalley-Eilenberg boundary map, and d is the Loday boundary map (see 10.6.2.1 in [CH]). The natural epimorphism

$$CL_*(\mathfrak{g}, M) \longrightarrow C_*(\mathfrak{g}, M)$$

is a morphism of chain complexes, which is an isomorphism in dimensions 0 and 1. We define $C_*^{\text{rel}}(\mathfrak{g}, M)$ such that $C_*^{\text{rel}}(\mathfrak{g}, M)[2]$ is the kernel of the above epimorphism. Let

$$H_*^{\text{rel}}(\mathfrak{g}, M) = H_*(C_*^{\text{rel}}(\mathfrak{g}, M)).$$

We recall that $HL_*(\mathfrak{g}, M) = H_*(CL_*(\mathfrak{g}, M))$ and $H_*(\mathfrak{g}, M) = H_*(C_*(\mathfrak{g}, M))$. Thus, by definition we have

PROPOSITION 1.1. — *One has an exact sequence*

$$\begin{aligned} 0 \longleftarrow H_2(\mathfrak{g}, M) \longleftarrow HL_2(\mathfrak{g}, M) \longleftarrow H_0^{\text{rel}}(\mathfrak{g}, M) \\ \longleftarrow H_3(\mathfrak{g}, M) \longleftarrow HL_3(\mathfrak{g}, M) \longleftarrow \dots \end{aligned}$$

and isomorphisms

$$HL_i(\mathfrak{g}, M) \cong H_i(\mathfrak{g}, M), \quad \text{for } i = 0, 1.$$

The product map $m : \mathfrak{g} \otimes \Lambda^n \mathfrak{g} \longrightarrow \Lambda^{n+1} \mathfrak{g}$ in the exterior algebra yields an epimorphism of chain complexes

$$C_*(\mathfrak{g}, \mathfrak{g}) \longrightarrow \overline{C}_*(\mathfrak{g}, k)[-1].$$

Here $\overline{C}_*(\mathfrak{g}, k)$ is the reduced chain complex :

$$\overline{C}_0(\mathfrak{g}, k) = 0, \overline{C}_i(\mathfrak{g}, k) = C_i(\mathfrak{g}, k), i > 0.$$

We define $CR_*(\mathfrak{g})$ such that $CR_*(\mathfrak{g})[1]$ is the kernel of $C_*(\mathfrak{g}, \mathfrak{g}) \longrightarrow \overline{C}_*(\mathfrak{g}, k)[-1]$. We denote the homology of $CR_*(\mathfrak{g})$ by $HR_*(\mathfrak{g})$. Then, by definition we have

PROPOSITION 1.2. — *One has an exact sequence*

$$\begin{aligned} 0 \longleftarrow H_2(\mathfrak{g}, k) \longleftarrow H_1(\mathfrak{g}, \mathfrak{g}) \longleftarrow HR_0(\mathfrak{g}) \longleftarrow H_3(\mathfrak{g}, k) \\ \longleftarrow H_2(\mathfrak{g}, \mathfrak{g}) \longleftarrow HR_1(\mathfrak{g}) \longleftarrow \dots \end{aligned}$$

As usual, we denote the kernel of $\otimes^2 \mathfrak{g} \rightarrow \Lambda^2 \mathfrak{g}$ by $\Gamma \mathfrak{g}$. Thus $\Gamma \mathfrak{g}$ is spanned by elements $x \otimes x$. Moreover, $\Gamma \mathfrak{g}$ is a \mathfrak{g} -submodule of $\mathfrak{g} \otimes \mathfrak{g}$ and the restriction of

$$1 \otimes m : \mathfrak{g} \otimes \mathfrak{g} \otimes \Lambda^i \mathfrak{g} \rightarrow \mathfrak{g} \otimes \Lambda^{i+1} \mathfrak{g}$$

to $\Gamma \mathfrak{g} \otimes \Lambda^i \mathfrak{g}$ gives an epimorphism of chain complexes

$$C_*(\mathfrak{g}, \Gamma \mathfrak{g}) \rightarrow CR_*(\mathfrak{g}),$$

which is an isomorphism in dimension 0. Hence $HR_0(\mathfrak{g}) \cong H_0(\mathfrak{g}, \Gamma \mathfrak{g})$ and for the cohomological version we have

$$HR^0(\mathfrak{g}) \cong \{\text{invariant quadratic forms on } \mathfrak{g}\}.$$

It follows from Theorem A that the same is also true for $H_0^{\text{rel}}(\mathfrak{g}, k)$ and $H_{\text{rel}}^0(\mathfrak{g}, k)$. Moreover, the corresponding map

$$\{\text{invariant quadratic forms on } \mathfrak{g}\} \rightarrow H^3(\mathfrak{g}, k)$$

coincides with the classical homomorphism of Koszul (see section 11 of [K]).

THEOREM A. — Let \mathfrak{g} be a Lie algebra and M be a right \mathfrak{g} -module. Then there exists a spectral sequence

$$E_{pq}^2 = HR_p(\mathfrak{g}) \otimes HL_q(\mathfrak{g}, M) \implies H_{p+q}^{\text{rel}}(\mathfrak{g}, M).$$

Proof. — We consider the following filtration on $C_*^{\text{rel}}(\mathfrak{g}, M)$. Let $\omega \in CL_n(\mathfrak{g}, M)$ and $x \in \mathfrak{g}$. Then it follows from 10.6.3.1 of [CH] that $d(\omega, x, x) = (d\omega, x, x)$. Here we write (α, y) instead of $\alpha \otimes y$. Thus the subspaces $\mathcal{F}_n^0 = C_n(\mathfrak{g}, M) \otimes \Gamma \mathfrak{g} \subset C_n^{\text{rel}}(\mathfrak{g}, M)$ give us a subcomplex and

$$H_*(\mathcal{F}_*^0) \cong HL_*(\mathfrak{g}, M) \otimes \Gamma \mathfrak{g}.$$

More generally, for $\rho = (x_1, \dots, x_i)$ and $\rho_j = (x_1, \dots, \hat{x}_j, \dots, x_i)$, $1 \leq j \leq i$, one has

$$\begin{aligned} d(\omega, x, x, \rho) &= (d\omega, x, x, \rho) + (-1)^{\deg \omega} (\omega, x, x, d\rho) \\ &+ \sum (-1)^{\deg \omega + j} ([\omega, x_j], x, x, \rho_j) \\ &+ \sum (-1)^{\deg \omega + j} (\omega, ([x, x_j], x) + (x, [x, x_j])), \rho_j). \end{aligned}$$

Since $(y, z) + (z, y) \in \Gamma \mathfrak{g}$, $y, z \in \mathfrak{g}$ we conclude that $\mathcal{F}_*^0 \subset \mathcal{F}_*^1 \subset \dots$ are subcomplexes of $C_*^{\text{rel}}(\mathfrak{g}, M)$ with $\mathcal{F}_n^n = C_n^{\text{rel}}(\mathfrak{g}, M)$ and

$$\mathcal{F}_*^s / \mathcal{F}_*^{s-1} \cong CL_*(\mathfrak{g}, M)[-s] \otimes (\text{Ker}(\otimes^{s+2} \mathfrak{g} \longrightarrow \Lambda^{s+2} \mathfrak{g}) / (\mathfrak{g} \otimes \text{Ker}(\otimes^{s+1} \mathfrak{g} \longrightarrow \Lambda^{s+1} \mathfrak{g}))).$$

Here $\mathcal{F}_n^s = CL_{n-s}(\mathfrak{g}, M) \otimes \text{Ker}(\otimes^{s+2} \mathfrak{g} \longrightarrow \Lambda^{s+2} \mathfrak{g})$ is spanned by $(\omega, x, x, x_1, \dots, x_i)$ with $i \leq s$. Thus the corresponding spectral sequence has the form :

$$E_{pq}^1 = HL_q(\mathfrak{g}, M) \otimes CR_p(\mathfrak{g})$$

because there is a natural isomorphism

$$\begin{aligned} \ker(\otimes^{n+2} \mathfrak{g} \longrightarrow \Lambda^{n+2} \mathfrak{g}) / \ker(\otimes^{n+2} \mathfrak{g} \longrightarrow \mathfrak{g} \otimes \Lambda^{n+1} \mathfrak{g}) \\ \cong \ker(\mathfrak{g} \otimes \Lambda^{n+1} \mathfrak{g} \longrightarrow \Lambda^{n+2} \mathfrak{g}). \end{aligned}$$

Hence the theorem follows from Proposition 10.1.7 of [CH] and from the above formula for d .

COROLLARY 1.3. — *If $H_*(\mathfrak{g}, M) = 0$, then $HL_*(\mathfrak{g}, M) = 0$.*

Proof. — It follows from Proposition 1.1 that $HL_i(\mathfrak{g}, M) = 0$ for $i = 0, 1$ and $HL_{n+2}(\mathfrak{g}, M) \cong H_n^{\text{rel}}(\mathfrak{g}, M)$ for $n \geq 0$. We prove by induction that $H_n^{\text{rel}}(\mathfrak{g}, M) = 0$. It holds when $n = 0$ because $E_{pq}^2 = 0$ for $q = 0, 1$ and Theorem A implies $H_n^{\text{rel}}(\mathfrak{g}, M) = 0$ for $n = 0, 1$. If we assume $H_i^{\text{rel}}(\mathfrak{g}, M) = 0$ for all $i \leq n$, then $HL_i(\mathfrak{g}, M) = 0$ for $i \leq n + 2$. Thus $E_{pq}^2 = 0$ for $q \leq n + 2$ and $H^{\text{rel}}(\mathfrak{g}, M) = 0$ for $i \leq n + 2$.

The cohomological versions of Theorem A and Corollary 3.3 are still true with the same proofs (of course now one will consider $C^*(\mathfrak{g}, \mathfrak{g}^*)$ instead of $C_*(\mathfrak{g}, \mathfrak{g})$, where \mathfrak{g}^* is the dual of \mathfrak{g}). There are well-known examples which satisfy the assumption of Corollary 1.3 (see for example Theorem 1 of [D] and the main result of [T]).

In the next section we consider the case when \mathfrak{g} is a finite dimensional semi-simple Lie algebra over a characteristic 0 field. Now as a sample application of Theorem A we compute

$$HL_* \mathfrak{g} := HL_*(\mathfrak{g}, k)$$

where \mathfrak{g} is two or three dimensional Lie algebra defined over the complex numbers.

Examples 1.4. — i) Let $\mathfrak{g} = \mathfrak{r}_2, \mathfrak{r}_3$ or $\mathfrak{r}_{3,\lambda}$, where $\lambda \neq -1, \frac{1}{2}$. Here \mathfrak{r}_2 is the nonabelian two dimensional Lie algebra, \mathfrak{r}_3 and $\mathfrak{r}_{3,\lambda}$ are three

dimensional Lie algebras with

$$\mathfrak{r}_3 : [x, y] = y, [x, z] = y + z$$

$$\mathfrak{r}_{3,\lambda} : [x, y] = y, [x, z] = \lambda z, \quad 0 < |\lambda| \leq 1.$$

Then $H_i(\mathfrak{g}) = 0$ for $i > 1$. Thus $HL_{n+2}(\mathfrak{g}) \cong H_n^{\text{rel}}(\mathfrak{g}), n \geq 0$. By the direct calculations we obtain : $HR_i(\mathfrak{g}) = 0$ for $i > 0$ and $\dim HR_0(\mathfrak{g}) = 1$. It follows from Theorem A that $H_n^{\text{rel}}(\mathfrak{g}) \cong HL_n(\mathfrak{g})$. Hence $\dim HL_n(\mathfrak{g}) = 1, n \geq 0$.

ii) Let $\mathfrak{g} = \mathfrak{r}_{3,\lambda}$ with $\lambda = -1$. In this case we still have $HR_i(\mathfrak{g})=0$ for $i > 0$ and $\dim HR_0(\mathfrak{g}) = 1$, but now $\dim H_i(\mathfrak{g}) = 1$, for $0 \leq i \leq 3$, and $H_i(\mathfrak{g}) = 0, i > 3$. Easy calculation shows that the connected homomorphism $H_3(\mathfrak{g}) \rightarrow H_0^{\text{rel}}(\mathfrak{g}) = HR_0(\mathfrak{g})$ is zero. Thus for $x_n := \dim HL_n(\mathfrak{g})$ one has $x_0 = 1, x_1 = 1, x_n = 2, n \geq 2$.

iii) Let $\mathfrak{g} = \mathfrak{r}_{3,\lambda}$ with $\lambda = \frac{1}{2}$. In this case $H_i(\mathfrak{g}) = 0$ for $i > 1$, $\dim HR_i(\mathfrak{g}) = 1$ for $i = 0, 1, 2$ and $HR_i(\mathfrak{g}) = 0$ for $i > 2$. Thus $HL_{n+2}(\mathfrak{g}) \cong H_n^{\text{rel}}(\mathfrak{g}), E_{pq}^2 = 0$ for $p > 2$ and $E_{pq}^2 = HL_q(\mathfrak{g})$ for $p = 0, 1, 2$. We claim that d_2 and hence all differentials are zero. Indeed, this follows from the fact that the class of $z \otimes (z \wedge x \wedge y)$ in $HR_2(\mathfrak{g})$ is a nonzero element and if ω is some element of $CL_*(\mathfrak{g})$ with $d\omega = 0$ then $d\rho = 0$. Here $\rho = (\omega, z, z, x, y) - (\omega, x, z, z, y) + (\omega, y, z, z, x) - (\omega, y, x, z, z)$. Thus $x_0 = 1, x_1 = 1, x_2 = 1, x_3 = 2, x_n = x_{n-2} + x_{n-3} + x_{n-4}, n \geq 4$, where $x_n = \dim HL_n(\mathfrak{g})$.

iv) Let $\mathfrak{g} = \mathfrak{n}_3$ be the three dimensional nilpotent Lie algebra : $[x, y] = z$. Then $\dim H_1(\mathfrak{g}) = 2, \dim H_2(\mathfrak{g}) = 2, \dim H_3(\mathfrak{g}) = 1$ and $\dim HR_0(\mathfrak{g}) = 3, \dim HR_1(\mathfrak{g}) = 3, \dim HR_2(\mathfrak{g}) = 1$. The connected homomorphism $H_3(\mathfrak{g}) \rightarrow HR_0(\mathfrak{g})$ as well as all differentials of the spectral sequence are still zero. To verify the last fact we remark that $z \otimes (z \wedge x \wedge y)$ gives a nonzero element in $HR_2(\mathfrak{g})$ and if $\omega \in CL_*(\mathfrak{g})$ is an element with $d\omega = 0$, then $d\rho = 0$, where $\rho = (\omega, z, z, x, y) - (\omega, x, z, z, y) + (\omega, y, z, z, x) + (\omega, x, y, z, z)$. Hence for $x_n := \dim HL_n(\mathfrak{g})$ we have $x_0 = 1, x_1 = 2, x_2 = 5, x_3 = 10$ and $x_n = 3x_{n-2} + 3x_{n-3} + x_{n-4}, n \geq 4$.

2. Semi-simple Lie algebras as Leibniz algebras.

In this section we assume that $\text{char } k = 0$ and \mathfrak{g} is a finite dimensional semi-simple Lie algebra. Proposition 2.1 was proved independently by Ntolo [N]. She used Casimir element to construct the explicit homotopy.

PROPOSITION 2.1. — *Let M be a finite dimensional right \mathfrak{g} -module and A be a finite dimensional corepresentation of \mathfrak{g} (see 1.5 and 1.11 of [LP]). Then*

$$\begin{aligned} HL_i(\mathfrak{g}, M) &= 0, \quad \text{if } i > 0 \\ HL_i(\mathfrak{g}, A) &= 0, \quad \text{if } i > 1. \end{aligned}$$

Proof. — It follows from the dual of Proposition 5.6 of [HS] and Corollary 1.3 that $HL_*(\mathfrak{g}, M) = 0$ if M is a nontrivial simple \mathfrak{g} -module. Thus, for arbitrary M , one has

$$HL_*(\mathfrak{g}, M) = HL_*(\mathfrak{g}, k) \otimes M_{\mathfrak{g}}.$$

But $HL_{i+1}(\mathfrak{g}, k) \cong HL_i(\mathfrak{g}, \mathfrak{g})$ (see Exercise E.10.6.1 of [CH] or take $C = k$ in Lemma 2.2) and $\mathfrak{g}_{\mathfrak{g}} = 0$. Thus $HL_i(\mathfrak{g}, k) = 0$ for $i \geq 1$. This completes the proof in the case of right modules. Let A be a corepresentation. Then there exists an extension (compare with the corresponding extension for representations in 1.10 of [LP])

$$0 \longrightarrow B^s \longrightarrow A \longrightarrow C^a \longrightarrow 0$$

such that B^s (resp. C^a) is an (anti)symmetric corepresentation and the proposition follows from Lemma 2.2.

LEMMA 2.2. — *Let \mathfrak{h} be a Leibniz algebra. Let B^s (resp. C^a) be an (anti)symmetric corepresentation of \mathfrak{h} . Let B and C be the underlying right \mathfrak{h} -modules. Then $HL_*(\mathfrak{h}, B^s) = HL_*(\mathfrak{h}, B)$, $HL_0(\mathfrak{h}, C^a) = C$ and*

$$HL_{i+1}(\mathfrak{h}, C^a) \cong HL_i(\mathfrak{h}, C \otimes \mathfrak{h}),$$

where $C \otimes \mathfrak{h}$ is considered as right \mathfrak{h} -module by the action :

$$[c \otimes h, x] = -[x, c] \otimes h + c \otimes [h, x], \quad h, x \in \mathfrak{h}, c \in C.$$

Proof. — One checks that the isomorphisms exist on the level of chain complexes.

The cohomological version of Proposition 2.1 is still true. As for Lie algebras one verifies that for an arbitrary Leibniz algebra \mathfrak{h} the space $H^2(\mathfrak{h}, \mathfrak{h})$ in the sense of 1.8 of [LP] is a space of obstructions for deformations in the category of Leibniz algebras. Thus we deduce from Proposition 2.1, Corollary 1.3 and [T] the following

PROPOSITION 2.3. — *Let \mathfrak{p} be a parabolic subalgebra of \mathfrak{g} . Then \mathfrak{p} as well as \mathfrak{g} is rigid in the category of all Leibniz algebras.*

PROPOSITION 2.4. — *Let $f : \mathfrak{h} \rightarrow \mathfrak{g}$ be an epimorphism from an arbitrary finite dimensional Leibniz algebra \mathfrak{h} to the semi-simple Lie algebra \mathfrak{g} . Then f has a section.*

Proof. — Let $\mathfrak{h}_{\text{Lie}} = \mathfrak{h}/([x, x] = 0)$ be the Liezation of \mathfrak{h} . Let $\text{ch}(\mathfrak{h}) \in H^2(\mathfrak{h}_{\text{Lie}}, \mathfrak{h}^{\text{ann}})$ be the characteristic element of \mathfrak{h} (see 1.10 in [LP]), corresponding to the extension

$$0 \longrightarrow \mathfrak{h}^{\text{ann}} \longrightarrow \mathfrak{h} \longrightarrow \mathfrak{h}_{\text{Lie}} \longrightarrow 0.$$

By definition f factors through $f_1 : \mathfrak{h}_{\text{Lie}} \rightarrow \mathfrak{g}$. It follows from Levi theorem that f_1 has a section s . By Proposition 2.1 we have

$$s^*(\text{ch}(\mathfrak{h})) \in HL^2(\mathfrak{g}, \mathfrak{h}^{\text{ann}}) = 0.$$

Thus s has a lifting $\mathfrak{g} \rightarrow \mathfrak{h}$.

3. Free Lie algebra approach.

Let \mathfrak{f} be a free Lie algebra and M be a right \mathfrak{f} -module. Since $H_i(\mathfrak{f}, -) = 0$, if $i > 1$, we deduce from Propositions 1.1 and 1.2 that $HL_{n+2}(\mathfrak{f}, M) \cong H_n^{\text{rel}}(\mathfrak{f}, M), n \geq 0$; $HR_i(\mathfrak{f}) = 0, i > 0$ and $HR_0(\mathfrak{f}) \cong H_1(\mathfrak{f}, \mathfrak{f})$. Then Theorem A gives

$$H_n^{\text{rel}}(\mathfrak{f}, M) \cong H_1(\mathfrak{f}, \mathfrak{f}) \otimes HL_n(\mathfrak{f}, M).$$

Thus we prove the following proposition.

PROPOSITION 3.1. — Let \mathbf{f} be a free Lie algebra and M be a right \mathbf{f} -module. Then

$$\begin{aligned} HL_n(\mathbf{f}, M) &\cong H_1(\mathbf{f}, \mathbf{f})^{\otimes k} \otimes H_0(\mathbf{f}, M), \quad \text{if } n = 2k \\ &\cong H_1(\mathbf{f}, \mathbf{f})^{\otimes k} \otimes H_1(\mathbf{f}, M), \quad \text{if } n = 2k + 1. \end{aligned}$$

THEOREM B. — Let \mathbf{g} be a Lie algebra and M be a right \mathbf{g} -module. Then there exists a spectral sequence $E_{pq}^r \implies HL_{p+q}(\mathbf{g}, M)$ with

$$\begin{aligned} E_{p,2k}^2 &= \bigoplus HR_{i_1}(\mathbf{g}) \otimes \dots \otimes HR_{i_k}(\mathbf{g}) \otimes H_0(\mathbf{g}, M) \\ E_{p,2k+1}^2 &= \bigoplus HR_{i_1}(\mathbf{g}) \otimes \dots \otimes HR_{i_k}(\mathbf{g}) \otimes H_{i_{k+1}+1}(\mathbf{g}, M) \end{aligned}$$

where the sum in the first (resp. second) formula is taken over all $i_1 + \dots + i_k = p$ (resp. $i_1 + \dots + i_{k+1} = p$).

Proof. — Let \mathbf{f}_* be a componentwise free simplicial Lie algebra, such that $\pi_0 \mathbf{f}_* = \mathbf{g}$ and $\pi_i \mathbf{f}_* = 0, i > 0$. Such an object exists because of [Q] and we obtain a spectral sequence

$$E_{pq}^1 = HL_q(\mathbf{f}_p, M) \implies HL_{p+q}(\mathbf{g}, M)$$

because $H_*(\text{Tot } CL_*(\mathbf{f}_*, M)) \cong HL_*(\mathbf{g}, M)$. Thus Theorem B is a consequence of Proposition 3.1, Künneth relation and Lemma 3.2.

LEMMA 3.2. — One has

$$\begin{aligned} \pi_i H_0(\mathbf{f}_*, M) &\cong H_0(\mathbf{g}, M), \quad \text{if } i = 0 \quad \text{and} \quad = 0, \quad \text{if } i > 0 \\ \pi_* H_1(\mathbf{f}_*, M) &\cong H_{*+1}(\mathbf{g}, M) \\ \pi_* H_1(\mathbf{f}_*, \mathbf{f}_*) &\cong HR_*(\mathbf{g}). \end{aligned}$$

Proof. — The first part follows from the fact that $H_0(\mathbf{f}_*, M)$ is a constant simplicial object. The second one is well-known (see for example [Q]). Since $H_*(\text{Tot } CR_*(\mathbf{f}_*)) \cong H_* CR_*(\mathbf{g})$, the last part is a consequence of Proposition 3.1.

4. The relation between $HL_*\mathfrak{h}$ and $HL_*\mathfrak{h}_{\text{Lie}}$.

Let $f : \mathfrak{h} \rightarrow \mathfrak{b}$ be an epimorphism of Leibniz algebras and let M be a right \mathfrak{b} -module. We may consider it also as an \mathfrak{h} -module through f . Let $C_*(\mathfrak{h}; \mathfrak{b}, M)$ be a chain complex, such that

$$CL_*(\mathfrak{h}; \mathfrak{b}, M)[1] = \text{Ker}(CL_*(\mathfrak{h}, M) \rightarrow CL_*(\mathfrak{b}, M))$$

and $HL_*(\mathfrak{h}; \mathfrak{b}, M) := H_*(CL_*(\mathfrak{h}; \mathfrak{b}, M))$. Then, by definition, we have

PROPOSITION 4.1. — *There exists an exact sequence*

$$\begin{aligned} 0 \leftarrow HL_1(\mathfrak{b}, M) \leftarrow HL_1(\mathfrak{h}, M) \leftarrow HL_0(\mathfrak{h}; \mathfrak{b}, M) \\ \leftarrow HL_2(\mathfrak{b}, M) \leftarrow HL_2(\mathfrak{h}, M) \leftarrow \dots \end{aligned}$$

THEOREM C. — *Let*

$$0 \rightarrow \mathfrak{a} \rightarrow \mathfrak{h} \rightarrow \mathfrak{b} \rightarrow 0$$

be a short exact sequence of Leibniz algebras. Assume $[h, a] = 0$, $h \in \mathfrak{h}$, $a \in \mathfrak{a}$. Then there exists a spectral sequence

$$E_{pq}^2 = HL_p(\mathfrak{b}, \mathfrak{a}) \otimes HL_q(\mathfrak{h}, M) \implies HL_{p+q}(\mathfrak{h}; \mathfrak{b}, M)$$

where a right \mathfrak{b} -module structure on \mathfrak{a} is given by $[a, x] := [a, f^{-1}(x)]$, $x \in \mathfrak{b}$, $a \in \mathfrak{a}$.

Proof. — Let $a \in \mathfrak{a}$ and $\omega \in CL_*(\mathfrak{h}, M)$. Then $d(\omega, a) = (d\omega, a)$, because $[\omega, a] = 0$. Thus

$$\mathcal{F}_*^0 = CL_*(\mathfrak{h}, M) \otimes \mathfrak{a} \subset CL_*(\mathfrak{h}; \mathfrak{b}, M)$$

is a subcomplex and $H_*(\mathcal{F}_*^0) = H_*(\mathfrak{h}, M) \otimes \mathfrak{a}$. More generally : for $\rho = (x_1, \dots, x_i)$ and $\rho_j = (x_1, \dots, \hat{x}_j, \dots, x_i)$, $1 \leq j \leq i$, one has

$$\begin{aligned} d(\omega, a, \rho) = (d\omega, a, \rho) + \sum (-1)^{\text{deg } \omega + j}([\omega, x_j], a, \rho_j) \\ + \sum (-1)^{\text{deg } \omega + j}(\omega, [a, x_j], \rho_j) + (-1)^{\text{deg } \omega}(\omega, a, d\rho). \end{aligned}$$

Since $[a, x] \in \mathfrak{a}$ we conclude that $\mathcal{F}_*^0 \subset \mathcal{F}_*^1 \subset \dots$ are subcomplexes of $CL_*(\mathfrak{h}; \mathfrak{b}, M)$ with $\mathcal{F}_n^s = CL_n(\mathfrak{h}; \mathfrak{b}, M)$. Here \mathcal{F}_*^s is spanned by (ω, a, ρ) with $\text{deg } \rho \leq s$. One checks that there exists an isomorphism

$$\mathcal{F}_*^s / \mathcal{F}_*^{s-1} \cong CL_*(\mathfrak{h}; \mathfrak{b}, M)[-s] \otimes \mathfrak{a} \otimes \mathfrak{b}^{\otimes s}$$

and the theorem follows from 10.1.7. of [CH] and the above formula for d .

Remark 4.2. — The assumptions of Theorem C hold for

$$0 \longrightarrow \mathfrak{h}^{\text{ann}} \longrightarrow \mathfrak{h} \longrightarrow \mathfrak{h}_{\text{Lie}} \longrightarrow 0.$$

Hence Theorem C gives a relation between $HL_*\mathfrak{h}$ and $HL_*\mathfrak{h}_{\text{Lie}}$. Since $HL_1(\mathfrak{h}, k) = HL_1(\mathfrak{h}_{\text{Lie}}, k)$, we obtain an exact sequence

$$0 \longleftarrow (\mathfrak{h}^{\text{ann}})_{\mathfrak{h}_{\text{Lie}}} \longleftarrow HL_2(\mathfrak{h}_{\text{Lie}}, k) \longleftarrow HL_2(\mathfrak{h}, k).$$

PROPOSITION 4.3. — *Let \mathfrak{h} be a finite dimensional Leibniz algebra, such that $\mathfrak{g} = \mathfrak{h}_{\text{Lie}}$ is a semi-simple Lie algebra over characteristic zero field or is a nilpotent Lie algebra with $\text{ch}(\mathfrak{h}) = 0$. Then $HL_*(\mathfrak{h}, k) = HL_*(\mathfrak{g}, k)$.*

Proof. — By Proposition 2.1 $\text{ch}(\mathfrak{h}) = 0$ in both cases. Hence the epimorphism $\mathfrak{h} \longrightarrow \mathfrak{g}$ has a section. It follows from Proposition 4.1 and Remark 4.2 that

$$HL_*(\mathfrak{h}, k) = HL_*(\mathfrak{g}, k) \oplus HL_{*-1}(\mathfrak{h}; \mathfrak{g}, k), \quad \text{and} \quad (\mathfrak{h}^{\text{ann}})_{\mathfrak{g}} = 0.$$

Theorem C gives the spectral sequence

$$E_{pq}^2 = HL_p(\mathfrak{g}, \mathfrak{h}^{\text{ann}}) \otimes HL_q(\mathfrak{h}, k) \implies HL_{p+q}(\mathfrak{h}; \mathfrak{g}, k).$$

By Theorem 1 of [D], Corollary 1.3 and Proposition 2.1 we have $HL_*(\mathfrak{g}, \mathfrak{h}^{\text{ann}}) = 0$. Thus $HL_*(\mathfrak{h}; \mathfrak{g}, k) = 0$, because $E_{*q}^2 = 0$ and $HL_*(\mathfrak{h}, k) = HL_*(\mathfrak{g}, k)$.

BIBLIOGRAPHY

- [D] J. DIXMIER, Cohomologie des algèbres de Lie nilpotentes. Acta Sci. Math. Szeged., 16 (1955), 246-250.
- [HS] P.J. HILTON, and U. STAMMBACH, A course in homological algebra (Grad. Texts in Math., vol. 4). Berlin, Heidelberg, New York, Springer Verlag, 1971.
- [K] J.L. KOSZUL, Homologie et cohomologie des algèbres de Lie. Bull. Soc. Math. France, 78 (1950), 65-127.
- [CH] J.L. LODAY, Cyclic homology. (Grund. Math. Wiss., Bd. 301), Berlin, Heidelberg, New York, Springer Verlag, 1992.
- [LP] J.L. LODAY, and T. PIRASHVILI, Universal enveloping algebras of Leibniz algebras and (co)homology, Math. Ann., 296 (1993), 139-158.

- [N] P. NTOLO, in preparation.
- [Q] D. QUILLEN, On the (co)-homology of commutative rings. Proc. Symp. Pure Math., 17 (1970), 65-87.
- [T] A.K. TOLPYGO, Cohomologies of parabolic Lie algebras, Mathematical notes of the Academy of Sciences of the USSR (Translations), 12 (1972), 585-588.

Manuscrit reçu le 10 septembre 1993.

Teimuraz PIRASHVILI,
Mathematical Institute
Rukhadze Str.1
Tbilisi, 380093 (Rep. of Georgia).