



ANNALES

DE

L'INSTITUT FOURIER

Plamen ILIEV

On the heat kernel and the Korteweg–de Vries hierarchy

Tome 55, n° 6 (2005), p. 2117-2127.

http://aif.cedram.org/item?id=AIF_2005__55_6_2117_0

© Association des Annales de l'institut Fourier, 2005, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

ON THE HEAT KERNEL AND THE KORTEWEG-DE VRIES HIERARCHY

by Plamen ILIEV

1. Introduction and examples.

Consider the one-dimensional Schrödinger (or Sturm-Liouville) operator

$$(1.1) \quad L = \frac{\partial^2}{\partial x^2} + u(x).$$

Its heat kernel $H(x, y, t)$ is the fundamental solution of the heat equation

$$(1.2) \quad \left(\frac{\partial}{\partial t} - L \right) f = 0.$$

It is well known that $H(t, x, y)$ has an asymptotic expansion of the form

$$(1.3) \quad H(x, y, t) \sim \frac{e^{-\frac{(x-y)^2}{4t}}}{\sqrt{4\pi t}} \left(1 + \sum_{n=1}^{\infty} H_n(x, y)t^n \right) \text{ as } t \rightarrow 0+.$$

The differential equation (1.2) for $H(x, y, t)$ implies the recursion-differential equations for the coefficients $H_n = H_n(x, y)$:

$$(1.4) \quad H_0 = 1$$

$$(1.5) \quad (x - y) \frac{\partial H_n}{\partial x} + nH_n = LH_{n-1} \text{ for } n \geq 1.$$

This system is known to admit unique smooth solutions $H_n = H_n(x, y)$ in some neighborhood of the diagonal $x = y$. The coefficients H_n are named after J. Hadamard [13], who constructed them for the first time.

Keywords: Heat kernel expansions, KdV hierarchy, tau functions.
Math. classification: 35Q53, 35K05, 37K10.

Computation of heat invariants of self-adjoint elliptic operators is a well known problem in spectral theory which has many applications, in particular to geometry and theoretical physics [5, 6, 10, 11, 17, 18]. The asymptotics of the one-dimensional Schrödinger operator are of particular interest due to their relations to the Korteweg-de Vries (KdV) hierarchy. More precisely, it is known that the restriction of the heat coefficients on the diagonal gives the right-hand sides of the KdV hierarchy, see [19, 21].

In the present paper we show that there are simple formulas for the Hadamard's coefficients $H_n(x, y)$ in terms of the τ -function of the KdV hierarchy (see the next section for a precise definition of the τ -function).

Remarkable explicit formulas for the coefficients of the Taylor expansion of $H_n(x, y)$ around the diagonal $x = y$ were previously constructed in [4], Theorem 1.3. However, these formulas have a rather complicated combinatorial structure and it is practically impossible to write a closed formula for the coefficients even for simple potentials u . One advantage of the formulas derived in this paper is that they give finite expressions for the heat coefficient if the τ -function is known (e.g. the solitons, or the more general algebro-geometric solutions of KdV).

To see the importance of the KdV equations, let us compute the first few coefficients using the defining relations (1.4)–(1.5). Anticipating the appearance of the τ -function, let us write $u(x)$ as

$$u(x) = 2 \frac{\partial^2 \log(\tau(x))}{\partial x^2}.$$

From (1.5) one can easily obtain simple formulas for H_1 and H_2

$$(1.6) \quad H_1(x, y) = \frac{2}{x-y} \left(\frac{\tau'(x)}{\tau(x)} - \frac{\tau'(y)}{\tau(y)} \right)$$

and

$$(1.7) \quad H_2(x, y) = \frac{2}{(x-y)^2} \left(\left(\frac{\tau''(x)}{\tau(x)} + \frac{\tau''(y)}{\tau(y)} \right) - H_1(x, y) - 2 \frac{\tau'(x)\tau'(y)}{\tau(x)\tau(y)} \right).$$

For the third coefficient we have the following formula

$$(1.8) \quad \begin{aligned} (x-y)^3 H_3(x, y) = & -6(x-y)H_2(x, y)2 \left(\frac{\tau'''(x)}{\tau(x)} - \frac{\tau'''(y)}{\tau(y)} \right) \\ & - 2 \left(\frac{\tau''(x)\tau'(x)}{\tau^2(x)} - \frac{\tau''(y)\tau'(y)}{\tau^2(y)} \right) \\ & + 4 \left(\frac{\tau'(x)}{\tau(x)} \frac{\tau''(y)}{\tau(y)} - \frac{\tau'(y)}{\tau(y)} \frac{\tau''(x)}{\tau(x)} \right) \\ & + \frac{4}{3} \left(\left(\frac{\tau'(x)}{\tau(x)} \right)^3 - \left(\frac{\tau'(y)}{\tau(y)} \right)^3 \right) + \int_y^x u^2(\xi) d\xi. \end{aligned}$$

Notice that the integral cannot be computed explicitly, unless something remarkable happens. This is the place where the KdV equation comes in. Assume that $u(x)$ depends on a additional parameter s_3 and it satisfies the KdV equation

$$(1.9) \quad 4\partial_3 u = u''' + 6uu',$$

where $\partial_3 = \partial/\partial s_3$ stands for the partial derivative with respect to s_3 , and u' is the derivative with respect to x . Then one can easily see that

$$(1.10) \quad \int_y^x u^2(\xi) d\xi = -\frac{2}{3} \left(\frac{\tau'''(x)}{\tau(x)} - \frac{\tau'''(y)}{\tau(y)} \right) + 2 \left(\frac{\tau''(x)\tau'(x)}{\tau^2(x)} - \frac{\tau''(y)\tau'(y)}{\tau^2(y)} \right) - \frac{4}{3} \left(\left(\frac{\tau'(x)}{\tau(x)} \right)^3 - \left(\frac{\tau'(y)}{\tau(y)} \right)^3 \right) + \frac{8}{3} \partial_3 \log \frac{\tau(x)}{\tau(y)},$$

which combined with (1.8) leads to a simple formula for $H_3(x, y)$.

We'll extend these computations by showing that if u is a solution of the KdV hierarchy and τ is the corresponding τ -function, then there are simple explicit formulas for $H_n(x, y)$ in terms of τ .

The paper is organized as follows. In the next section we recall some basic facts about the KdV hierarchy and Sato theory, which are needed for the formulation and the proof of the main result. In Section 3, we prove a general formula for $H_n(x, y)$. It is interesting that the smoothness of the coefficient $H_n(x, y)$ on the diagonal is related to the Gengenbauer polynomials. As a corollary of the main theorem, we see the symmetry of the coefficients about the diagonal $x = y$ as well as the connection between $H_n(x, x)$ and KdV equations.

As another application of the explicit formula, we show in [16] that the expansion is finite if and only if the potential $u(x)$ is a rational solution of the KdV hierarchy decaying at infinity studied in [1, 2]. Equivalently, one can characterize the corresponding operators as the rank one bispectral family in [9]. For related results concerning the finiteness property of the heat kernel expansion on the integers and rational solutions of the Toda lattice hierarchy see [12]. For solitons of the Toda lattice and purely discrete versions of the heat kernel see [14].

2. Korteweg-de Vries hierarchy and Sato theory.

In this section we recall some basic facts about KdV hierarchy and Sato theory. For more details on this and the more general Kadomtsev-Petviashvili hierarchy we refer the reader to the papers [20, 7] or the more detailed expositions [8, 22].

Let

$$L = \frac{\partial^2}{\partial x^2} + u(x)$$

be a second order differential operator. The KdV hierarchy is defined by the Lax equations

$$(2.1) \quad \frac{\partial L}{\partial s_j} = [(L^{j/2})_+, L],$$

where $j = 1, 3, 5, \dots$ is an odd positive integer and $(L^{j/2})_+$ is the differential part of the pseudo-differential operator $L^{j/2}$. The first equation (for $j = 1$) simply means that $u(x, s_1, s_3, s_5, \dots) = u(x + s_1, s_3, s_5, \dots)$, giving us the convenience to occasionally identify x and s_1 . The next equation (for $j = 3$) is exactly the KdV equation (1.9). Let us represent L in a dressing form

$$(2.2) \quad L = W \partial^2 W^{-1},$$

where W is a pseudo-differential operator of the form

$$(2.3) \quad W = \sum_{k=0}^{\infty} \psi_k \partial^{-k}, \quad \psi_0 = 1.$$

The wave (Baker) function $\Psi(x, s, z)$ and the adjoint wave function $\Psi^*(x, s, z)$ are defined as

$$(2.4) \quad \begin{aligned} \Psi(x, s, z) &= W \exp\left(xz + \sum_{i=1}^{\infty} s_{2i-1} z^{2i-1}\right) \\ &= \left(\sum_{k=0}^{\infty} \psi_k z^{-k}\right) \exp\left(xz + \sum_{i=1}^{\infty} s_{2i-1} z^{2i-1}\right) \end{aligned}$$

and

$$(2.5) \quad \begin{aligned} \Psi^*(x, s, z) &= (W^*)^{-1} \exp\left(-xz - \sum_{i=1}^{\infty} s_{2i-1} z^{2i-1}\right) \\ &= \left(\sum_{k=0}^{\infty} \psi_k^* z^{-k}\right) \exp\left(-xz - \sum_{i=1}^{\infty} s_{2i-1} z^{2i-1}\right), \end{aligned}$$

where W^* is the formal adjoint to the pseudo-differential operator W . Using (2.2) one can easily see that

$$(2.6) \quad L\Psi(x, s, z) = z^2\Psi(x, s, z) \text{ and } L\Psi^*(x, s, z) = z^2\Psi^*(x, s, z).$$

We shall also use the reduced wave function $\bar{\Psi}$ and the reduced adjoint wave function $\bar{\Psi}^*$ obtained from Ψ and Ψ^* , respectively, by omitting the exponential factor, *i.e.*

$$(2.7) \quad \bar{\Psi}(x, s, z) = \sum_{k=0}^{\infty} \psi_k z^{-k}$$

and

$$(2.8) \quad \bar{\Psi}^*(x, s, z) = \sum_{k=0}^{\infty} \psi_k^* z^{-k}.$$

Equations (2.6) imply

$$(2.9) \quad L\bar{\Psi}(x, s, z) + 2z\partial_x \bar{\Psi}(x, s, z) = 0 \text{ and } L\bar{\Psi}^*(x, s, z) - 2z\partial_x \bar{\Psi}^*(x, s, z) = 0.$$

Using equations (2.2)–(2.5) one can show that the wave and the adjoint wave function satisfy the following bilinear identities

$$(2.10) \quad \text{res}_z \left(z^{2n} \Psi^{(l)}(x, s, z) \Psi^*(x, s, z) \right) = 0,$$

for all nonnegative integers n and l , where $\Psi^{(l)}(x, s, z)$ is the l th derivative of Ψ with respect to x , and the residue is around $z = \infty$.

The remarkable discovery of the Kyoto school was that the KdV hierarchy (2.1) could be described by a function $\tau(x, s)$. This goes back to an earlier work of Hirota, see [15]. The reduced wave and the reduced adjoint wave functions can be expressed in terms of $\tau(x, s)$ by the following formulas

$$(2.11) \quad \bar{\Psi}(x, s, z) = \frac{\tau(x; s - [z^{-1}])}{\tau(x, s)} \quad \text{and} \quad \bar{\Psi}^*(x, s, z) = \frac{\tau(x; s + [z^{-1}])}{\tau(x, s)},$$

where $[z] = (z, z^3/3, z^5/5, \dots)$.

Finally, let us denote by $W_n(x, y)$ the coefficients of the function¹ $\bar{\Psi}(x, s, z)\bar{\Psi}^*(y, s, z)$, *i.e.*

$$(2.12) \quad \bar{\Psi}(x, s, z)\bar{\Psi}^*(y, s, z) = \sum_{n=0}^{\infty} W_n(x, y)z^{-n}.$$

¹ This function is closely related to the Green function for L .

Using (2.11) we can easily write an explicit formula for W_n in terms of the τ -function. If we denote by $\mathfrak{S}_k(s)$ the elementary Schur polynomials defined by

$$(2.13) \quad \sum_{k=0}^{\infty} \mathfrak{S}_k(s) z^k = \exp\left(\sum_{k=1}^{\infty} s_{2k-1} z^{2k-1}\right),$$

then we have

$$(2.14) \quad W_n(x, y) = \frac{\sum_{k=0}^n [\mathfrak{S}_k(-\tilde{\partial})\tau(x, s)] [\mathfrak{S}_{n-k}(\tilde{\partial})\tau(y, s)]}{\tau(x, s)\tau(y, s)},$$

where

$$(2.15) \quad \tilde{\partial} = (\partial_1, \partial_3/3, \dots, \partial_{2k-1}/(2k-1), \dots).$$

3. Explicit formulas for Hadamard’s coefficients.

The main result of the paper is the following theorem.

THEOREM 3.1. — *The Hadamard’s coefficients can be computed from the following relation*

$$(3.1) \quad H_n(x, y) = (-1)^n \sum_{k=0}^{n-1} \frac{2^{n-k}(n-k)_{2k}}{k!} \frac{W_{n-k}(x, y)}{(x-y)^{n+k}},$$

where $(\alpha)_k = \alpha(\alpha+1)\dots(\alpha+k-1)$ denotes the Pochhammer symbol, and $W_n(x, y)$ are defined by (2.14).

Proof. — To prove that the Hadamard’s coefficients are given by (3.1) we need to check that (1.5) holds and that $H_n(x, y)$ are smooth on the diagonal $x = y$.

To see that (1.5) holds, let us denote

$$(3.2) \quad f_n(x, y, z) = (-1)^n \sum_{k=0}^{n-1} \frac{2^{n-k}(n-k)_{2k}}{k!(x-y)^{n+k}} z^{n-k-1}.$$

Then (3.1) can be rewritten as

$$(3.3) \quad H_n(x, y) = \operatorname{res}_z [f_n(x, y, z) \bar{\Psi}(x, s, z) \bar{\Psi}^*(y, s, z)].$$

Using the last equation together with (2.9) one can easily see that $[(x - y)\partial_x + n] H_n(x, y) - L(x, \partial_x)H_{n-1}(x, y)$

$$= \operatorname{res}_z \left[((x - y)\partial_x f_n(x, y, z) + n f_n(x, y, z) - \partial_x^2 f_{n-1}(x, y, z)) \bar{\Psi}(x, s, z) \bar{\Psi}^*(y, s, z) + ((x - y)f_n(x, y, z) + 2z f_{n-1}(x, y, z) - 2\partial_x f_{n-1}(x, y, z)) \partial_x \bar{\Psi}(x, s, z) \bar{\Psi}^*(y, s, z) \right].$$

A direct computation now shows that

$$\begin{aligned} (x - y)\partial_x f_n(x, y, z) + n f_n(x, y, z) - \partial_x^2 f_{n-1}(x, y, z) &= 0 \\ (x - y)f_n(x, y, z) + 2z f_{n-1}(x, y, z) - 2\partial_x f_{n-1}(x, y, z) &= 0, \end{aligned}$$

which proves (1.5).

Next we need to show that $H_n(x, y)$ is well defined on the diagonal. Writing $H_n(x, y)$ as

$$(3.4) \quad H_n(x, y) = \frac{2(-1)^n}{(x - y)^{2n-1}} \sum_{k=0}^{n-1} \frac{2^k (x - y)^k (k + 1)_{2n-2k-2}}{(n - k - 1)!} W_{k+1}(x, y),$$

and applying L'Hôpital's rule we see that we need to prove that for $j = 0, 1, \dots, 2n - 2$ we have

$$(3.5) \quad \sum_{k=0}^{n-1} 2^k \binom{j}{k} \frac{(2n - 2 - k)!}{(n - k - 1)!} \partial_x^{j-k} W_{k+1}(x, y)|_{x=y} = 0.$$

Using (2.12) and

$$\partial_x^j \bar{\Psi}(x, s, z) = \exp\left(-xz - \sum_{i=1}^{\infty} s_{2i-1} z^{2i-1}\right) (\partial_x - z)^j \Psi(x, s, z)$$

we see that (3.5) is equivalent to the following identities

$$(3.6) \quad \operatorname{res}_z \left[\left(\sum_{k=0}^{n-1} 2^k \binom{j}{k} \frac{(2n - 2 - k)!}{(n - k - 1)!} z^k (\partial_x - z)^{j-k} \Psi(x, s, z) \right) \Psi^*(x, s, z) \right] = 0.$$

Equation (3.6) will follow from the bilinear identities (2.10) if we can show that the polynomial

$$(3.7) \quad P_{n,j}(w) = \sum_{k=0}^{n-1} 2^k \binom{j}{k} \frac{(2n - 2 - k)!}{(n - k - 1)!} (w - 1)^{j-k},$$

is an even/odd function when j is an even/odd number, respectively. It is a pleasant surprise to see that these polynomials are closely related to very

well known classical orthogonal polynomials – the so called Gegenbauer polynomials.

The Gegenbauer (or ultraspherical) polynomials are defined by

$$(3.8) \quad C_n^\lambda(w) = \sum_{k=0}^n 2^k \binom{\lambda + k - 1}{k} \binom{2\lambda + n + k - 1}{n - k} (w - 1)^k,$$

see for example [3], pages 302–303. Notice that this definition can be used for arbitrary λ . If $\lambda > -1/2$ and $\lambda \neq 0$ these polynomials are orthogonal on the interval $(-1, 1)$ with respect to $(1 - x^2)^{\lambda - \frac{1}{2}}$ which, in particular, implies that $C_n^\lambda(w)$ is an even/odd function when n is even/odd, respectively. However, we need these polynomials also for negative values of λ . In this case, we can use the three term recurrence relation

$$(3.9) \quad 2(n + \lambda)wC_n^\lambda(w) = (n + 1)C_{n+1}^\lambda(w) + (n + 2\lambda - 1)C_{n-1}^\lambda(w).$$

and $C_0^\lambda = 1$ and $C_1^\lambda = 2\lambda w$ to deduce that $C_n^\lambda(w)$ is an even/odd polynomial when n is even/odd, respectively.

Changing the summation index in (3.7) we can rewrite $P_{n,j}(w)$ as

$$(3.10) \quad P_{n,j}(w) = \sum_{k=\max(0, j-n+1)}^j 2^{j-k} \binom{j}{k} \frac{(2n - j - 2 + k)!}{(n - j + k - 1)!} (w - 1)^k.$$

From the last equation and the defining relation (3.8) for the Gegenbauer polynomials one can see that²

$$(3.11) \quad P_{n,j}(w) = j!2^{n-1}(2n - 2j - 3)!! C_j^{n-j-\frac{1}{2}}(w) \quad \text{for } 0 \leq j \leq n - 1,$$

$$(3.12) \quad P_{n,j}(w) = \frac{(-1)^{j-n+1}j!2^{n-1}}{(2j - 2n + 1)!!} C_j^{n-j-\frac{1}{2}}(w) \quad \text{for } n \leq j \leq 2n - 2,$$

which completes the proof. □

From (2.14) and (3.1) we obtain the following

COROLLARY 3.2. — *The Hadamard’s coefficients $H_n(x, y)$ are symmetric functions of x and y , i.e. we have*

$$H_n(x, y) = H_n(y, x).$$

Finally, we show that the heat coefficients $\{H_n(x, x)\}$ determine the right-hand sides of KdV equations (2.1).

² $(-1)!! = 1$ and $(2k - 1)!! = 1 \cdot 3 \cdots (2k - 1)$ for $k \geq 1$.

COROLLARY 3.3. — We have

$$(3.13) \quad H_n(x, x) = \frac{2^n}{(2n - 1)!!} W_{2n}(x, x)$$

and

$$(3.14) \quad [(L^{\frac{2n-1}{2}})_+, L] = 2\partial_x W_{2n}(x, x).$$

Thus, the KdV hierarchy (2.1) is equivalent to the following equations

$$(3.15) \quad \partial_{2n-1} u = \frac{(2n - 1)!!}{2^{n-1}} \partial_x H_n(x, x), \text{ for } n = 1, 2, \dots$$

Proof. — Using (3.1) and applying L'Hôpital's rule $2n - 1$ times we see that

$$(3.16) \quad \begin{aligned} H_n(x, x) &= (-1)^n \frac{2}{(2n - 1)!} \\ &\times \sum_{k=0}^{n-1} 2^k \binom{2n - 1}{k} \frac{(2n - k - 2)!}{(n - k - 1)!} \partial_x^{2n-1-k} W_{k+1}(y, x)|_{y=x} \\ &= (-1)^n \frac{2}{(2n - 1)!} \\ &\times \operatorname{res}_z [z^{2n-1} (P_{n,2n-1}(z^{-1} \partial_x) \Psi(x, s, z)) \Psi^*(x, s, z)], \end{aligned}$$

where $P_{n,2n-1}(w)$ is the polynomial defined by (3.10) for $j = 2n - 1$

$$(3.17) \quad P_{n,2n-1}(w) = \sum_{k=n}^{2n-1} 2^{2n-1-k} \binom{2n - 1}{k} \frac{(k - 1)!}{(k - n)!} (w - 1)^k.$$

Notice that this time we have

$$(3.18) \quad P_{n,2n-1}(w) = (-1)^n 2^{2n-1} (2n - 2)!! \left(C_{2n-1}^{-n+\frac{1}{2}}(w) + 1 \right),$$

and using the same argument (the bilinear identity and the fact that $C_{2n-1}^{-n+\frac{1}{2}}(w)$ is an odd polynomial) we obtain (3.13) from (3.16).

From (2.2), (2.4), (2.5) and (2.11) it follows that

$$L^{\frac{2n-1}{2}} = W \partial_x^{2n-1} W^{-1} = \sum_{i,j=0}^{\infty} \frac{\mathfrak{S}_i(-\tilde{\partial}) \tau(x, s)}{\tau(x, s)} \partial_x^{2n-1-i-j} \cdot \frac{\mathfrak{S}_j(\tilde{\partial}) \tau(x, s)}{\tau(x, s)}.$$

Combining this formula with (2.14) we find that the coefficient of ∂_x^{-1} in $L^{\frac{2n-1}{2}}$ is $W_{2n}(x, x)$. If we denote by $(L^{\frac{2n-1}{2}})_-$ the integral (Volterra) part of the pseudo-differential operator $L^{\frac{2n-1}{2}}$ we obtain

$$\begin{aligned} [(L^{\frac{2n-1}{2}})_+, L] &= [(L^{\frac{2n-1}{2}})_+, L]_+ \\ &= [L^{\frac{2n-1}{2}} - (L^{\frac{2n-1}{2}})_-, L]_+ = [L, (L^{\frac{2n-1}{2}})_-]_+ \\ &= [\partial_x^2 + u(x), W_{2n}(x, x) \partial_x^{-1} + O(\partial_x^{-2})]_+ = 2\partial_x(W_{2n}(x, x)), \end{aligned}$$

which gives (3.14) and completes the proof. □

BIBLIOGRAPHY

- [1] M. ADLER and J. MOSER, On a class of polynomials connected with the Korteweg-de Vries equation, *Comm. Math. Phys.*, 61 (1978), 1–30.
- [2] H. AIRAULT, H. P. MCKEAN and J. MOSER, Rational and elliptic solutions of the Korteweg-de Vries equation and a related many-body problem, *Comm. Pure Appl. Math.*, 30 (1977), 95–148.
- [3] G. ANDREWS, R. ASKEY and R. ROY, *Special Functions*, *Encyclopedia of Mathematics and Its Applications*, Vol. 71, Cambridge University Press (1990).
- [4] I. AVRAMIDI and R. SCHIMMING, A new explicit expression for the Korteweg-de Vries hierarchy, *Math. Nachr.*, 219 (2000), 45–64.
- [5] N. BERLINE, E. GETZLER and M. VERGNE, *Heat kernels and Dirac operators*, *Grundlehren der Mathematischen Wissenschaften*, 298, Springer-Verlag, Berlin, 1992.
- [6] M. BERGER, *Geometry of the Spectrum*, *Proc. Sympos. Pure Math.*, 27, Amer. Math. Soc., Providence (1975), 129–152.
- [7] E. DATE, M. JIMBO, M. KASHIWARA and T. MIWA, Transformation groups for soliton equations, in: M. Jimbo and T. Miwa (eds), *Proc. RIMS Symp. Nonlinear Integrable Systems - Classical and Quantum Theory (Kyoto 1981)*, World Scientific, Singapore (1983), 39–119.
- [8] L. A. DICKEY, *Soliton Equations and Hamiltonian Systems*, 2nd. Edition, *Advanced Series in Mathematical Physics*, Vol. 26, World Scientific, 2003.
- [9] J. J. DUISTERMAAT and F. A. GRÜNBAUM, Differential equations in the spectral parameter, *Comm. Math. Phys.*, 103 (1986), 177–240.
- [10] S. A. FULLING (ed.), *Heat kernel techniques and quantum gravity (Winnipeg, MB, 1994)*, *Discourses Math. Appl.*, 4, Texas A & M Univ., College Station, TX, 1995.
- [11] P. GILKEY, Heat equation asymptotics, *Differential geometry: Riemannian geometry (Los Angeles, CA, 1990)*, 317–326, *Proc. Sympos. Pure Math.*, 54, Part 3, Amer. Math. Soc., Providence, RI, 1993.
- [12] F. A. GRÜNBAUM and P. ILIEV, Heat kernel expansions on the integers, *Math. Phys. Anal. Geom.*, 5 (2002), 183–200, [math.CA/0206089](#).
- [13] J. HADAMARD, *Lectures on Cauchy's Problem*, New Haven, Yale Univ. Press, 1923.
- [14] L. HAINE, The spectral matrices of Toda solitons and the fundamental solution of some discrete heat equations, to appear in *Annales de l'Institut Fourier*.
- [15] R. HIROTA, *The direct method in soliton theory*, Translated from the 1992 Japanese original and edited by Atsushi Nagai, Jon Nimmo and Claire Gilson (with a foreword by Jarmo Hietarinta and Nimmo). *Cambridge Tracts in Mathematics*, 155. Cambridge University Press, Cambridge, 2004.
- [16] P. ILIEV, Finite heat kernel expansions on the real line, [math-ph/0504046](#).
- [17] M. KAC, Can one hear the shape of a drum?, *Amer. Math. Monthly*, 73 (1966), 1–23.
- [18] H. P. MCKEAN and I. SINGER, Curvature and the eigenvalues of the Laplacian, *J. Diff. Geom.*, 1 (1967), 43–69.
- [19] H. P. MCKEAN and P. van MOERBEKE, The spectrum of Hill's equation, *Invent. Math.*, 30 (1975), 217–274.

- [20] M. SATO and Y. SATO, Soliton equations as dynamical systems on infinite dimensional Grassmann manifolds, *Lect. Notes Num. Appl. Anal.*, 5 (1982), 259–271.
- [21] R. SCHIMMING, An explicit expression for the Korteweg-de Vries hierarchy, *Z. Anal. Anwendungen*, 7 (1988), 203–214.
- [22] P. van MOERBEKE, Integrable foundations of string theory, in: O. Babelon et al. (eds.), *Lectures on integrable systems*, CIMPA-Summer school at Sophia-Antipolis (1991), Singapore: World Scientific (1994) 163–267.

Plamen ILIEV,
Georgia Institute of Technology
School of Mathematics
Atlanta, GA 30332-0160 (USA)
iliev@math.gatech.edu

