



ANNALES

DE

L'INSTITUT FOURIER

Jean-Louis VERGER-GAUGRY

On gaps in Rényi β -expansions of unity for $\beta > 1$ an algebraic number

Tome 56, n° 7 (2006), p. 2565-2579.

http://aif.cedram.org/item?id=AIF_2006__56_7_2565_0

© Association des Annales de l'institut Fourier, 2006, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

ON GAPS IN RÉNYI β -EXPANSIONS OF UNITY FOR $\beta > 1$ AN ALGEBRAIC NUMBER

by Jean-Louis VERGER-GAUGRY

ABSTRACT. — Let $\beta > 1$ be an algebraic number. We study the strings of zeros (“gaps”) in the Rényi β -expansion $d_\beta(1)$ of unity which controls the set \mathbb{Z}_β of β -integers. Using a version of Liouville’s inequality which extends Mahler’s and Güting’s approximation theorems, the strings of zeros in $d_\beta(1)$ are shown to exhibit a “gappiness” asymptotically bounded above by $\log(M(\beta))/\log(\beta)$, where $M(\beta)$ is the Mahler measure of β . The proof of this result provides in a natural way a new classification of algebraic numbers > 1 with classes called $Q_i^{(j)}$ which we compare to Bertrand-Mathis’s classification with classes C_1 to C_5 (reported in an article by Blanchard). This new classification relies on the maximal asymptotic “quotient of the gap” value of the “gappy” power series associated with $d_\beta(1)$. As a corollary, all Salem numbers are in the class $C_1 \cup Q_0^{(1)} \cup Q_0^{(2)} \cup Q_0^{(3)}$; this result is also directly proved using a recent generalization of the Thue-Siegel-Roth Theorem given by Corvaja.

RÉSUMÉ. — Soit $\beta > 1$ un nombre algébrique. Nous étudions les plages de zéros (“lacunes”) dans le β -développement de Rényi $d_\beta(1)$ de l’unité qui contrôle l’ensemble \mathbb{Z}_β des β -entiers. En utilisant une version de l’inégalité de Liouville qui étend des théorèmes d’approximation de Mahler et de Güting, on montre que les plages de zéros dans $d_\beta(1)$ présentent une “lacunarité” asymptotiquement bornée supérieurement par $\log(M(\beta))/\log(\beta)$, où $M(\beta)$ est la mesure de Mahler de β . La preuve de ce résultat fournit de manière naturelle une nouvelle classification des nombres algébriques > 1 en classes appelées $Q_i^{(j)}$ que nous comparons à la classification de Bertrand-Mathis avec les classes C_1 à C_5 (reportée dans un article de Blanchard). Cette nouvelle classification repose sur la valeur asymptotique maximale du “quotient de lacune” de la série “lacunaire” associée à $d_\beta(1)$. Comme corollaire, tous les nombres de Salem sont dans la classe $C_1 \cup Q_0^{(1)} \cup Q_0^{(2)} \cup Q_0^{(3)}$; ce résultat est également obtenu par un théorème récent qui généralise le théorème de Thue-Siegel-Roth donné par Corvaja.

Keywords: Beta-integer, beta-numeration, PV number, Salem number, Perron number, Mahler measure, Diophantine approximation, Mahler’s series, mathematical quasicrystal.

Math. classification: 11B05, 11Jxx, 11J68, 11R06, 52C23.

1. Introduction

The exploration of the links between symbolic dynamics and number theory of β -expansions, when $\beta > 1$ is an algebraic number or more generally a real number, started with Bertrand-Mathis [6] [7]. Bertrand-Mathis, in Blanchard [8], reported a classification of real numbers according to their β -shift, using the properties of the Rényi β -expansion $d_\beta(1)$ of 1. A lot of questions remain open concerning the distribution of the algebraic numbers $\beta > 1$ in this classification. The Rényi β -expansion of 1 is important since it controls the β -shift [38] and the discrete and locally finite set $\mathbb{Z}_\beta \subset \mathbb{R}$ of β -integers [13] [18] [25] [26]. The aim of this note is to give a new Theorem (Theorem 1.1) on the gaps (strings of 0's) in $d_\beta(1)$ for algebraic numbers $\beta > 1$, and investigate how it provides (partial) answers to some questions of [8], in particular for Salem numbers (Corollary 1.2).

Theorem 1.1 provides an upper bound on the asymptotic quotient of the gap of $d_\beta(1)$ and is obtained by a version of Liouville's inequality extending Mahler's and Gütting's approximation theorems. The proof of Theorem 1.1 turns out to be extremely instructive in itself since it leads to a new classification of the algebraic numbers β as a function of the asymptotics of the gaps in $d_\beta(1)$ and "intrinsic features", namely the Mahler measure $M(\beta)$, of β (the definition of $M(\beta)$ is recalled in Section 3). The existence of this double parametrization, symbolic and algebraic, was guessed in [8] p 137. This new classification complements Bertrand-Mathis's (Blanchard [8] pp 137–139) and both are recalled below for comparison's sake. The question whether an algebraic number $\beta > 1$ is contained in one class or another has already been discussed by many authors [5] [6] [7] [8] [9] [10] [11] [12] [17] [22] [32] [33] [38] [39] [41] [42] and depends at least upon the distribution of the conjugates of β in the complex plane. Only the conjugates of β of modulus strictly greater than unity intervene in Theorem 1.1 via the Mahler measure of β . Corollary 1.2 is readily deduced from this remark. We deduce that Salem numbers belong to $C_1 \cup C_2 \cup Q_0$, whereas the Pisot numbers are in $C_1 \cup C_2$ [45].

Another proof of Corollary 1.2 consists of controlling the gaps of $d_\beta(1)$ by stronger Theorems of Diophantine Geometry which allow suitable collections of places of the number field $\mathbb{Q}(\beta)$ associated with the conjugates of β and the properties of $d_\beta(1)$ to be taken into account simultaneously. This alternative proof of Corollary 1.2, just sketched in Section 4, is obtained using the Theorem of Thue-Siegel-Roth given by Corvaja [1] [15].

THEOREM 1.1. — Let $\beta > 1$ be an algebraic number and $M(\beta)$ be its Mahler measure. Denote by $d_\beta(1) := 0.t_1t_2t_3\dots$, with $t_i \in A_\beta := \{0, 1, 2, \dots, \lceil \beta - 1 \rceil\}$, the Rényi β -expansion of 1. Assume that $d_\beta(1)$ is infinite and gappy in the following sense: there exist two sequences $\{m_n\}_{n \geq 1}$, $\{s_n\}_{n \geq 0}$ such that

$$1 = s_0 \leq m_1 < s_1 \leq m_2 < s_2 \leq \dots \leq m_n < s_n \leq m_{n+1} < s_{n+1} \leq \dots$$

with $(s_n - m_n) \geq 2$, $t_{m_n} \neq 0, t_{s_n} \neq 0$ and $t_i = 0$ if $m_n < i < s_n$ for all $n \geq 1$. Then

$$(1.1) \quad \limsup_{n \rightarrow +\infty} \frac{s_n}{m_n} \leq \frac{\log(M(\beta))}{\log(\beta)}.$$

Moreover, if $\liminf_{n \rightarrow +\infty} (m_{n+1} - m_n) = +\infty$, then

$$(1.2) \quad \limsup_{n \rightarrow +\infty} \frac{s_{n+1} - s_n}{m_{n+1} - m_n} \leq \frac{\log(M(\beta))}{\log(\beta)}.$$

As in Ostrowski [37] the quotient $s_n/m_n \geq 1$ is called *the quotient of the gap*, relative to the n th-gap (assuming $t_j \neq 0$ for all $s_n \leq j \leq m_{n+1}$ to characterize uniquely the gaps). Note that the term “lacunary” has often other meanings in literature and is not used here to describe “gappiness”. Denote by $\mathcal{L}(S_\beta)$ the language of the β -shift [8] [23] [24] [34]. Bertrand-Mathis’s classification ([8] pp 137–139) is as follows:

- C_1 : $d_\beta(1)$ is finite.
- C_2 : $d_\beta(1)$ is ultimately periodic but not finite.
- C_3 : $d_\beta(1)$ contains bounded strings of 0’s, but is not ultimately periodic.
- C_4 : $d_\beta(1)$ does not contain some words of $\mathcal{L}(S_\beta)$, but contains strings of 0’s with unbounded length.
- C_5 : $d_\beta(1)$ contains all words of $\mathcal{L}(S_\beta)$.

Present classes of algebraic numbers, with the notations of Theorem 1.1:

- $Q_0^{(1)}$: $1 = \lim_{n \rightarrow +\infty} \frac{s_n}{m_n}$ with $(m_{n+1} - m_n)$ bounded.
- $Q_0^{(2)}$: $1 = \lim_{n \rightarrow +\infty} \frac{s_n}{m_n}$ with $(s_n - m_n)$ bounded and $\lim_{n \rightarrow +\infty} (m_{n+1} - m_n) = +\infty$.
- $Q_0^{(3)}$: $1 = \lim_{n \rightarrow +\infty} \frac{s_n}{m_n}$ with $\limsup_{n \rightarrow +\infty} (s_n - m_n) = +\infty$.
- Q_1 : $1 < \limsup_{n \rightarrow +\infty} \frac{s_n}{m_n} < \frac{\log(M(\beta))}{\log(\beta)}$.
- Q_2 : $\limsup_{n \rightarrow +\infty} \frac{s_n}{m_n} = \frac{\log(M(\beta))}{\log(\beta)}$.

What are the relative proportions of each class in the whole set $\overline{\mathbb{Q}}_{>1}$ of algebraic numbers $\beta > 1$? Comparing C_2, C_3 and $Q_0^{(1)}$, what are the relative proportions in $Q_0^{(1)}$ of those β which give ultimate periodicity in

$d_\beta(1)$ and those for which $d_\beta(1)$ is not ultimately periodic ? Schmeling ([41] Theorem A) has shown that the class C_3 (of real numbers $\beta > 1$) has Hausdorff dimension one. We have:

- $\overline{Q}_{>1} \cap C_2 \subset Q_0^{(1)}$,
- $\overline{Q}_{>1} \cap C_3 \subset Q_0^{(1)} \cup Q_0^{(2)}$, with $C_3 \cap Q_0^{(3)} = \emptyset$,
- $\overline{Q}_{>1} \cap C_4 \subset Q_0^{(3)} \cup Q_1 \cup Q_2$.

The Pisot numbers β are contained in $C_1 \cup Q_0^{(1)}$ since they are such that $d_\beta(1)$ is finite or ultimately periodic (Parry [38], Bertrand-Mathis [5]). Recall that a Perron number is an algebraic integer $\beta > 1$ such that all the conjugates $\beta^{(i)}$ of β satisfy $|\beta^{(i)}| < \beta$. Conversely, as shown in Lind [32], Denker, Grillenberger, Sigmund [17] and Bertrand-Mathis [7], if $\beta > 1$ is such that $d_\beta(1)$ is ultimately periodic (finite or not), then β is a Perron number. Not all Perron numbers are attained in this way: a Perron number which possesses a real conjugate greater than 1 cannot be such that $d_\beta(1)$ is ultimately periodic ([8] p 138). Parry numbers belong to $C_1 \cup C_2$. Let $Q_0 = Q_0^{(1)} \cup Q_0^{(2)} \cup Q_0^{(3)}$.

COROLLARY 1.2. — *Let $\beta > 1$ be a Salem number which does not belong to C_1 . Then β belongs to the class Q_0 .*

The attribution of Salem numbers to C_1 , $Q_0^{(1)}$, $Q_0^{(2)}$ and $Q_0^{(3)}$ is an open problem in general, except in low degree. Boyd [9] [12] has shown that Salem numbers of degree 4 belong to C_2 , hence to $Q_0^{(1)}$. It is also the case of some Salem numbers of degree 6 and ≥ 8 in the framework of a probabilistic model [11] [12]. In Section 5 we ask the question whether Corollary 1.2 could still be true for Perron numbers.

The definition of the class Q_0 does not make any allusion to β , i.e. to $M(\beta)$, to the conjugates of β , to the minimal polynomial of β or to its length, etc, but takes only into account the quotients of the gaps in $d_\beta(1)$. Hence this class Q_0 can be applied to real numbers $\beta > 1$ in full generality instead of only to algebraic numbers > 1 . The question whether there exist transcendental numbers $\beta > 1$ which belong to the class Q_0 was asked in [8]; what proportion appears in each subclass ? Examples of transcendental numbers (Komornik-Loreti constant [2] [29], Sturmian numbers [14]) in Q_0 are given in Section 5.

In the present note, we deal with the algebraicity of values of "gappy" series, deduced from $d_\beta(1)$, at the algebraic point β^{-1} . In a related context, more related to transcendency, Nishioka [36] and Corvaja Zannier [16] have followed different paths and applied the Subspace Theorem [43] to deduce different results.

2. Definitions

For $x \in \mathbb{R}$ the integer part of x is $[x]$ and its fractional part $\{x\} = x - [x]$. The smallest integer larger than or equal to x is denoted by $\lceil x \rceil$. For $\beta > 1$ a real number and $z \in [0, 1]$ we denote by $T_\beta(z) = \beta z \pmod{1}$ the β -transform on $[0, 1]$ associated with β [38] [40], and iteratively, for all integers $j \geq 0$, $T_\beta^{j+1}(z) := T_\beta(T_\beta^j(z))$, where by convention $T_\beta^0 = Id$.

Let $\beta > 1$ be a real number. A beta-representation (or β -representation, or representation in base β) of a real number $x \geq 0$ is given by an infinite sequence $(x_i)_{i \geq 0}$ and an integer $k \in \mathbb{Z}$ such that $x = \sum_{i=0}^{+\infty} x_i \beta^{-i+k}$, where the digits x_i belong to a given alphabet $(\subset \mathbb{N})$ [23] [24] [34]. Among all the beta-representations of a real number $x \geq 0$, $x \neq 1$, there exists a particular one called Rényi β -expansion, which is obtained via the greedy algorithm: in this case, k satisfies $\beta^k \leq x < \beta^{k+1}$ and the digits

$$(2.1) \quad x_i := \lfloor \beta T_\beta^i \left(\frac{x}{\beta^{k+1}} \right) \rfloor \quad i = 0, 1, 2, \dots,$$

belong to the finite canonical alphabet $\mathbb{A}_\beta := \{0, 1, 2, \dots, \lceil \beta - 1 \rceil\}$. If β is an integer, then $\mathbb{A}_\beta := \{0, 1, 2, \dots, \beta - 1\}$; if β is not an integer, then $\mathbb{A}_\beta := \{0, 1, 2, \dots, \lfloor \beta \rfloor\}$. We denote by

$$(2.2) \quad \langle x \rangle_\beta := x_0 x_1 x_2 \dots x_k . x_{k+1} x_{k+2} \dots$$

the couple formed by the string of digits $x_0 x_1 x_2 \dots x_k x_{k+1} x_{k+2} \dots$ and the position of the dot, which is at the k -th position (between x_k and x_{k+1}). By definition the integer part (in base β) of x is $\sum_{i=0}^k x_i \beta^{-i+k}$ and its fractional part (in base β) is $\sum_{i=k+1}^{+\infty} x_i \beta^{-i+k}$. If a Rényi β -expansion ends in infinitely many zeros, it is said to be finite and the ending zeros are omitted. If it is periodic after a certain rank, it is said to be eventually periodic (the period is the smallest finite string of digits possible, assumed not to be a string of zeros); for the substitutive approach see [19] [39].

The Rényi β -expansion which plays an important role in the theory is the Rényi β -expansion of 1, denoted by $d_\beta(1)$ and defined as follows: since $\beta^0 \leq 1 < \beta$, the value $T_\beta(1/\beta)$ is set to 1 by convention. Then using the formulae (2.1)

$$(2.3) \quad t_1 = \lfloor \beta \rfloor, t_2 = \lfloor \beta \{ \beta \} \rfloor, t_3 = \lfloor \beta \{ \beta \{ \beta \} \} \rfloor, \dots$$

The writing

$$d_\beta(1) = 0.t_1 t_2 t_3 \dots$$

corresponds to

$$1 = \sum_{i=1}^{+\infty} t_i \beta^{-i}.$$

Links between the set \mathbb{Z}_β of beta-integers and $d_\beta(1)$ are evoked in [18] [21] [27] [26] [46]. A real number $\beta > 1$ such that $d_\beta(1)$ is finite or eventually periodic is called a *beta-number* or more recently a *Parry number* (this recent terminology appears in [18]). In particular, it is called a *simple beta-number* or a *simple Parry number* when $d_\beta(1)$ is finite. Beta-numbers (Parry numbers) are algebraic integers ([38]) and all their conjugates lie within a compact subset which looks like a fractal in the complex plane [44]. The conjugates of beta-numbers are all bounded above in modulus by the golden mean $\frac{1}{2}(1 + \sqrt{5})$ ([20] [44]).

3. Proof of Theorem 1.1

Since algebraic numbers $\beta > 1$ for which the Rényi β -expansion $d_\beta(1)$ of 1 is finite are excluded, we may consider that β does not belong to \mathbb{N} . Indeed, if $\beta = h \in \mathbb{N}$, then $d_h(1) = 0.h$ is finite (Lothaire [34], Chap. 7). If $\beta \notin \mathbb{N}$, then $\lceil \beta - 1 \rceil = \lfloor \beta \rfloor$ and the alphabet A_β equals $\{0, 1, 2, \dots, \lfloor \beta \rfloor\}$, where $\lfloor \beta \rfloor$ denotes the greatest integer smaller than or equal to β .

Let $f(z) := \sum_{i=1}^{+\infty} t_i z^i$ be the “gappy” power series deduced from the representation $d_\beta(1) = 0.t_1 t_2 t_3 \dots$ associated with the β -shift (*gappy* in the sense of Theorem 1.1). Since $d_\beta(1)$ is assumed to be infinite, its radius of convergence is 1. By definition, it satisfies

$$(3.1) \quad f(\beta^{-1}) = 1,$$

which means that the function value $f(\beta^{-1})$ is algebraic, equal to 1, at the real algebraic number β^{-1} in the open disk of convergence $D(0, 1)$ of $f(z)$ in the complex plane. This fact is a general intrinsic feature of the Rényi expansion process which leads to the following important consequence by the theory of admissible power series of Mahler [35].

PROPOSITION 3.1. —

$$(3.2) \quad \limsup_{n \rightarrow +\infty} \frac{s_n}{m_n} < +\infty.$$

Proof. — This is a consequence of Theorem 1 in [35]. Indeed, if we assume that there exists a sequence of integers (n_i) which tends to infinity such that $\lim_{i \rightarrow +\infty} s_{n_i}/m_{n_i} = +\infty$, then $f(z)$ would be *admissible* in the sense of [35]. Since $f(z)$ is a power series with nonnegative coefficients, which is not a polynomial, the function value $f(\beta^{-1})$ should not be algebraic. But it equals 1. Contradiction. \square

Let us improve Proposition 3.1. Assume that

$$(3.3) \quad \limsup \frac{s_n}{m_n} > \frac{\log(M(\beta))}{\log(\beta)}$$

and show the contradiction with (1.1) and (1.2). Recall that, if

$$P_\beta(X) = \sum_{i=0}^d \alpha_i X^i = \alpha_d \prod_{i=0}^{d-1} (X - \beta^{(i)})$$

with $d \geq 1$, $\alpha_0 \alpha_d \neq 0$, denotes the minimal polynomial of $\beta = \beta^{(0)} > 1$, having $\beta^{(1)}, \beta^{(2)}, \dots, \beta^{(d-1)}$ as conjugates, the Mahler measure of β is by definition

$$M(\beta) := |\alpha_d| \prod_{i=0}^{d-1} \max\{1, |\beta^{(i)}|\}.$$

Gütting [28] has shown that the approximation of algebraic numbers by algebraic numbers is fairly difficult to realize by polynomials. In the present proof, we use a version of Liouville’s inequality which generalizes approximation theorems obtained by Gütting [28], and apply it to the values of the “polynomial tails” of the power series $f(z)$ at the algebraic number β^{-1} , to obtain the contradiction. Let us write

$$(3.4) \quad f(z) = \sum_{n=0}^{+\infty} Q_n(z)$$

with

$$(3.5) \quad Q_n(z) := \sum_{i=s_n}^{m_{n+1}} t_i z^i, \quad n = 0, 1, 2, \dots$$

By construction the polynomials $Q_n(z)$, of degree m_{n+1} , are not identically zero and $Q_n(1) > 0$ is an integer for all $n \geq 0$.

Denote by $S_n(z) = -1 + \sum_{i=1}^{m_n} t_i z^i$ the m_n th-section polynomial of the power series $f(z) - 1$ for all $n \geq 1$. Recall that, for $R(X) = \sum_{i=0}^v \alpha_i X^i \in \mathbb{Z}[X]$, $L(R) := \sum_{i=0}^v |\alpha_i|$ denotes the length of the polynomial $R(X)$. We have: $L(S_n) = 1 + \sum_{i=1}^{m_n} t_i = 1 + \sum_{j=0}^{n-1} Q_j(1)$. From Theorem 5 in [28] we deduce that only one of the following cases (G-i) or (G-ii) holds, for all $n \geq 1$:

$$(3.6) \quad (G - i) \quad S_n(\beta^{-1}) = 0,$$

$$(3.7) \quad (G - ii) \quad |S_n(\beta^{-1})| \geq \frac{1}{\left(1 + \sum_{j=0}^{n-1} Q_j(1)\right)^{d-1} \left(L(P_\beta^*)\right)^{m_n}},$$

where $P_\beta^*(X) = X^d P_\beta(1/X)$ is the reciprocal polynomial of the minimal polynomial of β , for which $L(P_\beta) = L(P_\beta^*) \in \mathbb{N} \setminus \{0, 1\}$.

Case (G-i) is impossible for any n . Indeed, if there exists an integer $n_0 \geq 1$ such that (G-i) holds, then, since all the digits t_i are positive and that $\beta^{-1} > 0$, we would have $t_i = 0$ for all $i \geq s_{n_0}$. This would mean that the Rényi expansion of 1 in base β is finite, which is excluded by assumption. Contradiction. Therefore, the only possibility is (G-ii), which holds for all integers $n \geq 1$. From Lemma 3.10 and Liouville's inequality (Proposition 3.14) in Waldschmidt [47] the inequality (G-ii) can be improved to

$$(3.8) \quad (L - ii) \quad |S_n(\beta^{-1})| \geq \frac{1}{\left(1 + \sum_{j=0}^{n-1} Q_j(1)\right)^{d-1} (M(\beta))^{m_n}}.$$

This improvement may be important; recall the well-known inequalities:

$$M(\beta) \leq L(P_\beta) \leq 2^{\deg(\beta)} M(\beta)$$

and see [47] p. 113 for comparison with different heights. On the other hand, since $|S_n(\beta^{-1})| = \sum_{i=s_n}^{+\infty} t_i \beta^{-i}$ for all integers $n \geq 1$, we deduce

$$(3.9) \quad |S_n(\beta^{-1})| \leq \frac{\lfloor \beta \rfloor}{1 - \beta^{-1}} \beta^{-s_n} \quad n = 1, 2, \dots$$

Putting together (3.8) and (3.9), we deduce that

$$(3.10) \quad \frac{\beta^{s_n}}{\left(1 + \sum_{j=0}^{n-1} Q_j(1)\right)^{d-1} M(\beta)^{m_n}} \leq \frac{\lfloor \beta \rfloor}{1 - \beta^{-1}}$$

should be satisfied for $n = 1, 2, 3, \dots$. Denote

$$u_n := \frac{\beta^{s_n}}{\left(1 + \sum_{j=0}^{n-1} Q_j(1)\right)^{d-1} M(\beta)^{m_n}} \quad \text{for all } n \geq 1.$$

Proof of (1.1): from (3.3) assumed to be true there exists a sequence of integers (n_i) which tends to infinity and an integer i_0 such that

$$\frac{s_{n_i}}{m_{n_i}} > \frac{\log(M(\beta))}{\log(\beta)} \quad \text{for all } i \geq i_0.$$

Now,

$$(3.11) \quad \left(\frac{1}{1 + \lfloor \beta \rfloor m_{n_i}} \right)^{d-1} \left(\frac{\beta^{\frac{s_{n_i}}{m_{n_i}}}}{M(\beta)} \right)^{m_{n_i}} \leq \frac{1}{\left(1 + \sum_{j=0}^{n_i-1} Q_j(1) \right)^{d-1}} \left(\frac{\beta^{\frac{s_{n_i}}{m_{n_i}}}}{M(\beta)} \right)^{m_{n_i}} \leq u_{n_i}.$$

For $i \geq i_0$ the inequality

$$(3.12) \quad 1 = \frac{\beta^{\frac{\log(M(\beta))}{\log(\beta)}}}{M(\beta)} < \frac{\beta^{\frac{s_{n_i}}{m_{n_i}}}}{M(\beta)}$$

holds. This implies that the left-hand side member of (3.11) tends exponentially to infinity when i tends to infinity. By (3.11) this forces u_{n_i} to tend to infinity. The contradiction now comes from (3.10) since the sequence (u_n) should be uniformly bounded.

Proof of (1.2): for $n = 1, 2, \dots$, let us rewrite the n -th quotient

$$(3.13) \quad \frac{u_{n+1}}{u_n} = \frac{\beta^{s_{n+1}-s_n}}{M(\beta)^{m_{n+1}-m_n}} \frac{\left(1 + \sum_{j=0}^{n-1} Q_j(1) \right)^{d-1}}{\left(1 + \sum_{j=0}^n Q_j(1) \right)^{d-1}}$$

as

$$(3.14) \quad \frac{u_{n+1}}{u_n} = \frac{\left(\frac{\beta^{\frac{s_{n+1}-s_n}{m_{n+1}-m_n}}}{M(\beta)} \right)^{m_{n+1}-m_n}}{(m_{n+1} - m_n + 1)^{(d-1)}} \left[\frac{(m_{n+1} - m_n + 1)^{(d-1)} \left(1 + \sum_{j=0}^{n-1} Q_j(1) \right)^{d-1}}{\left(1 + \sum_{j=0}^n Q_j(1) \right)^{d-1}} \right]$$

and denote

$$(3.15) \quad U_n := \frac{1}{(m_{n+1} - m_n + 1)^{(d-1)}} \left(\frac{\beta^{\frac{s_{n+1}-s_n}{m_{n+1}-m_n}}}{M(\beta)} \right)^{m_{n+1}-m_n}$$

and

$$(3.16) \quad W_n := (m_{n+1} - m_n + 1)^{(d-1)} \left(\frac{1 + \sum_{j=0}^{n-1} Q_j(1)}{1 + \sum_{j=0}^n Q_j(1)} \right)^{d-1}$$

so that $u_{n+1}/u_n = U_n W_n$.

LEMMA 3.2. —

$$(3.17) \quad 0 < \liminf_{n \rightarrow +\infty} W_n$$

Proof. — Assume the contrary. Then there exists a subsequence (n_i) of integers which tends to infinity such that $\lim_{i \rightarrow +\infty} W_{n_i} = 0$. In other terms, for all $\epsilon > 0$, there exists i_1 such that $i \geq i_1$ implies $W_{n_i} \leq \epsilon$, equivalently

$$(3.18) \quad (m_{n_i+1} - m_{n_i} + 1) \left(1 + \sum_{j=0}^{n_i-1} Q_j(1)\right) \leq \epsilon^{\frac{1}{d-1}} \times \left(1 + \sum_{j=0}^{n_i} Q_j(1)\right).$$

Since, by hypothesis, $t_{s_n} \geq 1$ and $t_{m_n+1} \geq 1$ for all $n \geq 1$, we have: $n_i \leq 1 + \sum_{j=0}^{n_i-1} Q_j(1)$. On the other hand, $Q_{n_i}(1) \leq \lfloor \beta \rfloor (m_{n_i+1} - m_{n_i} + 1)$. Then, from (3.18) with ϵ taken equal to 1, we would have

$$(3.19) \quad n_i \leq 1 + \sum_{j=0}^{n_i-1} Q_j(1) \leq \frac{Q_{n_i}(1)}{(m_{n_i+1} - m_{n_i} + 1) - 1} \leq \lfloor \beta \rfloor \times \frac{m_{n_i+1} - m_{n_i} + 1}{m_{n_i+1} - m_{n_i}} \leq \frac{3}{2} \lfloor \beta \rfloor.$$

But the left-hand side member of (3.19) tends to infinity which is impossible. Contradiction. □

Let us assume that (1.2) does not hold and show the contradiction ; that is, assume that $\liminf_{n \rightarrow +\infty} (m_{n+1} - m_n) = +\infty$ and $\limsup_{n \rightarrow +\infty} (s_{n+1} - s_n) / (m_{n+1} - m_n) > \log(M(\beta)) / \log(\beta)$ hold. Then

$$(3.20) \quad 1 = \frac{\beta^{\frac{\log(M(\beta))}{\log(\beta)}}}{M(\beta)} < \frac{\beta^{\frac{s_{n_i+1} - s_{n_i}}{m_{n_i+1} - m_{n_i}}}}{M(\beta)}$$

for some sequence of integers (n_i) which tends to infinity. This proves that $\limsup_{n \rightarrow +\infty} U_n = +\infty$ since $\lim_{i \rightarrow +\infty} U_{n_i} = +\infty$ exponentially, by (3.15) and (3.20).

By Lemma 3.2 there exists $r > 0$ such that $W_n \geq r$ for all n large enough. Therefore, $u_{n+1}/u_n = U_n W_n \geq r U_n$ for all n large enough. Since $\limsup_{n \rightarrow +\infty} U_n = +\infty$ we conclude that $\limsup_{n \rightarrow +\infty} u_{n+1}/u_n = +\infty$, hence that $\limsup_{n \rightarrow +\infty} u_n = +\infty$. This contradicts (3.10) and proves (1.2).

4. A direct proof of Corollary 1.2

Let $\beta > 1$ be a Salem number such that $\beta \notin C_1$. Using the notations of Theorem 1.1 we show that the assumption

$$(4.1) \quad \limsup_{n \rightarrow +\infty} \frac{s_n}{m_n} > 1$$

leads to a contradiction.

Denote by \mathbb{K} the algebraic number field $\mathbb{Q}(\beta)$, considered as a multivalued field with the product formula [15] [43] (see also [30]).

The present proof is merely an adaptation of that of Theorem 1 in [1], though the aims are different, and therefore does not merit publication. We simply point out a few hints for the interested reader.

The main result which is used is Corollary 1 of the Main Theorem in [15], as in [1]. This is a version of the Thue-Siegel-Roth Theorem given by Corvaja which is stronger than Roth Theorem for number fields [31] [43] to the extent it allows us to introduce a *missing proportion of places* of \mathbb{K} by considering the projective approximation of the point at infinity in $\mathbb{P}^1(\mathbb{K})$. Since β is a Salem number, it is a unit [4]. Hence, this missing proportion has just to be chosen among the pairwise distinct Archimedean places of \mathbb{K} .

5. On the class \mathcal{Q}_0

5.1. Perron numbers

Let us give, after Solomyak ([44], p 483), the example of a Perron number which is not a beta-number and therefore which is not in the class \mathcal{C}_2 , without knowing whether it is in the class \mathcal{Q}_0 . This example allows us to estimate the sharpness of the upper bound $\log(M(\beta))/\log(\beta)$ in (1.1). Recall that a real number $\beta > 1$ is a beta-number if the orbit of $x = 1$ under the transformation $T_\beta : x \rightarrow \beta x \pmod{1}$ is finite [34] [39]. The set of all conjugates of all beta-numbers is the union of the closed unit disc in the complex plane and the set of reciprocals of zeros of the function class $\{f(z) = 1 + \sum a_j z^j \mid 0 \leq a_j \leq 1\}$. The closure of this domain, say Φ , is compact and was studied by Flatto, Lagarias and Poonen [20] and Solomyak [44]. After [44], the Perron number $\beta = \frac{1}{2}(1 + \sqrt{13})$, dominant root of $P_\beta(X) = X^2 - X - 3$, is not a beta-number, though its only conjugate $\beta' = \frac{1}{2}(1 - \sqrt{13})$ lies in the interior $\text{int}(\Phi)$. We have $M(\beta) = 3$. By Theorem 1.1 the "quotients of the gaps" are asymptotically bounded above by $\log(3)/\log(\beta) = 1.3171\dots$, a much better bound than the degree $d = 2$ of β (see Lemma 5.1). This does not suffice to conclude that $\frac{1}{2}(1 + \sqrt{13})$ belongs to \mathcal{Q}_0 .

Do all Perron numbers belong to \mathcal{Q}_0 ? Let $\beta > 1$ be a Perron number of degree $d \geq 2$ and denote by $\beta^{(1)}, \beta^{(2)}, \dots, \beta^{(d-1)}$ the conjugates of $\beta = \beta^{(0)}$, roots of the minimal polynomial $P_\beta(X)$ of β . Let $K_\beta := \max\{|\beta^{(i)}| \mid i = 1, 2, \dots, d-1\}$.

LEMMA 5.1. — Let $n = n_\beta$ (with $2 \leq n_\beta \leq d$) be the number of conjugates of β of modulus strictly greater than unity (including β). Then

$$(5.1) \quad \frac{\log(M(\beta))}{\log(\beta)} \leq n - \frac{n-1}{(d\beta)^{6d^3} \log \beta}.$$

Proof. — Obvious since (Lemma 2 in [33]): $K_\beta < \beta(1 - \frac{1}{(d\beta)^{6d^3}})$. \square

The upper bound (5.1) does not allow us to give a positive answer to the question and has probably to be improved.

5.2. Transcendental numbers

Let us show that the Komornik-Loreti constant [2] [29] belongs to $\mathbb{Q}_0^{(1)}$.

THEOREM 5.2. — There exists a smallest $q \in (1, 2)$ for which there exists a unique expansion of 1 as $1 = \sum_{n=1}^{\infty} \delta_n q^{-n}$, with $\delta_n \in \{0, 1\}$. Furthermore, for this smallest q , the coefficient δ_n is equal to 0 (respectively, 1) if the sum of the binary digits of n is even (respectively, odd). This number q can then be obtained as the unique positive solution of $1 = \sum_{n=1}^{\infty} \delta_n q^{-n}$. It is equal to 1.787231650...

This constant q is named Komornik-Loreti constant. Allouche and Cosnard [2] have shown the following result.

THEOREM 5.3. — The constant q is a transcendental number, where the sequence of coefficients $(\delta_n)_{n \geq 1}$ is the Prouhet-Thue-Morse sequence on the alphabet $\{0, 1\}$.

The uniqueness of the development of 1 in base q given by Theorem 5.2 allows us to write

$$d_q(1) = 0.\delta_1\delta_2\delta_3\dots,$$

the coefficients δ_n being the digits of the Rényi q -expansion of 1. Since the strings of zeros and 1's in the Prouhet-Thue-Morse sequence are known (Thue, 1906/1912; [3]) and uniformly bounded, the constant q belongs to the class $\mathbb{Q}_0^{(1)}$.

As second example, let us show that Sturmian numbers in the interval $(1, 2)$ (in the sense of [14]) belong to $\mathbb{Q}_0^{(1)}$.

A real number $\beta > 1$ is called a Sturmian number if $d_\beta(1)$ is a Sturmian word over a binary alphabet $\{a, b\}$, with $0 \leq a < b = \lfloor \beta \rfloor$. Chi and Kwon [14] have shown the following theorem.

THEOREM 5.4. — Every Sturmian number is transcendental.

Let us consider all the Sturmian numbers $\beta \in (1, 2)$ for which the two-letter alphabet is $\{0, 1\}$. For such numbers gappiness appears in $d_\beta(1)$ (in the sense of Theorem 1.1). By Theorem 3.3 in [14] strings of zeros, resp. of 1's, cannot be arbitrarily long. This gives the claim.

Acknowledgments

The author thanks J.-P. Allouche, B. Adamczewski, J. Bernat, V. Berthé, C. Frougny, J.-P. Gazeau and the anonymous referee for valuable comments, discussions and remarks. The author would like to thank Catriona MacLean for her careful rereading of the manuscript.

BIBLIOGRAPHY

- [1] B. ADAMCZEWSKI, “Transcendance “à la Liouville” de certains nombres réels”, *C. R. Acad. Sci. Paris* **338** (2004), no. I, p. 511-514.
- [2] J.-P. ALLOUCHE & M. COSNARD, “The Komornik-Loreti constant is transcendental”, *Amer. Math. Monthly* **107** (2000), p. 448-449.
- [3] J.-P. ALLOUCHE & J. SHALLIT, “The ubiquitous Prouhet-Thue-Morse sequence”, in *Sequences and Their Applications* (C. Ding, T. Helleseth & H. Niederreiter, eds.), Proceedings of SETA'98, Springer-Verlag, 1999, p. 1-16.
- [4] M.-J. BERTIN, A. DECOMPS-GUILLOUX, M. GRANDET-HUGOT, M. PATHIAUX-DELEFOSSE & J.-P. SCHREIBER, *Pisot and Salem Numbers*, Birkhäuser, 1992.
- [5] A. BERTRAND-MATHIS, “Questions diverses relatives aux systèmes codés : applications au θ -shift”, preprint.
- [6] ———, “Développements en base Pisot et répartition modulo 1”, *C. R. Acad. Sci. Paris* **285** (1977), no. A, p. 419-421.
- [7] ———, “Développements en base θ et répartition modulo 1 de la suite $(x\theta^n)$ ”, *Bull. Soc. Math. Fr.* **114** (1986), p. 271-324.
- [8] F. BLANCHARD, “ β -expansions and Symbolic Dynamics”, *Theoret. Comput. Sci.* **65** (1989), p. 131-141.
- [9] D. BOYD, “Salem numbers of degree four have periodic expansions”, in *Théorie des Nombres - Number Theory* (Berlin and New York), Walter de Gruyter & Co., Eds. J.M. de Koninck and C. Levesque, 1989, p. 57-64.
- [10] ———, “On beta expansions for Pisot numbers”, *Math. Comp.* **65** (1996), p. 841-860.
- [11] ———, “On the beta expansion for Salem numbers of degree 6”, *Math. Comp.* **65** (1996), p. 861-875.
- [12] ———, “The beta expansions for Salem numbers”, in *Organic Mathematics* (Providence, RI), Canad. Math. Soc. Conf. Proc. 20, A.M.S., 1997, p. 117-131.
- [13] C. BURDIK, C. FROUGNY, J.-P. GAZEAU & R. KREJCAR, “Beta-integers as natural counting systems for quasicrystals”, *J. Phys. A: Math. Gen.* **31** (1998), p. 6449-6472.
- [14] D. P. CHI & D. KWON, “Sturmian words, β -shifts, and transcendence”, *Theoret. Comput. Sci.* **321** (2004), p. 395-404.
- [15] P. CORVAJA, “Autour du Théorème de Roth”, *Monath. Math.* **124** (1997), p. 147-175.

- [16] P. CORVAJA & U. ZANNIER, “Some New Applications of the Subspace Theorem”, *Compositio Mathematica* **131** (2002), p. 319-340.
- [17] M. DENKER, C. GRILLENBERGER & K. SIGMUND, *Ergodic Theory on compact spaces*, Springer Lecture Notes in Math. 527, 1976.
- [18] A. ELKHARRAT, C. FROUGNY, J.-P. GAZEAU & J.-L. VERGER-GAUGRY, “Symmetry groups for beta-lattices”, *Theor. Comp. Sci.* **319** (2004), p. 281-305.
- [19] S. FABRE, “Substitutions et β -systèmes de numération”, *Theoret. Comput. Sci.* **137** (1995), p. 219-236.
- [20] L. FLATTO, J. LAGARIAS & B. POONEN, “The zeta function of the beta transformation”, *Ergod. Th. and Dynam. Sys.* **14** (1994), p. 237-266.
- [21] C. FROUGNY, J.-P. GAZEAU & R. KREJCAR, “Additive and multiplicative properties of point sets based on beta-integers”, *Theoret. Comput. Sci.* **303** (2003), p. 491-516.
- [22] C. FROUGNY & B. SOLOMYAK, “Finite beta-expansions”, *Ergod. Theor. Dynam. Sys.* **12** (1992), p. 713-723.
- [23] C. FROUGNY, “Number Representation and Finite Automata”, *London Math. Soc. Lecture Note Ser.* **279** (2000), p. 207-228.
- [24] ———, “Algebraic Combinatorics on Words”, chap. Numeration systems, 7, Cambridge University Press, 2003.
- [25] J.-P. GAZEAU, “Pisot-Cyclotomic Integers for Quasilattices”, in *The Mathematics of Long-Range Aperiodic Order* (Dordrecht), Ed. R. V. Moody, Kluwer Academic Publisher, 1997, p. 175-198.
- [26] J.-P. GAZEAU & J.-L. VERGER-GAUGRY, “Geometric study of the set of β -integers for a Perron number and mathematical quasicrystals”, *J. Th. Nombres Bordeaux* **16** (2004), p. 1-25.
- [27] ———, “Diffraction spectra of weighted Delone sets on β -lattices with β a quadratic unitary Pisot number”, *Ann. Inst. Fourier*, 2006.
- [28] R. GÜTING, “Approximation of algebraic numbers by algebraic numbers”, *Michigan Math. J.* **8** (1961), p. 149-159.
- [29] V. KOMORNIK & P. LORETI, “Unique developments in non-integer bases”, *Amer. Math. Monthly* **105** (1998), p. 636-639.
- [30] S. LANG, “Fundamentals of Diophantine Geometry”, *Springer-Verlag, New York* (1983), p. 158-187.
- [31] W. J. LEVEQUE, “Topics in Number Theory”, *Addison-Wesley II* (1956), p. 121-160.
- [32] D. LIND, “The entropies of topological Markov shifts and a related class of algebraic integers”, *Erg. Th. Dyn. Syst.* **4** (1984), p. 283-300.
- [33] ———, “Matrices of Perron numbers”, *J. Number Theory* **40** (1992), p. 211-217.
- [34] M. LOTHAIRE, *Algebraic Combinatorics on Words*, Cambridge University Press, 2003.
- [35] K. MAHLER, “Arithmetic properties of lacunary power series with integral coefficients”, *J. Austr. Math. Soc.* **5** (1965), p. 56-64.
- [36] K. NISHIOKA, “Algebraic independence by Mahler’s method and S-units equations”, *Compositio Math.* **92** (1994), p. 87-110.
- [37] A. OSTROWSKI, “On representation of analytical functions by power series”, *J. London Math. Soc.* **1** (1926), p. 251-263, (*Addendum*), *ibid* **4** (1929), p. 32.
- [38] W. PARRY, “On the β -expansions of real numbers”, *Acta Math. Acad. Sci. Hung.* **11** (1960), p. 401-416.
- [39] N. PYTHÉAS FOGG, *Substitutions in dynamics, arithmetics and combinatorics*, Springer Lecture Notes in Math. 1794, 2003.
- [40] A. RÉNYI, “Representations for real numbers and their ergodic properties”, *Acta Math. Acad. Sci. Hung.* **8** (1957), p. 477-493.

- [41] J. SCHMELING, “Symbolic dynamics for β -shift and self-normal numbers”, *Ergod. Th. & Dynam. Sys.* **17** (1997), p. 675-694.
- [42] K. SCHMIDT, “On periodic expansions of Pisot numbers and Salem numbers”, *Bull. London Math. Soc.* **12** (1980), p. 269-278.
- [43] W. M. SCHMIDT, *Diophantine Approximations and Diophantine Equations*, Springer Lecture Notes in Math. 1467, 1991.
- [44] B. SOLOMYAK, “Conjugates of beta-numbers and the zero-free domain for a class of analytic functions”, *Proc. London Math. Soc. (3)* **68** (1993), p. 477-498.
- [45] W. P. THURSTON, “Groups, tilings, and finite state automata”, A.M.S. Colloquium Lectures, Boulder, Summer 1989.
- [46] J.-L. VERGER-GAUGRY, “On self-similar finitely generated uniformly discrete (SFU-) sets and sphere packings”, in *Number Theory and Physics* (L. Nyssen, ed.), IRMA Lectures in Mathematics and Theoretical Physics, E.M.S. Publishing House, 2006.
- [47] M. WALDSCHMIDT, *Diophantine Approximation on Linear Algebraic Groups. Transcendence Properties of the Exponential Function in Several Variables*, Springer-Verlag, Berlin, 2000.

Jean-Louis VERGER-GAUGRY
Université de Grenoble I
Institut Fourier
UMR CNRS 5582
BP 74 - Domaine Universitaire,
38402 Saint-Martin d'Hères (France)
jlverger@ujf-grenoble.fr