



ANNALES

DE

L'INSTITUT FOURIER

Nicola PAGANI

Chen–Ruan Cohomology of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$

Tome 63, n° 4 (2013), p. 1469-1509.

http://aif.cedram.org/item?id=AIF_2013__63_4_1469_0

© Association des Annales de l'institut Fourier, 2013, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

CHEN–RUAN COHOMOLOGY OF $\mathcal{M}_{1,n}$ AND $\overline{\mathcal{M}}_{1,n}$

by Nicola PAGANI (*)

ABSTRACT. — In this work we compute the Chen–Ruan cohomology of the moduli spaces of smooth and stable n -pointed curves of genus 1. In the first part of the paper we study and describe stack theoretically the twisted sectors of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$. In the second part, we study the orbifold intersection theory of $\overline{\mathcal{M}}_{1,n}$. We suggest a definition for an orbifold tautological ring in genus 1, which is a subring of both the Chen–Ruan cohomology and of the stringy Chow ring.

RÉSUMÉ. — Dans ce travail on calcule la cohomologie de Chen–Ruan de l'espace de modules des courbes lisses et stables de genre 1 avec n points marqués. Dans la première partie on étudie et on décrit les secteurs tordus de $\mathcal{M}_{1,n}$ et $\overline{\mathcal{M}}_{1,n}$, en tant que champs.

Dans la deuxième partie, on étudie la théorie d'intersection orbifold de $\overline{\mathcal{M}}_{1,n}$. On donne une définition possible de l'anneau tautologique orbifold en genre 1, comme sous-anneau simultanément de la cohomologie de Chen–Ruan et de l'anneau de Chow orbifold.

1. Introduction

Motivated by physics, Chen–Ruan cohomology was introduced in the paper [6] in the analytic category, and in the two papers by Abramovich–Graber–Vistoli [1] and [2] in the algebraic category. This has produced two parallel objects: the Chen–Ruan cohomology and the stringy Chow ring, which provide the basis to develop the quantum cohomology ring of an orbifold. This cohomology ring recovers as a subalgebra the ordinary rational

Keywords: moduli spaces, Gromov–Witten, orbifold, cohomology, tautological ring.

Math. classification: 14H10, 14N35, 55N32, 14D23, 14H37, 55P50.

(*) I would like to acknowledge my advisor, Barbara Fantechi, for suggesting to me the topic of this work and for helpful discussions and suggestions. I would like to thank Gilberto Bini, Carel Faber and Orsola Tommasi for sharing with me their ideas devoted to further developments of the present work. A special thank to Sebastian Krug, who discovered some mistakes in the previous version of this preprint, and to the anonymous referee, who improved the quality of the manuscript. This work was supported by SISSA, GNSAGA, the Clay Mathematical Institute and KTH, Royal Institut of Technology.

cohomology ring of the topological space that underlies the orbifold. As a vector space, the Chen–Ruan cohomology of X is simply the cohomology of the inertia stack of X . If X is an orbifold, its *inertia stack* $I(X)$ is, loosely speaking, constructed as the disjoint union, for g in the stabilizer of some point x of X , of the locus stabilized by g in X (see Definition 3.1). As an example, the orbifold X itself appears as a connected component of $I(X)$, as the locus fixed by the identity automorphism, which is trivially in the stabilizer group of every point. All the other connected components of the inertia stack $I(X)$ are usually called *twisted sectors*. In this paper we use the algebraic language, and whenever the word “orbifold” is mentioned, it stands for smooth Deligne–Mumford stack.

Among the first examples of smooth Deligne–Mumford stacks in the literature there are the moduli of smooth pointed curves $\mathcal{M}_{g,n}$ and their compactifications $\overline{\mathcal{M}}_{g,n}$. It seems thus interesting to study their Chen–Ruan cohomology. This has been done so far for $\overline{\mathcal{M}}_{1,1}$ (a special case of weighted projective space) and for \mathcal{M}_2 and $\overline{\mathcal{M}}_2$ by Spencer [26] (see also [27]).

In the present work, we investigate the Chen–Ruan cohomology ring for $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$ with rational coefficients, assuming knowledge of the cohomology of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$. We show how it is possible to describe the stringy Chow ring in a similar fashion. Indeed we show that for each twisted sector, the cycle map from the Chow ring to cohomology is an isomorphism.

The main results of this paper are the complete stack-theoretic description of the twisted sectors, and the explicit computation of the Chen–Ruan product as an extension of the usual cup product.

THEOREM 1.1. — (*Theorem 3.22, Corollary 3.23*) *Each twisted sector of $\overline{\mathcal{M}}_{1,n}$ is isomorphic to*

$$A \times \overline{\mathcal{M}}_{0,n_1} \times \overline{\mathcal{M}}_{0,n_2} \times \overline{\mathcal{M}}_{0,n_3} \times \overline{\mathcal{M}}_{0,n_4},$$

where the $n_i \geq 3$ are integers and A is in the set

$$\{B\mu_3, B\mu_4, B\mu_6, \mathbb{P}(4, 6), \mathbb{P}(2, 4), \mathbb{P}(2, 2)\}.$$

Here BG is the classifying stack of principal G -bundles, and $\mathbb{P}(a, b)$ is a weighted projective stack.

THEOREM 1.2. — (*Theorem 7.2*) *The Chen–Ruan cohomology ring of $\overline{\mathcal{M}}_{1,n}$ is generated as an algebra over the ordinary cohomology ring of $\overline{\mathcal{M}}_{1,n}$ by the fundamental classes of the twisted sectors with explicit relations.*

The theory developed in this manuscript can be seen as an extension of the study of the cohomological properties of the moduli spaces of curves, initiated by Harer and Mumford ([19]) in the eighties. The latter framework has produced important results in algebraic geometry, topology, mathematical and theoretical physics, representation theory and number theory. On the other hand, our results are the first steps towards the study and the understanding of more recent topics of investigation, such as the stringy topology and the Gromov-Witten theory of the moduli spaces of curves (see, for example, [3]).

This paper is part of the PhD thesis [20], where the case of genus g bigger than 1 is also discussed. However, we believe that the genus 1 case can be conveniently described within a more explicit and elementary framework, mainly thanks to the fact that the automorphism groups of stable genus 1 marked curves are cyclic. In general we make an effort to limit to a minimum the use of technicalities, whereas various aspects of the theory could be developed in bigger generality to produce higher genera results, see [21], [22] and [23].

1.1. Description of the sections

In Section 2 we recall some known results that we will use and fix our notation. The complete, stack-theoretic description of the twisted sectors of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$ is given in Section 3, where we prove the first main result 1.1. This result allows us to compute the generating series of the orbifold Poincaré polynomials for $\overline{\mathcal{M}}_{1,n}$. In Section 4, we compute the Chen–Ruan cohomology of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$ as a graded vector space. To do so, we introduce the unconventional rational grading on the cohomology of the inertia stack, usually referred to as *age*, or *degree shifting number*, or *fermionic shift*. In Section 5, we describe the twisted sectors of the second inertia stack of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$. Here a simplification occurs, indeed we show that every double twisted sector is canonically isomorphic to a twisted sector. In Section 6 we begin the study of the orbifold intersection theory on $\overline{\mathcal{M}}_{1,n}$, we compute all excess intersection bundles, and their top Chern classes. Finally, in Section 7, we determine the Chen–Ruan cup product and we prove the second main result 1.2. In this section a proposal for an orbifold tautological ring for $\overline{\mathcal{M}}_{1,n}$ is motivated and advanced.

2. Foundation

2.1. General notation

We work in the category of schemes of finite type over \mathbb{C} . Although we treat only this case, some of our results can easily be extended to the case of an arbitrary field of characteristic different from 2 and 3. In the paper, algebraic stack means Deligne–Mumford stack. Intersection theory on schemes is defined in [10], on Deligne–Mumford stacks it is defined in [28]. We refer to these texts for the definitions and first properties of the Chow groups A_* . In this manuscript we work with cohomology and Chow ring with rational coefficients. Since all the spaces we consider are smooth, there is a standard identification of A^* with the dual of A_* .

We use \mathbb{G}_m to denote the group scheme of invertible multiplicative elements of \mathbb{C} . The discrete group subscheme of \mathbb{G}_m of the N -th roots of 1 is called μ_N . The generators of μ_2 , μ_4 and μ_6 are conventionally chosen to be respectively -1 , i and ϵ . Since we work over the complex numbers, we can fix $\epsilon = e^{\frac{2i\pi}{6}}$. We denote by S_n the group of permutations of $[n] := \{1, 2, \dots, n\}$.

If G is a finite abelian group, $G^\vee = \text{Hom}(G, \mathbb{C}^*)$ is the group of characters of G . We call the trivial G -gerbe over a point $BG = [\text{Spec}(\mathbb{C})/G]$; it is the classifying stack of the group G . If X is a scheme, an element in $\text{Pic}(X \times BG)$ is a pair (L, χ) where $L \in \text{Pic}(X)$ and $\chi \in G^\vee$. For a complete treatment on gerbes, we refer to the book [14], or to [5].

2.2. Notation for $\mathcal{M}_{g,n}$ and $\overline{\mathcal{M}}_{g,n}$, and some cohomological results

If I is a finite set $\overline{\mathcal{M}}_{g,I}$ is the moduli stack of stable genus g curves with marked points in the set I . If $I = \{i, \bullet\}$ then we conventionally define $\overline{\mathcal{M}}_{0,I}$ as a point labeled by i . When the set I is $[n]$, the set of the first n natural numbers, we write $\mathcal{M}_{g,n}$ and $\overline{\mathcal{M}}_{g,n}$ instead of $\mathcal{M}_{g,[n]}$ and $\overline{\mathcal{M}}_{g,[n]}$.

If $I \subset J$, then $\pi_I : \mathcal{M}_{g,J} \rightarrow \mathcal{M}_{g,I}$ is the morphism that remembers only the sections inside I . We give the same name to the morphism $\pi_I : \overline{\mathcal{M}}_{g,J} \rightarrow \overline{\mathcal{M}}_{g,I}$ that forgets all the sections but the ones in I and stabilizes. With this notation, let s_i be the i -th section of $\pi_{[n]} : \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$. By definition, the *cotangent line bundle* \mathbb{L}_i is the line bundle $s_i^*(\omega_{\pi_{[n]}})$, where $\omega_{\pi_{[n]}}$ is the relative dualizing sheaf. We also define $\psi_i := c_1(\mathbb{L}_i)$.

Let $k > 0$ and let (I_1, \dots, I_k) be a partition of $[n]$. We define $j_{g,k}$ as the morphism gluing the marked points labeled with the same symbol:

$$j_{g,k} : \overline{\mathcal{M}}_{g, \coprod_{i=1}^k \bullet_i} \times \overline{\mathcal{M}}_{g_1, I_1 \sqcup \bullet_1} \times \dots \times \overline{\mathcal{M}}_{g_k, I_k \sqcup \bullet_k} \rightarrow \overline{\mathcal{M}}_{g+\sum g_i, n},$$

note that $j_{g,k}$ depends on the partition I_1, \dots, I_k and on the choice of g_1, \dots, g_k , although we do not make this explicit in our notation. We also define:

$$j : \overline{\mathcal{M}}_{g,n \sqcup \bullet_1 \sqcup \bullet_2} \rightarrow \overline{\mathcal{M}}_{g+1,n}$$

as the morphism gluing together \bullet_1 and \bullet_2 . In this paper, we will be dealing with the case of genus 1 curves. We will be using several times the map $j_{1,k}$, where all the g_i are set equal to 0. We call this map simply j_k , so if I_1, \dots, I_k is a partition of $[n]$ we have the gluing map:

$$j_k : \overline{\mathcal{M}}_{1, \coprod_{i=1}^k \bullet_i} \times \overline{\mathcal{M}}_{0, I_1 \sqcup \bullet_1} \times \dots \times \overline{\mathcal{M}}_{0, I_k \sqcup \bullet_k} \rightarrow \overline{\mathcal{M}}_{1,n}.$$

The product space on the left admits projection maps onto each factor. We denote by p the projection map onto the first factor p , and with p_i the projection maps onto the genus 0 component with marked points in the set I_i .

We recall the definition and main properties of the tautological ring for the moduli spaces of curves.

DEFINITION 2.1. — ([8, Section 0.1]) *The system of tautological rings $R^*(\overline{\mathcal{M}}_{g,n})$ is defined to be the set of smallest \mathbb{Q} -subalgebras of the Chow rings*

$$R_{g,n}^* = R^*(\overline{\mathcal{M}}_{g,n}) \subset A^*(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$$

that is closed under push-forward via all forgetful and gluing maps.

Remark 2.2. — The system of tautological rings is closed under pull-back via the forgetful and the gluing maps. Each tautological ring is an S_n -representation via the action that permutes the points. We denote by $RH^*(\overline{\mathcal{M}}_{g,n})$ the image of $R^*(\overline{\mathcal{M}}_{g,n})$ under the cycle map to the ring of even cohomology classes.

DEFINITION 2.3. — *We define $B_{g,n}^*$ to be the smallest system of vector subspaces of the Chow rings $A^*(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ that contain the fundamental classes, and that are stable under push-forward via all gluing maps (see Definition 2.1). A boundary strata class is an element in $B_{g,n}^*$ that corresponds to a closed irreducible proper substack of $\overline{\mathcal{M}}_{g,n}$.*

Obviously, the tautological ring contains all boundary strata classes.

Notation 2.4. — If $I \subset [n], |I| \geq 2$, we denote by d_I the closure of the substack of $\overline{\mathcal{M}}_{1,n}$ of reducible nodal curves with two smooth components, where the marked points in the set I are on the genus 0 component and the marked points on the genus 1 curve are in the complement. The closure of the substack of $\overline{\mathcal{M}}_{1,n}$ of irreducible curves of geometric genus 0 is called

d_{irr} . We will sometimes indicate by d_I also the class $[d_I] \in H^2(\overline{\mathcal{M}}_{1,n})$ represented by the divisor d_I . These elements form a basis for $B_{1,n}^1$.

Analogously, given $I \subset [n]$, such that $|I| \geq 2$ and $[n] \setminus I \geq 2$, $\Delta_I = \Delta_{[n] \setminus I}$ is the sublocus of $\overline{\mathcal{M}}_{0,n}$ whose general element has two genus 0 components with marked points in I in the first one and in $[n] \setminus I$ in the second one. These elements generate $B_{0,n}^1$.

In general, $B_{g,n}^* \subset R_{g,n}^* \subset A^*(\overline{\mathcal{M}}_{g,n}, \mathbb{Q}) \rightarrow H^*(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ are all distinct. By [8, p.2], the ψ -classes defined in Section 2.2 are in the tautological ring. In genus 0 we have the equalities

$$(2.1) \quad B_{0,n}^* = R_{0,n}^* = A^*(\overline{\mathcal{M}}_{0,n}, \mathbb{Q}) = H^{2*}(\overline{\mathcal{M}}_{0,n}, \mathbb{Q});$$

moreover the cohomology is generated by the boundary divisor classes (see [17]). In the remainder of this section, we will see what we can say for the cohomology of $\overline{\mathcal{M}}_{1,n}$, in analogy with (2.1). The next proposition is a straightforward consequence of Theorem *, [16, Theorem 1.1].

PROPOSITION 2.5. — *The tautological ring $R^*(\overline{\mathcal{M}}_{1,n})$ is additively generated by boundary strata classes (see 2.3), so that $B_{1,n}^* = R_{1,n}^*$.*

So let us now introduce some results concerning the tautological ring of $\overline{\mathcal{M}}_{1,n}$. These results were originally claimed by Getzler in [12], and they were recently proved by Petersen in [24].

THEOREM 2.6. — *(claimed for the first time in [12], proved in [24]) The boundary strata classes span the even cohomology of $\overline{\mathcal{M}}_{1,n}$. Moreover the cycle map is injective when restricted to the tautological algebra $R^*(\overline{\mathcal{M}}_{1,n})$.*

Note that the second sentence of the statement follows, in Petersen’s proof, from the stronger fact that all relations among the generators of the even cohomology come from genus 0 relations, and from Getzler’s relation [12, Theorem 1.8], see [24] for more details. The tautological ring of $\overline{\mathcal{M}}_{1,n}$ has been studied in detail by Belorousski [4]. When $n \leq 10$, the picture is similar to the genus 0 case (2.1).

PROPOSITION 2.7. — *([4, Theorem 3.1.1, Theorem 3.6.3]) For $n \leq 10$ the following two equalities also hold*

$$R_{1,n}^* = A^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q}) = H^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q}).$$

It is well known that the eleventh cohomology group of $\overline{\mathcal{M}}_{1,11}$ is non-zero. It follows that the second and third equalities of the proposition above are no longer true for $n \geq 11$ (see for instance [15, p.2]).

Anyway, after Proposition 2.5, and Theorem 2.6, we obtain a decomposition of the cohomology in boundary strata classes and odd cohomology

$$(2.2) \quad H^*(\overline{\mathcal{M}}_{1,n}) = B_{1,n}^* \oplus H^{odd}(\overline{\mathcal{M}}_{1,n}).$$

The rings $R_{1,n}^* = B_{1,n}^*$ are, in general, not multiplicatively generated by the boundary divisors (as it happens in genus 0), by a result of Belorousski.

THEOREM 2.8. — [4, Chapter 3] *The Chow ring of $\overline{\mathcal{M}}_{1,n}$ is generated by the divisors precisely when $n \leq 5$.*

Anyway, it is possible to give a simple and geometric description of the additive generators of $R_{1,n}^*$ besides the product of boundary divisors. We denote by $R_{1,n}^{div}$ the subring of $R_{1,n}^*$ generated by the classes of the divisors.

DEFINITION 2.9. — *The banana locus (cf. [4, p.49]) is the locus in $\overline{\mathcal{M}}_{1,n}$ of curves whose general element has two rational components joined in two nodes. A subbanana cycle is a boundary strata class in $\overline{\mathcal{M}}_{1,n}$ that is contained in the banana locus. Let $R_{1,n}^{ban}$ be the vector subspace of $R_{1,n}^*$ generated by subbanana cycles.*

Then we have⁽¹⁾

$$(2.3) \quad R_{1,n}^* = B_{1,n}^* = R_{1,n}^{div} + R_{1,n}^{ban}.$$

Indeed, let us consider the complement of the banana locus in $\overline{\mathcal{M}}_{1,n}$. Reasoning by induction on the codimension, it is simple to see that the boundary strata classes on this complement can all be written as products of divisor classes.

3. The Chen–Ruan cohomology of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$ as vector spaces

3.1. Definition of Chen–Ruan cohomology as a vector space

The following is a natural stack associated to a stack X , which points to where X fails to be an algebraic space.

DEFINITION 3.1. — ([28, Definition 1.12]) *Let X be an algebraic stack. The inertia stack $I(X)$ of X is defined as the fiber product $X \times_{X \times X} X$ where both morphisms $X \rightarrow X \times X$ are the diagonal morphisms. There is a natural map $f : I(X) \rightarrow X$.*

⁽¹⁾We learned this from Belorousski's thesis [4], although it is not explicitly written there.

The construction of Chen–Ruan cohomology based on the definition of inertia orbifold was given for the first time in [6, Definition 3.2.3]. As observed in [1, Section 4.4], the latter is nothing but the coarse moduli space of the inertia stack we have just introduced. In [2, 7.3] the algebraic counterpart of Chen–Ruan cohomology is introduced, under the name of stringy Chow ring. It is built on the rigidification of the cyclotomic inertia stack introduced in [2, Section 3]. In this paper we work over \mathbb{C} , and all cohomologies are taken with rational coefficients. Therefore, the cohomologies of the inertia stack, of the cyclotomic inertia stack, of the rigidified cyclotomic inertia stack and of the inertia orbifold are all canonically isomorphic, since all of them share the same coarse moduli space.

Remark 3.2. — If Y is a twisted sector of $I(X)$, then the map f of Definition 3.1 restricts to a map $f|_Y : Y \rightarrow X$. In general $f|_Y : Y \rightarrow X$ is a composition of a stack covering and a closed embedding, as easily follows for instance from [28, Lemma 1.13], or from [6, 3.1.3]. In the present paper however, since all the stacks we consider are abelian orbifolds, the map $f|_Y$ is a closed embedding. So if Y is a twisted sector, it can be written as $Y = (Z, g)$, where Z is a closed substack of X and g is an automorphism in the generic stabilizer of Z .

DEFINITION 3.3. — *If X is an algebraic stack, the connected component of the inertia stack associated with the identity automorphism is called the untwisted sector of the inertia stack. All the remaining connected components are called the twisted sectors of $I(X)$. The latter are sometimes called also the twisted sectors of X .*

PROPOSITION 3.4. — [2, Corollary 3.1.4] *Let X be a smooth algebraic stack. Then the inertia stack $I(X)$ is smooth.*

DEFINITION 3.5. — *Let X be a smooth algebraic stack. Let T be a set of indices in bijection with the twisted sectors of $I(X)$. We say that the equality*

$$I(X) = X \sqcup \coprod_{i \in T} (X_i, g_i),$$

is a decomposition of the inertia stack of X in twisted sectors.

Notation 3.6. — In order to simplify the notation, if (A, g) , (A, g') are two twisted sectors, we shall write $(A, g/g')$ to denote the disjoint union of the two twisted sectors (A, g) and (A, g') in the inertia stack. When we write A we refer to the image of the closed embedding of the twisted sector inside the original stack X (see Remark 3.2).

We can then define the Chen–Ruan cohomology vector space.

DEFINITION 3.7. — ([6, Definition 3.2.3]) *Let X be a smooth algebraic stack. Then the Chen–Ruan cohomology is by definition*

$$H_{CR}^*(X, \mathbb{Q}) := H^*(I(X), \mathbb{Q})$$

as a rational vector space.

The Chen–Ruan cohomology decomposes as in Definition 3.5

$$H_{CR}^*(X, \mathbb{Q}) = H^*(X, \mathbb{Q}) \oplus \bigoplus_{i \in T} H^*(X_i, \mathbb{Q}).$$

3.2. The inertia stack of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$

The twisted sectors in case $n = 1$ are well known as a direct consequence of the Weierstrass Theorem. We refer to [25, III.1] for the basic material on this topic. First of all, recall that every curve of the form:

$$C_{a,b} = \{[x : y : z] \mid zy^2 = x^3 + az^2x + bz^3, \Delta := 4a^3 + 27b^2 \neq 0\} \subset \mathbb{P}^2$$

is a smooth genus 1 curve. If, instead:

$$C_{a,b} = \{[x : y : z] \mid zy^2 = x^3 + az^2x + bz^3, \Delta := 4a^3 + 27b^2 = 0, (a, b) \neq (0, 0)\},$$

then $C_{a,b}$ is a nodal curve of arithmetic genus 1, geometric genus 0 and one node. All genus 1 curves with a marked point admit this description.

THEOREM 3.8. — [25, III.1] (*Weierstrass representation*) *Let (C, P) be an elliptic curve, possibly nodal. Then there exist $(a, b) \in \mathbb{C}^2$ such that (C, P) is isomorphic to $(C_{a,b}, [0 : 1 : 0])$, where $C_{a,b}$ is as above. If α is an isomorphism of (C, P) with (D, Q) then there exists $\lambda \in \mathbb{G}_m$ such that, up to the isomorphism above, α is*

$$\alpha : \begin{cases} a \rightarrow \lambda^4 a \\ b \rightarrow \lambda^6 b \\ x \rightarrow \lambda^2 x \\ y \rightarrow \lambda^3 y \\ z \rightarrow z. \end{cases}$$

From this it follows that the moduli stack $\overline{\mathcal{M}}_{1,1}$ is isomorphic to the weighted projective stack $\mathbb{P}(4, 6)$.

Notation 3.9. — There are two elements of $\overline{\mathcal{M}}_{1,1}$ that are stabilized by the action of a group respectively isomorphic to μ_4 and μ_6 , we call them respectively C_4 and C_6 . These are classes of curves whose Weierstrass representation can be chosen respectively as:

$$C_4 := \{[x : y : z] \mid y^2z = x^3 + xz^2\} \subset \mathbb{P}^2,$$

$$C_6 := \{[x : y : z] \mid y^2z = x^3 + z^3\} \subset \mathbb{P}^2.$$

If (C, P) is an elliptic curve, and G is its automorphism group, then it can be identified canonically with μ_N for a certain $N \in \{2, 4, 6\}$.

Notation 3.10. — If (C, P) is an elliptic curve, and G is its automorphism group, then G acts effectively on $T_P^\vee(C)$, the cotangent space in C to P , which is canonically isomorphic to \mathbb{C} . We identify G with μ_N under this isomorphism.

The decomposition of the inertia stack of $\mathcal{M}_{1,1}$ and $\overline{\mathcal{M}}_{1,1}$ in twisted sectors (Definition 3.5), is a simple way to summarize the well-known facts that we have exposed in this section.

COROLLARY 3.11. — *With the notation introduced in Notation 3.6 and 3.10, the decomposition of the inertia stack of $\mathcal{M}_{1,1}$ in twisted sectors is:*

$$I(\mathcal{M}_{1,1}) = (\mathcal{M}_{1,1}, 1) \sqcup (\mathcal{M}_{1,1}, -1) \sqcup (C_4, i / -i) \sqcup (C_6, \epsilon / \epsilon^2 / \epsilon^4 / \epsilon^5)$$

and that of $\overline{\mathcal{M}}_{1,1}$ is:

$$I(\overline{\mathcal{M}}_{1,1}) = (\overline{\mathcal{M}}_{1,1}, 1) \sqcup (\overline{\mathcal{M}}_{1,1}, -1) \sqcup (C_4, i / -i) \sqcup (C_6, \epsilon / \epsilon^2 / \epsilon^4 / \epsilon^5).$$

3.2.1. The case of $\mathcal{M}_{1,n}$

We now study the inertia stack of $\mathcal{M}_{1,n}$. Note that if $n > 4$, the objects of $\mathcal{M}_{1,n}$ are rigid, and therefore in that range $I(\mathcal{M}_{1,n}) = \mathcal{M}_{1,n}$. A simple analysis of the fixed points of the action of μ_3, μ_4 and μ_6 on the curves C_4 and C_6 (see Notation 3.9) by Theorem 3.8 leads to three special points of $\overline{\mathcal{M}}_{1,n}, n \leq 3$.

Notation 3.12. — We call the point in $\mathcal{M}_{1,2}$ stabilized by i or $-i$ C'_4 , the point in $\mathcal{M}_{1,2}$ stabilized by ϵ^2 or ϵ^4 C'_6 , and the point in $\mathcal{M}_{1,3}$ stabilized by ϵ^2 or ϵ^4 C''_6 .

To complete the study of the loci fixed by automorphisms in $\mathcal{M}_{1,n}$, we shall need the loci fixed by the elliptic involution (according to Notation 3.10 we write it as (-1)). We give a special name to them.

DEFINITION 3.13. — Let $1 \leq i \leq 4$. We define A_i as the closed substack of $\mathcal{M}_{1,i}$ whose objects $A_i(S)$ are i -marked smooth genus 1 curves over S such that the sections are stabilized by the elliptic involution.

We shall see, as a consequence of Lemma 3.17, that A_i is connected for all i (note that $A_1 = \mathcal{M}_{1,1}$). What we have just discussed, leads to the following description:

COROLLARY 3.14. — The decomposition of the inertia stack of $\mathcal{M}_{1,n}$ (Notation 3.5, 3.6, 3.10) is:

- $I(\mathcal{M}_{1,1}) = \mathcal{M}_{1,1} \sqcup (\mathcal{M}_{1,1}, -1) \sqcup (C_4, i/ - i) \sqcup (C_6, \epsilon/\epsilon^2/\epsilon^4/\epsilon^5)$;
- $I(\mathcal{M}_{1,2}) = \mathcal{M}_{1,2} \sqcup (A_2, -1) \sqcup (C'_4, i/ - i) \sqcup (C'_6, \epsilon^2/\epsilon^4)$;
- $I(\mathcal{M}_{1,3}) = \mathcal{M}_{1,3} \sqcup (A_3, -1) \sqcup (C''_6, \epsilon^2/\epsilon^4)$;
- $I(\mathcal{M}_{1,4}) = \mathcal{M}_{1,4} \sqcup (A_4, -1)$;
- $I(\mathcal{M}_{1,n}) = \mathcal{M}_{1,n}$ if $n \geq 5$.

We collect the twisted sectors of $\mathcal{M}_{1,n}$ in the following table. Different rows correspond to different automorphisms, while the i -th column corresponds to the twisted sectors of $\mathcal{M}_{1,i}$.

	1	2	3	4
-1	A_1	A_2	A_3	A_4
ϵ^2/ϵ^4	C_6	C'_6	C''_6	\emptyset
$i/ - i$	C_4	C'_4	\emptyset	\emptyset
ϵ/ϵ^5	C_6	\emptyset	\emptyset	\emptyset

We now investigate the geometry of the spaces A_i introduced in Definition 3.13. In particular, this will give their cohomology.

Remark 3.15. — Using analytic methods (see [7, Chapter 3]), it is known that the coarse moduli space of A_i is a genus 0 quasiprojective curve, and also how many points are needed to compactify it. In the literature, the coarse moduli space for A_2 is known under the name of $Y_1(2) = Y_0(2)$. The coarse moduli space for $A_3 \cong A_4$ is usually called $Y(2)$. We here want to give an algebraic and stack-theoretic description of those spaces, that we could not find anywhere.

DEFINITION 3.16. — We define \overline{A}_i as the closure of A_i , in $\overline{\mathcal{M}}_{1,i}$.

We have already observed that the stack $\overline{A}_1 \cong \overline{\mathcal{M}}_{1,1}$ is isomorphic to $\mathbb{P}(4, 6)$ as a consequence of Theorem 3.8. Following the same strategy that can be used to prove the latter isomorphism, we can obtain the following result:

LEMMA 3.17. — *The stack \overline{A}_2 is isomorphic to the weighted projective stack $\mathbb{P}(2, 4)$. The stacks \overline{A}_3 and \overline{A}_4 are isomorphic to the weighted projective stack $\mathbb{P}(2, 2)$.*

Proof. — We first study the case of \overline{A}_2 . Let us define the following space:

$$B_1 := \{((a, b), [x : y : z]) \mid (a, b) \neq (0, 0), zy^2 = x^3 + az^2x + bz^3\} \subset \mathbb{A}_0^2 \times \mathbb{P}^2.$$

The projection onto the first factor, with the section $\sigma_1(a, b) := ((a, b), [0 : 1 : 0])$, describes this space as an elliptic fibration over \mathbb{A}_0^2 , so it determines a unique map $\phi : \mathbb{A}_0^2 \rightarrow \overline{\mathcal{M}}_{1,1}$, (here $\overline{\mathcal{C}}_{1,1}$ is the universal curve):

$$\begin{array}{ccc} B_1 & \longrightarrow & \overline{\mathcal{C}}_{1,1} \xrightarrow{\sim} \overline{\mathcal{M}}_{1,2} \\ \downarrow \left. \begin{array}{l} \sigma_1 \\ \downarrow \end{array} \right) & & \downarrow \left. \begin{array}{l} x_1 \\ \downarrow \end{array} \right) \\ \mathbb{A}_0^2 & \xrightarrow{\phi} & \overline{\mathcal{M}}_{1,1}. \end{array}$$

making the diagram cartesian. It is a well-known consequence of the Weierstrass theorem (3.8) made in families that the map ϕ factors via the quotient $[\mathbb{A}_0^2/\mathbb{G}_m]$, where \mathbb{G}_m acts with weights 4 and 6, and that the resulting map $\tilde{\phi} : [\mathbb{A}_0^2/\mathbb{G}_m] \rightarrow \overline{\mathcal{M}}_{1,1}$ is an isomorphism of stacks. The locus in B_1 cut out by the equation $y = 0$ surjects onto $\overline{A}_2 \subset \overline{\mathcal{M}}_{1,2}$. This locus is isomorphic to \mathbb{A}_0^2 with parameters (a, x) . Again as a consequence of the Weierstrass theorem, the action of \mathbb{G}_m with weights 4 and 2 can be factored out, to obtain an isomorphism of stacks $[\mathbb{A}_0^2/\mathbb{G}_m] \rightarrow \overline{A}_2$. The forgetful map $\overline{A}_2 \rightarrow \overline{A}_1$ lifts to the map of the charts $\mathbb{A}_0^2 \rightarrow \mathbb{A}_0^2$

$$(3.1) \quad (a, x) \rightarrow (a, -ax - x^3).$$

Now we study the case of \overline{A}_3 . Let:

$$B_2 := \{((a, x_1), [x : y : z]) \mid (a, x_1) \neq (0, 0), zy^2 = x^3 + az^2x + (-ax_1 - x_1^3)z^3\}.$$

In this case, the projection onto the first factor with the two sections

$$\begin{cases} \sigma_1(a, x_1) = ((a, x_1), [0 : 1 : 0]) \\ \sigma_2(a, x_1) = ((a, x_1), [x_1 : 0 : 1]) \end{cases}$$

does not give a map to $\overline{\mathcal{M}}_{1,2}$ since the image of the second section intersects the singular locus. We define

$$\Lambda := \{((a, x_1), [x : y : z]) \mid x = x_1, y = 0, 4a^3 + 27(-ax_1 - x_1^3)^2 = 0\} \subset B_2.$$

Let $p : \tilde{B}_2 \rightarrow B_2$ be the blow-up of B_2 in Λ . Now the projection of \tilde{B}_2 onto \mathbb{A}_0^2 admits two distinct sections $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$ that do not intersect the

singular locus, and such that $p \circ \tilde{\sigma}_i = \sigma_i$. In this way, we obtain the cartesian diagram:

$$\begin{array}{ccccc}
 \tilde{B}_2 & \longrightarrow & \tilde{\mathcal{C}}_{1,2} & \xrightarrow{\sim} & \overline{\mathcal{M}}_{1,3} \\
 \downarrow & \nearrow \tilde{\sigma}_1 & \downarrow & \nearrow x_1 & \\
 \mathbb{A}_0^2 & \xrightarrow{\psi} & \mathcal{M}_{1,2} & & \\
 & & \downarrow \tilde{\sigma}_2 & & \\
 & & & & \mathbb{A}_0^2 \xrightarrow{x_2}
 \end{array}$$

We denote by $\tilde{\psi}$ the smooth map $\tilde{B}_2 \rightarrow \overline{\mathcal{M}}_{1,3}$. Let F be the locus in B_2 cut out by the equation $y = 0$ and \tilde{F} its strict transform under $p : \tilde{B}_2 \rightarrow B_2$. The map $\tilde{\psi}$ restricted to \tilde{F} surjects onto \overline{A}_3 . There is an isomorphism λ from \mathbb{A}_0^2 (parameters (x_2, x_1)) to F

$$\lambda : (x_2, x_1) \rightarrow ((-x_1^2 - x_1x_2 - x_2^2, x_1), [x_2 : 0 : 1]).$$

Since F is smooth, the restriction of the map $p : \tilde{F} \rightarrow F$ is an isomorphism, and therefore λ lifts to an isomorphism $\tilde{\lambda} : \mathbb{A}_0^2 \rightarrow \tilde{F}$. So we have a surjection:

$$\tilde{\psi} \circ \tilde{\lambda} : \mathbb{A}_0^2 \rightarrow \overline{A}_3.$$

Again, as a consequence of Weierstrass theorem, this map factors via the quotient $[\mathbb{A}^2/\mathbb{G}_m]$, where the action has weights 2 and 2, thus inducing an isomorphism of stacks $[\mathbb{A}_0^2/\mathbb{G}_m] \rightarrow \overline{A}_3$. The forgetful map $\overline{A}_3 \rightarrow \overline{A}_2$ lifts to the map of the charts $\mathbb{A}_0^2 \rightarrow \mathbb{A}_0^2$

$$(3.2) \quad (x_1, x_2) \rightarrow (-x_1^2 - x_1x_2 - x_2^2, x_1).$$

To conclude the proof, we observe that the restriction of the forgetful map, $\overline{A}_4 \rightarrow \overline{A}_3$, is an equivalence of categories. Indeed, when three among the four 2-torsion points of an elliptic curve have been chosen, the fourth is uniquely determined. □

Note that as a consequence of the proof, we deduce from (3.1) and (3.2) a description of the forgetful maps $\overline{A}_3 \rightarrow \overline{A}_2 \rightarrow \overline{A}_1$ in terms of maps of weighted projective stacks.

In Figure 3.1 we show the two points in $\overline{A}_2 \setminus A_2$, the three points in $\overline{A}_3 \setminus A_3$ are in Figure 3.2. The irreducible components of these curves are all rational: the geometric genus of each component is written at one extreme of the curve itself. The first marked point is not pictured and is at infinity. If coordinates are chosen on the vertical genus 0 curve, in such a way that the two intersection points with the other component are $0, \infty$, the marked points (2 and) 2, 3 are (chosen among) the points with coordinates $1, -1$.




Figure 3.1. The two points that compactify A_2




Figure 3.2. The three points that compactify A_3

We have thus described the spaces A_i as open dense substacks of weighted projective stacks of dimension 1. In particular, their rational cohomology follows from the following corollary.

COROLLARY 3.18. — *The coarse moduli spaces of \overline{A}_i is isomorphic to \mathbb{P}^1 . If $i < 4$, the coarse moduli spaces of A_i is \mathbb{P}^1 minus i points. The moduli stack A_4 coincides with A_3 .*

3.2.2. The case of $\overline{\mathcal{M}}_{1,n}$

Up to now, we have determined the decomposition in twisted sectors of the inertia stack (3.5) of $\mathcal{M}_{1,n}$ (Corollary 3.14). As we have observed in Remark 3.2, each twisted sector of $\overline{\mathcal{M}}_{1,n}$ admits a closed embedding inside $\overline{\mathcal{M}}_{1,n}$ itself. So, if (Z, g) is a twisted sector of $\mathcal{M}_{1,n}$, we can consider the Deligne–Mumford compactification $\overline{Z} \subset \overline{\mathcal{M}}_{1,n}$. It is easily seen that (\overline{Z}, g) is a twisted sector of $\overline{\mathcal{M}}_{1,n}$. In the last section, we have studied all such twisted sectors (\overline{Z}, g) . We will see in this section that there are further twisted sectors of $\mathcal{M}_{1,n}$, whose image inside $\overline{\mathcal{M}}_{1,n}$ is completely contained in the boundary.

Let (Z, g) be a twisted sector of $\mathcal{M}_{1,k}$. The twisted sector (\overline{Z}, g) has k marked points that we rename \bullet_i , where $1 \leq i \leq k$. Let now (I_1, \dots, I_k) be a partition of $[n]$, such that $I_i \neq \emptyset$. Let now j_k be the morphism gluing together the same symbols (defined in Section 2.2):

DEFINITION 3.19. — Let Z be a twisted sector inside $\mathcal{M}_{1,k}$. We say \overline{Z} is a base twisted sector. We define $\overline{Z}^{(I_1, \dots, I_k)}$ as:

$$\overline{Z}^{(I_1, \dots, I_k)} := j_k(\overline{Z} \times \overline{\mathcal{M}}_{0, I_1 \sqcup \bullet_1} \times \dots \times \overline{\mathcal{M}}_{0, I_k \sqcup \bullet_k}).$$

We then say \overline{Z} is the base twisted sector associated with $\overline{Z}^{(I_1, \dots, I_k)}$.

THEOREM 3.20. — If (Z, α) is a twisted sector in $\mathcal{M}_{1,k}$, and (I_1, \dots, I_k) is a partition of $[n]$, $(\overline{Z}^{(I_1, \dots, I_k)}, \alpha)$ is a twisted sector of the inertia stack of $\overline{\mathcal{M}}_{1,n}$.

Proof. — The automorphism α lifts to an automorphism α' of $\overline{Z}^{(I_1, \dots, I_k)}$ that acts as α on the base and as the identity on the components $\overline{\mathcal{M}}_{0,n}$. We can call with the same name α and α' , and represent them by the same element in μ_N (see Notation 3.10). It is easy to check that $\overline{Z}^{(I_1, \dots, I_k)}$ is a connected component of the inertia stack of $\overline{\mathcal{M}}_{1,n}$. \square

Notation 3.21. — Let $\sigma \in S_k$. Then $\overline{Z}^{(I_1, \dots, I_k)} = \overline{Z}^{(I_{\sigma(1)}, \dots, I_{\sigma(k)})}$. The twisted sector is identified up to isomorphism by Z and the partition $\{I_1, \dots, I_k\}$ where the ordering of the I_i s does not matter. From now on we will simply denote this twisted sector in $\overline{\mathcal{M}}_{1,n}$ by $\overline{Z}^{\{I_1, \dots, I_k\}}$ or $\overline{Z}^{I_1, \dots, I_k}$: the set of parameters for the twisted sectors whose base space is Z is the set of the k partitions of $[n]$. Note also that \overline{Z} is identified with $\overline{Z}^{\{1\}, \dots, \{k\}}$ for every twisted sector Z of $\overline{\mathcal{M}}_{1,k}$.

With the notation just introduced, we state the main result of this section.

THEOREM 3.22. — The decomposition of $I(\overline{\mathcal{M}}_{1,n})$ in twisted sectors is (see Notation 3.5, 3.6, 3.10):

$$\begin{aligned} & (\overline{\mathcal{M}}_{1,n}, 1) \sqcup (\overline{A}_1^{[n]}, -1) \sqcup (\overline{A}_2^{I_1, I_2}, -1) \sqcup (\overline{A}_3^{I_1, I_2, I_3}, -1) \\ & \sqcup (\overline{A}_4^{I_1, I_2, I_3, I_4}, -1) \sqcup (C_4^{[n]}, i/ - i) \sqcup (C_4^{I_1, I_2}, i/ - i) \\ & \sqcup (C_6^{I_1, I_2}, \epsilon^2/\epsilon^4) \sqcup (C_6^{I_1, I_2, I_3}, \epsilon^2/\epsilon^4) \sqcup (C_6^{[n]}, \epsilon/\epsilon^2/\epsilon^4/\epsilon^5), \end{aligned}$$

where each disjoint union is taken over the set of all possible decompositions of the set $[n]$ in 1, 2, 3 or 4 non-empty subsets: $[n] = \coprod I_i$.

Proof. — We have just seen in Theorem 3.20 that all the components that appear in the decomposition are twisted sectors of $\overline{\mathcal{M}}_{1,n}$. We have to prove that there are no further connected components in the inertia stack

of $\overline{\mathcal{M}}_{1,n}$. To see that there are no further twisted sectors, one can work by induction using the fact that

$$\pi_{[n]} : \overline{\mathcal{M}}_{1,n+1} \rightarrow \overline{\mathcal{M}}_{1,n}$$

is the universal curve. □

From this, we obtain the following corollary that describes all the possible twisted sectors of $\overline{\mathcal{M}}_{1,n}$ stack-theoretically.

COROLLARY 3.23. — *Each twisted sector of $\overline{\mathcal{M}}_{1,n}$ is isomorphic to a product*

$$A \times \overline{\mathcal{M}}_{0,n_1} \times \overline{\mathcal{M}}_{0,n_2} \times \overline{\mathcal{M}}_{0,n_3} \times \overline{\mathcal{M}}_{0,n_4},$$

where $n_1, \dots, n_4 \geq 3$ are integers and A is in the set

$$\{B\mu_3, B\mu_4, B\mu_6, \mathbb{P}(4, 6), \mathbb{P}(2, 4), \mathbb{P}(2, 2)\}.$$

Proof. — It is a consequence of Theorem 3.22, Lemma 3.17, and the fact that $C_4 \cong B\mu_4$, $C_6 \cong B\mu_6$, $C'_6 \cong C''_6 \cong B\mu_3$ (as a consequence of Theorem 3.8). □

3.3. The cohomology of the inertia stack of $\overline{\mathcal{M}}_{1,n}$

We can use the results established in Theorem 3.22 and Corollary 3.23 to compute the dimension of the vector space $H^*_{CR}(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$ (see Definition 3.7). We write the formula for the dimension as a function of the dimension of $H^*(\overline{\mathcal{M}}_{0,n})$, which is well known after Keel [17]. Then, let

$$h(n) := \dim H^*(\overline{\mathcal{M}}_{0,n+1}, \mathbb{Q}) = \sum_k a^k(n)$$

(the latter notation is the one of [17, p. 550] shifted by 1).

COROLLARY 3.24. — *The dimension of the Chen–Ruan cohomology vector space of $\overline{\mathcal{M}}_{1,n}$ is:*

$$\begin{aligned} \dim(H^*_{CR}(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})) &= \dim(H^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})) + 8h(n) + 3 \sum \binom{n}{i, j} h(i)h(j) \\ &+ \frac{2}{3} \sum \binom{n}{i, j, k} h(i)h(j)h(k) + \frac{1}{12} \sum \binom{n}{i, j, k, l} h(i)h(j)h(k)h(l), \end{aligned}$$

where the sum is over indices $1 \leq i, j, k, l \leq n$ such that their sum is n .

Proof. — This result is obtained from Theorem 3.22 and Corollary 3.23, using the fact that the dimension of the cohomology of a point is 1 and the dimension of the cohomology of the projective line is 2. □

We introduce the generating polynomials:

$$(3.3) \quad P_0(s) := \sum_{n=0}^{\infty} \frac{Q_0(n)}{n!} s^n$$

$$(3.4) \quad P_1(s) := \sum_{n=0}^{\infty} \frac{Q_1(n)}{n!} s^n$$

$$(3.5) \quad P_1^{CR}(s) := \sum_{n=0}^{\infty} \frac{Q_1^{CR}(n)}{n!} s^n$$

where:

$$\begin{aligned} Q_0(n) &:= \dim H^*(\overline{\mathcal{M}}_{0,n+1}) = h(n) \\ Q_1(n) &:= \dim H^*(\overline{\mathcal{M}}_{1,n}) \\ Q_1^{CR}(n) &:= \dim H_{CR}^*(\overline{\mathcal{M}}_{1,n}) \end{aligned}$$

with the convention that when the right hand side is not defined, the left hand side equals 1. Formula 3.24 can now be written compactly.

THEOREM 3.25. — *The following equality between power series relates the dimensions of the cohomology group of $\overline{\mathcal{M}}_{0,n}$ and $\overline{\mathcal{M}}_{1,n}$ with the dimension of the Chen–Ruan cohomology group of $\overline{\mathcal{M}}_{1,n}$.*

$$(3.6) \quad P_1^{CR}(s) = P_1(s) + 8P_0(s) + 3P_0(s)^2 + \frac{2}{3}P_0(s)^3 + \frac{1}{12}P_0(s)^4.$$

3.4. The classes of the twisted sectors in $\overline{\mathcal{M}}_{1,n}$

In this section, we want to express the classes $[Y]$ for all Y a twisted sector of $\overline{\mathcal{M}}_{1,n}$, as linear combinations of elements in $R^*(\overline{\mathcal{M}}_{1,n})$. It is possible to express them as linear combinations of products of divisor classes in $\overline{\mathcal{M}}_{1,n}$. This is due to the fact that there are base twisted sectors (Definition 3.19) in genus 1 only up to 4 marked points, and Belorousski’s Theorem 2.8. In fact, we manage to compute these classes as linear combinations of products of S_n -invariant boundary divisor classes.

Notation 3.26. — If Y is a base twisted sector (3.19), we can write

$$[Y] \in A^*(\overline{\mathcal{M}}_{1,n}) = R^*(\overline{\mathcal{M}}_{1,n}) = H^{ev}(\overline{\mathcal{M}}_{1,n}) :$$

the two equalities hold when $n \leq 10$, see 2.7. If $i : Y \rightarrow \overline{\mathcal{M}}_{1,n}$ is the restriction of the map from the inertia stack, $[Y]$ is the push-forward via i of the fundamental class of the twisted sector Y (see 3.2).

We use the notation for the divisors introduced in Notation 2.4.

THEOREM 3.27. — *Let Y be a base twisted sector of $\overline{\mathcal{M}}_{1,n}$ (Definition 3.19). We express its class in the cohomology ring as a linear combination of products of S_n -invariant divisor classes.*

- Base space classes coming from $\overline{\mathcal{M}}_{1,1}$:
 - (1) $[\overline{A}_1] = 1$, the fundamental class of $\overline{\mathcal{M}}_{1,1}$;
 - (2) $[C_4] = \frac{1}{2}d_{irr}$
 - (3) $[C_6] = \frac{1}{3}d_{irr}$.
- Base space classes coming from $\overline{\mathcal{M}}_{1,2}$:
 - (1) $[\overline{A}_2] = \frac{1}{4}d_{irr} + 3d_{\{1,2\}}$;
 - (2) $[C'_4] = \frac{1}{2}d_{irr}d_{\{1,2\}}$;
 - (3) $[C'_6] = \frac{2}{3}d_{irr}d_{\{1,2\}}$.
- Base space classes coming from $\overline{\mathcal{M}}_{1,3}$:
 - (1) $[\overline{A}_3] = \frac{1}{4}d_{irr} \left(\sum_{\{i,j\} \subset \{1,2,3\}} d_{\{i,j\}} \right) + \frac{1}{4}d_{irr}d_{\{1,2,3\}} + 2d_{\{1,2,3\}} \left(\sum_{\{i,j\} \subset \{1,2,3\}} d_{\{i,j\}} \right)$;
 - (2) $[C''_6] = \frac{2}{9}d_{irr}d_{\{1,2,3\}} \left(\sum_{\{i,j\} \subset \{1,2,3\}} d_{\{i,j\}} \right)$.
- Base space classes coming from $\overline{\mathcal{M}}_{1,4}$:
 - (1) $[\overline{A}_4] = \frac{1}{2}d_{\{1,2,3,4\}} \sum_{\{i,j,k\} \subset \{1,2,3,4\}} d_{\{i,j,k\}} \left(\sum_{\{l,m\} \subset \{i,j,k\}} d_{\{l,m\}} \right) + \frac{1}{12}d_{irr} \sum_{\{i,j,k\} \subset \{1,2,3,4\}} d_{\{i,j,k\}} \left(\sum_{\{l,m\} \subset \{i,j,k\}} d_{\{i,j\}} \right) + \frac{1}{12}d_{irr}d_{\{1,2,3,4\}} \left(\sum_{\{i,j\} \subset \{1,2,3,4\}} d_{\{i,j\}} \right)$.

Proof. — We use the methods first established by Mumford [19, Section III]. For the classes of the points the result is trivial. We show how to obtain the result for the classes of the spaces \overline{A}_i . We refer to [4] for all the bases of the Chow groups of $\overline{\mathcal{M}}_{1,n}$ that we use in the following.

First of all, a basis of $A^1(\overline{\mathcal{M}}_{1,2})$ is given by d_{irr} and $d_{\{1,2\}}$. Therefore:

$$(3.7) \quad [\overline{A}_2] = ad_{irr} + bd_{\{1,2\}}.$$

Taking the push-forward via $\pi_1 : \overline{\mathcal{M}}_{1,2} \rightarrow \overline{\mathcal{M}}_{1,1}$, and using that the forgetful morphism restricted to \overline{A}_2 is of degree three (Lemma 3.17), gives that $b = 3$. Now we observe that \overline{A}_2 does not intersect $d_{\{1,2\}}$; by using that

$$d_{\{1,2\}}d_{\{1,2\}} = -\frac{1}{24}, \quad d_{\{1,2\}}d_{irr} = 1/2,$$

we obtain $a = \frac{1}{4}$.

A basis of the S_3 -invariants of $A^2(\overline{\mathcal{M}}_{1,3})$ is given by:

$$d_{irr} (d_{\{1,2\}} + d_{\{1,3\}} + d_{\{2,3\}}), d_{irr}d_{\{1,2,3\}}, d_{\{1,2,3\}} \left(\sum_{\{i,j\} \subset \{1,2,3\}} d_{\{i,j\}} \right),$$

therefore $[\overline{A}_3]$ can be uniquely written as:

$$ad_{irr} (d_{\{1,2\}} + d_{\{1,3\}} + d_{\{2,3\}}) + bd_{irr}d_{\{1,2,3\}} + cd_{\{1,2,3\}} \sum_{\{i,j\} \subset \{1,2,3\}} d_{\{i,j\}}.$$

Taking the push-forwards via the map forgetting one marked point, and using that these forgetful morphisms restricted to \overline{A}_3 are of degree two (Lemma 3.17), gives:

$$2a = \frac{1}{2}, \quad c = 2.$$

Now to determine b , intersect the class of \overline{A}_3 with $d_{\{1,2\}}$ to find $b = a$.

The class of \overline{A}_4 is computed similarly. The dimension of the S_4 -invariants of $A^3(\overline{\mathcal{M}}_{1,4})$ is four; the four coefficients of the class of \overline{A}_4 are obtained by intersecting with the four S_4 -invariant divisors of $\overline{\mathcal{M}}_{1,4}$. Another way to obtain three relations among the four coefficients is by considering the push-forward via the map that forgets the last marked point, this provides a nontrivial check of the result. □

COROLLARY 3.28. — *Let (Y, g) be a twisted sector of $\overline{\mathcal{M}}_{1,n}$. Then $[Y]$ is in the subalgebra generated by the divisors of $\overline{\mathcal{M}}_{1,n}$.*

Proof. — As a consequence of Theorem 3.22, every twisted sector class is $j_{k*}p^*([Z])$, where Z is a base twisted sector in $\overline{\mathcal{M}}_{1,k}$, $k \leq 4$, and the maps fit into the diagram:

$$\begin{array}{ccc} \overline{\mathcal{M}}_{1,k} \times \overline{\mathcal{M}}_{0,I_1+1} \times \dots \times \overline{\mathcal{M}}_{0,I_k+1} & \xrightarrow{j_k} & \overline{\mathcal{M}}_{1,n} \\ \downarrow p & & \\ \overline{\mathcal{M}}_{1,k} & & \end{array}$$

where j_k is the gluing map defined in Section 2.2, and p is the projection onto the first factor.

Suppose without loss of generality that $1 \in I_1, \dots, k \in I_k$, and $\pi_{\{1, \dots, k\}}$ be the forgetful map. Then the class of Y is just the transversal intersection of

$$\pi_{\{1, \dots, k\}}^*([Z]) \quad \text{and} \quad j_{k*}([\overline{\mathcal{M}}_{1,k} \times \overline{\mathcal{M}}_{0,I_1+1} \times \dots \times \overline{\mathcal{M}}_{0,I_k+1}]).$$

The second class is clearly in the subalgebra generated by the divisors, and so is the first since the class of $[Z]$ is in the subalgebra generated by the divisors by Theorem 3.27. □

4. The age grading

4.1. Definition of Chen–Ruan degree

Let X be a smooth algebraic stack of dimension n , and $x \in X$ a point. Let $T(X)$ be the tangent bundle of X . For any g in the stabilizer group of x of order k , there is a basis of $T_x(X)$ consisting of eigenvectors for the g -action. In terms of such a basis, the g -action is given by a diagonal matrix $M = \text{diag}(\xi^{a_1}, \dots, \xi^{a_n})$ where $\xi = e^{\frac{2\pi i}{k}}$ and $a_i < k$. For any pair (x, g) , define a function $a(x, g) := \frac{1}{k} \sum a_i$. This function is nonnegative and takes rational values. Moreover, this function is constant on each connected component of the inertia stack ([6, Chapter 3.2]).

DEFINITION 4.1. — ([6, Chapter 3.2]) *Let X be an algebraic smooth stack. The age of a twisted sector $X_{(g)}$ is defined to be $a(x, g)$ for any point $(x, g) \in X_{(g)}$.*

The age is also referred to as degree shifting number, or fermionic shift. The algebraic definition of age was given in [1, 7.1] and [2, 7.1].

PROPOSITION 4.2. — *Let (Y, g) , (Y, g^{-1}) be two connected components of the inertia stack of an algebraic stack X which are exchanged by the involution of the inertia stack. Then*

$$a((Y, g)) + a((Y, g^{-1})) = \text{codim}_Y X.$$

Remark 4.3. — If $i : Y \rightarrow X$ is a twisted sector, and $x \in Y$ is a point, then the following splitting holds:

$$T_x X = T_x Y \oplus N_Y X|_x.$$

If $G := \langle g \rangle$ is in the stabilizer group of x , then $T_x X$ is a representation of G which splits as a sum of two representations: $T_x Y$ and $N_Y X|_x$. The first of such representations is trivial by the definition of twisted sector. Therefore what is needed in order to compute the age of a twisted sector is to study the action of G on $N_Y X|_x$.

In conclusion of this subsection we define the orbifold degree.

DEFINITION 4.4. — We define the d -th degree Chen–Ruan cohomology group as follows:

$$H_{CR}^d(X, \mathbb{Q}) := \bigoplus_i H^{d-2a(X_i, g_i)}(X_i, \mathbb{Q}),$$

where the sum is over all the connected components of the inertia stack of X . If α is an element of this vector space, its orbifold degree is d .

4.2. Age of $\mathcal{M}_{1,n}$ and $\overline{\mathcal{M}}_{1,n}$

We start our computations in the smooth case. The result for the age of $\mathcal{M}_{1,1}$ is well known, so we work out the other cases.

If ϕ_N is a generator of μ_N (and therefore of μ_N^\vee since we work over \mathbb{C}), we denote by $\langle \phi_N^k \rangle$ the one dimensional complex vector space with the action of μ_N , where ϕ_N acts as the multiplication by ϕ_N^k . Recall that if $G = \text{Aut}(C, P)$ is the automorphism group of an elliptic curve, it is canonically identified with μ_N for a certain N (Notation 3.10).

LEMMA 4.5. — Let $Z \subset \mathcal{M}_{1,k}$ be the closed embedding of a twisted sector (Z, α) of $\mathcal{M}_{1,k}$, with $Z \in \{C_4, C'_4, C_6, C'_6, C''_6\}$. The normal bundle $N_Z \mathcal{M}_{1,k}$ is a representation of μ_N on a k -dimensional vector space:

- as a representation of μ_4 , $N_{C_4} \mathcal{M}_{1,1}$ is $\langle i^2 \rangle$,
- as a representation of μ_4 , $N_{C'_4} \mathcal{M}_{1,2}$ is $\langle i^2 \rangle \oplus \langle i^3 \rangle$,
- as a representation of μ_6 , $N_{C_6} \mathcal{M}_{1,1}$ is $\langle \epsilon^4 \rangle$,
- as a representation of μ_3 generated by ϵ^2 , $N_{C'_6} \mathcal{M}_{1,2}$ is $\langle \epsilon^2 \rangle \oplus \langle \epsilon^4 \rangle$,
- as a representation of μ_3 generated by ϵ^2 , $N_{C''_6} \mathcal{M}_{1,3}$ is $\langle \epsilon^2 \rangle \oplus \langle \epsilon^4 \rangle \oplus \langle \epsilon^4 \rangle$.

Proof. — The age for the twisted sectors of $\mathcal{M}_{1,1}$ is known from the description of it as an open substack of $\mathbb{P}(4, 6)$ (see for instance [18]). It is easily checked that i acts on the normal bundle as the multiplication by $i^6 = i^2$ and ϵ acts as multiplication by ϵ^4 .

We study the tangent bundle to C'_4 in $\mathcal{M}_{1,2}$, the other cases follow through. Since the forgetful morphism $\overline{\mathcal{M}}_{1,2} \rightarrow \overline{\mathcal{M}}_{1,1}$ is the universal curve, we have that the following diagram is cartesian (see 3.9 for the definition of C_4):

$$\begin{array}{ccccccc} & & g & & & & \\ & \curvearrowright & & \curvearrowleft & & & \\ C'_4 & \longrightarrow & C_4 & \longrightarrow & [C_4/\mu_4] & \longrightarrow & \overline{\mathcal{M}}_{1,2} \\ & & & & \downarrow & & \downarrow \\ & & & & C_4 & \longrightarrow & \overline{\mathcal{M}}_{1,1}. \end{array}$$

The normal bundle $N_{C'_4} \overline{\mathcal{M}}_{1,2}$ is therefore isomorphic as a representation of μ_4 to the direct sum: $g^*(N_{[C_4/\mu_4]} \overline{\mathcal{M}}_{1,2}) \oplus N_{C'_4} \mathcal{C}_4$, since $C_4 \rightarrow [C_4/\mu_4]$ is a finite étale map. The first term in the direct sum is equivariantly isomorphic to $N_{C_4} \overline{\mathcal{M}}_{1,1}$ (since the diagram is cartesian and the forgetful map is flat). The normal bundle $N_{C'_4} \mathcal{C}_4$ is equivariantly isomorphic to the tangent space to \mathcal{C}_4 in the second marked point. In the Weierstrass representation of \mathcal{C}_4 (3.9), this is the point with projective coordinates: $[0 : 0 : 1]$. The tangent space in these coordinates is then parametrized by $[0 : t : 1]$ and i acts on it as the multiplication by i^3 (Theorem 3.8). \square

With this result, it is straightforward to compute the age of all the twisted sectors of $\mathcal{M}_{1,n}$, by using the fact that the twisted sector corresponding to an involutive automorphism have age equal to half the codimension (Proposition 4.2).

Remark 4.6. — We observe that, in the proof of the above proposition, we had to use a different argument for the case $n = 1$ and $n > 1$. This is due to the fact that there is no forgetful map from $\mathcal{M}_{1,1}$. In particular, if $(C, p) \in \mathcal{M}_{1,1}$, and $\mu_N = \text{Aut}(C, P)$ then the actions of μ_N on $N_{(C,p)} \mathcal{M}_{1,1}$ and on $T_p \mathcal{C}$ do not necessarily coincide.

In the following Proposition, we give a description of the normal bundle $N_Z \overline{\mathcal{M}}_{1,n}$ where Z is a twisted sector. We restrict to the cases where the base space of Z (Definition 3.19) has dimension 0, since the automorphism element of all others is involutive. Hence, let Z be in $\{C_4, C'_4, C_6, C'_6, C''_6\}$. The normal bundle $N_Z \mathcal{M}_{1,k}$ is a k -dimensional vector space with an action of μ_N on it which we studied in Lemma 4.5 (N can be 3, 4 or 6). For the sake of simplicity, we identify Z^{I_1, \dots, I_k} with $Z \times \overline{\mathcal{M}}_{0, I_1+1} \times \dots \times \overline{\mathcal{M}}_{0, I_k+1}$ (see Definition 3.19). We denote by p, p_1, \dots, p_k the projections onto each factor. We indicate by $\underline{\mathbb{C}}$ the trivial bundle of rank 1.

PROPOSITION 4.7. — *Let I_1, \dots, I_k be a partition of $[n]$ in $k \leq 3$ subsets. Suppose that $|I_i| > 1$ for all i s. Then the normal bundle $N_{Z^{I_1, \dots, I_k}} \overline{\mathcal{M}}_{1,n}$ is of rank $2k$ and splits as a direct sum of line bundles.* ⁽²⁾

- (1) $N_{C'_4} \overline{\mathcal{M}}_{1,n}$ is isomorphic to: $(i^2, \underline{\mathbb{C}}) \oplus (i^3, p_1^*(\mathbb{L}_{n+1}^\vee))$,
- (2) $N_{C'_4} \overline{\mathcal{M}}_{1,n}$ is isomorphic to: $(i^2, \underline{\mathbb{C}}) \oplus (i^3, \underline{\mathbb{C}}) \oplus (i^3, p_1^*(\mathbb{L}_{I_1+1}^\vee)) \oplus (i^3, p_2^*(\mathbb{L}_{I_2+1}^\vee))$,
- (3) $N_{C_6} \overline{\mathcal{M}}_{1,n}$ is isomorphic to: $(\epsilon^4, \underline{\mathbb{C}}) \oplus (\epsilon^5, p_1^*(\mathbb{L}_{n+1}^\vee))$,

(2) see last paragraph of Section 2.1 for a line bundle on a trivial gerbe

- (4) $N_{C_6^{I_1, I_2} \overline{\mathcal{M}}_{1,n}}$ is isomorphic to: $(\epsilon^2, \underline{\mathbb{C}}) \oplus (\epsilon^4, \underline{\mathbb{C}}) \oplus (\epsilon^4, p_1^*(\mathbb{L}_{I_1+1}^\vee)) \oplus (\epsilon^4, p_2^*(\mathbb{L}_{I_2+1}^\vee))$,
- (5) $N_{C_6^{I_1, I_2, I_3} \overline{\mathcal{M}}_{1,n}}$ is isomorphic to: $(\epsilon^2, \underline{\mathbb{C}}) \oplus (\epsilon^4, \underline{\mathbb{C}}) \oplus (\epsilon^4, \underline{\mathbb{C}}) \oplus (\epsilon^4, p_1^*(\mathbb{L}_{I_1+1}^\vee)) \oplus (\epsilon^4, p_2^*(\mathbb{L}_{I_2+1}^\vee)) \oplus (\epsilon^4, p_3^*(\mathbb{L}_{I_3+1}^\vee))$.

If some of the I_i s has cardinality 1, the normal bundle has the same description after removing the corresponding components $p_i^*(\mathbb{L}_{I_i+1}^\vee)$.

We postpone the proof of this proposition, in order to immediately see that as a consequence of it we can compute the age of all the twisted sectors of $\overline{\mathcal{M}}_{1,n}$. We use the convention that $\delta(I) = \delta_{1,|I|}$, the Kronecker delta.

COROLLARY 4.8. — In this table we recollect the age, or degree shifting number, of the twisted sectors of $\overline{\mathcal{M}}_{1,n}$.

Component	Aut	Codimension	Age
$\overline{A}_1^{[n]}$	-1	$1 - \delta(n)$	$\frac{1}{2}(1 - \delta(n))$
$\overline{A}_2^{I_1, I_2}$	-1	$3 - \delta(I_1) - \delta(I_2)$	$\frac{1}{2}(3 - \delta(I_1) - \delta(I_2))$
$\overline{A}_3^{I_1, I_2, I_3}$	-1	$5 - \delta(I_1) - \delta(I_2) - \delta(I_3)$	$\frac{1}{2}(5 - \delta(I_1) - \delta(I_2) - \delta(I_3))$
$\overline{A}_4^{I_1, I_2, I_3, I_4}$	-1	$7 - \delta(I_1) - \delta(I_2) - \delta(I_3) - \delta(I_4)$	$\frac{1}{2}(7 - \delta(I_1) - \delta(I_2) - \delta(I_3) - \delta(I_4))$
$C_4^{[n]}$	i	$2 - \delta(n)$	$\frac{5}{4} - \frac{3}{4}\delta(n)$
$C_4^{[n]}$	$-i$	$2 - \delta(n)$	$\frac{3}{4} - \frac{1}{4}\delta(n)$
$C_4^{I_1, I_2}$	i	$4 - \delta(I_1) - \delta(I_2)$	$\frac{11}{4} - \frac{3}{4}(\delta(I_1) + \delta(I_2))$
$C_4^{I_1, I_2}$	$-i$	$4 - \delta(I_1) - \delta(I_2)$	$\frac{5}{4} - \frac{1}{4}(\delta(I_1) + \delta(I_2))$
$C_6^{I_1, I_2}$	ϵ^2	$4 - \delta(I_1) - \delta(I_2)$	$\frac{7}{3} - \frac{2}{3}(\delta(I_1) + \delta(I_2))$
$C_6^{I_1, I_2}$	ϵ^4	$4 - \delta(I_1) - \delta(I_2)$	$\frac{5}{3} - \frac{1}{3}(\delta(I_1) + \delta(I_2))$
$C_6^{I_1, I_2, I_3}$	ϵ^2	$6 - \delta(I_1) - \delta(I_2) - \delta(I_3)$	$\frac{11}{3} - \frac{2}{3}(\delta(I_1) + \delta(I_2) + \delta(I_3))$
$C_6^{I_1, I_2, I_3}$	ϵ^4	$6 - \delta(I_1) - \delta(I_2) - \delta(I_3)$	$\frac{7}{3} - \frac{1}{3}(\delta(I_1) + \delta(I_2) + \delta(I_3))$
$C_6^{[n]}$	ϵ	$2 - \delta(n)$	$\frac{3}{2} - \frac{5}{6}\delta(n)$
$C_6^{[n]}$	ϵ^2	$2 - \delta(n)$	$1 - \frac{2}{3}\delta(n)$
$C_6^{[n]}$	ϵ^4	$2 - \delta(n)$	$1 - \frac{1}{3}\delta(n)$
$C_6^{[n]}$	ϵ^5	$2 - \delta(n)$	$\frac{1}{2} - \frac{1}{6}\delta(n)$

Proof. — The age of the sectors with an involutive automorphism is half the codimension by Lemma 4.2. They are all the twisted sectors whose associated base twisted sector (3.19) has dimension 1. For the remaining twisted sectors, we simply apply Proposition 4.7.

□

To prove Proposition 4.7, we use the following result due to Mumford:

LEMMA 4.9. — ([19]) *Let I_1, \dots, I_k be a partition of $[n]$, and j_k the gluing map defined in Section 2.2*

$$j_k : \overline{\mathcal{M}}_{1, \coprod_{i=1}^k \bullet_i} \times \overline{\mathcal{M}}_{0, I_1 \sqcup \bullet_1} \times \dots \times \overline{\mathcal{M}}_{0, I_k \sqcup \bullet_k} \rightarrow \overline{\mathcal{M}}_{1, n}.$$

Let p be the projection onto the first factor, and p_1, \dots, p_k the projections onto the moduli spaces of genus 0 curves. Then the normal bundle of the map j_k is isomorphic to:

$$N_{j_k} = \bigoplus_{i=1}^k p^*(\mathbb{L}_{\bullet_i}^\vee) \otimes p_i^*(\mathbb{L}_{\bullet_i}^\vee),$$

where \mathbb{L}_{\bullet_i} are the cotangent line bundles defined in Section 2.2, the first one is on $\overline{\mathcal{M}}_{1, \coprod_{i=1}^k \bullet_i}$, and the second one is on $\overline{\mathcal{M}}_{0, I_i \sqcup \bullet_i}$

Proof. — (of Proposition 4.7) If j_k denotes as usual the gluing morphism defined in Section 2.2, the following diagram is cartesian by definition of Z^{I_1, \dots, I_k} (p is the projection onto the first factor)

$$\begin{array}{ccc} Z^{I_1, \dots, I_k} \hookrightarrow & \xrightarrow{\lambda} & \overline{\mathcal{M}}_{1, k} \times \overline{\mathcal{M}}_{0, I_1 \sqcup \bullet_1} \times \dots \times \overline{\mathcal{M}}_{0, I_k \sqcup \bullet_k} \hookrightarrow \\ \downarrow p & \square & \downarrow p \\ Z \hookrightarrow & \longrightarrow & \overline{\mathcal{M}}_{1, k}. \end{array}$$

In this case the following isomorphism holds

$$p^*(N_Z \overline{\mathcal{M}}_{1, k}) \cong N_{Z^{I_1, \dots, I_k}} \overline{\mathcal{M}}_{1, k} \times \overline{\mathcal{M}}_{0, I_1 \sqcup \bullet_1} \times \dots \times \overline{\mathcal{M}}_{0, I_k \sqcup \bullet_k} = N_\lambda$$

since the diagram is cartesian and p is flat. This is the trivial bundle with a certain representation of μ_N , that we studied in Lemma 4.5.

The map $j_k \circ \lambda$ is the restriction to $\overline{Z}^{I_1, \dots, I_k}$ of the map $I(\overline{\mathcal{M}}_{1, n}) \rightarrow \overline{\mathcal{M}}_{1, n}$. Therefore, the normal bundle $N_{Z^{I_1, \dots, I_k}} \overline{\mathcal{M}}_{1, n}$ is isomorphic to the direct sum: $\lambda^* N_{j_k} \oplus N_\lambda$. To conclude the proof, we have now to study the first term $\lambda^* N_{j_k}$.

According to Lemma 4.9, we have:

$$\lambda^* N_{j_k} = \left(\bigoplus_{i=1}^k p^*(\mathbb{L}_{\bullet_i}^\vee) \otimes p_i^*(\mathbb{L}_{\bullet_i}^\vee) \right) \Big|_{Z^{I_1, \dots, I_k}}.$$

The term $p^*(\mathbb{L}_{\bullet_i}^\vee)|_{Z^{I_1, \dots, I_k}}$ is the trivial bundle whose constant fiber is the tangent space to Z in the i -th marked point. It carries the representation of μ_N , as the latter group acts on the tangent space to Z in the i -th marked point (this action can be computed explicitly, see Lemma 4.5). On the

other hand, the term $p_i^*(\mathbb{L}_{\bullet_i}^\vee)|_{Z^{I_1, \dots, I_k}}$ carries the trivial representation of μ_N , but it is non trivial as a line bundle. This last observation concludes the proof. \square

We can now write a formula analogous to (3.6), adding a new variable to separate the different degrees in the Chen–Ruan cohomology of $\overline{\mathcal{M}}_{1,n}$. We define:

$$(4.1) \quad P_0(s, t) := \sum_{n=0}^{\infty} \frac{Q_0(n, m)}{n!} s^n t^m,$$

$$(4.2) \quad P_1(s, t) := \sum_{n=0}^{\infty} \frac{Q_1(n, m)}{n!} s^n t^m,$$

$$(4.3) \quad P_{1,\alpha}^{CR}(s, t) := \sum_{n=0}^{\infty} \frac{Q_{1,\alpha}^{CR}(n, m)}{n!} s^n t^m,$$

where

$$\begin{aligned} Q_0(n, m) &:= \dim H^{2m}(\overline{\mathcal{M}}_{0,n+1}) = a^m(n), \\ Q_1(n, m) &:= \dim H^{2m}(\overline{\mathcal{M}}_{1,n}), \\ Q_{1,\alpha}^{CR}(n) &:= \dim H_{CR}^{2m+\alpha}(\overline{\mathcal{M}}_{1,n}). \end{aligned}$$

The relevant values of α are $\{0, \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}\} =: A$. Summing over α and shifting by α the degree of t , we have the generating series of the orbifold Poincaré polynomials:⁽³⁾

$$P_1^{CR}(s, t) := \sum_{\alpha \in A} t^\alpha P_{1,\alpha}^C R(s, t).$$

In other words, the coefficient of degree n in the variable s of P_1^{CR} is the orbifold Poincaré polynomial of $\overline{\mathcal{M}}_{1,n}$ divided by $n!$. The power series (4.1) are described in [11, Theorem 5.9], [13, Theorem 2.6]. There, the author describes the cohomology of the moduli of genus 0 and genus 1, n -pointed stable curves as an S_n -representation.

THEOREM 4.10. — *The following equalities between power series relate the dimension of the m -th Chen–Ruan cohomology group of $\overline{\mathcal{M}}_{1,n}$ ($n > 4$) with the Betti numbers of the moduli of pointed stable curves of genus 0*

⁽³⁾Note that the orbifold Poincaré polynomials are, strictly speaking, not polynomials.

and 1.

$$\begin{aligned}
 P_{1,0}^{CR}(s, t) &= P_1 + (t + t^2)P_0 + 3s(t^2 + t^3)P_0^2 + \frac{s}{24}(t^3 + t^4)P_0^3 \\
 &\quad + 2s(t + t^2)\frac{\partial}{\partial s}(sP_0) + \frac{s^3}{6}(t^2 + t^3)\frac{\partial^3}{\partial s^3}(s^3P_0) \\
 P_{1,\frac{1}{4}}^{CR}(s, t) &= tP_0 + \frac{t^2}{2}P_0^2 \\
 P_{1,\frac{1}{3}}^{CR}(s, t) &= \frac{t^2}{2}P_0^2 + \frac{t^2}{6}P_0^3 + ts\frac{\partial}{\partial s}(sP_0) + \frac{s^2t^2}{2}\frac{\partial^2}{\partial s^2}(s^2P_0) \\
 P_{1,\frac{1}{2}}^{CR}(s, t) &= 2(1 + t)P_0 + \frac{t + t^2}{2}P_0^2 + \frac{t^2 + t^3}{6}P_0^3 + \frac{t^3 + t^4}{24}P_0^4 \\
 &\quad + \frac{s^2}{2}(t + t^2)\frac{\partial^2}{\partial s^2}(s^2P_0) \\
 P_{1,\frac{2}{3}}^{CR}(s, t) &= \frac{t}{2}P_0^2 + \frac{t^3}{6}P_0^3 + st\frac{\partial}{\partial s}(sP_0) + \frac{s^2}{2}t\frac{\partial^2}{\partial s^2}(s^2P_0) \\
 P_{1,\frac{3}{4}}^{CR}(s, t) &= P_0 + \frac{t}{2}P_0^2
 \end{aligned}$$

5. The second inertia stack

The definition of the Chen–Ruan product involves the second inertia stack.

DEFINITION 5.1. — *Let X be an algebraic stack. The second inertia stack $I_2(X)$ is defined as:*


$$I_2(X) = I(X) \times_X I(X).$$

Remark 5.2. — Like the inertia stack, the second inertia stack is smooth (cfr. 3.4, see also [1, p. 15], after noticing that $\mathcal{K}_{0,3}(X, 0) \cong I_2(X)$).

A point in $I_2(X)$ is a triplet (x, g, h) where x is a point of X and $g, h \in \text{Aut}(x)$. It can equivalently be given as $(x, g, h, (gh)^{-1})$.

Remark 5.3. — The second inertia stack comes with three natural morphisms to $I(X)$: p_1 and p_2 , the two projections of the fiber product, and p_3 which acts on points sending (x, g, h) to (x, gh) . This gives the following diagram, where $(Y, g, h, (gh)^{-1})$ is a double twisted sector and (X_1, g) ,

(X_2, h) , $(X_3, (gh))$ are twisted sectors:



Let us now study the double twisted sectors in the case when $X = \overline{\mathcal{M}}_{1,n}$. From now on we focus on the compact case, since the case of $\mathcal{M}_{1,n}$ follows through analogously and much more simply. The following follows easily from the fact that all isotropy groups of $\overline{\mathcal{M}}_{1,n}$ are cyclic (cf. also Remark 3.2).

PROPOSITION 5.4. — *Let $(Z, g, h, (gh)^{-1})$ be a double twisted sector of $\overline{\mathcal{M}}_{1,n}$. Then either (Z, g) or (Z, h) or $(Z, (gh)^{-1})$ is a twisted sector of the inertia stack of $\overline{\mathcal{M}}_{1,n}$.*

Remark 5.5. — We label each sector of $I_2(X)$ via the triplet $(g, h, (gh)^{-1})$. There are two automorphism groups acting on $I_2(X)$: an involution sending a sector labeled with $(g, h, (gh)^{-1})$ into $(g^{-1}, h^{-1}, (gh))$, and S_3 which permutes the three automorphisms. Up to permutations and involution, the following are all the possible labels of the sectors in $I_2(X)$ that correspond to non-empty connected sectors:

- $(1, 1, 1)$, generated group μ_1 ;
- $(1, -1, -1)$, generated group μ_2 ;
- $(\epsilon^2, \epsilon^2, \epsilon^2)$, generated group μ_3 ;
- $(1, \epsilon^2, \epsilon^4)$, generated group μ_3 ;
- $(1, i, -i)$ generated group μ_4 ;
- $(i, i, -1)$, generated group μ_4 ;
- $(\epsilon, \epsilon, \epsilon^4)$, generated group μ_6 ;
- $(\epsilon, \epsilon^2, -1)$, generated group μ_6 ;
- $(1, \epsilon, \epsilon^5)$, generated group μ_6 .

We now describe the sectors of the double inertia stack. We do so up to the automorphisms described in the previous remark, and up to the permutations of the marked points. The next proposition is easily obtained after Theorem 3.22.

PROPOSITION 5.6. — *Up to permutation of the automorphisms, and up to involution, the following are the twisted sectors of $I_2(\mathcal{M}_{1,n})$:*

$$\begin{aligned} & \left(\overline{A_1}^{[n]}, (1, -1, -1)\right), \quad \left(\overline{A_2}^{I_1, I_2}, (1, -1, -1)\right), \quad \left(\overline{A_3}^{I_1, I_2, I_3}, (1, -1, -1)\right) \\ & \quad \left(\overline{A_4}^{I_1, I_2, I_3, I_4}, (1, -1, -1)\right), \quad \left(C_6^{I_1, I_2}, (1, \epsilon^2, \epsilon^4)/(\epsilon^2, \epsilon^2, \epsilon^2)\right) \\ & \quad \left(C_6^{I_1, I_2, I_3}, (1, \epsilon^2, \epsilon^4)/(\epsilon^2, \epsilon^2, \epsilon^2)\right), \quad \left(C_4^{[n]}, (1, i, -i)/(i, i, -1)\right) \\ & \left(C_4^{I_1, I_2}, (1, i, -i)/(i, i, -1)\right), \quad \left(C_6^{[n]}, (1, \epsilon, \epsilon^5)/(\epsilon, \epsilon, \epsilon^4)/(\epsilon, \epsilon^2, -1)/(\epsilon^2, \epsilon^2, \epsilon^2)\right), \end{aligned}$$

where I_1, I_2, I_3, I_4 form a partition of $[n]$ in non-empty subsets.

6. The excess intersection bundle

6.1. Definition of the Chen–Ruan product

We review the definition of the excess intersection bundle over $I_2(X)$, for X an algebraic smooth stack. Let $(Y, g, h, (gh)^{-1})$ be a twisted sector of $I_2(X)$. Let $H := \langle g, h \rangle$ be the group generated by g and h .

CONSTRUCTION 1. — Let $\gamma_0, \gamma_1, \gamma_\infty$ be three small loops around $0, 1, \infty \subset \mathbb{P}^1$. Any map $\pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}) \rightarrow H$ corresponds to an H -principal bundle on $\mathbb{P}^1 \setminus \{0, 1, \infty\}$. Let $\pi^0 : C^0 \rightarrow \mathbb{P}^1 \setminus \{0, 1, \infty\}$ be the H -principal bundle which corresponds to the map $\gamma_0 \rightarrow g, \gamma_1 \rightarrow h, \gamma_\infty \rightarrow (gh)^{-1}$. It can be uniquely extended to a ramified H -Galois covering $C \rightarrow \mathbb{P}^1$ (see [9, Appendix]), where C is a smooth compact curve. Note that H acts on C (and, by definition, \mathbb{P}^1 is the quotient C/H as varieties), and hence on $H^1(C, \mathcal{O}_C)$.

Let $f : Y \rightarrow X$ be the restriction of the canonical map $I_2(X) \rightarrow X$ to the twisted sector Y . Note that H acts on $f^*(T_X)$.

DEFINITION 6.1. — [6] *With the same notation as in the previous paragraph, the excess intersection bundle over Y is defined as:*

$$E_Y = \left(H^1(C, \mathcal{O}_C) \otimes_{\mathbb{C}} f^*(T_X)\right)^H,$$

i.e. the H -invariant subbundle of the expression between parenthesis.

Remark 6.2. — Since $H^1(C, \mathcal{O}_C)^H = 0$, it is the same to consider in the previous definition:

$$\left(H^1(C, \mathcal{O}_C) \otimes N_Y X\right)^H,$$

where N_Y is the coker of $T_Y \rightarrow f^*(T_X)$.

We now review the definition of the Chen–Ruan product.

DEFINITION 6.3. — *Let $\alpha \in H_{CR}^*(X)$, $\beta \in H_{CR}^*(X)$. We define:*

$$\alpha *__{CR} \beta = p_{3*} (p_1^*(\alpha) \cup p_2^*(\beta) \cup c_{top}(E)).$$

THEOREM 6.4. — ([6]) *With the age grading defined in the previous section, $(H_{CR}^*(X, \mathbb{Q}), *__{CR})$ is a graded $(H^*(X, \mathbb{Q}), \cup)$ -algebra.*

Theorem 6.4 allows us to compute the rank of the excess intersection bundle in terms of the already computed age grading. If $(Y, (g, h, (gh)^{-1}))$ is a sector of the second inertia stack, the rank of the excess intersection bundle is (here we stick to the notation introduced in Remark 5.3):

$$(6.1) \quad \text{rk}(E_{(Y,g,h)}) = a(X_1, g) + a(X_2, h) + a(X_3, (gh)^{-1}) - \text{codim}_Y X.$$

COROLLARY 6.5. — *The excess intersection bundle over double twisted sectors when either g, h , or $(gh)^{-1}$ is the identity, is the zero bundle.*

One other useful consequence of Proposition 4.2 and Theorem 6.4 relates the rank of the excess bundle over a double twisted sector and the rank of the excess bundle over the double twisted sector obtained inverting the automorphisms that label the sector

$$(6.2) \quad \text{rk}(E_{(Y,g^{-1},h^{-1})}) = \sum_{i=1}^3 \text{codim}_{X_i} X - 2 \text{codim}_Y X - \text{rk}(E_{(Y,g,h)}).$$

Let $(Y, (g, h, (gh)^{-1}))$ be a double twisted sector in $I_2(\overline{\mathcal{M}}_{1,n})$, and let H be the group generated by $(g, h, (gh)^{-1})$. We will study $N_Y X$ and $H^1(C, \mathcal{O}_C)$ as H -representations.

6.2. The excess intersection bundle for $\overline{\mathcal{M}}_{1,n}$

We have seen in Remark 5.5 what are all the possible couples of automorphisms that correspond to non-empty connected substacks of $I_2(\overline{\mathcal{M}}_{1,n})$. Thanks to Corollary 5.4, the double twisted sectors whose excess intersection bundles have non-zero rank are those labelled by:

$$(6.3) \quad (\epsilon^2, \epsilon^2, \epsilon^2), (i, i, -1), (\epsilon, \epsilon, \epsilon^4), (\epsilon, \epsilon^2, -1)$$

up to permutation and involution. The top Chern classes of the excess intersection bundles for $\mathcal{M}_{1,n}$ are always 0 or 1, since the coarse moduli spaces of the double twisted sectors labeled by these automorphisms are points.

The ranks of the excess intersection bundles for the twisted sectors labeled by 6.3 can be computed thanks to formulas (6.1) and (6.2).

PROPOSITION 6.6. — *In the following table we list the ranks of the excess intersection bundles over all the double twisted sectors (Z, g, h) of $\overline{\mathcal{M}}_{1,n}$, such that g, h and $gh \neq 1$:*

(g, h)	Double twisted sector	$\text{rk}(E)$	(g^{-1}, h^{-1})	$\text{rk}(E)$
(ϵ^2, ϵ^2)	$C_6^{[n]}$	1	(ϵ^4, ϵ^4)	1
(ϵ^2, ϵ^2)	$C_6^{I_1, I_2}$	3	(ϵ^4, ϵ^4)	1
(ϵ^2, ϵ^2)	$C_6^{I_1, I_2, I_3}$	5	(ϵ^4, ϵ^4)	1
(i, i)	$C_4^{[n]}$	1	$(-i, -i)$	0
(i, i)	$C_4^{I_1, I_2}$	3	$(-i, -i)$	0
(ϵ, ϵ)	$C_6^{[n]}$	2	(ϵ^5, ϵ^5)	0
(ϵ, ϵ^2)	$C_6^{[n]}$	1	(ϵ^4, ϵ^5)	0

Remark 6.7. — The above proposition and Corollary 6.5 imply that many top Chern classes of excess intersection bundles are 1 (those of the rank 0 excess intersection bundles). Moreover it is obvious, after Proposition 4.7, that all the rank 3 or 5 bundles of Corollary 6.5 contain at least one trivial subbundle of rank 1, this implying that their top Chern classes is zero.

Now we want to compute explicitly the remaining, non trivial, excess intersection bundles and their top Chern classes for $\overline{\mathcal{M}}_{1,n}$. We study the decomposition of $H^1(C, \mathcal{O}_C)$ as a representation of H in the cases corresponding to non-zero ranks in Proposition 6.6. Here ϕ_N is a generator for μ_N^\vee , the same chosen in Section 4.2. As in that section, we indicate by $\langle \phi_N^k \rangle$ the one dimensional complex vector space endowed with the action of $\phi_N \in \mu_N^\vee$ given by the product by ϕ_N^k .

PROPOSITION 6.8. — *Let H be generated by two elements g, h as in 6.6. Let $C \rightarrow \mathbb{P}^1$ be the H -covering associated with the generators g, h (see Construction 1). We study $H^1(C, \mathcal{O}_C)$ as an H -representation:*

- $H = \mu_3: g = \epsilon^2, h = \epsilon^2$, then $H^1(C, \mathcal{O}_C) = \langle \epsilon^2 \rangle$,
- $H = \mu_3: g = \epsilon^4, h = \epsilon^4$, then $H^1(C, \mathcal{O}_C) = \langle \epsilon^4 \rangle$,
- $H = \mu_4: g = i, h = i$, then $H^1(C, \mathcal{O}_C) = \langle i \rangle$,
- $H = \mu_6: g = \epsilon, h = \epsilon^2$, then $H^1(C, \mathcal{O}_C) = \langle \epsilon \rangle$,
- $H = \mu_6: g = \epsilon, h = \epsilon$, then $H^1(C, \mathcal{O}_C) = \langle \epsilon \rangle \oplus \langle \epsilon^2 \rangle$.

Proof. — By Serre duality, we compute the action of H on $H^0(C, \Omega_C^1)$. In the first four cases, the curve C has genus 1, thus Ω_C^1 is trivial, therefore the action of H coincides with its action on the cotangent space of a

point of total ramification. In the fifth case the curve C has genus 2; by choosing suitable coordinates one can represent respectively C and a basis for $H^0(C, \Omega_C^1)$ as

$$y^2 = x^6 - 1, \quad \left(\frac{dx}{y}, x \frac{dx}{y} \right).$$


Here the automorphism acts by fixing y and mapping $x \rightarrow \epsilon x$. From this description the result for the last case follows. \square

With all this, and thanks to Proposition 4.7, we can compute the excess intersection bundles and their respective top Chern classes. We know already that, among the list of couples of automorphisms of Proposition 6.6, the rank 3 and 5 bundles have top Chern class zero (see 6.7). Among the vector bundles having rank greater than zero, we can prove:

COROLLARY 6.9. — *In Table 6.6, the top Chern classes of all the excess intersection bundles (which are all line bundles) corresponding to the couple (ϵ^4, ϵ^4) are zero. The top Chern class of the excess intersection bundle that corresponds to the couple (ϵ, ϵ) (of rank 1), is also zero.*

Proof. — From Proposition 4.7 and Proposition 6.8, it is straightforward to see that all the excess bundles mentioned in the statement contain a trivial subbundle, forcing their top Chern class to be zero. \square

So we are now left with three rank 1 excess intersection bundles, whose top Chern class is non-zero and not 1. In the following diagram and in the following lemma, we identify the isomorphic spaces in order to simplify the notation for the projection maps.



where λ can be 4 or 6. Here \bullet is the gluing point.

COROLLARY 6.10. — *The only top Chern classes of the excess intersection bundles over double twisted sectors of $\overline{\mathcal{M}}_{1,n}$ that are not 0 nor 1 are:*

- (1) $(C_6^{[n]}, (\epsilon^2, \epsilon^2, \epsilon^2)) \cong B_{\mu_6} \times \overline{\mathcal{M}}_{0,n,\bullet}$, where the top Chern class of the excess intersection bundle is $-p_1^*(\psi_\bullet) = -p_1^*(\psi_{n+1})$;

- (2) $(C_4^{[n]}, (i, i, -1)) \cong B_{\mu_4} \times \overline{\mathcal{M}}_{0,n\sqcup\bullet}$, where the top Chern class of the excess intersection bundle is $-p_1^*(\psi_\bullet) = -p_1^*(\psi_{n+1})$;
- (3) $(C_6^{[n]}, (\epsilon, \epsilon^2, -1)) \cong B_{\mu_6} \times \overline{\mathcal{M}}_{0,n\sqcup\bullet}$, where the top Chern class of the excess intersection bundle is $-p_1^*(\psi_\bullet) = -p_1^*(\psi_{n+1})$.

Proof. — The fact that all other top Chern classes are zero or one follows from all the considerations in this section. In particular, we have observed in the beginning of the section that the excess intersection bundle that may have top Chern class different from 1 are listed in 6.6. In Remark 6.7 and in Corollary 6.9 we have computed the top Chern class of all the remaining cases to be zero or 1.

The result stated then follows as a consequence of Proposition 4.7, Proposition 6.8 (and from the very definition of the ψ classes, see Section 2.2). \square

Note that when $n = 2$ the top Chern classes in Corollary above are 0 too, because the sectors involved are all points.

To conclude, we summarize the result we have obtained in this section:

THEOREM 6.11. — *All top Chern classes of the excess intersection bundles over all double twisted sectors are explicitly given. They can be:*

- (1) either 1, for all the sectors listed in Proposition 5.6 such that one of the three automorphisms of the labeling is 1,
- (2) or again 1, for some of the sectors in the list 6.6, more precisely those mentioned in Remark 6.7,
- (3) or 0, for some of the sectors listed in 6.6, as discussed in Remark 6.7 and in Corollary 6.9;
- (4) or a pullback of a ψ class over a component $\overline{\mathcal{M}}_{0,n}$, for the remaining elements of the list 6.6, as in Corollary 6.10.

7. Pull-backs and push-forwards of strata to the twisted sectors

In order to compute the Chen–Ruan product, one has to compute pull-backs from the twisted sectors to the double twisted sectors and push-forwards from the double twisted sectors to the twisted sectors. Thanks to Corollary 6.5, it is necessary and sufficient to only compute push-forwards and pull-backs between twisted sectors of the inertia stack.

In this section we fix n and call $X := \overline{\mathcal{M}}_{1,n}$. Let (Y, g) be a twisted sector of X , and $f : Y \rightarrow X$ be the closed embedding of the twisted sector.

LEMMA 7.1. — *The cycle map*

$$A^*(Y, \mathbb{Q}) \rightarrow H^{2*}(Y, \mathbb{Q})$$

is a graded ring isomorphism. Moreover the Chow ring of each twisted sector is generated by divisors.

Proof. — All factors of each twisted sector have Chow ring isomorphic to the even cohomology. The cohomology ring of $\overline{\mathcal{M}}_{0,n}$ is generated by divisors due to the work of Keel [17]. Each of the spaces $\overline{A}_1, \overline{A}_2, \overline{A}_3, \overline{A}_4$ has coarse moduli space isomorphic to \mathbb{P}^1 . \square

We can now state and prove the result announced in the introduction. For some of the results needed in the proof we refer to the following two subsections on pull-backs and push-forwards.

THEOREM 7.2. — *The Chen–Ruan cohomology ring of $\overline{\mathcal{M}}_{1,n}$ is generated as an $H^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$ -algebra by the fundamental classes of the twisted sectors with explicit relations.*

Proof. — We will prove in Corollary 7.3 that f^* , the induced pull-back in cohomology, is surjective. This suffices to prove the generation part of the statement. Indeed, let (X_i, g_i) be two twisted sectors, and $f_i : (X_i, g_i) \rightarrow X$ be the canonical projection maps. Let $\alpha_i \in H^*((X_i, g_i), \mathbb{Q})$ for $i = 1, 2$. Let $\tilde{\alpha}_i$ be two liftings of α_i to $H^*(\overline{\mathcal{M}}_{1,n})$ obtained by using the surjectivity of f_i^* . Then we have

$$\alpha_1 *_{CR} \alpha_2 = (\tilde{\alpha}_1 \cup \tilde{\alpha}_2) *_{CR} 1_{(X_1, g)} *_{CR} 1_{(X_2, h)},$$

this proves the generation claim.

Then, as a consequence of Proposition 5.4, $(X_1, g) \times_X (X_2, h)$ is connected, and hence a double twisted sector of $I_2(X)$. Let $(X_3, g_1 g_2)$ be such a twisted sector of $I(X)$, and denote by f_3 the closed embedding of $(X_3, g_1 g_2)$ in X . Let $p_3 : (X_1, g_1) \times_X (X_2, g_2) \rightarrow (X_3, g_1 g_2)$ be the third projection of the double twisted sector as in 5.3. Let E be the excess intersection bundle on $(X_1, g_1) \times_X (X_2, g_2)$, and $\gamma := p_{3*}(c_{top}(E))$.

In 7.10 we fix a candidate, for every couple X_1, X_2 of twisted sectors, of a cohomology class $\beta = \beta((X_1, g_1), (X_2, g_2)) \in H^*(X, \mathbb{Q})$ such that $p_{3*}(c_{top}(E)) = f_3^*(\beta)$. In Section 7.1 we fix a set of multiplicative generators for $H^*(X, \mathbb{Q})$: boundary divisors, subbanana cycles (2.9) and any fixed set of additive generators for the odd cohomology of X . Our relations depend upon these choices. Finally, we obtain the formula for the Chen–Ruan product

$$(7.1) \quad \alpha_1 *_{CR} \alpha_2 = f_3^*(\tilde{\alpha}_1 \cup \tilde{\alpha}_2 \cup \beta) = \tilde{\alpha}((X_1, g_1), (X_2, g_2)) *_{CR} 1_{(X_3, g_1 g_2)},$$

where we posed $\tilde{\alpha}((X_1, g), (X_2, g_2)) := \tilde{\alpha}_1 \cup \tilde{\alpha}_2 \cup \beta((X_1, g_1), (X_2, g_2))$.

From (7.1), the relations appear naturally divided into two sets. The first set of relations, presented in Section 7.3 comes from the Chen–Ruan product of all couples of twisted sectors. The second set of relations comes, for each twisted sector Y , from the classes of $\overline{\mathcal{M}}_{1,n}$ that are in the kernel of f^* . These are all the relations of $H^*(\overline{\mathcal{M}}_{1,n})$ as a module over $H_{CR}^*(\overline{\mathcal{M}}_{1,n})$ generated by the fundamental classes of the twisted sectors. These relations are determined in Section 7.1. □

Note that we actually obtain finitely many generators of the even part of $H_{CR}^*(\overline{\mathcal{M}}_{1,n})$ as a \mathbb{Q} -algebra as a consequence of Section 7.1. What we mean by “even part” here is in the original grading on $I(\overline{\mathcal{M}}_{1,n})$, without considering the age grading shift. The Chen–Ruan cohomology of $\overline{\mathcal{M}}_{1,n}$ as a \mathbb{Q} -algebra is generated by the fundamental classes of the twisted sectors, the boundary divisors of $\overline{\mathcal{M}}_{1,n}$ and the subbanana cycles (see Definition 2.9). In this description the relations are the two sets of relations described in the proof of Theorem 7.2, and the set of relations for $R^*(\overline{\mathcal{M}}_{1,n})$ (see [24]).

We observe that the first part of the proof of Theorem 7.2 shows that the fundamental classes of the twisted sectors generate the stringy Chow ring of $\overline{\mathcal{M}}_{1,n}$ (see [1], [2]) as an algebra over the ordinary Chow ring of $\overline{\mathcal{M}}_{1,n}$.

7.1. Pull-backs

Let now (Y, g) be a twisted sector of $\overline{\mathcal{M}}_{1,n}$. Let $f : Y \rightarrow \overline{\mathcal{M}}_{1,n}$ be the restriction to Y of the natural map from the inertia stack to $\overline{\mathcal{M}}_{1,n}$. In this section we study the pull-back morphism

$$f^* : H^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q}) \rightarrow H^*(Y, \mathbb{Q}).$$

The main results of this section are:

- (1) the explicit description of the pull-back of the divisor classes of $\overline{\mathcal{M}}_{1,n}$;
- (2) the pull-back morphism f^* is determined by its restriction to the subalgebra of the cohomology generated by the divisors (Proposition 7.5).

Point (1) is enough for proving the generation part of Theorem 7.3. Anyway it is only as a consequence of point (2) that we know all relations of $H_{CR}^*(\overline{\mathcal{M}}_{1,n})$ as an $H^*(\overline{\mathcal{M}}_{1,n})$ -module (and, together with those of Section 7.3, all relations as an $H^*(\overline{\mathcal{M}}_{1,n})$ -algebra).

Let us fix a twisted sector $(Y, g) = (\overline{Z}^{I_1, \dots, I_k}, g)$ (see Theorem 3.20); and identify it with the product of $\overline{Z} \times \overline{\mathcal{M}}_{0, I_1+1} \times \dots \times \overline{\mathcal{M}}_{0, I_k+1}$. Therefore we have⁽⁴⁾

$$\begin{aligned} A^* \left(\overline{Z}^{I_1, \dots, I_k} \right) &= H^{2*}(\overline{Z}^{I_1, \dots, I_k}) \\ &= A^*(\overline{Z}) \times A^*(\overline{\mathcal{M}}_{0, I_1+1}) \times \dots \times A^*(\overline{\mathcal{M}}_{0, I_k+1}). \end{aligned}$$

We call p, p_1, \dots, p_k the projections onto the factors.

The relations are obtained as follows

- take all divisor classes that are in the kernel of f^* ,
- for all $1 \leq i \leq k$, and each relation R in $A^1(\overline{\mathcal{M}}_{0, I_i+1})$ (see [17]), take all linear combinations of divisor classes that pull-back via f to R ,
- take any finite set generating $H^{odd}(\overline{\mathcal{M}}_{1,n})$,
- take all the subbanana cycles (see Definition 2.9).

Let us first deal with point (1), and thus compute the pull-back morphism for the divisor classes. The notation for the divisors in $\overline{\mathcal{M}}_{1,n}$ was introduced in Notation 2.4. The pull-back $f^*(d_{irr})$ is zero when the base space Z is a point. Otherwise,

- (1) it is $\frac{1}{2}[pt] \times [\overline{\mathcal{M}}_{0, n+1}]$, when the space is $\overline{A}_1^{[n]}$;
- (2) it is $\frac{3}{2}[pt] \times [\overline{\mathcal{M}}_{0, I_1+1}] \times [\overline{\mathcal{M}}_{0, I_2+1}]$, when the space is $\overline{A}_2^{I_1, I_2}$;
- (3) it is $3[pt] \times [\overline{\mathcal{M}}_{0, I_1+1}] \times [\overline{\mathcal{M}}_{0, I_2+1}] \times [\overline{\mathcal{M}}_{0, I_3+1}]$, when the space is $\overline{A}_3^{I_1, I_2, I_3}$;
- (4) it is $3[pt] \times [\overline{\mathcal{M}}_{0, I_1+1}] \times [\overline{\mathcal{M}}_{0, I_2+1}] \times [\overline{\mathcal{M}}_{0, I_3+1}] \times [\overline{\mathcal{M}}_{0, I_4+1}]$, when the space is $\overline{A}_4^{I_1, I_2, I_3, I_4}$.

This is a simple consequence of Theorem 3.27.

The pullback $f^*([d_M])$ is zero when M is not contained in any of the I_i s. Assume now that the base space Z is a point. If M is contained in (wlog) I_1 , then there are two cases. If M is a proper subset of I_1 , then

$$f^*([d_M]) \cong [Z] \times \Delta_M \times [\overline{\mathcal{M}}_{0, I_2+1}] \times \dots \times [\overline{\mathcal{M}}_{0, I_k+1}].$$

Otherwise, if $I = M$, then

$$f^*([d_M]) = p_1^*(-\psi_{I+1}).$$

Finally, if \overline{Z} is one of the one-dimensional spaces \overline{A}_i , then the pullback $f^*([d_M])$ is computed by applying Theorem 3.27, similarly to $f^*(d_{irr})$.

A very important theoretical result follows as a corollary of our description of the pull-back morphism.

⁽⁴⁾ see Lemma 7.1 for the next equality

COROLLARY 7.3. — *The morphisms $f^* : R^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q}) \rightarrow A^*(Y, \mathbb{Q})$ are surjective. The same holds for the induced map in cohomology.*

Proof. — Thanks to Lemma 7.1, it is sufficient to prove that the morphism $f^* : R^1(\overline{\mathcal{M}}_{1,n}) \rightarrow A^1(Y)$ is surjective. The Kunneth decomposition reduces the problem to proving that one can obtain all divisors of each single factor of each twisted sector by pull-back from $R^1(\overline{\mathcal{M}}_{1,n})$. The above discussion shows that the set of divisor classes $\{d_{irr}, d_M\}_{M \subset [n]}$ surjects via f^* onto $A^1(Y, \mathbb{Q})$. □

Here is another way to express this result.

COROLLARY 7.4. — *If Y is a twisted sector, then the cohomology $H^*(Y, \mathbb{Q})$ is an $H^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$ -module generated by the fundamental class $[Y]$.⁽⁵⁾*

Following the plan established at the beginning of this section, we now deal with point (2). As a combination of 2.5, 2.6 and (2.3), we have that all cohomology classes of $\overline{\mathcal{M}}_{1,n}$ can be written as sums of odd cohomology classes, of subbanana cycles (Definition 2.9), and of products of divisors. This leads to the decomposition

$$(7.2) \quad H^*(\overline{\mathcal{M}}_{1,n}) = H^{odd}(\overline{\mathcal{M}}_{1,n}) \oplus (R_{1,n}^{div} + R_{1,n}^{ban}),$$

where the sum of the last two terms is the vector space generated by boundary strata classes. It is clear that the pull-back morphism f^* is zero when restricted to the odd cohomology classes. We now prove that f^* is zero when restricted to the vector subspace $R_{1,n}^{ban}$ generated by the subbanana cycles. Using Corollary 3.28 we explicitly express Y as a linear combination of product of divisor classes in $\overline{\mathcal{M}}_{1,n}$. The product of a subbanana cycle with all the summands that contain a factor d_{irr} is zero, because $d_{irr}^2 = 0$, and because a subbanana cycle is, in particular, a closed substack of d_{irr} . All the other summands have product zero with the subbanana cycles, because the set theoretic intersection of the substacks of $\overline{\mathcal{M}}_{1,n}$ that they describe is empty. We can now conclude.

PROPOSITION 7.5. — *The pull-back morphism f^* is determined by its restriction to the subalgebra of $H^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$ generated by the divisors.*

⁽⁵⁾ Indeed $H^*(Y, \mathbb{Q})$ is cyclic also as an $R^*(\overline{\mathcal{M}}_{1,n})$ -module, or as a module over the subring of the tautological ring generated by the divisors.

7.2. Push–forwards

We now start the study of the push–forward morphism. Let

$$g : Z \rightarrow Y, \quad f : Y \rightarrow X$$

be respectively the inclusion of a double twisted sector in a twisted sector and of a twisted sector inside $X = \overline{\mathcal{M}}_{1,n}$. We study the push–forward morphism induced in cohomology by f and g . Here is an easy corollary of Corollary 3.28 and Corollary 7.3.

COROLLARY 7.6. — *The push–forward morphism*

$$f_* : A^*(Z, \mathbb{Q}) \rightarrow A^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$$

has image in the tautological ring. The same holds for the push–forward map in cohomology.

Lemma 7.3 and Corollary 7.6 make it possible to define an orbifold tautological ring in genus 1.

DEFINITION 7.7. — *We define the orbifold tautological ring in genus 1*

- *Let $R_{CR}^*(\overline{\mathcal{M}}_{1,n})$ be defined as $R^*(\overline{\mathcal{M}}_{1,n}) \oplus \bigoplus H^*((X_i, g_i), \mathbb{Q})$ as a vector space, where X_i are all the twisted sectors and the grading is inherited from $H_{CR}^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$;*
- *the product is the product $*_{CR}$ restricted to this rationally graded vector space.*

Note that, as a consequence of Theorem 2.6, $R_{CR}^*(\overline{\mathcal{M}}_{1,n})$ is a Gorenstein ring (a Poincaré duality ring).

We show how $g_*([Z])$ can be obtained as a pull–back of a class in X in a canonical way.

DEFINITION 7.8. — *If $\lambda = 4$ or $\lambda = 6$, we define C_λ^* via the following pull–back diagram*

$$\begin{array}{ccc} C_\lambda^* & \hookrightarrow & \overline{\mathcal{M}}_{1,n} \\ \downarrow & \square & \downarrow \pi_1 \\ C_\lambda & \hookrightarrow & \overline{\mathcal{M}}_{1,1}. \end{array}$$

Note that the equality: $[C_\lambda^*] = \frac{2}{\lambda} d_{irr}$ holds in the tautological ring of $\overline{\mathcal{M}}_{1,n}$.

PROPOSITION 7.9. — *With the notation introduced in this section, for Z a double twisted sector and Y a twisted sector, there is a canonical choice of W closed substack of $\overline{\mathcal{M}}_{1,n}$, such that $g_*([Z]) = f_*([W])$.*

Proof. — The only cases are, thanks to Proposition 5.6:

- (1) it happens that $Z = Y$ or $Y = X$. In all these cases we choose $W := X$;
- (2) either $Z = C_\lambda^{[n]}$ for $\lambda = 4, 6$ and $Y = \overline{A_1}^{[n]}$. In these cases we choose $W := C_\lambda^*$;
- (3) or $Z = C_\lambda^{I_1, I_2}$ and $Y = \overline{A_2}^{I_1, I_2}$. In these cases we choose $W = C_\lambda^*$.

One can easily check that these are all the cases that occur, and that the intersections are transversal. □

We have just fixed the cohomology classes that represent via pull– back the push–forward of all the fundamental classes. This choice determines the top Chern class of the excess intersection bundles via the projection formula.

COROLLARY 7.10. — *Let now E be the excess intersection bundle over the double twisted sector Z . Once the choice of 7.9 is fixed, a cohomology class β on $\overline{\mathcal{M}}_{1,n}$ is determined such that:*

$$g_*(c_{top}(E)) = f^*(\beta).$$

Proof. — If the top Chern class of E is zero, we fix β to be zero. When the top Chern class is 1, the choice of Proposition 7.9 determines the class β of this corollary too. The list of non trivial top Chern classes of excess intersection bundles (non-zero and not 1), is given in 6.10. So, if the top Chern class is a ψ class, there are only two possibilities: either the double twisted sector Z is isomorphic to the twisted sector Y (case 1 of Corollary 6.10), or $Z = C_\lambda^{[n]}$ and $Y = \overline{A_1}^{[n]}$ (cases 2 and 3 of Corollary 6.10). In the first case, we choose β to be $d_{[n]}$, the divisor with all markings on the genus 0 component, and in the second case we choose

$$\beta := d_{[n]} \cup [C_\lambda^*],$$

where C_λ^* is defined in 7.8. □

7.3. Products of the fundamental classes of the twisted sectors

If X_i, X_j are twisted sectors, in this section we compute all the products $1_{X_i} *_{CR} 1_{X_j}$. In the perspective of Theorem 3.23, this section and Section 7.1, provide the relations of $H_{CR}^*(\overline{\mathcal{M}}_{1,n})$ as an $H^*(\overline{\mathcal{M}}_{1,n})$ -algebra generated by the fundamental classes of the twisted sectors.

Remark 7.11. — An explicit computation of all intersections of the twisted sectors, shows that besides the orbifold intersections of the kinds $(1_{X_i}, \alpha) *_{CR} (1_{X_i}, \beta)$, and besides the trivial products $1_{\overline{\mathcal{M}}_{1,n}} *_{CR} 1_{X_i}$, the only pairs of twisted sectors whose fundamental classes give rise to non-zero Chen–Ruan products are in the following list:

- (1) $((\overline{A_1}^{[n]}, -1), (C_4^{[n]}, i/ - i)),$
- (2) $((\overline{A_1}^{[n]}, -1), (C_6^{[n]}, \epsilon/\epsilon^2/\epsilon^4/\epsilon^5)),$
- (3) $((\overline{A_2}^{I_1, I_2}, -1), (C_4^{I_1, I_2}, i/ - i)).$

We now compute the products of the pairs just described. Here if (X, α) is a twisted sector, we write $H^*((X, \alpha), \mathbb{Q})$, which is a direct summand of $H^*_{CR}(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$ with its own grading. In other words, we implicitly assume the inclusion

$$i : H^*((X, \alpha), \mathbb{Q}) \subset H^*_{CR}(\overline{\mathcal{M}}_{1,n}, \mathbb{Q})$$

shifts the degree by twice the age of (X, α) . As usual, $-p_1^*(\psi_\bullet)$ is the Chern class of Corollary 6.10 and Theorem 6.11.

COROLLARY 7.12. — *With our usual notation for the twisted sectors, and with the notation introduced above, here is the explicit result of all the Chen–Ruan products described in Remark 7.11:*

- (1) $[(C_4^{[n]}, i)] *_{CR} [(\overline{A_1}^{[n]}, -1)] = p_1^*(-\psi_\bullet) \in H^2((C_4^{[n]}, -i), \mathbb{Q}),$
- (2) $[(C_4^{[n]}, -i)] *_{CR} [(\overline{A_1}^{[n]}, -1)] = [C_4^{[n]}] \in H^0((C_4^{[n]}, i), \mathbb{Q}),$
- (3) $[(C_6^{[n]}, \epsilon)] *_{CR} [(\overline{A_1}^{[n]}, -1)] = p_1^*(-\psi_\bullet) \in H^2((C_6^{[n]}, \epsilon^4), \mathbb{Q}),$
- (4) $[(C_6^{[n]}, \epsilon^2)] *_{CR} [(\overline{A_1}^{[n]}, -1)] = p_1^*(-\psi_\bullet) \in H^2((C_6^{[n]}, \epsilon^5), \mathbb{Q}),$
- (5) $[(C_6^{[n]}, \epsilon^4)] *_{CR} [(\overline{A_1}^{[n]}, -1)] = [C_6^{[n]}] \in H^0((C_6^{[n]}, \epsilon), \mathbb{Q}),$
- (6) $[(C_6^{[n]}, \epsilon^5)] *_{CR} [(\overline{A_1}^{[n]}, -1)] = [C_6^{[n]}] \in H^0((C_6^{[n]}, \epsilon^2), \mathbb{Q}),$
- (7) $[(\overline{A_2}^{I_1, I_2}, -1)] *_{CR} [(C_4^{I_1, I_2}, i)] = 0 \in H^2((C_4^{I_1, I_2}, -i), \mathbb{Q}),$
- (8) $[(\overline{A_2}^{I_1, I_2}, -1)] *_{CR} [(C_4^{I_1, I_2}, -i)] = [C_4^{I_1, I_2}] \in H^2((C_4^{I_1, I_2}, i), \mathbb{Q}),$

COROLLARY 7.13. — *With our usual notation for the twisted sectors, and with the notation introduced above, here we recollect all the products of the kind $(1_{X_i}, \alpha) *_{CR} (1_{X_i}, \beta)$:*

- (1) $[(X, \alpha)] *_{CR} [(X, \alpha^{-1})] = [X] \in H^*(\overline{\mathcal{M}}_{1,n}, \mathbb{Q}),$
- (2) $[(C_4^{[n]}, i)] *_{CR} [(C_4^{[n]}, i)] = p_1^*(-\psi_\bullet) \cap [C_4^{[n]}] \in H^4((\overline{A_1}^{[n]}, -1), \mathbb{Q}),$
- (3) $[(C_4^{[n]}, -i)] *_{CR} [(C_4^{[n]}, -i)] = [C_4^{[n]}] \in H^2((\overline{A_1}^{[n]}, -1), \mathbb{Q}),$
- (4) $[(C_4^{I_1, I_2}, i)] *_{CR} [(C_4^{I_1, I_2}, i)] = 0 \in H^4((\overline{A_2}^{I_1, I_2}, -1), \mathbb{Q}),$
- (5) $[(C_4^{I_1, I_2}, -i)] *_{CR} [(C_4^{I_1, I_2}, -i)] = [C_4^{I_1, I_2}] \in H^2((\overline{A_2}^{I_1, I_2}, -1), \mathbb{Q}),$
- (6) $[(C_6^{[n]}, \epsilon)] *_{CR} [(C_6^{[n]}, \epsilon)] = 0 \in H^4((C_6^{[n]}, \epsilon^2), \mathbb{Q}),$

- (7) $[(C_6^{[n]}, \epsilon)] *_{CR} [(C_6^{[n]}, \epsilon^2)] = p_1^*(-\psi_\bullet) \cap [C_6^{[n]}] \in H^4((\overline{A_1}^{[n]}, -1), \mathbb{Q})$,
- (8) $[(C_6^{[n]}, \epsilon)] *_{CR} [(C_6^{[n]}, \epsilon^4)] = 0 \in H^4((C_6^{[n]}, \epsilon^5), \mathbb{Q})$,
- (9) $[(C_6^{[n]}, \epsilon^2)] *_{CR} [(C_6^{[n]}, \epsilon^2)] = p_1^*(-\psi_\bullet) \in H^2((C_6^{[n]}, \epsilon^4), \mathbb{Q})$,
- (10) $[(C_6^{[n]}, \epsilon^2)] *_{CR} [(C_6^{[n]}, \epsilon^5)] = p_1^*(-\psi_\bullet) \in H^2((C_6^{[n]}, \epsilon), \mathbb{Q})$,
- (11) $[(C_6^{[n]}, \epsilon^4)] *_{CR} [(C_6^{[n]}, \epsilon^4)] = 0 \in H^2((C_6^{[n]}, \epsilon^2), \mathbb{Q})$,
- (12) $[(C_6^{[n]}, \epsilon^4)] *_{CR} [(C_6^{[n]}, \epsilon^5)] = [C_6^{[n]}] \in H^2((\overline{A_1}^{[n]}, -1), \mathbb{Q})$,
- (13) $[(C_6^{[n]}, \epsilon^5)] *_{CR} [(C_6^{[n]}, \epsilon^5)] = [C_6^{[n]}] \in H^0((C_6^{[n]}, \epsilon^4), \mathbb{Q})$,
- (14) $[(C_6^{I_1, I_2, I_3}, \epsilon^2)] *_{CR} [(C_6^{I_1, I_2, I_3}, \epsilon^2)] = 0 \in H^4((C_6^{I_1, I_2, I_3}, \epsilon^4), \mathbb{Q})$ and
the result still holds when $I_3 = \emptyset$,
- (15) $[(C_6^{I_1, I_2, I_3}, \epsilon^4)] *_{CR} [(C_6^{I_1, I_2, I_3}, \epsilon^4)] = 0 \in H^2((C_6^{I_1, I_2, I_3}, \epsilon^2), \mathbb{Q})$ and
the result still holds when $I_3 = \emptyset$,

Moreover, the product of two fundamental classes of twisted sectors that do not belong to this list, nor to the one of Corollary 7.12, is zero.

BIBLIOGRAPHY

- [1] D. ABRAMOVICH, T. GRABER & A. VISTOLI, “Algebraic orbifold quantum products”, in *Orbifolds in mathematics and physics (Madison, WI, 2001)*, Contemp. Math., vol. 310, Amer. Math. Soc., Providence, RI, 2002, p. 1-24.
- [2] ———, “Gromov–Witten theory of Deligne–Mumford stacks”, *Amer. J. Math.* **130** (2008), no. 5, p. 1337-1398.
- [3] A. ADEM, J. LEIDA & Y. RUAN, *Orbifolds and stringy topology*, Cambridge Tracts in Mathematics, vol. 171, Cambridge University Press, Cambridge, 2007.
- [4] P. BELOROUSKI, “Chow Rings of moduli spaces of pointed elliptic curves”, PhD Thesis, Chicago, 1998.
- [5] L. BREEN, *On the classification of 2-gerbes and 2-stacks*, Astérisque, no. 225, 1994, 160 pp.
- [6] W. CHEN & Y. RUAN, “A new cohomology theory of orbifold”, *Comm. Math. Phys.* **248** (2004), no. 1, p. 1-31.
- [7] F. DIAMOND & J. SHURMAN, *A first course in modular forms*, Graduate Texts in Mathematics, vol. 228, Springer-Verlag, New York, 2005.
- [8] C. FABER & R. PANDHARIPANDE, “Relative maps and tautological classes”, *J. Eur. Math. Soc. (JEMS)* **7** (2005), no. 1, p. 13-49.
- [9] B. FANTECHI & L. GÖTTSCHE, “Orbifold cohomology for global quotients”, *Duke Math. J.* **117** (2003), no. 2, p. 197-227.
- [10] W. FULTON, *Intersection Theory*, Springer-Verlag, Berlin, 1984.
- [11] E. GETZLER, “Operads and moduli of genus 0 Riemann surfaces”, in *The moduli space of curves (Texel Island, 1994)*, Progr. Math., vol. 129, Birkhäuser Boston, Boston, MA, 1995, p. 199-230.
- [12] ———, “Intersection theory on $\overline{M}_{1,4}$ and elliptic Gromov-Witten invariants”, *J. Amer. Math. Soc.* **10** (1997), no. 4, p. 973-998.
- [13] ———, “The semi-classical approximation for modular operads”, *Comm. Math. Phys.* **194** (1998), no. 2, p. 481-492.
- [14] J. GIRAUD, *Cohomologie non abélienne (French)*, Die Grundlehren der mathematischen Wissenschaften, Band, vol. 179, Springer-Verlag, Berlin-New York, 1971.

- [15] T. GRABER & R. PANDHARIPANDE, “Constructions of nontautological classes on moduli spaces of curves”, *Michigan Math. J.* **51** (2003), no. 1, p. 93-109.
- [16] T. GRABER & R. VAKIL, “Relative virtual localization and vanishing of tautological classes on moduli spaces of curves”, *Duke Math. J.* **130** (2005), no. 1, p. 1-37.
- [17] S. KEEL, “Intersection theory of moduli space of stable n -pointed curves of genus zero”, *Trans. Amer. Math. Soc.* **330** (1992), no. 2, p. 545-574.
- [18] É. MANN, “Orbifold quantum cohomology of weighted projective spaces”, *J. Algebraic Geom.* **17** (2008), no. 1, p. 137-166.
- [19] D. MUMFORD, “Towards an enumerative geometry of the moduli space of curves”, in *Arithmetic and geometry Vol II*, Progr. Math., vol. 36, Birkhäuser Boston, Boston, MA, 1983, p. 271-328.
- [20] N. PAGANI, “Chen–Ruan cohomology of moduli of curves”, PhD Thesis, SISSA (Trieste), 2009.
- [21] ———, “The Chen-Ruan cohomology of moduli of curves of genus 2 with marked points”, *Adv. Math.* **229** (2012), no. 3, p. 1643-1687.
- [22] ———, “The orbifold cohomology of moduli of hyperelliptic curves”, *Int. Math. Res. Not. IMRN* (2012), no. 10, p. 2163-2178.
- [23] N. PAGANI & O. TOMMASI, “The orbifold cohomology of moduli of genus 3 curves”, <http://arxiv.org/abs/1103.0151>.
- [24] D. PETERSEN, “The structure of the tautological ring in genus 1”, <http://arxiv.org/abs/1205.1586v1>.
- [25] J. H. SILVERMAN, *The arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, New York, 1992.
- [26] J. E. SPENCER, “The stringy Chow ring of the moduli stack of genus-two curves and its Deligne-Mumford compactification”, PhD Thesis, Boston, 2004.
- [27] ———, “The orbifold cohomology of the moduli of genus-two curves”, in *Gromov-Witten theory of spin curves and orbifolds*, Contemp. Math., vol. 403, Amer. Math. Soc., Providence, RI, 2006, p. 167-184.
- [28] A. VISTOLI, “Intersection theory on algebraic stacks and on their moduli spaces”, *Invent. Math.* **97** (1989), no. 3, p. 613-670.

Manuscrit reçu le 1^{er} avril 2011,
révisé le 10 juin 2012,
accepté le 30 août 2012.

Nicola PAGANI
Institut für Algebraische Geometrie,
Leibniz Universität Hannover,
Welfengarten 1, D-30167 Hannover, Germany
npagani@math.uni-hannover.de