



ANNALES

DE

L'INSTITUT FOURIER

Gilbert LEVITT

Generalized Baumslag–Solitar groups: rank and finite index subgroups

Tome 65, n° 2 (2015), p. 725-762.

http://aif.cedram.org/item?id=AIF_2015__65_2_725_0



© Association des Annales de l'institut Fourier, 2015,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

GENERALIZED BAUMSLAG–SOLITAR GROUPS: RANK AND FINITE INDEX SUBGROUPS

by Gilbert LEVITT

ABSTRACT. — A generalized Baumslag–Solitar (GBS) group is a finitely generated group acting on a tree with infinite cyclic edge and vertex stabilizers. We show how to determine effectively the rank (minimal cardinality of a generating set) of a GBS group; as a consequence, one can compute the rank of the mapping torus of a finite order outer automorphism of a free group F_n . We also show that the rank of a finite index subgroup of a GBS group G cannot be smaller than the rank of G . We determine which GBS groups are large (some finite index subgroup maps onto F_2), and we solve the commensurability problem (deciding whether two groups have isomorphic finite index subgroups) in a particular family of GBS groups.

RÉSUMÉ. — Un groupe de Baumslag–Solitar généralisé (groupe GBS) est un groupe de type fini agissant sur un arbre avec stabilisateurs de sommets et d’arêtes infinis cycliques. Nous déterminons explicitement le rang (nombre minimal de générateurs) d’un groupe GBS, et en déduisons le rang de la suspension d’un automorphisme d’ordre fini d’un groupe libre F_n . Nous montrons aussi que le rang ne peut pas diminuer quand on passe à un sous-groupe d’indice fini d’un groupe GBS. Nous déterminons quels groupes GBS sont larges (un sous-groupe d’indice fini se surjecte sur F_2), et nous résolvons le problème de commensurabilité (décider si deux groupes ont des sous-groupes d’indice fini isomorphes) dans une certaine famille de groupes GBS.

1. Introduction and statement of results

This paper studies generalized Baumslag–Solitar (GBS) groups. These are finitely generated groups G acting on a tree T with infinite cyclic edge and vertex stabilizers (equivalently, fundamental groups of finite graphs of groups Γ with all vertex and edge groups \mathbb{Z}). Basic examples are provided by the Baumslag–Solitar groups $BS(m, n) = \langle a, t \mid ta^mt^{-1} = a^n \rangle$, known for sometimes being non-Hopfian [3, 10], and one-relator groups with non-trivial center [30].

Keywords: Group, Baumslag–Solitar, rank, finite index.

Math. classification: 20E06, 20E08, 20F05, 20F65.

GBS groups have been studied in particular in relation with JSJ decompositions [18], quasi-isometries [33], automorphisms [6, 25], cohomological dimension [23], the Haagerup property [11], Bredon cohomology [13], one-relator groups [9, 27, 30]. Solving the conjugacy problem in GBS groups is possible but non-trivial [4], while the isomorphism problem for GBS groups is open in general (see [7, 19, 25] for partial results).

Our first result is a computation of the rank (minimal cardinality of a generating set). The rank is a basic invariant of groups, still there are very few families of groups for which the rank is known (and the rank of a (Gromov)–hyperbolic group is not even computable [2]). It is therefore somewhat surprising that there is a complete and explicit way of computing the rank $\text{rk}(G)$ of a GBS group G .

A GBS group has standard generating sets, with one generator a_v per vertex of the graph of groups Γ , and one stable letter t_e for each edge outside of a given maximal subtree. It turns out that the rank is achieved by a subset of any standard generating set, and that there is a simple combinatorial criterion to decide which subsets of a standard set are themselves generating sets.

The basic notion involved in this criterion is that of a *plateau*, which we shall explain after considering two simple examples (see also Definition 3.1).




Figure 1.1. G_1 has rank 2 if n is odd, 3 if n is even. $\{v_a\}$ is a 3–plateau, $\{v_b\}$ is a 2–plateau for n even, $\{v_c\}$ is a 5–plateau, e_1 is a p –plateau if $p > 3$ divides n , e_2 is a 2–plateau if n is odd, and Γ is a p –plateau for $p > 5$ not dividing n . If n is odd, every plateau meets $\{v_a, v_c\}$, so $G_1 = \langle a, c \rangle$.

Figure 1.1 represents the labelled graph Γ associated to $G_1 = \langle a, b, c \mid a^3 = b^2, b^n = c^5 \rangle$ (the graph is labelled by 3, 2, n , 5, which are indices of edge groups in vertex groups). If n is even, G_1 has rank 3 because adding the relation $b^2 = 1$ maps G_1 onto $\mathbb{Z}/3\mathbb{Z} * \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/5\mathbb{Z}$, which has rank 3 by Grushko’s theorem. On the other hand, if n is odd, $G_1 = \langle a, c \rangle$ has rank 2.

The group $G_2 = \langle a, b, c, d \mid a^3 = b^2, b^3 = c^7, c^{10} = d^5 \rangle = \langle a, b, d \rangle = \langle a, c, d \rangle$ of Figure 1.2 has rank 3: adding the relations $b^2 = c^2 = 1$ maps G_2




Figure 1.2. $\{v_a\}$ is a 3-plateau, $\{v_d\}$ is a 5-plateau, e_2 is a 2-plateau, $e_1 \cup e_2$ and $\{v_d\}$ are 5-plateaux, e_3 is a 7-plateau, and Γ is a p -plateau for $p > 7$. Every plateau meets $\{v_a, v_b, v_d\}$ and $\{v_a, v_c, v_d\}$, so $G_2 = \langle a, b, d \rangle = \langle a, c, d \rangle$.

onto $\mathbb{Z}/3\mathbb{Z} * \langle b, c \mid b^3 = c^7, b^2 = c^2 = 1 \rangle * \mathbb{Z}/5\mathbb{Z}$, which has rank 3 because the middle group is $\mathbb{Z}/2\mathbb{Z}$.

This $\mathbb{Z}/2\mathbb{Z}$ is associated to the edge e_2 , which we call a 2-plateau: 2 is a prime, e_2 is a connected non-empty subgraph P , the labels carried by edges in P (3 and 7) are not divisible by 2, but to exit the plateau one must pass by a label (2 or 10) which is divisible by 2 (see Figures 1.1 and 1.2 for examples, and Definition 3.1 for a precise definition). A plateau P is proper if $P \neq \Gamma$.

We will show that one obtains a generating set of minimal cardinality $\text{rk}(G)$ by removing certain generators a_v from any standard generating set $\{(a_v), (t_\varepsilon)\}$. The set remains a generating set provided one keeps at least one v in each plateau (see Figures 1.1 and 1.2). Thus:

THEOREM 1.1 (Theorem 3.2). — *Let G be a GBS group represented by a labelled graph Γ . The rank of G is $\beta(\Gamma) + \mu(\Gamma)$, where $\beta(\Gamma)$ is the first Betti number and $\mu(\Gamma)$ is the minimal cardinality of a set of vertices meeting every plateau.*

On Figure 1.3, $G = \langle a, b, c, t \mid a^3 = b^5, b^2 = c^3, tc^5t^{-1} = a^2 \rangle = \langle a, b, t \rangle = \langle a, c, t \rangle = \langle b, c, t \rangle$. There are four plateaux: Γ and each of the edges, so $\mu(\Gamma) = 2$ and $\text{rk}(G) = 1 + 2 = 3$ by the theorem. In this example Grushko’s theorem is not sufficient to prove $\text{rk}(G) \geq 3$, and we have to use Dunwoody’s folding sequences [15].

COROLLARY 1.2 (Corollary 4.4). — *One may compute the rank of the mapping torus of a finite order outer automorphism of a finitely generated free group F_n .*

Indeed, these mapping tori are exactly the GBS groups with a non-trivial center (see Proposition 4.1).

Using Theorem 1.1, we can show:




Figure 1.3. e_1 is a 2-plateau, e_2 is a 5-plateau, e_3 is a 3-plateau, Γ is a p -plateau for $p > 5$, and $G = \langle a, b, t \rangle = \langle a, c, t \rangle = \langle b, c, t \rangle$.

THEOREM 1.3 (Theorem 5.1). — *If \bar{G} is a finite index subgroup of a GBS group G , then $\text{rk}(\bar{G}) \geq \text{rk}(G)$.*

This property holds in free groups and surface groups, but not in arbitrary hyperbolic groups. As pointed out by I. Kapovich, the Rips construction [31] (applied to finite groups) yields hyperbolic groups of arbitrarily large rank containing a 2-generated subgroup of finite index.

A group \bar{G} as in the theorem is represented by a labelled graph $\bar{\Gamma}$, with a map $\pi : \bar{\Gamma} \rightarrow \Gamma$ sending edge to edge and vertex to vertex. It is easy to show $\beta(\bar{\Gamma}) \geq \beta(\Gamma)$. If some component of $\pi^{-1}(P)$ is a plateau whenever $P \subset \Gamma$ is a plateau, the result is clear. Unfortunately this is not true in general, so that $\mu(\bar{\Gamma})$ may be smaller than $\mu(\Gamma)$, and we have to show that this is compensated for by an increase in β . This is not too hard to do when Γ contains no proper 2-plateau, but 2-plateaux bring technical complications. See for instance Figure 1.4 for a subgroup of index 2 of $G = \langle a, b, t \mid a^2 = b^2, tb^3t^{-1} = b^5 \rangle$. The graph Γ representing G contains two proper 2-plateaux (the terminal vertex and the ellipse), but $\bar{\Gamma}$ contains no proper plateau.

If Γ represents G , and $\pi : \Gamma' \rightarrow \Gamma$ is a finite covering, one may lift the labels from Γ to Γ' , and Γ' represents a finite index subgroup of G . When Γ contains no proper plateau, every finite index subgroup of G is isomorphic to one obtained by this construction (Proposition 6.5). On the




Figure 1.4.

other hand, when Γ contains a proper p -plateau P , there are other finite index subgroups, obtained by taking a covering of degree p branched over P (on Figure 1.4, the branching is over the union of two 2-plateaux). As an application, we identify which GBS groups are large (i.e. they have a finite index subgroup mapping onto a free group F_2):

THEOREM 1.4 (Theorem 6.7). — *A non-cyclic GBS group G is large if and only if it cannot be represented by a labelled graph homeomorphic to a circle and containing no proper plateau.*

When Γ is a circle, with labels denoted x_i and y_i alternatively, the absence of a proper plateau is equivalent to $\prod x_i$ being coprime with $\prod y_i$. Theorem 1.4 was proved independently by T. Mecham [28]; see [16] for the case of Baumslag–Solitar groups. We show in [26] that G is large if and only if it is not a quotient of $BS(m, n)$ with m, n coprime.

There may be infinitely many different labelled graphs Γ representing a given GBS group G (this is why the isomorphism problem is difficult). Uniqueness (up to sign changes and restricting to reduced graphs) holds when Γ is strongly slide-free [17]: no two labels near a given vertex divide each other. Using this, we show:

THEOREM 1.5 (Theorem 6.9). — *The commensurability problem is solvable among GBS groups represented by labelled graphs which are strongly slide-free and contain no proper plateau.*

The commensurability problem consists in deciding whether two given groups contain isomorphic finite index subgroups. Among Baumslag–Solitar groups $BS(m, n)$, it is solved when m and n are coprime [33], or equal in absolute value, but the general case is open. Note that all GBS groups which are virtually $F_n \times \mathbb{Z}$ for some n are commensurable (provided $n > 1$); these groups are called unimodular (see Section 2), in particular every GBS group with non-trivial center is unimodular.

Theorem 1.5 applies in particular to the non-large groups of Theorem 1.4. We show that any GBS group has a finite index subgroup represented by a labelled graph with no proper plateau (Proposition 6.8), but unfortunately this graph is not strongly slide-free in general.

Deciding commensurability in Theorem 1.5 amounts to deciding whether the labelled graphs have a common finite covering; Leighton’s graph covering theorem [24, 29] ensures that this is possible.

I am grateful to M. Forester for explaining how to visualize finite index subgroups, to R. Weidmann for suggesting the use of folding sequences, and to W. Dicks for mentioning Leighton’s theorem and the reference [29]. This work was partly supported by ANR-07-BLAN-0141-01 and ANR-2010-BLAN-116-03.

2. Preliminaries

We refer to [18, 19, 25] for basic facts about GBS groups.

GBS groups are represented by labelled graphs. A *labelled graph* is a finite graph Γ where each oriented edge e has a label λ_e , a nonzero integer (possibly negative). We denote by V the set of vertices of Γ , and by \mathcal{E} the set of non-oriented edges. We view a non-oriented edge as $\varepsilon = (e, \tilde{e})$, where \tilde{e} is e with the opposite orientation. We denote by $v = o(e)$ the origin of e , and by E_v the set of oriented edges with origin v . The cardinality $|E_v|$ of E_v is the *valence* of v . A vertex is *terminal* if it has valence one. We say that λ_e is the *label of e near the vertex $o(e)$* , and that λ_e is a label *carried by e or ε* . There are $|E_v|$ labels near a vertex v .

We write $m \wedge n$ for the greatest common divisor (gcd) of two integers, with the convention that $m \wedge n > 0$ regardless of the sign of m, n . On the other hand, $lcm(m, n) := \frac{mn}{m \wedge n}$ may be negative.

A connected labelled graph defines a graph of groups. All edge and vertex groups are \mathbb{Z} , and the inclusion from the edge group G_e to the vertex group $G_{o(e)}$ is multiplication by λ_e . The fundamental group G of the graph of groups is the *GBS group represented by Γ* (we do not always assume that Γ is connected, but we implicitly do so whenever we refer to the group it represents). The group G may be presented as follows.

Choose a maximal subtree $\Gamma_0 \subset \Gamma$. There is one generator a_v for each vertex $v \in V$, and one generator t_ε (stable letter) for each $\varepsilon \in \mathcal{E}_0$, the set of non-oriented edges not in Γ_0 . Each non-oriented edge $\varepsilon = (e, \tilde{e})$ of Γ contributes a relation R_ε . The relation is $(a_{o(e)})^{\lambda_e} = (a_{o(\tilde{e})})^{\lambda_{\tilde{e}}}$ if ε is in Γ_0 , and $t_\varepsilon(a_{o(e)})^{\lambda_e} t_\varepsilon^{-1} = (a_{o(\tilde{e})})^{\lambda_{\tilde{e}}}$ if ε is not in Γ_0 (replacing e by \tilde{e} amounts to replacing t_ε by its inverse). This will be called a *standard presentation* of G , and the generating set will be called a *standard generating set* (associated to Γ and Γ_0).

A GBS group is *elementary* if it is isomorphic to \mathbb{Z} , or \mathbb{Z}^2 , or the Klein bottle group $K = \langle x, y \mid x^2 = y^2 \rangle = \langle a, t \mid tat^{-1} = a^{-1} \rangle$. These are the only virtually abelian GBS groups, and they have very special properties. Unless mentioned otherwise, our results apply to all GBS groups, but we do not always provide proofs for elementary groups.

A labelled graph Γ is *minimal* if its Bass–Serre tree T contains no proper G -invariant subtree; this is equivalent to no label near a terminal vertex being equal to ± 1 . The graph Γ is *reduced* [17] if any edge e such that $\lambda_e = \pm 1$ is a loop (e and \tilde{e} have the same origin). Any labelled graph may be made reduced by a sequence of elementary collapses (see [17]); these collapses do not change G .

The group G represented by Γ does not change if one changes the sign of all labels near a given vertex v , or if one changes the sign of the labels $\lambda_e, \lambda_{\tilde{e}}$ carried by a given non-oriented edge ε . These will be called *admissible sign changes*.

In general, there may be infinitely many reduced labelled graphs representing a given G . One case when uniqueness holds (up to admissible sign changes) is when Γ is *strongly slide-free* ([17], see also [21]): if e and e' are edges with the same origin, λ_e does not divide $\lambda_{e'}$.

We denote by $\beta(\Gamma)$ the first Betti number of Γ . If G is non-elementary, all labelled graphs representing G have the same β , and we sometimes denote it by $\beta(G)$.

Let G be represented by Γ . An element $g \in G$ is *elliptic* if it fixes a point in the Bass–Serre tree T associated to Γ , or equivalently if some conjugate of g belongs to a vertex group of Γ . All elliptic elements are

pairwise *commensurable* (they have a common power). An element which is not elliptic is *hyperbolic*. If G is non-elementary, ellipticity or hyperbolicity does not depend on the choice of Γ representing G .

The quotient of G by the subgroup generated by all elliptic elements may be identified with the topological fundamental group $\pi_1^{top}(\Gamma)$ of the graph Γ , a free group of rank $\beta(\Gamma)$. In particular, any generating set of G contains at least $\beta(\Gamma)$ hyperbolic elements.

If G is non-elementary, there is a *modular homomorphism* $\Delta_G : G \rightarrow \mathbb{Q}^*$ associated to G . It may be characterized as follows: given any non-trivial elliptic element a , there is a non-trivial relation $ga^mg^{-1} = a^n$, and $\Delta_G(g) = \frac{m}{n}$ (the numbers m, n may depend on the choice of a , but m/n does not).

Elliptic elements have modulus 1, so Δ_G factors through $\pi_1^{top}(\Gamma)$ for any Γ representing G . One may define Δ_G directly in terms of loops in Γ : if $\gamma \in \pi_1^{top}(\Gamma)$ is represented by an edge-loop (e_1, \dots, e_m) , then $\Delta_G(\gamma) = \prod \frac{\lambda_{e_i}}{\lambda_{e_i}}$. Note that Δ_G is trivial if Γ is a tree.

Δ_G is trivial if and only if the center of G is non-trivial. In this case the center is cyclic and only contains elliptic elements (see [25]). Moreover, there is an epimorphism $G \rightarrow \mathbb{Z}$ whose kernel contains no non-trivial elliptic element (see Proposition 3.3 of [25]).

A non-elementary G is *unimodular* if the image of Δ_G is contained in $\{1, -1\}$. This is equivalent to G having a normal infinite cyclic subgroup, and also to G being virtually $F_n \times \mathbb{Z}$ for some $n \geq 2$.

Given any Γ , one may perform admissible sign changes so that at most $\beta(\Gamma)$ labels are negative (edges in some maximal subtree only carry positive labels, edges not in the subtree carry at most one negative label). If Δ_G only takes positive values, one may make all labels positive.

3. Computing the rank

Let G be a GBS group. In this section we fix a (connected) labelled graph Γ representing G . We assume that it is finite, but not necessarily minimal or reduced.

DEFINITION 3.1 (plateau, plateaunic number). — *Let p be a prime number. A non-empty connected subgraph $P \subset \Gamma$ is a p -plateau if the following condition holds for every oriented edge e with $v = o(e)$ belonging to P : the label λ_e of e near v is divisible by p if and only if e is not contained in P . We say that P is a plateau if it is a p -plateau for some p . It is proper if $P \neq \Gamma$.*

The plateaunic number $\mu(\Gamma)$ is the minimal cardinality of a set of vertices meeting every plateau.

Let P be a p -plateau. Given $e = vw$, there are four possibilities. If $e \subset P$, none of the labels $\lambda_e, \lambda_{\bar{e}}$ is divisible by p . If $v, w \in P$ but e is not contained in P , both labels $\lambda_e, \lambda_{\bar{e}}$ are divisible by p . If $v \in P$ and $w \notin P$ (or vice versa), the label of e near v is divisible by p . If $v, w \notin P$, there is no restriction.

If v is a vertex, $\{v\}$ is a p -plateau if and only if p divides every label near v . A terminal vertex with label $\neq \pm 1$ is a plateau. Two p -plateaux are disjoint or equal. The whole graph Γ is a p -plateau for all but finitely many p , so $\mu(\Gamma) \geq 1$.

THEOREM 3.2. — *Let G be a GBS group represented by a labelled graph Γ .*

- *Every standard generating set contains a generating set of cardinality $\text{rk}(G)$.*
- *The rank of G equals $\beta(\Gamma) + \mu(\Gamma)$, where $\beta(\Gamma)$ is the first Betti number of Γ and $\mu(\Gamma)$ is its plateaunic number (see Definition 3.1).*

In particular, the rank of G is computable from any labelled graph representing G . It follows from Proposition 3.9 below that one obtains a generating set of cardinality $\text{rk}(G)$ by finding a set of vertices $V_1 \subset V$ of cardinality $\mu(\Gamma)$ meeting every plateau, and deleting generators a_v for $v \notin V_1$ from any standard generating set $\{(a_v)_{v \in V}, (t_\varepsilon)_{\varepsilon \in \mathcal{E}_0}\}$.

Remark 3.3. — $\beta(\Gamma) + \mu(\Gamma)$ is an invariant: it only depends on G . For G non-elementary it is well-known that $\beta(\Gamma)$ is an invariant, and it is not hard to deduce directly from [17] or [8] that $\mu(\Gamma)$ is also an invariant.

The remainder of this section is devoted to the proof of Theorem 3.2, so we fix G and Γ .

Given a prime number p and a p -plateau $P \subset \Gamma$, fix a maximal subtree $P_0 \subset P$, and a maximal subtree $\Gamma_0 \subset \Gamma$ with $\Gamma_0 \cap P = P_0$. Consider the associated standard presentation, with generators a_v and t_ε . We define a quotient G' of G by adding the relations $a_v = 1$ for $v \notin P$, and $(a_v)^p = 1$ for $v \in P$.

LEMMA 3.4. — *The rank of G' is $\beta(\Gamma) + 1$.*

Proof. — Consider an edge $\varepsilon = (e, \bar{e})$ not contained in P . The elements $(a_{o(e)})^{\lambda_e}$ and $(a_{o(\bar{e})})^{\lambda_{\bar{e}}}$ are killed in G' since λ_e is divisible by p if $o(e) \in P$, so we may remove the relation R_ε from the presentation of G' .

For simplicity, we first consider the case when Γ is a tree. Then G' is generated by the elements a_v for $v \in P$, and $(a_v)^p = 1$ in G' . The other relations come from edges of P , they are of the form $(a_{o(e)})^{\lambda_e} = (a_{o(\bar{e})})^{\lambda_{\bar{e}}}$. The key point here (as in the group G_2 of Figure 1.2) is that no exponent λ_e (or $\lambda_{\bar{e}}$) is divisible by p , so multiplication by λ_e defines an automorphism of $\mathbb{Z}/p\mathbb{Z}$. Thus G' is the fundamental group of a graph of groups with underlying graph P , vertex and edge groups $\mathbb{Z}/p\mathbb{Z}$, and inclusions given by multiplication by the λ_e 's. It follows that G' is isomorphic to $\mathbb{Z}/p\mathbb{Z}$, so has rank $1 = \beta(\Gamma) + 1$.

The general case is similar. Since $\Gamma_0 \cap P = P_0$, there are $\beta(\Gamma) - \beta(P)$ edges not contained in either P or Γ_0 . The associated stable letters are not involved in any relation, so G' is the free product of a free group of rank $\beta(\Gamma) - \beta(P)$ with $G'' = \langle (a_v)_{v \in P}, (t_\varepsilon)_{\varepsilon \subset P \setminus P_0} \rangle$. As in the previous case, G'' is the fundamental group of a graph of $\mathbb{Z}/p\mathbb{Z}$'s based on P . It maps onto the topological fundamental group of P with kernel $\mathbb{Z}/p\mathbb{Z}$, so has rank $\beta(P) + 1$. By Grushko's theorem, G' has rank $\beta(\Gamma) - \beta(P) + \beta(P) + 1 = \beta(\Gamma) + 1$. \square

DEFINITION 3.5 (Minimally hyperbolic). — *A generating set S of G is minimally hyperbolic if it contains exactly $\beta(\Gamma)$ hyperbolic elements.*

Recall that any generating set contains at least $\beta(\Gamma)$ hyperbolic elements. Standard generating sets are minimally hyperbolic.

COROLLARY 3.6. — *Let S be a minimally hyperbolic generating set. For each elliptic element $s \in S$, fix a vertex v_s of Γ such that s has a conjugate contained in the vertex group G_{v_s} . Then every plateau contains some v_s . In particular, S has at least $\beta(\Gamma) + \mu(\Gamma)$ elements.*

Proof. — Let P be a p -plateau, and let G' be as above. If P contains no v_s , every a_{v_s} is killed in G' , so every elliptic element of S is killed in G' . Since S is minimally hyperbolic, G' may be generated by $\beta(\Gamma)$ elements. This contradicts Lemma 3.4. \square

The following fact is used in [26].

COROLLARY 3.7. — *Let S be a minimally hyperbolic generating set. Given a p -plateau P , there exist $s \in S$ and $v \in P$ such that s is conjugate to a power $(a_v)^q$ with q not divisible by p .*

Proof. — As in the previous proof, some elliptic $s \in S$ must survive in G' . Being elliptic, s is conjugate to some power $(a_v)^q$ with $v \in V$. Survival implies that v is in P and q is not divisible by p . \square

Remark 3.8. — If S is not minimally hyperbolic, let $h(S)$ be the number of hyperbolic elements in S . If $P_1, \dots, P_i, \dots, P_k$ are disjoint p_i -plateaux

with $k > h(S) - \beta(\Gamma)$, there exist $s \in S$, and i , such that s is conjugate to $(a_v)^q$ with $v \in P_i$ and q not divisible by p_i . To prove this, one defines G' by adding relations $a_v = 1$ for $v \notin \cup_i P_i$ and $(a_v)^{p_i} = 1$ for $v \in P_i$. It has rank $\beta(\Gamma) + k$, so some elliptic $s \in S$ survives in G' .

PROPOSITION 3.9. — *Let $S = \{(a_v)_{v \in V}, (t_\varepsilon)_{\varepsilon \in \mathcal{E}_0}\}$ be a standard generating set. For $V_1 \subset V$, define $S_1 \subset S$ by keeping only the t_ε 's and the a_v 's for $v \in V_1$. Then S_1 generates G if and only if V_1 meets every plateau.*

Proof. — If S_1 generates, V_1 meets every plateau by Corollary 3.6. We now suppose that $G_1 = \langle S_1 \rangle$ is not G , and we construct a plateau P disjoint from V_1 . We may assume $V_1 \neq \emptyset$ and $V_1 \neq \Gamma$.

For $v \in V$, define the integer $n_v \geq 1$ by $G_1 \cap \langle a_v \rangle = \langle a_v^{n_v} \rangle$; it exists because $V_1 \neq \emptyset$ and the a_v 's are all commensurable. Note that $n_v = 1$ if $v \in V_1$. Since $G_1 \neq G$, we have $n_v > 1$ for some $v \notin V_1$, so we may fix a prime p dividing some n_v . For each $v \notin V_1$, let p^{δ_v} be the maximal power dividing n_v , and let $\delta \geq 1$ be the maximal value of δ_v (for $v \notin V_1$).

We now define a subgraph $\Gamma_1 \subset \Gamma$ as follows. Its vertices are the vertices $v \notin V_1$ such that $\delta_v = \delta$. An edge $\varepsilon = (e, \tilde{e})$ is in Γ_1 if and only if its vertices are in Γ_1 and none of $\lambda_e, \lambda_{\tilde{e}}$ is divisible by p . We complete the proof by showing that every component of Γ_1 is a p -plateau.

This is equivalent to the following fact: if e is an edge with $\delta_{o(e)} = \delta$ such that the label λ_e of e near $o(e)$ is not divisible by p , then $\delta_{o(\tilde{e})} = \delta$ and the label $\lambda_{\tilde{e}}$ near $o(\tilde{e})$ is not divisible by p . Let us write $v = o(e)$ and $w = o(\tilde{e})$.

Let $a_e = (a_v)^{\lambda_e}$. Since p does not divide λ_e , and $\delta_v = \delta$, the smallest n such that $(a_e)^n \in G_1$ is divisible by p^δ . The standard presentation of G contains a relation R_ε expressing a_e as $(a_w)^{\lambda_{\tilde{e}}}$ (possibly conjugated by a stable letter t_ε). Let $\theta = \frac{lcm(n_w, \lambda_{\tilde{e}})}{\lambda_{\tilde{e}}} = \frac{n_w}{n_w \wedge \lambda_{\tilde{e}}}$. Since G_1 contains $(a_w)^{n_w}$ and all stable letters, it contains $(a_w)^{\lambda_{\tilde{e}}\theta}$ and $(a_e)^\theta$. It follows that p^δ divides θ , hence n_w , so $\delta_w = \delta$. Since n_w is not divisible by $p^{\delta+1}$, we deduce that p does not divide $\lambda_{\tilde{e}}$. □

Corollary 3.6 and Proposition 3.9 imply:

COROLLARY 3.10.

- Every standard generating set contains a generating set of cardinality $\beta(\Gamma) + \mu(\Gamma)$. In particular, $\beta(\Gamma) + \mu(\Gamma) \geq \text{rk}(G)$.
- The minimal cardinality of a minimally hyperbolic generating set is $\beta(\Gamma) + \mu(\Gamma)$.

This is clear, recalling that standard generating sets are minimally hyperbolic. The following proposition now completes the proof of Theorem 3.2.

PROPOSITION 3.11. — *There exists a minimally hyperbolic generating set of cardinality $\text{rk}(G)$.*

Proof. — The proof is by induction on $r = \text{rk}(G)$, with the result clear for $r = 1$. We fix a generating set X of cardinality r , which we view as an epimorphism π from the free group $F(X)$ to G . Let T_0 be the Cayley tree of $F(X)$, with the natural action of $F(X)$. Let T be the Bass–Serre tree of Γ , with the natural action of G . Trimming Γ if necessary, we may assume that G acts minimally on T .

By Theorem 2.1 of [15], there exists a sequence $T_0 \xrightarrow{f_0} T_1 \xrightarrow{f_1} \dots \rightarrow T_i \rightarrow \dots \xrightarrow{f_{n-1}} T_n = T$ of simplicial trees T_i with an action of a group G_i such that:

- (1) $G_0 = F(X)$, $G_n = G$, and there are epimorphisms $\pi_i : G_i \rightarrow G_{i+1}$, with $\pi_{n-1} \circ \dots \circ \pi_0 = \pi$;
- (2) the restriction of $\pi_{n-1} \circ \dots \circ \pi_i$ to each vertex stabilizer of T_i is injective;
- (3) the map $f_i : T_i \rightarrow T_{i+1}$ is equivariant with respect to π_i ; it is either a subdivision or a basic fold followed by a vertex morphism (see [15] for definitions).

The action of G_i on T_i is not necessarily minimal. By condition 2 (and equivariance), all vertex and edge stabilizers are cyclic (possibly trivial). In particular, G_i is a free product of GBS groups. Note that all groups G_i have rank r . If a group acts on a tree T , we say that a generating set is minimally T -hyperbolic if no other generating set contains fewer hyperbolic elements.

Consider the smallest k such that G_{k+1} is a GBS group.

- We first show that G_{k+1} has a minimally T_{k+1} -hyperbolic generating set of cardinality r .

The previous group G_k is a nontrivial free product of (possibly cyclic) GBS groups of rank $< r$. Indeed, define a graph of groups $\hat{\Gamma}_k$ by collapsing to a point each edge of $\Gamma_k = T_k/G_k$ with stabilizer \mathbb{Z} . We can write G_k as the free product of GBS vertex groups H_v of $\hat{\Gamma}_k$, together with $\beta(\hat{\Gamma}_k)$ infinite cyclic groups generated by stable letters. The free product is nontrivial because G_k is not a GBS group.

By induction on r , each H_v has a minimally T_k -hyperbolic generating set S_v of cardinality $\text{rk}(H_v)$. Consider a generating set S_k of G_k obtained by adjoining $\beta(\hat{\Gamma}_k)$ stable letters to the union of the S_v 's. By Grushko's theorem, it has r elements. Since each S_v is minimally T_k -hyperbolic, exactly $\beta(\Gamma_k)$ elements of S_k are hyperbolic in T_k .

If passing from T_k to T_{k+1} involves no basic fold of type III (see Figure 1 in [Du]), one has $\beta(\Gamma_{k+1}) = \beta(\Gamma_k)$. Since $\pi_k(g)$ is elliptic in T_{k+1} if g is elliptic in T_k , the image of S_k in G_{k+1} is the desired minimally T_{k+1} -hyperbolic generating set. If there is a fold of type III, so that $\beta(\Gamma_{k+1}) = \beta(\Gamma_k) - 1$, the stable letter associated to the folded loop (the element g in Figure 1 of [15]) becomes elliptic in T_{k+1} . Once again $\pi_k(S_k)$ is a minimally T_{k+1} -hyperbolic generating set of G_{k+1} .

• We now show by induction on $i \geq k + 1$ that G_i has a minimally T_i -hyperbolic generating set of cardinality r . This proves the proposition since $G_n = G$.

The result is true for $i = k + 1$. Assume it is true for some i . Let S_i be a minimally T_i -hyperbolic generating set of G_i of cardinality r , obtained by removing elliptic elements from any standard generating set (such an S_i exists by the induction hypothesis and Corollary 3.10). As above, the image of S_i in G_{i+1} is the desired generating set: if there is a basic fold of type III, we have $\beta(\Gamma_{i+1}) = \beta(\Gamma_i) - 1$, but the definition of S_i guarantees that some stable letter in S_i becomes elliptic in T_{i+1} . □

4. Finite order automorphisms of free groups

We denote by F_n the free group of rank n . Given $\Phi \in \text{Out}(F_n)$, we denote by M_Φ the mapping torus $M_\Phi = F_n \rtimes_\Phi \mathbb{Z} = \langle F_n, t \mid tgt^{-1} = \alpha(g) \rangle$, where $\alpha \in \text{Aut}(F_n)$ is any representative of Φ .

PROPOSITION 4.1. — *Given a group G , the following are equivalent:*

- G is a GBS group with non-trivial center;
- there exist n and $\Phi \in \text{Out}(F_n)$ of finite order such that $G = M_\Phi$.

Proof. — We may assume that G is not \mathbb{Z} , \mathbb{Z}^2 , or the Klein bottle group.

Let G be a GBS group with non-trivial center Z . Then (see Section 2) G is virtually $F_r \times \mathbb{Z}$ for some $r \geq 2$, and there is an epimorphism $\tau : G \rightarrow \mathbb{Z}$ which is non-trivial on Z . The restriction of τ to $F_r \times \mathbb{Z}$ is non-trivial on the factor \mathbb{Z} , so its kernel is free of finite rank (projecting onto F_r identifies the kernel with a finite index subgroup of F_r). It follows that $\ker \tau$ is finitely generated and virtually free, hence equal to some F_n since G is torsion-free. This shows that $G = M_\Phi$ for some $\Phi \in \text{Out}(F_n)$. View the center Z as a subgroup of M_Φ and write a generator as $t^k g$ with $k \in \mathbb{Z}$ and $g \in F_n$. One has $k \neq 0$, and Φ^k is the identity because Z is central.

Conversely, suppose that $\Phi \in \text{Out}(F_n)$ has finite order. We first show that M_Φ is a GBS group. The following simple argument is due to V.

Guirardel. By [12, 22, 34], Φ is induced by an automorphism of a finite graph Λ with fundamental group isomorphic to F_n . The group M_Φ is the fundamental group of a 2-complex made of circles (associated to vertices of Λ) and annuli (associated to edges), so is a GBS group. We now show that the center of M_Φ is non-trivial.

Let $\alpha \in \text{Aut}(F_n)$ be a representative of Φ . There exists $k \geq 1$ such that α^k is conjugation by some $g \in F_n$. If g is trivial, t^k is central. If not, we consider the subgroup of M_Φ generated by F_n and t^k . It has finite index and non-trivial center (generated by $t^k g^{-1}$). It follows that the modular map Δ_{M_Φ} (see Section 2) has finite image, so M_Φ has a normal infinite cyclic subgroup Z (containing $t^k g^{-1}$). This subgroup is not contained in F_n , so is in fact central: otherwise any non-trivial $z \in Z$ would be conjugate to z^{-1} , a contradiction since z maps non-trivially under the natural projection from M_Φ to $\mathbb{Z} = \langle t \rangle$. \square

Example 4.2. — $\text{Out}(F_2)$ contains an element Φ of order 6. It is represented by an automorphism f of the theta graph (it has 2 vertices joined by 3 edges, and f is a symmetry of order 6). One considers midpoints of edges as vertices, so that there is one orbit of edges (with period 6) and two orbits of vertices (one with period 3, one with period 2). It follows that $M_\Phi = \langle a, b \mid a^3 = b^2 \rangle$.

Remark. — It may be shown that M_Φ is not a GBS group if $\Phi \in \text{Out}(F_n)$ has infinite order: M_Φ is residually finite (as a semi-direct product of finitely generated residually finite groups), so cannot be a GBS group unless it is virtually $F_r \times \mathbb{Z}$ (see [26] for a proof); but then Φ has finite order (argue as in the proof of the proposition).

COROLLARY 4.3. — *Let $\alpha \in \text{Aut}(F_n)$ and $k \geq 2$. If α^k is conjugation by some $g \in F_n$, then $\alpha(g) = g$.*

Proof. — We have seen in the proof of Proposition 4.1 that $t^k g^{-1}$ is central in M_Φ . In particular it commutes with t , so t commutes with g . This means $\alpha(g) = g$. \square

Remark. — This result is not specific to $G = F_n$. It holds whenever the outer automorphism defined by α may be represented by a homeomorphism of finite order of a space X with $\pi_1(X) = G$.

COROLLARY 4.4. — *One may compute the rank of the mapping torus of a finite order automorphism of a free group F_n : there is an algorithm which, given $\Phi \in \text{Out}(F_n)$ of finite order, computes the rank of $M_\Phi = F_n \rtimes_\Phi \mathbb{Z}$.*

Proof. — Given Φ of finite order, we know that M_Φ is a GBS group. We may find a standard GBS presentation of M_Φ by applying Tietze transformations to the presentation as a semidirect product, or by arguing as in Example 4.2. We then apply Theorem 3.2. \square

Remark. — Given $\Phi \in \text{Out}(F_n)$ and $\Psi \in \text{Out}(F_q)$, both of finite order, one may decide whether M_Φ and M_Ψ are isomorphic [19].

5. The rank of finite index subgroups

This section is devoted to the proof of the following result:

THEOREM 5.1. — *If G is a GBS group, and $\overline{G} \subset G$ has finite index, then $\text{rk}(\overline{G}) \geq \text{rk}(G)$.*

5.1. A reduction

The first step in the proof is to reduce the theorem to a result about graphs (Proposition 5.4).

We say that a map between graphs is a *morphism* if it sends vertex to vertex and edge to edge.

We represent G by a labelled graph Γ . In this section we will assume that Γ is reduced (if $\lambda_e = \pm 1$, then e is a loop). The group \overline{G} acts on the Bass–Serre tree T of Γ and this yields a graph of groups $\overline{\Gamma} = T/\overline{G}$, with a morphism $\pi : \overline{\Gamma} \rightarrow \Gamma$. We describe $\overline{\Gamma}$ and π using topology. More conceptually, π (and the admissible maps of Section 6) are coverings of graphs of groups in the sense of Bass [1]; our point of view is closer to that of Scott–Wall [32].

It is standard to associate a foliated 2–complex Θ to Γ . One associates a circle C_v to each vertex v , an annulus $A_\varepsilon = [0, 1] \times S^1$ foliated by circles $\{*\} \times S^1$ to each non-oriented edge ε , and boundaries of annuli are glued to circles by maps whose degree (positive or negative) is given by the labels of Γ . One recovers Γ from Θ by collapsing each circle to a point. The fundamental group of Θ is G , and \overline{G} defines a finite covering $\rho : \overline{\Theta} \rightarrow \Theta$ whose degree is the index of \overline{G} . The complex $\overline{\Theta}$ is also made of circles and annuli, $\overline{\Gamma}$ is the corresponding graph, and π is induced by ρ .

DEFINITION 5.2 (multiplicity $m_x, m_{\overline{x}}$). — *A point x of $\overline{\Gamma}$ corresponds to a circle of $\overline{\Theta}$, and the restriction of ρ to this circle is a covering map*

whose degree we call the multiplicity m_x of x : we thus associate a positive integer m_x to each point x of $\bar{\Gamma}$. All points belonging to the interior of a given edge have the same multiplicity, so we also define the multiplicity $m_{\bar{e}}$ of an edge \bar{e} of $\bar{\Gamma}$. Given a point u in Γ , the sum $\sum_{x \in \pi^{-1}(u)} m_x$ is constant, equal to the index of \bar{G} ; we call it the total multiplicity of π .

Algebraically, multiplicities are indices $[G_p : G_p \cap \bar{G}]$, with G_p the stabilizer of a point $p \in T$.

LEMMA 5.3. — *The morphism $\pi : \bar{\Gamma} \rightarrow \Gamma$ satisfies the following condition (*) :*

Given an edge e of Γ , with origin v and label λ_e near v , and $x \in \pi^{-1}(v)$, define $k_{x,e}$ as the gcd $m_x \wedge \lambda_e$. Then (see Figure 5.1) there are $k_{x,e}$ edges of $\bar{\Gamma}$ with origin x mapping to e ; they each have multiplicity $m_x/k_{x,e}$, and their label near x is $\lambda_e/k_{x,e}$.




Figure 5.1. condition (*)

Proof. — A neighborhood of C_v in the annulus A_e corresponding to e has fundamental group \mathbb{Z} , and one simply studies its preimage by ρ . The group G_x carried by x in $\bar{\Gamma}$ has index m_x in G_v , and G_e has index λ_e in G_v . Their intersection has index $lcm(m_x, \lambda_e) = m_x \lambda_e / k_{x,e}$ in G_v , index $m_x / k_{x,e}$ in G_e , and index $\lambda_e / k_{x,e}$ in G_x . □

Remark. — One may also prove this lemma algebraically, by considering intersections of \bar{G} with vertex and edge stabilizers of T .

PROPOSITION 5.4. — *Let Γ and $\bar{\Gamma}$ be connected labelled graphs. Assume that Γ is reduced (if $\lambda_e = \pm 1$, then Γ is a loop), and a positive multiplicity is assigned to each vertex and edge of $\bar{\Gamma}$. If there exists a morphism $\pi : \bar{\Gamma} \rightarrow \Gamma$ satisfying $(*)$, then $\beta(\bar{\Gamma}) + \mu(\bar{\Gamma}) \geq \beta(\Gamma) + \mu(\Gamma)$.*

Recall that β is the first Betti number, and μ is the plateaunic number (Definition 3.1). It follows from Theorem 3.2 that this proposition implies Theorem 5.1. The remainder of this section is devoted to its proof.

5.2. The main idea

Let π be as in the proposition. It is open: the image of a neighborhood of a vertex x is a neighborhood of $\pi(x)$. If $\Gamma' \subset \Gamma$ is a connected subgraph, every component of $\pi^{-1}(\Gamma')$ maps onto Γ' , and the restriction satisfies $(*)$. Note that no label near a terminal vertex of $\bar{\Gamma}$ is ± 1 ($\bar{\Gamma}$ is minimal, but not necessarily reduced).

In the situation of Lemma 5.3, let \bar{e} be an edge of $\bar{\Gamma}$ with origin x mapping onto e . We call \bar{e} a lift of e (at x).

Let p be a prime.

If λ_e is not divisible by p (in particular if e is contained in a p -plateau), the label $\lambda_{\bar{e}}$ of \bar{e} near x is not divisible by p . Moreover $k_{x,e}$ is not divisible by p , so m_x and $m_{\bar{e}}$ are both divisible by p or both non divisible by p .

If λ_e is divisible by p , note that $k_{x,e}$ is divisible by p if and only if m_x is. If m_x is not divisible by p , then $\lambda_{\bar{e}}$ is divisible by p .

It is easy to check that $\beta(\bar{\Gamma}) \geq \beta(\Gamma)$ (see Lemma 5.6). Suppose that, for each plateau $P \subset \Gamma$, some component of $\pi^{-1}(P)$ is a plateau. Then $\mu(\bar{\Gamma}) \geq \mu(\Gamma)$ and the result is clear. We therefore consider a p -plateau P , and we study $\pi^{-1}(P)$. We may assume $P \neq \Gamma$.

If there exists a vertex $x \in \pi^{-1}(P)$ with m_x not divisible by p , it follows from previous observations that the same holds for all points in the component \bar{P} of $\pi^{-1}(P)$ containing x , and \bar{P} is a p -plateau.

We therefore assume that m_x is divisible by p for every vertex $x \in \pi^{-1}(P)$. Since $P \neq \Gamma$, there is an edge e with origin $v \in P$ which is not contained in P . If $x \in \pi^{-1}(v)$, the number of lifts of e at x is divisible by p since both m_x and λ_e are. In particular, denoting by d_v the valence of a vertex, we have $d_x \geq d_v + p - 1$.

We conclude that, if P is a p -plateau such that no component of $\pi^{-1}(P)$ is a p -plateau, there exist $v \in P$ and $x \in \pi^{-1}(v)$ such that the valence of x satisfies $d_x \geq d_v + p - 1$.

The main idea of the proof now is the following: the existence of such P 's may cause $\mu(\bar{\Gamma})$ to be smaller than $\mu(\Gamma)$, but the increase in valence will force $\beta(\bar{\Gamma})$ to be larger than $\beta(\Gamma)$, so that $\beta(\bar{\Gamma}) + \mu(\bar{\Gamma}) \geq \beta(\Gamma) + \mu(\Gamma)$ does hold (recall the formula $\beta(\Gamma) = 1 + \frac{1}{2} \sum_v (d_v - 2)$ for the first Betti number of a connected graph Γ).

If $p > 2$ for all P 's as above, we have $d_x \geq d_v + 2$ and the proof is not too hard, as we shall now explain. On the other hand, if $p = 2$, we only have $d_x \geq d_v + 1$ and this makes the proof much more complicated.

5.3. The simple case

Notation.

- β : first Betti number (we write $\bar{\beta}$, β' for the first Betti number of $\bar{\Gamma}$, Γ' , etc.)
- μ : plateaunic number
- t : number of terminal vertices
- d_v : valence of a vertex
- V : vertex set
- E_v : oriented edges with origin v
- $\bar{\Gamma}_v$: preimage $\pi^{-1}(v)$
- $m_x, m_{\bar{e}}$: multiplicity of a vertex, an edge

The following lemma is left to the reader.

LEMMA 5.5. — *Let Γ be a finite connected graph, with vertex set V . Then:*

$$\begin{aligned} \beta &= 1 + \sum_{v \in V} \left(\frac{d_v}{2} - 1 \right) \\ \beta + t &= 1 + \sum_{v \in V} \left\lfloor \frac{d_v}{2} - 1 \right\rfloor \\ \beta + \frac{t}{2} &= 1 + \sum_{v \in V} \max\left(\frac{d_v}{2} - 1, 0 \right). \end{aligned}$$

□

LEMMA 5.6. — Let $\pi : \bar{\Gamma} \rightarrow \Gamma$ be an open morphism between finite connected graphs. Then:

$$\begin{aligned} \beta &\leq \bar{\beta} \\ \beta + \frac{t}{2} &\leq \bar{\beta} + \frac{\bar{t}}{2} \end{aligned}$$

(β, t refer to Γ , and $\bar{\beta}, \bar{t}$ refer to $\bar{\Gamma}$).

Proof. — Consider any loop γ in Γ . Since π is open, there exists a loop in $\bar{\Gamma}$ projecting onto a power of γ . This proves the first inequality.

For the other inequality, write

$$\max\left(\frac{d_v}{2} - 1, 0\right) \leq \sum_{x \in \bar{\Gamma}_v} \max\left(\frac{d_x}{2} - 1, 0\right)$$

for $v \in V$ and use the third equality of the previous lemma. □

Let π be as in Proposition 5.4.

DEFINITION 5.7 (Δ_v). — For $v \in V$, define

$$\Delta_v = \sum_{x \in \bar{\Gamma}_v} \left| \frac{d_x}{2} - 1 \right| - \left| \frac{d_v}{2} - 1 \right|.$$

This is non-negative, unless v is a terminal vertex whose preimages all have valence 2 (in this case $\Delta_v = -\frac{1}{2}$). Note that such a v is a 2-plateau, since the label near v must be even. Thus Δ_v is always non-negative if Γ contains no proper 2-plateau.

Also note that the second equality of Lemma 5.5 yields

$$\sum_{v \in V} \Delta_v = (\bar{\beta} + \bar{t}) - (\beta + t).$$

Remark. — Since Δ_v may be negative, it is not always true that $\beta + t \leq \bar{\beta} + \bar{t}$ (for instance, $\bar{\Gamma}$ may be a circle subdivided into two intervals, mapping onto an interval). See Lemma 5.20 for a complete discussion.

DEFINITION 5.8 (boundary, frontier point, interior plateau). —

The boundary ∂P of a p -plateau is the set of oriented edges e with origin v in P and terminal point not in P ; the vertex v is a frontier point of P . Note that the label λ_e of the edge e is divisible by p .

A plateau P is interior if P is not the whole graph and P contains no terminal vertex.

Every terminal vertex of Γ (or $\bar{\Gamma}$) is a plateau, so μ is the sum of t and the minimal cardinality of a set meeting every interior plateau (unless $t = 0$ and Γ is the only plateau).

DEFINITION 5.9 (totally unfolded). — A plateau $P \subset \Gamma$ is p -totally unfolded if P is a p -plateau and, given any edge $e \in \partial P$ with origin $v \in P$, and $x \in \bar{\Gamma}_v$, the number of lifts of e at x is divisible by p . A plateau is totally unfolded if it is p -totally unfolded for some p .

It may happen that P is both a p -plateau and a p' -plateau, totally unfolded as a p -plateau but not as a p' -plateau. We consider such a P as totally unfolded.

The arguments in the previous subsection show:

LEMMA 5.10. — If no component of $\pi^{-1}(P)$ is a plateau, then P is totally unfolded. □

DEFINITION 5.11 (minimal plateau, c). — A minimal plateau is a plateau $P \subset \Gamma$ which is interior, totally unfolded, and minimal for these properties. If a minimal plateau is p -totally unfolded, we say that it is a minimal p -plateau.

Let c be the minimal cardinality of a set of vertices meeting every minimal plateau.

LEMMA 5.12. — If Γ contains no proper 2-plateau, one has $\beta + t + c \leq \bar{\beta} + \bar{t}$.

Proof. — We have seen that $\sum_{v \in V} \Delta_v = (\bar{\beta} + \bar{t}) - (\beta + t)$, so it suffices to show $\sum_{v \in V} \Delta_v \geq c$. Since Δ_v is non-negative at terminal vertices (because there is no 2-plateau), we may restrict the sum to non-terminal vertices.

For each minimal plateau P_i , we fix an edge $e_i \in \partial P_i$ with origin $v_i \in P_i$, and an odd prime p_i such that P_i is p_i -totally unfolded.

Given a non-terminal vertex v , consider the minimal plateaux P_i such that $v_i = v$. Let n_v be their number (possibly 0). Note that the associated p_i 's are distinct. We shall show $\Delta_v \geq n_v$. Assuming this, $\sum_{v \in V} \Delta_v$ is bounded below by the number of minimal plateaux, hence by c as required.

Fix $x \in \bar{\Gamma}_v$. Since v is not terminal, the terms $\frac{d_x}{2} - 1$ and $\frac{d_v}{2} - 1$ are non-negative and

$$2\Delta_v = d_x - d_v + \sum_{y \in \bar{\Gamma}_v \setminus \{x\}} (d_y - 2) \geq d_x - d_v.$$

Given an edge e with origin v , consider the plateaux P_j such that $e_j = e$. Let n_e be their number (possibly 0). The number $d_{x,e}$ of lifts of e at x is divisible by the product of the corresponding p_j 's, so is bounded below by $2n_e + 1$ (because the product of n odd prime numbers is at least $2n + 1$).

Summing over all edges e with origin v we get

$$d_x = \sum_{e \in E_v} d_{x,e} \geq 2 \sum_{e \in E_v} n_e + d_v = 2n_v + d_v,$$

so $2\Delta_v \geq d_x - d_v \geq 2n_v$ as required. □

COROLLARY 5.13. — *Proposition 5.4 is true if Γ contains no proper 2-plateau.*

Proof. — One has $\bar{\mu} = \bar{t} + |A|$, where $A \subset \bar{\Gamma}$ is a subset of minimal cardinality such that every plateau of $\bar{\Gamma}$ meets A or contains a terminal vertex. If $C \subset V$ is a set of minimal cardinality c meeting every minimal plateau of Γ , the union of $C \cup \pi(A)$ with the terminal vertices meets every plateau of Γ (totally unfolded or not) by Lemma 5.10, so $\mu \leq c + |A| + t$. Using $\beta + t + c \leq \bar{\beta} + \bar{t}$, we get

$$\beta + \mu \leq \beta + c + |A| + t \leq \bar{\beta} + \bar{t} + |A| = \bar{\beta} + \bar{\mu}.$$

□

Proposition 5.4 thus follows fairly directly from the inequality of Lemma 5.12. Unfortunately, that inequality does not always hold, and in the general case we will have to deduce the proposition from the weaker lemma 5.24.

5.4. The general case

Unless mentioned otherwise, π is as in Proposition 5.4.

The assumption that there is no proper 2-plateau was used in the previous subsection to ensure $\Delta_v \geq 0$ and $\prod_{j=1}^{n_e} p_j \geq 2n_e + 1$. This motivates the following definitions.

DEFINITION 5.14 (bad vertex, good vertex, V_g). — *A vertex v of Γ is bad if $\Delta_v < 0$, i.e. if v is terminal and all its preimages in $\bar{\Gamma}$ have valence 2. The label near a bad vertex is even.*

Let V_g be the set of good (i.e. not bad) vertices of Γ .

DEFINITION 5.15 (bad plateau). — *A minimal plateau $P \subset \Gamma$ is bad if it is 2-totally unfolded, its boundary consists of a single edge e , and e has exactly 2 lifts in $\bar{\Gamma}$ (so the origin of e has a single preimage since P is totally unfolded).*

DEFINITION 5.16 (t_g, c_g). — *Let t_g be the number of good terminal vertices, and c_g the number of good minimal plateaux.*

LEMMA 5.17. — *One has $\beta + t_g + c_g \leq \bar{\beta} + \bar{t}$.*

Proof. — We argue as in the proof of Lemma 5.12. Since $\Delta_v = -\frac{1}{2}$ if v is bad, we have $\sum_{v \in V \setminus V_g} \Delta_v = -\frac{1}{2}(t - t_g)$ and

$$\sum_{v \in V_g} \Delta_v = \sum_{v \in V} \Delta_v + \frac{1}{2}(t - t_g) = (\bar{\beta} + \bar{t}) - (\beta + t) + \frac{1}{2}(t - t_g) \leq (\bar{\beta} + \bar{t}) - (\beta + t_g),$$

and we reduce to showing $\sum_{v \in V_g} \Delta_v \geq c_g$. We define p_i, e_i, v_i, n_v, n_e as above (with $p_i = 2$ now allowed), except that we restrict to good plateaux, and we try to prove $\Delta_v \geq n_v$ for $v \in V_g$ (we will not succeed in all cases).

In the proof of Lemma 5.12, we deduced $\Delta_v \geq n_v$ from the inequalities

$$\begin{aligned} 2\Delta_v &\geq d_x - d_v \\ d_x = \sum_{e \in E_v} d_{x,e} &\geq \sum_{e \in E_v} \left(\prod_{j=1}^{n_e} p_j \right) \\ \prod_{j=1}^{n_e} p_j &\geq 2n_e + 1. \end{aligned}$$

The first two inequalities are still true, but the third inequality may be wrong. This happens precisely when $n_e = 1$ and $p_1 = 2$ (recall that the p_j 's are distinct), so we have $\Delta_v \geq n_v$ unless v is the chosen frontier point v_i of a 2-totally unfolded plateau P_i . In this case we have only proved $2\Delta_v \geq 2n_v - 1$ (this does not imply $\Delta_v \geq n_v$ because Δ_v is not necessarily an integer). To conclude, we shall use goodness of P_i to find an edge in ∂P_i with a lift not taken into account in the previous estimates.

If e has more than 2 lifts in $\bar{\Gamma}$, or if there are 2 edges with origin v in ∂P_i , we still have $\Delta_v \geq n_v$ because one of the first two inequalities displayed above is strict. Otherwise, since P_i is good, it has another frontier point w_i . Every edge with origin w_i in ∂P_i has at least 2 lifts, and we have $\Delta_v + \Delta_{w_i} \geq n_v + n_{w_i}$ even if $\Delta_v < n_v$. The w_i 's are distinct because 2-plateaux are equal or disjoint, so we get $\sum_{v \in V_g} \Delta_v \geq \sum_{v \in V_g} n_v = c_g$. \square

In the following definitions, and in Lemma 5.20, the graphs do not have to be labelled, and π is just an open morphism between finite connected graphs.

DEFINITION 5.18 (accordion). — $\pi : \bar{\Gamma} \rightarrow \Gamma$ is an accordion if $\bar{\Gamma}$ is homeomorphic to a circle and Γ to an interval (see Figure 5.2). Thus the terminal vertices of Γ are bad, and $\bar{\Gamma}$ is the union of $2n$ intervals (possibly subdivided), each mapped homeomorphically to Γ . We call n the size of the accordion.




Figure 5.2. an accordion of size 2

DEFINITION 5.19 (branched covering). — $\pi : \bar{\Gamma} \rightarrow \Gamma$ is a branched 2-covering of a tree (or simply a branched covering) if Γ is a tree, and for $u \in \Gamma$ the preimage $\bar{\Gamma}_u$ consists of a single point if u is a terminal vertex of Γ , of 2 points otherwise (see Figure 5.3). In particular, all terminal vertices of Γ are bad.

An accordion of size 1 is a branched covering.

The following topological lemma may be seen as a warm-up for the proof of Lemma 5.24.

LEMMA 5.20. — Let $\pi : \bar{\Gamma} \rightarrow \Gamma$ be an open morphism between finite connected graphs. One has $\beta + t = \bar{\beta} + \bar{t} + 1$ if π is an accordion or a branched 2-covering of a tree, $\beta + t \leq \bar{\beta} + \bar{t}$ otherwise.

Proof. — If π is an accordion, one has $\beta + t = 0 + 2 = 2$ and $\bar{\beta} + \bar{t} = 1 + 0 = 1$. If it is a branched 2-covering of a tree, then $\beta = \bar{t} = 0$ and $\bar{\beta} = t - 1$. We now consider the general case.

As a preliminary observation, note that a connected graph satisfies $\beta + t \geq 2$ unless it is homeomorphic to a point or a circle. If $\bar{\Gamma}$ is a circle, Γ is a circle or an interval. Also note that the lemma is true if Γ has only one edge. We will argue by induction on the number of edges of Γ .

Let v be a bad vertex of Γ (the result follows from Lemma 5.17 if there is none). We define a vertex w as follows (see Figure 5.4).

Let v_1 be the vertex adjacent to the terminal vertex v . We let $w = v_1$ if the valence of v_1 is different from 2, or if the valence is 2 and every $x \in \bar{\Gamma}_{v_1}$ is the origin of at least 2 edges not mapping onto v_1v . Otherwise,




Figure 5.3. a branched 2-covering of a tree




Figure 5.4. the vertex w (w cannot be v_1 because of x_0)

we consider the vertex $v_2 \neq v$ adjacent to v_1 and we iterate. We obtain a vertex $w = v_q$ such that w has valence $\neq 2$, or w has valence 2 and every $x \in \bar{\Gamma}_w$ is the origin of at least 2 edges not mapping onto wv_{q-1} .

If w is terminal, then Γ is a segment, $\beta + t = 2$, and $\bar{\beta} + \bar{t} \geq 2$ since π is not an accordion, so the lemma is proved.

Otherwise, we consider the graph Γ' obtained from Γ by removing the segment vw (but not the vertex w). No vertex of $\bar{\Gamma}_w$ is terminal in $\bar{\Gamma}' = \pi^{-1}(\Gamma')$. With obvious notations, one has $\beta' = \beta$ and $t' \geq t - 1$, with equality if and only if w has valence ≥ 3 in Γ . We distinguish two cases.

- First case: $\bar{\Gamma}'$ is connected. We then consider $\bar{\beta}'$ and \bar{t}' . The second equality of Lemma 5.5 implies $\bar{\beta}' + \bar{t}' < \bar{\beta} + \bar{t}$, because any $x \in \bar{\Gamma}_w$ has valence ≥ 2 in $\bar{\Gamma}'$, so $|\frac{d_x}{2} - 1|$ is larger when computed in $\bar{\Gamma}$ than when computed in $\bar{\Gamma}'$.

The restriction $\pi' : \bar{\Gamma}' \rightarrow \Gamma'$ of π is open, so we may use induction and write

$$\beta + t \leq \beta' + t' + 1 \leq \bar{\beta}' + \bar{t}' + 2 \leq \bar{\beta} + \bar{t} + 1.$$

The lemma is proved if one of the inequalities is strict. If not, w has valence ≥ 3 in Γ , the map π' is an accordion or a branched covering, and $\bar{\beta}' + \bar{t}' = \bar{\beta} + \bar{t} - 1$.

This equality drastically limits the possibilities for the preimage of the segment vw (see Figure 5.5): either w has two preimages x, x' , and $\pi^{-1}(vw)$ is an arc joining them, or w has a single preimage x , and $\pi^{-1}(vw)$ is a circle containing x or a lollipop (an arc xy with a circle attached to y).




Figure 5.5. possible preimages of vw

We complete the proof by showing that π must be a branched covering. Since π' is an accordion or a branched covering, w has at least 2 preimages because it is not terminal in Γ' . It follows that w has two preimages, and $\pi^{-1}(vw)$ is an arc joining them, so π is a branched covering of the tree $\Gamma = \Gamma' \cup vw$.

• Second case: $\bar{\Gamma}'$ has several components $\bar{\Gamma}'_i$. By induction, $\bar{\beta}'_i + \bar{t}'_i \geq \beta' + t' - 1$. Also, $\bar{\beta} + \bar{t} \geq \sum_i (\bar{\beta}'_i + \bar{t}'_i)$ since no vertex in $\bar{\Gamma}_w$ is terminal in $\bar{\Gamma}'$. We then write

$$\bar{\beta} + \bar{t} \geq \sum_i (\bar{\beta}'_i + \bar{t}'_i) \geq 2(\beta' + t' - 1) \geq 2(\beta + t - 2) = \beta + t + (\beta + t - 4).$$

We are done if $\beta + t \geq 4$. Since Γ is not a circle, one cannot have $\beta + t = 1$. If $\beta + t = 2$, one has $\bar{\beta} + \bar{t} \geq 2$ since $\bar{\Gamma}$ is not a circle. There remains the possibility that $\beta + t = 3$ and $\bar{\beta} + \bar{t} = 2$. If this happens, one has $\sum_i (\bar{\beta}'_i + \bar{t}'_i) = 2$, so $\bar{\Gamma}'$ consists of two disjoint circles and $\bar{\Gamma}$ is their union with an arc. It follows that Γ is homeomorphic to an arc or a lollipop, so $\beta + t = 2$. □

LEMMA 5.21. — *Let $\pi : \bar{\Gamma} \rightarrow \Gamma$ be as in Proposition 5.4. If P is a 2-plateau, the parity of the cardinality of $\pi^{-1}(u)$ is the same for all $u \in P$. If P is a 2-totally unfolded plateau, and $P \neq \Gamma$, all points in $\pi^{-1}(P)$ have even multiplicity; in particular, the total multiplicity of π is even.*

Proof. — This follows easily from condition (*). With the notations of Lemma 5.3, if $e \subset P$, then λ_e is odd, so $k_{x,e}$ is odd. This proves the first assertion.

If P is 2-totally unfolded, points of $\bar{\Gamma}$ mapping to a frontier point of P have even multiplicity (because m_x is even if $k_{x,e}$ is), and this propagates to all points of $\pi^{-1}(P)$ since all numbers $k_{x,e}$ associated to edges in P are odd. □

DEFINITION 5.22 (generalized branched covering). — *We say that π is a generalized branched 2-covering of a tree if there exist minimal 2-plateaux P_i such that:*

- (1) *the graph obtained from Γ by collapsing each P_i to a point is a tree;*
- (2) *if $u \in \Gamma$ is not a terminal vertex and does not belong to $\cup_i P_i$, its preimage $\bar{\Gamma}_u$ consists of 2 points;*
- (3) *if v is a terminal vertex of Γ , or a frontier point of some P_i , its preimage is a single point (in particular, terminal vertices are bad).*

The P_i 's are the branching plateaux of π . The union of the P_i 's and the terminal vertices is the branching locus.

See a simple example on Figure 1.4. The ellipse is the unique branching plateau.

When there is no P_i , we recover the definition of a branched covering (Definition 5.19).

Recall (Definition 5.11) that a minimal 2-plateau is interior, 2-totally unfolded, and minimal among all interior totally unfolded plateaux (not necessarily 2-plateaux). Any two 2-plateaux are equal or disjoint, so the branching plateaux are disjoint. It follows from Lemma 5.21 that all points in a branching plateau have an odd number of preimages, so the branching locus is the set of points with an odd number of preimages.

DEFINITION 5.23 (exceptional). — π is exceptional if it is an accordion or a generalized branched 2-covering of a tree.

LEMMA 5.24. — If $\pi : \bar{\Gamma} \rightarrow \Gamma$ is as in Proposition 5.4, then:

$$\beta + t + c \leq \bar{\beta} + \bar{t} \quad \text{if } \pi \text{ is not exceptional}$$

$$\beta + t + c \leq \bar{\beta} + \bar{t} + 1 \quad \text{if } \pi \text{ is exceptional.}$$

Recall that c is the minimal cardinality of a set of vertices meeting every minimal plateau.

Proof. — We elaborate on the proof of Lemma 5.20. If Γ' is a connected subgraph, the restriction π' of π to the preimage $\bar{\Gamma}'$ satisfies (*), and Γ' is reduced, so we can argue inductively, defining minimal plateaux of Γ' , and a number c' , as in Subsection 5.3 (Definition 5.11).

The lemma follows from Lemma 5.20 if $c = 0$, in particular if Γ has only one edge or if π is an accordion. It follows from Lemma 5.17 if there is no bad vertex or bad plateau.

- First suppose that there is a bad vertex v . We define $w, \Gamma', \bar{\Gamma}'$ as in the proof of Lemma 5.20.

It follows from the definition of w that any interior totally unfolded plateau which meets the segment vw must contain w . In particular $c = 0$ if w is terminal, so the result is true in this case. We therefore assume that w is not terminal (in particular, $\bar{\Gamma}$ is not a circle).

The intersection of a minimal plateau of Γ with Γ' is totally unfolded, but it is not necessarily interior: it may be equal to Γ' , and w may be a terminal vertex of Γ' (if it has valence 2 in Γ).

If $A \subset \Gamma'$ meets every minimal plateau of Γ' , then $A \cup \{w\}$ meets every minimal plateau of Γ , so $c \leq c' + 1$. Equality is possible only if Γ' is contained in a minimal plateau of Γ , or w has valence 2 in Γ .

1) First assume that no minimal plateau of Γ contains Γ' . Then

$$\beta + t + c \leq \beta' + t' + c' + 1$$

since $t \leq t' + 1$ with equality only if w has valence > 2 in Γ (see the proof of Lemma 5.20), and $c \leq c' + 1$ with equality only if w has valence 2 (and of

course $\beta = \beta'$). We now argue as in the proof of Lemma 5.20, distinguishing two subcases.

1a) If $\bar{\Gamma}'$ is connected, we have

$$\beta + t + c \leq \beta' + t' + c' + 1 \leq \bar{\beta}' + \bar{t}' + 1 \leq \bar{\beta} + \bar{t}$$

if π' is not exceptional (as above, the last inequality follows from Lemma 5.5). If π' is exceptional, we only get $\beta + t + c \leq \bar{\beta} + \bar{t} + 1$. We assume that equality holds, and we show that π is exceptional or $c = 0$.

We have $\bar{\beta}' + \bar{t}' = \bar{\beta} + \bar{t} - 1$, so $\bar{\Gamma} \setminus \bar{\Gamma}'$ is very restricted (as pointed out in the proof of Lemma 5.20, $\pi^{-1}(vw)$ is an arc, a circle, or a lollipop). In particular, w has at most 2 preimages; if π' is an accordion, it has size (in the sense of Definition 5.18) at most 2.

If w has 2 preimages, then π is exceptional with the same P_i 's as π' if π' is a generalized branched covering, while $c = 0$ if π' is an accordion of size 2. If w has 1 preimage, let w_1w be the maximal subsegment of vw consisting of points with a single preimage ($w_1 = \pi(y)$ if y is as on Figure 5.5). Points between v and w_1 have 2 preimages ($w_1 \neq v$ since v is a bad vertex).

Define the branching locus of π as the union of that of π' with w_1w and $\{v\}$. The component containing w is a 2-totally unfolded plateau of Γ because, if e is an oriented edge with origin in w_1w , its label λ_e has the same parity as the number of lifts of e (the total multiplicity of π is even by Lemma 5.21). It is minimal because the branching plateaux of π' are. The conditions of Definition 5.22 are satisfied, so π is exceptional.

1b) If $\bar{\Gamma}'$ has several components $\bar{\Gamma}'_i$, then

$$\begin{aligned} \bar{\beta} + \bar{t} &\geq \sum_i (\bar{\beta}'_i + \bar{t}'_i) \\ &\geq 2(\beta' + t' + c' - 1) \\ &\geq 2(\beta + t + c - 2) \\ &= \beta + t + c + (\beta + t + c - 4). \end{aligned}$$

As in the proof of Lemma 5.20, we just have to rule out the possibility that $\beta + t + c = 3$ and $\bar{\beta} + \bar{t} = 2$. If this happens, $\bar{\Gamma}'$ is the union of two disjoint circles and an arc, mapping onto an arc or a lollipop, and $c = 0$.

2) If some minimal plateau P of Γ contains Γ' , we have $t = 1$ (the only terminal vertex of Γ is v), $c = 1$ (every minimal plateau contains w), and $\bar{t} = \bar{t}' = 0$.

2a) If $\bar{\Gamma}'$ is connected, we have

$$\beta + c + t = \beta' + 2 \leq \bar{\beta}' + 2 \leq \bar{\beta} + 1.$$

We assume that both inequalities are equalities, and we show that π is exceptional.

The equality $\beta' + 2 = \bar{\beta} + 1$ implies that one gets $\bar{\Gamma}$ from $\bar{\Gamma}'$ by attaching an arc, a circle, or a lollipop. It cannot be an arc because of the totally unfolded interior plateau P containing Γ' . It follows that w has a single preimage.

If π' is exceptional, one shows that π is exceptional as in case 1a. If not, we have $\beta' + t' \leq \bar{\beta}' + \bar{t}' = \bar{\beta}'$ by Lemma 5.20, so $t' = 0$ since we assume $\beta' = \bar{\beta}'$. Thus Γ' and $\bar{\Gamma}'$ are graphs with no terminal vertices, and $\beta' = \bar{\beta}'$ implies that π' is an isomorphism by Lemma 5.5. It follows that π is exceptional, with branching locus $\Gamma' \cup w_1w \cup \{v\}$.

2b) If $\bar{\Gamma}'$ is not connected, we have

$$\beta + c + t - 1 = \beta' + 1 \leq 2\beta' \leq \bar{\beta}'_1 + \bar{t}'_1 + \bar{\beta}'_2 + \bar{t}'_2 \leq \bar{\beta} + \bar{t}.$$

The first inequality holds because $\beta' \geq 1$: since Γ' is contained in a minimal plateau of Γ , no vertex other than w may be terminal in Γ' . The second inequality comes from Lemma 5.6. The third inequality was used in the proof of Lemma 5.20; as in the previous subcase, the existence of P implies that $\bar{\Gamma} \setminus \bar{\Gamma}'$ cannot be an open segment, so the inequality is strict. We deduce $\beta + c + t \leq \bar{\beta} + \bar{t}$.

- We now assume that there is a bad plateau P . Let v be its unique frontier point. Let Γ' be the graph obtained from Γ by removing $P \setminus \{v\}$. The preimages $\bar{\Gamma}'$ of Γ' and \bar{P} of P are connected because $\bar{\Gamma}_v$ is a single point.

Denoting by $\beta(P)$ and $\beta(\bar{P})$ the first Betti numbers of P and its preimage \bar{P} , we have $\beta(\bar{P}) \geq \beta(P)$ by Lemma 5.6 and

$$\begin{aligned} t' &= t + 1 \\ \bar{t}' &= \bar{t} \\ \beta' &= \beta - \beta(P) \\ \bar{\beta}' &= \bar{\beta} - \beta(\bar{P}). \end{aligned}$$

By minimality of P , any minimal plateau of Γ either contains v , or is contained in $\Gamma' \setminus \{v\}$ and is a minimal plateau of Γ' . This shows $c \leq c' + 1$. By induction we get

$$\beta + t + c \leq \beta' + \beta(P) + t' - 1 + c' + 1 \leq \beta' + \beta(\bar{P}) + t' + c' \leq \bar{\beta}' + \bar{t}' + \beta(\bar{P}) = \bar{\beta} + \bar{t}$$

if π' is not exceptional. If π' is exceptional, so is π (with P added to the branching locus), and we get $\beta + t + c \leq \bar{\beta} + \bar{t} + 1$ as required. □

We can now conclude.

Proof of Proposition 5.4. — If π is not exceptional, we write

$$\beta + \mu \leq \beta + c + |A| + t \leq \bar{\beta} + \bar{t} + |A| = \bar{\beta} + \bar{\mu}$$

as in the proof of Corollary 5.13, with $A \subset \bar{\Gamma}$ a subset of minimal cardinality such that every plateau of $\bar{\Gamma}$ meets A or contains a terminal vertex.

If π is exceptional, we only have $\beta + c + t \leq \bar{\beta} + \bar{t} + 1$. In this case we get the required inequality $\beta + \mu \leq \bar{\beta} + \bar{\mu}$ by showing $\mu < c + |A| + t$. We fix $x \in A$, noting that A is nonempty because $\bar{\Gamma}$ has no terminal vertices.

If π is an accordion, we claim that $\pi(A \setminus \{x\})$ meets every interior plateau $P \subset \Gamma$. This implies $\mu < |A| + t$, hence the result since $c = 0$. To prove the claim, consider the preimage of P . It consists of $2n$ disjoint arcs, each a plateau of $\bar{\Gamma}$ (with $n \geq 1$ as in Definition 5.18). Each of these plateaux meets A , so one of them contains a point of $A \setminus \{x\}$.

In the case of a generalized branched covering, c is bounded below by k , defined as the number of branching plateaux P_i . We prove $\mu < c + |A| + t$ by constructing a set $C \subset \Gamma$ of cardinality at most $k + |A| + t - 1$ meeting every plateau. There are two cases.

First suppose that $\pi(x)$ belongs to a branching plateau P_{i_0} . For $i \neq i_0$, let v_i be the point of P_i closest to P_{i_0} ; it is well-defined because one gets a tree by collapsing each P_i to a point. Let C consist of the v_i 's, the terminal vertices, and $\pi(A)$. We show that C meets every interior plateau $P \subset \Gamma$.

If P is not totally unfolded, it meets $\pi(A)$ by Lemma 5.10, so assume that it is. If it is contained in some P_i , it equals P_i by minimality of P_i , so contains $\pi(x)$ or v_i . Otherwise, because of condition (2) in Definition 5.22, P contains the closure of a component of $\Gamma \setminus \cup_i P_i$. Since one gets a tree by collapsing each P_i to a point, this closure contains a v_i or a terminal vertex.

Now suppose that $\pi(x)$ belongs to no P_i . For each i we let v_i be the point of P_i closest to $\pi(x)$, and we let C consist of the v_i 's, the terminal vertices, and $\pi(A \setminus \{x\})$.

Arguing as in the previous case, we see that C meets every totally unfolded interior plateau. We just have to check that it meets interior plateaux P contained in the component of $\Gamma \setminus \cup_i P_i$ containing $\pi(x)$. As in the case of an accordion, $\pi^{-1}(P)$ has two components, each a plateau of $\bar{\Gamma}$. At least one of them does not contain x , so P meets $\pi(A \setminus \{x\})$. \square

6. Plateaux and finite index subgroups

Let Γ be a labelled graph representing a GBS group G . We have seen in Subsection 5.1 that any finite index subgroup \overline{G} yields a map $\pi : \overline{\Gamma} \rightarrow \Gamma$ between labelled graphs. We formalize the properties of π .

DEFINITION 6.1 (Admissible map). — *Let $\overline{\Gamma}$ and Γ be labelled graphs (not necessarily connected). An admissible map from $\overline{\Gamma}$ to Γ is a pair (π, m) where $\pi : \overline{\Gamma} \rightarrow \Gamma$ is a morphism, and m assigns a positive multiplicity to each vertex and edge of $\overline{\Gamma}$ so that condition $(*)$ of Lemma 5.3 is satisfied. We usually denote an admissible map simply by π , keeping m implicit.*

If Γ is connected, the total multiplicity of π is $\sum_{x \in \pi^{-1}(u)} m_x$; it does not depend on $u \in \Gamma$.

Recall condition $(*)$: if $\pi(x) = o(e)$, then e has $k_{x,e} = m_x \wedge \lambda_e$ lifts with origin x , each with multiplicity $m_x/k_{x,e}$ and label $\lambda_e/k_{x,e}$.

Remark 6.2. — Suppose $\overline{\Gamma}$ is connected. Given an admissible $\pi : \overline{\Gamma} \rightarrow \Gamma$, there are infinitely many m 's such that (π, m) is admissible. One may show that all of them are multiples of a single m_0 for which no prime divides all edge multiplicities.

If $\pi : \overline{\Gamma} \rightarrow \Gamma$ and $\pi' : \Gamma \rightarrow \Gamma'$ are admissible, so is $\pi' \circ \pi$, with multiplicity function $m(m' \circ \pi)$; this is easy to check, using the formula $\lambda \wedge \mu\mu' = (\lambda \wedge \mu')(\frac{\lambda}{\lambda \wedge \mu'} \wedge \mu)$.

LEMMA 6.3. — *Let Γ be a labelled graph representing a GBS group G .*

- (1) *If \overline{G} is a subgroup of finite index, there is a labelled graph $\overline{\Gamma}$ representing \overline{G} and an admissible map $\pi : \overline{\Gamma} \rightarrow \Gamma$ whose total multiplicity is the index of \overline{G} .*
- (2) *Conversely, any admissible map $\pi : \overline{\Gamma} \rightarrow \Gamma$ with $\overline{\Gamma}$ connected may be obtained from a finite index subgroup \overline{G} as in (1). In particular, the group represented by $\overline{\Gamma}$ embeds into G as a finite index subgroup.*

Proof. — The first assertion was proved in Subsection 5.1. The second assertion may be deduced from [1], but we provide a direct proof. We consider the 2-complex Θ associated to Γ as in Subsection 5.1 and we construct a covering map $\rho : \overline{\Theta} \rightarrow \Theta$ inducing π . The desired group \overline{G} is the fundamental group of $\overline{\Theta}$.

Let $C = \mathbb{R}/\mathbb{Z}$ be the standard circle. We view Θ as the union of circles $C_v = \{v\} \times C$ and annuli $A_\varepsilon = \varepsilon \times C$. The complex $\overline{\Theta}$ is made of circles $C_x = \{x\} \times C$ associated to vertices of $\overline{\Gamma}$ and annuli $A_{\overline{\varepsilon}} = \overline{\varepsilon} \times C$ associated to edges. If x is an endpoint of $\overline{\varepsilon}$, we attach the corresponding boundary

circle of $A_{\bar{\varepsilon}}$ to C_x by the map $(x, \theta) \mapsto (x, \lambda\theta)$, where λ is the label carried by $\bar{\varepsilon}$ near x . To define ρ , we map each circle $\{y\} \times C$, for $y \in \bar{\Gamma}$, to $\{\pi(y)\} \times C$ by $(y, \theta) \mapsto (\pi(y), m_y\theta)$. Condition $(*)$ ensures that this is compatible with the attaching maps of Θ and $\bar{\Theta}$. \square

If $\pi : \Gamma' \rightarrow \Gamma$ is a covering map between finite graphs (in the topological sense), and Γ is labelled, then labelling Γ' so that π is label-preserving, and letting m be constant (coprime with all labels) on each component of Γ' , makes π admissible. Such an admissible map may be characterized as follows.

LEMMA 6.4. — *Let $\pi : \bar{\Gamma} \rightarrow \Gamma$ be an admissible map between labelled graphs. The following conditions are equivalent:*

- (1) π is a covering map (in the topological sense);
- (2) all numbers $k_{x,e} = m_x \wedge \lambda_e$ are equal to 1;
- (3) π preserves labels: $\lambda_{\pi(\bar{\varepsilon})} = \lambda_{\bar{\varepsilon}}$;
- (4) the multiplicity m is constant on every component of $\bar{\Gamma}$.

If these conditions are satisfied, we say that π is a *topological covering*, or that $\bar{\Gamma}$ is a topological covering of Γ . Note that every admissible π is a covering of graphs of groups in the sense of [1].

Proof. — By $(*)$, condition (2) is equivalent to π being a covering, to π being label-preserving, and also to every edge of $\bar{\Gamma}$ having the same multiplicity as its endpoints. \square

PROPOSITION 6.5. — *Given a connected labelled graph Γ , the following conditions are equivalent:*

- every admissible $\pi : \bar{\Gamma} \rightarrow \Gamma$ is a topological covering;
- Γ contains no proper plateau.

Proof.

• We first suppose that π is not a covering, and we construct a proper plateau as in the proof of Proposition 3.9. Using Lemma 6.4, fix a prime p dividing some k_{x_0, e_0} . It divides m_{x_0} , so let p^δ with $\delta \geq 1$ be the maximal power of p dividing the multiplicity of a vertex.

We define a subgraph $\Gamma_1 \subset \Gamma$ as follows. A vertex is in Γ_1 if and only if it has a preimage whose multiplicity is divisible by p^δ ; an edge ε is in Γ_1 if its endpoints are, and none of the two labels carried by ε is divisible by p . Note that $\Gamma_1 \neq \Gamma$ because p divides λ_{e_0} . We show that every component of Γ_1 is a p -plateau.

Consider an edge e with origin $v \in \Gamma_1$ such that p does not divide λ_e . We have to check that its other endpoint w is in Γ_1 , and the label near w is not

divisible by p . Let x be a preimage of v with multiplicity divisible by p^δ , and \bar{e} a lift of e with origin x . Since p does not divide λ_e , the multiplicity of \bar{e} is divisible by p^δ . So is the multiplicity of the terminal point y of \bar{e} . In particular, $w \in \Gamma_1$. By our choice of δ , the multiplicity of y cannot be divisible by $p^{\delta+1}$. This implies that the label near w is not divisible by p .

• Conversely, given a proper p -plateau P , we construct a *branched covering ramified over P* . It is an admissible map $\pi : \bar{\Gamma} \rightarrow \Gamma$ which is not a topological covering: points in P have one preimage, points not in P have p preimages. Compare Figure 1.4, which represents a covering ramified over the union of two 2-plateaux (the terminal vertex and the ellipse).

To construct $\bar{\Gamma}$, we start with $\Gamma \times \{1, \dots, p\}$, and we identify (x, i) with (x, j) whenever $x \in P$. The map from $\bar{\Gamma}$ to Γ is the natural projection. The multiplicity of (the image of) (x, i) is defined as 1 if $x \notin P$, as p if $x \in P$. The label of an edge \bar{e} of $\bar{\Gamma}$ is the same as the label of its projection e , except if the origin of e is in P but e is not contained in P ; in this case λ_e is divisible by p and we define $\lambda_{\bar{e}} = \lambda_e/p$. One checks that (*) is satisfied. □

COROLLARY 6.6. — *Let Γ be a labelled graph representing a GBS group G . Suppose that Γ contains no proper plateau. Then a GBS group \bar{G} is isomorphic to a finite index subgroup of G if and only if it may be represented by a labelled graph $\bar{\Gamma}$ which is a topological covering of Γ .* □

Recall that a group is *large* if some finite index subgroup maps onto the free group F_2 . A GBS group represented by a labelled graph Γ maps onto F_2 if and only if $\beta(\Gamma) \geq 2$ (see [25], p. 483).

THEOREM 6.7. — *Let G be a non-cyclic GBS group. The following are equivalent:*

- (1) G is not large;
- (2) G may be represented by a labelled graph Γ homeomorphic to a circle and containing no proper plateau.

This was proved in [16] for $G = BS(m, n)$ (in this case the absence of a proper plateau is equivalent to $m \wedge n = 1$), and independently by T. Mecham [28] in general. When Γ is a circle, fix an orientation and denote by x_i (resp. y_i) the labels of edges whose orientation agrees (resp. disagrees) with that of the circle; the absence of a proper plateau is equivalent to $\prod x_i \wedge \prod y_i = 1$. When G is not large, finite index subgroups \bar{G} of G are determined, up to isomorphism, by the degree of the topological covering $\bar{\Gamma} \rightarrow \Gamma$ (see [14, 33] for the case of $G = BS(m, n)$ with m and n coprime). It would be interesting to generalize the results of [5, 14, 20] to these groups.

Proof. — We may assume that G is non-elementary. Let G be represented by Γ . It is large if $\beta(\Gamma) \geq 2$. If Γ is a tree, then G is virtually $F_n \times \mathbb{Z}$ so is large (one may also use branched coverings to see this). Assume therefore $\beta(\Gamma) = 1$. If Γ contains a proper plateau P (in particular, if Γ is minimal and has a terminal vertex), a branched covering ramified over P as in the proof of Proposition 6.5 yields an admissible map $\bar{\Gamma} \rightarrow \Gamma$ with $\beta(\bar{\Gamma}) \geq 2$, so G is large. On the other hand, since every connected space covering a circle is a circle, Corollary 6.6 implies that G is not large if Γ is a circle with no proper plateau. \square

PROPOSITION 6.8. — *Given a connected labelled graph Γ , there exists an admissible map $\pi : \bar{\Gamma} \rightarrow \Gamma$ such that $\bar{\Gamma}$ is connected and contains no proper plateau.*

Every GBS group has a finite index subgroup represented by a labelled graph with no proper plateau.

Proof. — The second assertion follows from the first one and Lemma 6.3, so we concentrate on the first. Given p , we shall construct an admissible π such that $\bar{\Gamma}$ is connected, $\bar{\Gamma}$ contains no proper p -plateau, and all multiplicities are powers of p . Noting that, for $p' \neq p$, there is no proper p' -plateau in $\bar{\Gamma}$ if there is none in Γ , the proposition follows by induction on the number of primes p such that Γ contains a proper p -plateau.

The idea is the following. Consider the union of all proper p -plateaux in Γ , and construct a branched covering ramified over it as in the proof of Proposition 6.5. Then iterate. Since the graph grows, it is not obvious that this process terminates, so we prefer to work purely within Γ .

We construct graphs Γ_i which are identical to Γ , but with different labels. Let \mathcal{P}_1 be the (disjoint) union of the proper p -plateaux of $\Gamma_1 = \Gamma$. If $\mathcal{P}_1 \neq \emptyset$, define a new labelled graph Γ_2 by dividing by p the label of each oriented edge e in $\partial\mathcal{P}_1$ (see Definition 5.8). Define \mathcal{P}_2 as the union of the proper p -plateaux of Γ_2 , and iterate until obtaining Γ_{r+1} containing no proper p -plateau (the process terminates because labels decrease).

Given a vertex v of Γ , define $P(v)$ as the number of $i \in \{1, \dots, r\}$ such that $v \in \mathcal{P}_i$. Define $P(e)$ similarly for an edge e by counting the number of times that $e \subset \mathcal{P}_i$. Note that $P(\bar{e}) = P(e)$ if \bar{e} is the opposite edge. If e has origin v , then $P(v) - P(e)$ is the number of times that the label of e gets divided by p when passing from Γ to Γ_{r+1} . In particular, $p^{P(v)-P(e)}$ divides the label λ_e of e in Γ .

We now describe π by describing preimages. Its total multiplicity is p^r . The preimage of a vertex v consists of $p^{r-P(v)}$ points, each of multiplicity $p^{P(v)}$. The preimage of an edge e consists of $p^{r-P(e)}$ edges of multiplicity

$p^{P(e)}$, each with label $\lambda_e/p^{P(v)-P(e)}$. If e has origin v , exactly $p^{P(v)-P(e)}$ lifts of e are attached to each preimage of v (there is a choice in the way lifts of e are attached to preimages of v).

This defines a graph $\bar{\Gamma}$ and a map π . We check that $(*)$ is satisfied, and $\bar{\Gamma}$ has no proper p -plateau.

Condition $(*)$ is equivalent to the equality $p^{P(v)-P(e)} = \lambda_e \wedge p^{P(v)}$, for e an edge with origin v . There are two cases. If $P(e) = 0$, then λ_e is divisible by $p^{P(v)}$, so $\lambda_e \wedge p^{P(v)} = p^{P(v)}$. If $P(e) > 0$, there exists $i \in \{1, \dots, r\}$ such that p does not divide the label of e in Γ_i , so λ_e is divisible by $p^{P(v)-P(e)}$ but not by $p^{P(v)-P(e)+1}$. Thus $\lambda_e \wedge p^{P(v)} = p^{P(v)-P(e)}$.

The label of an edge of $\bar{\Gamma}$ depends only on its image in Γ , and it is the label of that image in the graph Γ_{r+1} . Since Γ_{r+1} has no proper p -plateau, it follows that the only possible proper p -plateaux of $\bar{\Gamma}$ are its connected components. Restricting to a component of $\bar{\Gamma}$ yields the desired map π (one may show that $\bar{\Gamma}$ is connected, but this is not needed). □

THEOREM 6.9. — *Given two GBS groups G_1, G_2 represented by labelled graphs Γ_1, Γ_2 which are strongly slide-free and contain no proper plateau, one may decide whether G_1 and G_2 have isomorphic finite index subgroups or not.*

Γ is *strongly slide-free* [17] if, whenever two edges e, f have the same origin, then λ_e does not divide λ_f . There is at most one minimal strongly slide-free labelled graph representing G [17].

The group $G = BS(m, n)$ satisfies the hypotheses of the theorem if and only if m and n are coprime (unless G is solvable). The theorem also applies to GBS groups represented by a circle with no proper plateau.

Proof. — Consider G_1, G_2 as in the theorem. After replacing them by a subgroup of index 2 if needed, the modulus maps Δ_{G_i} only take positive values, so we may assume that all labels are positive (see Section 2). We also assume that Γ_1, Γ_2 are minimal. Note that they have no terminal vertex (such a vertex would be a plateau). By a covering, we always mean a label-preserving topological covering between labelled graphs.

Suppose that H is isomorphic to a finite index subgroup of both G_1 and G_2 . By Corollary 6.6, it is represented by labelled graphs $\bar{\Gamma}_i$ which are coverings of Γ_i . These graphs are minimal and strongly slide-free, so by Theorem 1.2 of [17] (see also [21]) they are the same. This means that the labelled graphs Γ_1 and Γ_2 have a common finite covering if G_1 and G_2 are commensurable. The converse is also true by Assertion 2 of Lemma 6.3, so we are reduced to deciding whether Γ_1 and Γ_2 have a common finite covering (as labelled graphs).

We claim that, given Γ_1 and Γ_2 , we can find a number M such that, if Γ_1 and Γ_2 have a common finite covering, then they have one with at most M edges. Assuming the claim, we can decide existence of a common finite covering by inspection.

We deduce the claim from the fact that the proof of Leighton's graph covering theorem [24] is constructive. We follow the account given in [29], proof of Theorem 1.1.

Assume that Γ_1 and Γ_2 have a common finite covering, and denote by $\tilde{\Gamma}$ their universal covering. We first subdivide edges e of Γ_1 and Γ_2 having the same label at both ends, placing labels 1 near the created midpoint. This ensures that $\text{Aut}(\tilde{\Gamma})$ acts on $\tilde{\Gamma}$ without inversions.

As in [29], we let $C = \tilde{\Gamma} / \text{Aut}(\tilde{\Gamma})$ be the graph of colors. It is a quotient of Γ_i , so belongs to an explicit finite list. For each graph in this list, we define numbers n_i, m_k, r_k using Γ_1 as in [29], and we compute numbers a_i and b_k . As shown in [29], Γ_1 and Γ_2 have a common covering whose number of edges may be explicitly bounded in terms of $C, \Gamma_1, \Gamma_2, b_k$. Since there are only finitely many possibilities for C , the existence of M follows. \square

BIBLIOGRAPHY

- [1] H. BASS, "Covering theory for graphs of groups", *J. Pure Appl. Algebra* **89** (1993), no. 1-2, p. 3-47.
- [2] G. BAUMSLAG, C. F. MILLER, III & H. SHORT, "Unsolvable problems about small cancellation and word hyperbolic groups", *Bull. London Math. Soc.* **26** (1994), no. 1, p. 97-101.
- [3] G. BAUMSLAG & D. SOLITAR, "Some two-generator one-relator non-Hopfian groups", *Bull. Amer. Math. Soc.* **68** (1962), p. 199-201.
- [4] B. BEEKER, "Multiple conjugacy problem in graphs of free abelian groups", <http://arxiv.org/abs/1106.3978>, to appear in *Groups, geometry, dynamics*.
- [5] J. O. BUTTON, "A formula for the normal subgroup growth of Baumslag-Solitar groups", *J. Group Theory* **11** (2008), no. 6, p. 879-884.
- [6] M. CLAY, "Deformation spaces of G -trees and automorphisms of Baumslag-Solitar groups", *Groups Geom. Dyn.* **3** (2009), no. 1, p. 39-69.
- [7] M. CLAY & M. FORESTER, "On the isomorphism problem for generalized Baumslag-Solitar groups", *Algebr. Geom. Topol.* **8** (2008), no. 4, p. 2289-2322.
- [8] ———, "Whitehead moves for G -trees", *Bull. Lond. Math. Soc.* **41** (2009), no. 2, p. 205-212.
- [9] D. J. COLLINS, "Generation and presentation of one-relator groups with centre", *Math. Z.* **157** (1977), no. 1, p. 63-77.
- [10] D. J. COLLINS & F. LEVIN, "Automorphisms and Hopficity of certain Baumslag-Solitar groups", *Arch. Math. (Basel)* **40** (1983), no. 5, p. 385-400.
- [11] Y. CORNUILLER & A. VALETTE, "On equivariant embeddings of generalized Baumslag-Solitar groups", *Geom. Dedicata* **175** (2015), p. 385-401.
- [12] M. CULLER, "Finite groups of outer automorphisms of a free group", in *Contributions to group theory*, Contemp. Math., vol. 33, Amer. Math. Soc., Providence, RI, 1984, p. 197-207.

- [13] D. DEGRUISE & N. PETROSYAN, “Bredon cohomological dimensions for groups acting on CAT(0)-spaces”, <http://arxiv.org/abs/1203.3884>.
- [14] F. A. DUDKIN, “Subgroups of finite index in Baumslag-Solitar groups”, *Algebra Logika* **49** (2010), no. 3, p. 331-345, 427, 429.
- [15] M. J. DUNWOODY, “Folding sequences”, in *The Epstein birthday schrift*, Geom. Topol. Monogr., vol. 1, Geom. Topol. Publ., Coventry, 1998, p. 139-158 (electronic).
- [16] M. EDJVET & S. J. PRIDE, “The concept of “largeness” in group theory. II”, in *Groups—Korea 1983 (Kyoungju, 1983)*, Lecture Notes in Math., vol. 1098, Springer, Berlin, 1984, p. 29-54.
- [17] M. FORESTER, “Deformation and rigidity of simplicial group actions on trees”, *Geom. Topol.* **6** (2002), p. 219-267 (electronic).
- [18] ———, “On uniqueness of JSJ decompositions of finitely generated groups”, *Comment. Math. Helv.* **78** (2003), no. 4, p. 740-751.
- [19] ———, “Splittings of generalized Baumslag-Solitar groups”, *Geom. Dedicata* **121** (2006), p. 43-59.
- [20] E. GELMAN, “Subgroup growth of Baumslag-Solitar groups”, *J. Group Theory* **8** (2005), no. 6, p. 801-806.
- [21] V. GUIRARDEL, “A very short proof of Forester’s rigidity result”, *Geom. Topol.* **7** (2003), p. 321-328 (electronic).
- [22] D. G. KHRAMTSOV, “Finite groups of automorphisms of free groups”, *Mat. Zametki* **38** (1985), no. 3, p. 386-392, 476.
- [23] P. H. KROPHOLLER, “Baumslag-Solitar groups and some other groups of cohomological dimension two”, *Comment. Math. Helv.* **65** (1990), no. 4, p. 547-558.
- [24] F. T. LEIGHTON, “Finite common coverings of graphs”, *J. Combin. Theory Ser. B* **33** (1982), no. 3, p. 231-238.
- [25] G. LEVITT, “On the automorphism group of generalized Baumslag-Solitar groups”, *Geom. Topol.* **11** (2007), p. 473-515.
- [26] ———, “Quotients and subgroups of Baumslag-Solitar groups”, *J. Group Theory* **18** (2015), no. 1, p. 1-43.
- [27] J. MCCOOL, “A class of one-relator groups with centre”, *Bull. Austral. Math. Soc.* **44** (1991), no. 2, p. 245-252.
- [28] T. MECHAM, *Largeness of graphs of abelian groups*, ProQuest LLC, Ann Arbor, MI, 2009, Thesis (Ph.D.)—The University of Oklahoma, 72 pages.
- [29] W. D. NEUMANN, “On Leighton’s graph covering theorem”, *Groups Geom. Dyn.* **4** (2010), no. 4, p. 863-872.
- [30] A. PIETROWSKI, “The isomorphism problem for one-relator groups with non-trivial centre”, *Math. Z.* **136** (1974), p. 95-106.
- [31] E. RIPS, “Subgroups of small cancellation groups”, *Bull. London Math. Soc.* **14** (1982), no. 1, p. 45-47.
- [32] P. SCOTT & T. WALL, “Topological methods in group theory”, in *Homological group theory (Proc. Sympos., Durham, 1977)*, London Math. Soc. Lecture Note Ser., vol. 36, Cambridge Univ. Press, Cambridge-New York, 1979, p. 137-203.
- [33] K. WHYTE, “The large scale geometry of the higher Baumslag-Solitar groups”, *Geom. Funct. Anal.* **11** (2001), no. 6, p. 1327-1343.
- [34] B. ZIMMERMANN, “Über Homöomorphismen n -dimensionaler Henkelkörper und endliche Erweiterungen von Schottky-Gruppen”, *Comment. Math. Helv.* **56** (1981), no. 3, p. 474-486.

Manuscrit reçu le 30 mai 2013,
 accepté le 14 mai 2014.

Gilbert LEVITT
Laboratoire de Mathématiques Nicolas Oresme
Université de Caen et CNRS (UMR 6139)
BP 5186
F-14032 Caen Cedex, (France)
levitt@unicaen.fr