



ANNALES

DE

L'INSTITUT FOURIER

Daniel SCHLIEBNER

On Lorentzian manifolds with highest first Betti number

Tome 65, n° 4 (2015), p. 1423-1436.

http://aif.cedram.org/item?id=AIF_2015__65_4_1423_0



© Association des Annales de l'institut Fourier, 2015,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

ON LORENTZIAN MANIFOLDS WITH HIGHEST FIRST BETTI NUMBER

by Daniel SCHLIEBNER (*)

ABSTRACT. — We consider Lorentzian manifolds with parallel light-like vector field V . Being parallel and light-like, the orthogonal complement of V induces a codimension one foliation. Assuming compactness of the leaves and non-negative Ricci curvature on the leaves it is known that the first Betti number is bounded by the dimension of the manifold or the leaves if the manifold is compact or non-compact, respectively. We prove in the case of the maximality of the first Betti number that every such Lorentzian manifold is – up to finite cover – diffeomorphic to the torus (in the compact case) or the product of the real line with a torus (in the non-compact case) and has very degenerate curvature, i.e. the curvature tensor induced on the leaves is light-like.

RÉSUMÉ. — On considère des variétés lorentziennes avec un champ de vecteurs de genre lumière parallèle. Comme il est parallèle et de genre lumière, son complément orthogonal induit un feuilletage de codimension un. Si on suppose les feuilles compactes et la courbure de Ricci positive (ou nulle) sur les feuilles, on sait que le premier nombre de Betti est borné par la dimension de la variété ou de la feuille, suivant la compacité ou non-compacité de la variété. Nous démontrons que dans le cas où le nombre de Betti est maximal, quitte à la remplacer par un revêtement fini, toute telle variété lorentzienne est soit diffeomorphe au tore (si elle est compacte) ou alors au produit d'une droite avec un tore (autrement), et que sa courbure est très dégénérée, c'est-à-dire le tenseur de courbure induit sur les feuilles est de genre lumière.

1. Introduction

We consider $(n + 2)$ -dimensional Lorentzian manifolds⁽¹⁾ (\mathcal{M}, g) with parallel light-like vector field V , i.e. with $g(V, V) = 0$ and $\nabla^g V = 0$ for the Levi-Civita connection ∇^g of g . Since the vector field V is parallel and

Keywords: Lorentzian manifolds, holonomy groups, Betti number.

Math. classification: 53C15, 53C29, 53C12.

(*) The author is funded by the Berlin Mathematical School (BMS).

⁽¹⁾In this paper, all manifolds are assumed to be smooth, connected and without boundary.

light-like we obtain a parallel line bundle $\mathbb{V} := \mathbb{R}V$, while the orthogonal complement distribution

$$\mathbb{V}^\perp := \{X \in \Gamma(T\mathcal{M}) \mid g(X, V) = 0\}$$

defines a parallel sub-distribution of the tangent bundle of codimension one and $\mathbb{V} \subset \mathbb{V}^\perp$. Being parallel, \mathbb{V}^\perp induces a codimension one foliation on \mathcal{M} (which we will, in abuse of notation, also denote by \mathbb{V}^\perp) into light-like hypersurfaces which are the leaves of the foliation, each of which we usually will denote with L , i.e. $TL = \mathbb{V}^\perp$. The holonomy⁽²⁾ of each such Lorentzian manifold is non-irreducible and contained in the stabilizer $O(n) \times \mathbb{R}^n$ of V in $O(1, n + 1)$. If, moreover, the projection of the holonomy group onto \mathbb{R}^n is surjective it is even indecomposable. An overview over this topic can be found for example in [6, 1]. A general interest in this area is to find topological consequences on the manifold \mathcal{M} if it admits such a Lorentzian metric. In light of this motivation, K. Lärz studied in his dissertation [10], among other aspects of global holonomy theory, topological questions on Lorentzian manifolds which admit a parallel light-like vector field. One result he obtained was the boundedness of the first Betti number⁽³⁾ $b_1(\mathcal{M})$ by the dimension of \mathcal{M} under additional assumptions on the structure of the foliation and the Ricci curvature, cf. [10, Theorem 2.82], see also Lemma 3.2 in Section 3. To be precise, he proved that every Lorentzian manifold with parallel light-like vector field⁽⁴⁾ such that the leaves of \mathbb{V}^\perp are compact⁽⁵⁾ and the Ricci curvature of g is non-negative on $\mathbb{V}^\perp \times \mathbb{V}^\perp$, fulfills $\varepsilon \leq b_1(\mathcal{M}) \leq \dim(\mathcal{M}) - 1 + \varepsilon$, where $\varepsilon = 1$ if \mathcal{M} is compact, and $\varepsilon = 0$ if \mathcal{M} is non-compact.

A natural question occurring in this context is, if one can describe the cases in which the upper bound for the first Betti number is actually reached. This is basically motivated by the classical Bochner result by which any compact, oriented Riemannian manifold \mathcal{N} with non-negative Ricci curvature has $b_1(\mathcal{N}) \leq \dim \mathcal{N}$ and $b_1(\mathcal{N}) = \dim \mathcal{N}$ if and only if it is isometric to the flat torus [16, Ch. 7, Corollary 19].

(2) By the *holonomy* of (\mathcal{M}, g) we mean the group $\text{Hol}_x(\mathcal{M}, g) := \{\mathcal{P}_\gamma \mid \gamma \text{ loop in } x\} \subset O(T_x\mathcal{M}, g_x)$ of parallel displacements along loops closed in $x \in \mathcal{M}$.

(3) We define the first Betti number of any manifold \mathcal{M} to be the rank of $H^1(\mathcal{M}, \mathbb{R})$.

(4) Essentially his assumption on the light-like vector field V was weaker. Namely he just assumed V to be *recurrent*, i.e. such that $\nabla^g V = \omega \otimes V$, with $\omega \in \Omega^1(\mathcal{M})$ and $\ker \omega = \mathbb{V}^\perp$.

(5) Note that by [3], all leaves are diffeomorphic and either dense or closed. Indeed, a time-orientable indecomposable, non-irreducible Lorentzian manifold is obviously transversally parallelizable, see e.g. [10, Lemma 2.47].

The main intention of this paper is to give a full answer to this question in the Lorentzian case under the assumptions that the leaves of \mathbb{V}^\perp are compact and the Ricci curvature of g is non-negative on $\mathbb{V}^\perp \times \mathbb{V}^\perp$. Beside the ideas already used in [10] our main contribution is to prove, along a leaf of \mathbb{V}^\perp , the existence of a horizontal and integrable realization of the screen bundle $\Sigma = \mathbb{V}^\perp/\mathbb{V}$, which is a subbundle \mathbb{S} of $T\mathcal{M}$ isomorphic to Σ . This is done by exploiting the maximality assumption on the Betti number (Proposition 3.3) and it relates the Ricci curvature of g along a leaf with the Ricci curvature of a Riemannian metric (Lemma 2.2). We point out that horizontal and integrable screen distributions are a crucial tool in the study of Lorentzian manifolds with indecomposable holonomy, see e.g. [11, 10, 12], since they provide a link between Riemannian and Lorentzian geometry. As a result we find out that under the assumptions above, the consequence for the Lorentzian metric g is not to be flat but still having very degenerate curvature. Indeed, the obtained metrics have light-like hypersurface curvature. These metrics are in some sense generalizations of pp-waves and were in light of this motivation already studied by T. Leistner in [11]. Topologically, the manifolds turn out to be diffeomorphic (homeomorphic in dimension four) to a finite cover of the torus (in the compact case) or the product of the real line with the torus (in the non-compact case).

DEFINITION. — A Lorentzian manifold (\mathcal{M}, g) with a global non-trivial parallel light-like vector field $V \in \Gamma(T\mathcal{M})$, i.e. $V \neq 0$, $g(V, V) = 0$ and $\nabla^g V = 0$, has light-like hypersurface curvature, iff the curvature R satisfies $R(X, Y)W \in \Gamma(\mathbb{V})$ for all $X, Y, W \in \Gamma(\mathbb{V}^\perp)$.

THEOREM 1.1. — Let (\mathcal{M}, g) be an orientable $(n + 2)$ -dimensional Lorentzian manifold with parallel light-like vector field V . Assume that the leaves of the codimension one foliation induced by the distribution \mathbb{V}^\perp are compact and $\text{Ric}|_{\mathbb{V}^\perp \times \mathbb{V}^\perp} \geq 0$. Then:

- (i) if \mathcal{M} is compact, then $b_1(\mathcal{M}) \leq n + 2$ and $b_1(\mathcal{M}) = n + 2$ if and only if \mathcal{M} is – up to finite cover – diffeomorphic (homeomorphic⁽⁶⁾ if $\dim \mathcal{M} = 4$) to the torus and g has light-like hypersurface curvature;

⁽⁶⁾ In dimension four, \mathcal{M} is only known to be homeomorphic to the torus and to our best knowledge it seems to be an open problem in geometric topology if \mathcal{M} must be also diffeomorphic to the torus.

- (ii) if \mathcal{M} is non-compact, then $b_1(\mathcal{M}) \leq n+1$ and $b_1(\mathcal{M}) = n+1$ if and only if \mathcal{M} is isometric to $\mathbb{R} \times \mathbb{T}^{n+1}$ and g has light-like hypersurface curvature.

In both cases, the leaves of \mathbb{V}^\perp are all diffeomorphic to the torus \mathbb{T}^{n+1} .

Concerning non-orientability of \mathcal{M} we give the following

Remark. — If \mathcal{M} is non-orientable, the topological statements asserted in the Main Theorem inherit to the 2-fold orientation covering $\widehat{\mathcal{M}}$ for \mathcal{M} . In particular we can still conclude $b_1(\mathcal{M}) \leq n+2$ since $b_1(\widehat{\mathcal{M}}) = b_1(\mathcal{M}) + b_{n-1}(\mathcal{M})$, cf. [2]. Moreover, since a covering is a local isometry, (\mathcal{M}, g) has also light-like hypersurface curvature.

Acknowledgements. — We would like to thank Helga Baum for motivating us to investigate the problem considered in this paper.

2. Preliminaries

Throughout this section, let (\mathcal{M}, g) be a Lorentzian manifold with parallel light-like vector field. The techniques used within the present paper involve cohomology theory for foliations. Hence, we will present the required definitions and facts used in this paper but with just giving references to the statements in the literature. Indeed, the book [20] is a very good reference for the most of the theory used here.

2.1. Screen distributions and screen bundles

By the parallel light-like vector field V on (\mathcal{M}, g) , we obtain a filtration

$$\mathbb{V} \subset \mathbb{V}^\perp \subset T\mathcal{M}$$

of the tangent bundle and a codimension two vector bundle

$$\Sigma := \mathbb{V}^\perp / \mathbb{V} \longrightarrow \mathcal{M}$$

which is called the *screen bundle* of (\mathcal{M}, g) . It comes with a naturally given connection ∇^Σ , induced by the Levi-Civita connection ∇^g of g : $\nabla_X^\Sigma \varphi := [\nabla_X^g Y]$ for some $Y \in \Gamma(\mathbb{V}^\perp)$ s.t. $[Y] = \varphi$, where $[\cdot] : \mathbb{V}^\perp \rightarrow \Sigma$ is the projection. Taking into account a non-canonical splitting $s : \Sigma \rightarrow \mathbb{V}^\perp$ of the short exact sequence $0 \rightarrow \mathbb{V} \rightarrow \mathbb{V}^\perp \rightarrow \Sigma \rightarrow 0$ one obtains a codimension two distribution $\mathbb{S} := s(\Sigma) \subset T\mathcal{M}$, called a *screen distribution* or *realization of the screen bundle* Σ . Each such realization of the screen bundle is in 1-to-1 correspondence with a *screen vector field* $Z \in \Gamma(T\mathcal{M})$ with $g(Z, Z) = 0$ and $g(V, Z) = 1$.

2.2. Riemannian foliations and Riemannian flows

Let us begin with mentioning that we do not need to distinguish between the leaves topologically. Since \mathbb{V}^\perp is defined by the closed one form $\sigma := g(V, \cdot)$, all leaves are diffeomorphic [20, Corollary 3.31] if \mathcal{M} is closed. Moreover, all leaves have trivial leaf holonomy (for a definition we refer to [20, Section 3]), cf. [17] (see [20, Theorem 3.29] for an English version). Hence, if \mathcal{M} is non-compact, then the assumption on all the leaves being compact implies that they are all diffeomorphic, [19, Ch. 2, Corollary 8.6]. In particular, in case (i) of the Main Theorem we have that if one leaf L of \mathbb{V}^\perp is compact then so are all leaves.

Given a foliated manifold $(\mathcal{N}, \mathcal{F})$ and a Riemannian metric h on \mathcal{N} , then h is said to be *bundle-like* for $(\mathcal{N}, \mathcal{F})$, iff $(\mathcal{L}_X h)(Z_1, Z_2) = 0$ for all $X \in \Gamma(T\mathcal{F})$ and $Z_i \in \Gamma(T\mathcal{F}^\perp)$, where \mathcal{L} denotes the Lie-derivative. In this case, $(\mathcal{N}, \mathcal{F}, h)$ is called a *Riemannian foliation*. Given a foliated manifold $(\mathcal{N}, \mathcal{F}, h)$ with Riemannian metric h , consider the normal bundle $Q := T\mathcal{N}/T\mathcal{F}$ and the exact sequence $0 \rightarrow T\mathcal{F} \rightarrow T\mathcal{N} \xrightarrow{\pi} Q \rightarrow 0$. Then, for a splitting $s : Q \rightarrow T\mathcal{F}^\perp$, the metric h induces a *transversal metric* h^T on Q by $h^T := s^*h|_{T\mathcal{F}^\perp}$. We define a connection on Q by

$$(2.1) \quad \nabla_X^T \varphi := \begin{cases} \pi(\nabla_X^h Y_\varphi), & X \in \Gamma(T\mathcal{F}^\perp), \\ \pi([X, Y_\varphi]), & X \in \Gamma(T\mathcal{F}), \end{cases}$$

for any $\varphi \in \Gamma(Q)$ and $s(\varphi) = Y_\varphi$. It is torsion-free [20, Proposition 3.8] and if h is bundle-like, it is metric [20, Theorem 5.8]. Moreover, if h_1 and h_2 are two bundle-like metrics w.r.t. $(\mathcal{N}, \mathcal{F})$ such that $h_1^T = h_2^T$, then $\nabla_1^T = \nabla_2^T$ [20, Theorem 5.9]. For bundle-like h , ∇^T is called the *transversal Levi-Civita connection* of the Riemannian foliation $(\mathcal{N}, \mathcal{F}, h)$. Given the transversal Levi-Civita connection ∇^T of a Riemannian foliation one obtains the corresponding curvature tensors R^T . Considering the bundle isomorphism $s(Q) \simeq Q$ one obtains the transversal Ricci curvature Ric^T defined through $\text{Ric}^T(e_i, e_j) := \sum_{k=1}^{\dim Q} R^T(e_i, e_k, e_k, e_j)$ for an h^T -orthonormal frame $\varphi_\ell \in \Gamma(Q)$ and $s(\varphi_\ell) = e_\ell$.

Given a realization \mathbb{S} of the screen bundle, we obtain a naturally associated Riemannian metric g^R corresponding to \mathbb{S} by

$$(2.2) \quad g^R(V, \cdot) := g(Z, \cdot), \quad g^R(Z, \cdot) := g(V, \cdot), \quad g^R(X, \cdot) := g(X, \cdot) \text{ for } X \in \Gamma(\mathbb{S})$$

and extension by linearity. This Riemannian metric is of certain interest since it relates aspects of Riemannian geometry naturally to the Lorentzian geometry. An important example is the following: assume that the

realization \mathbb{S} of Σ is *integrable* (i.e. $[\Gamma(\mathbb{S}), \Gamma(\mathbb{S})] \subset \Gamma(\mathbb{S})$) and *horizontal* (i.e. $[\Gamma(\mathbb{S}), \Gamma(\mathbb{V})] \subset \Gamma(\mathbb{S})$) at least along L . Then $\nabla^g|_L$ (the induced connection on a leaf L) and $\nabla^R|_L$ (the Levi-Civita connection of the induced metric by g^R on L) only differ by a section $T\mathcal{N} \rightarrow \mathbb{V}$, $\nabla^R|_L V = 0$ and $\text{Ric}^g|_{\mathbb{V}^\perp \times \mathbb{V}^\perp} = \text{Ric}^{g^R}|_{\mathbb{V}^\perp \times \mathbb{V}^\perp}$. We will use this as a key step in the proof of our Proposition 3.3. Another important property of g^R is the fact that it constitutes a *Riemannian flow* on L . Let us explain this in more detail. A *Riemannian flow* $(\mathcal{N}, \mathcal{F}, h)$ is a one-dimensional foliation \mathcal{F} s.t. h is bundle-like w.r.t. \mathcal{F} . Now, since the parallel light-like vector field V on \mathcal{M} has no zero, it constitutes a one-dimensional foliation \mathcal{F} on \mathcal{M} and in particular on each leaf L by its flow. The important property for any g^R as in (2.2) associated to some realization \mathbb{S} of the screen bundle is now:

LEMMA 2.1 ([10, Lemma 2.48]). — *For each leaf L of \mathbb{V}^\perp and any realization \mathbb{S} of the screen bundle, (L, \mathcal{F}, g^R) is a Riemannian flow, where \mathcal{F} is given by the flow of V restricted to L .*

For a Riemannian flow $(\mathcal{N}, \mathcal{F}, h)$ defined by a non-singular vector field $V \in \Gamma(T\mathcal{N})$ such that $h(V, V) = 1$, we define the *mean curvature one form* $\kappa \in \Omega^1(\mathcal{N})$ to be $\kappa := h(\nabla_V^h V, \cdot)$.

Finally the following property of g^R will be important:

LEMMA 2.2. — *Let (\mathcal{M}, g) be a Lorentzian manifold with parallel light-like vector field V . If along L there exists a horizontal and integrable realization \mathbb{S} of the screen bundle, then $\nabla^h V = 0$ and*

$$(2.3) \quad [R^g(X, Y) - R^h(X, Y)]W \in \Gamma(\mathbb{V}) \text{ for all } X, Y, W \in \Gamma(\mathbb{S}),$$

where h denotes the Riemannian metric on L defined by $h(X, \cdot) := g(X, \cdot)$ for $X \in \Gamma(\mathbb{S})$, $h(V, V) = 1$ and extension by linearity. In particular we have that $\text{Ric}^g|_{\mathbb{V}^\perp \times \mathbb{V}^\perp} = \text{Ric}^h$.

Proof. — Since $h(V, V) = 1$, we obtain $h(\nabla^h V, V) = 0$. Moreover, applying the horizontality and involutivity property of \mathbb{S} , we see by the Koszul formula for h that

$$h(\nabla_{S_1}^h V, S_2) = g(\nabla_{S_1}^g V, S_2) = 0$$

for all $S_1, S_2 \in \Gamma(\mathbb{S})$ since $\nabla^g V = 0$. This proves $\nabla^h V = 0$. Moreover, as

$$h(\nabla_{S_1}^h S_2, S_3) = g(\nabla_{S_1}^g S_2, S_3)$$

for all $S_i \in \Gamma(\mathbb{S})$ we obtain $\nabla^g|_{\mathcal{L}^\perp} - \nabla^h \in \Gamma(\mathbb{L})$ for all $X, Y \in \Gamma(T\mathcal{L}^\perp)$ which proves (2.3). □

2.3. Basic and twisted cohomology of foliations

In this paragraph we deal with the term of basic cohomology. A comprehensive introduction can be found, e.g. in [20, Chapter 4, Chapter 7]. Let $(\mathcal{N}, \mathcal{F})$ be an arbitrary foliated manifold.

DEFINITION 2.3. — *A k -form $\alpha \in \Omega^k(\mathcal{N})$ is called basic iff $X \lrcorner \alpha = 0$ and $\mathcal{L}_X \alpha = 0$ for all $X \in \Gamma(T\mathcal{F})$. The set of basic k -forms is denoted by $\Omega_B^k(\mathcal{F})$.*

Since $\mathcal{L}_X d\alpha = d\mathcal{L}_X \alpha = 0$ and $X \lrcorner d\alpha = \mathcal{L}_X \alpha - d(X \lrcorner \alpha) = 0$ for all $X \in \Gamma(T\mathcal{F})$ and $\alpha \in \Omega_B^k(\mathcal{F})$, we obtain a subcomplex $(\Omega_B^*(\mathcal{F}), d_B)$ of the de Rham-complex with $d_B := d|_{\Omega_B^*(\mathcal{F})}$ and hence a corresponding cohomology $H_B^*(\mathcal{F})$, the *basic cohomology*. Given d_B there is also a formal L^2 -adjoint δ_B [20, Theorem 7.10] and hence a transversal Laplacian $\Delta_B = d_B \delta_B + \delta_B d_B$ on $\Omega_B^*(\mathcal{F})$. Naturally, we then say that a basic form $\alpha \in \Omega_B^k(\mathcal{F})$ is *basic-harmonic* iff $\Delta_B \alpha = 0$. For Riemannian flows we have the following important result relating the mean curvature one form with the basic-harmonic forms:

THEOREM 2.4 ([4, 13]). — *Let $(\mathcal{N}, \mathcal{F}, g)$ be a Riemannian flow on a compact manifold \mathcal{N} . Then there exists a bundle-like metric \hat{g} on \mathcal{N} such that κ is basic-harmonic and $g^T = \hat{g}^T$.*

Not enough, we need another definition of cohomology for foliations. Namely, since for the basic cohomology, Poincaré duality not necessarily holds, one defines another cohomology theory. The so called *twisted cohomology* for a foliation $(\mathcal{N}, \mathcal{F}, h)$ with $\kappa \in \Omega_B^1(\mathcal{N})$ is defined by the subcomplex $(\Omega_B^*(\mathcal{F}), d_\kappa := d_B - \kappa \wedge \cdot)$ of the de Rham-complex.⁽⁷⁾ We will denote its cohomology groups by $H_\kappa^*(\mathcal{F})$. It behaves in a nice way as to obtain a type of Poincaré duality when comparing the basic cohomology and the twisted cohomology [20, Theorem 7.54].

The orientability assumption in the Main Theorem has the following background. Since most of the results about Hodge theory in basic cohomology require the foliation to be transversally oriented (i.e. there is an orientation of the normal bundle), we need the following

LEMMA 2.5. — *Let (\mathcal{M}, g) be an oriented Lorentzian manifold with parallel light-like vector field. Then, for each leaf L of \mathbb{V}^\perp and any realization*

⁽⁷⁾ Here, κ is in general the mean curvature form of the foliation [20, (3.20)]. However, we will use this only in the setting of Riemannian flows s.t. κ is the mean curvature one form.

\mathbb{S} of the screen bundle, the Riemannian flow (L, \mathcal{F}, g^R) is transversally orientable. Moreover, L is orientable.

Proof. — The argument can be found in [10, p. 78]. Namely, one can prove that $\text{Hol}(\nabla^T) \subset \text{SO}(\dim \mathbb{S})$ and hence Σ is orientable. In particular, each leaf L is orientable since any screen vector field $Z \in \Gamma(T\mathcal{M})$ defines a unit normal vector field. \square

3. Main Results

Let (\mathcal{M}, g) be an oriented Lorentzian manifold with parallel light-like vector field V inducing an integrable codimension one distribution $\mathbb{V}^\perp \subset T\mathcal{M}$ and hence a codimension one foliation on \mathcal{M} . Recall from the previous section that, by fixing a realization \mathbb{S} of the screen bundle and denoting by g^R the associated Riemannian metric on \mathcal{M} , we obtain a Riemannian flow \mathcal{F} on L by the flow of V and g^R restricted to L (Lemma 2.1). Throughout this section we will denote this Riemannian flow by (L, \mathcal{F}, g^R) .

LEMMA 3.1. — *Let L be a compact $(n+1)$ -dimensional leaf of \mathbb{V}^\perp . Then $H_{\text{dR}}^1(L) = H_B^1(\mathcal{F}) \oplus H$ for H a subgroup of $H_B^n(\mathcal{F}) \in \{0, \mathbb{R}\}$.*

Proof. — Since (L, \mathcal{F}, g^R) is a Riemannian flow, by Theorem 2.4, there exists a bundle-like metric \widehat{g}^R such that the mean-curvature one form κ of $(L, \mathcal{F}, \widehat{g}^R)$ is basic-harmonic. By the Gysin sequence for (L, \widehat{g}^R) , cf. [18, Theorem 3.2], we obtain an exact sequence

$$(3.1) \quad 0 \longrightarrow H_B^1(\mathcal{F}) \longrightarrow H_{\text{dR}}^1(L) \longrightarrow H_\kappa^0(\mathcal{F}) \longrightarrow H_B^2(\mathcal{F}) \longrightarrow \dots$$

and by taking into account that $H_\kappa^0(\mathcal{F}) \cong H_B^n(\mathcal{F})$, cf. [20, Theorem 7.54], this translates into the long exact sequence

$$(3.2) \quad 0 \longrightarrow H_B^1(\mathcal{F}) \longrightarrow H_{\text{dR}}^1(L) \xrightarrow{\Phi} H_B^n(\mathcal{F}) \longrightarrow H_B^2(\mathcal{F}) \longrightarrow \dots$$

and thus we obtain the short exact sequence

$$(3.3) \quad 0 \longrightarrow H_B^1(\mathcal{F}) \longrightarrow H_{\text{dR}}^1(L) \longrightarrow H \longrightarrow 0$$

for $H := \text{im } \Phi \subset H_B^n(\mathcal{F})$. In particular this is a short exact sequence of vector spaces and hence splits as a direct sum. Since $H_B^n(\mathcal{F}) \in \{0, \mathbb{R}\}$ by [20, Corollary 7.57] this completes the proof. \square

By assuming non-negativity of the Ricci curvature on $TL \times TL$, we obtain the following estimation for the dimensions of $H_{\text{dR}}^1(L)$ and $H_B^1(\mathcal{F})$. This is precisely the [10, Proposition 2.81] of K. Lärz. However, to be self-contained, make the upcoming proofs more precise and to fix notation we will present its full proof here.

LEMMA 3.2. — *Let L be a compact $(n + 1)$ -dimensional leaf of \mathbb{V}^\perp and $\text{Ric}^g|_{TL \times TL} \geq 0$. Then $b_1(L) \leq \dim H_B^1(\mathcal{F}) + 1 \leq \dim L$.*

Proof. — Again, consider the Riemannian flow (L, \mathcal{F}, g^R) and the bundle-like metric \widehat{g}^R such that the mean-curvature one form κ of $(L, \mathcal{F}, \widehat{g}^R)$ is basic-harmonic, constituted by Theorem 2.4. In particular, the induced metrics on $TL/T\mathcal{F} =: \Sigma|_L$ coincide and hence so do the induced transversal connections. Defining by $\widehat{\mathbb{S}} := \ker \widehat{g}^R(V, \cdot)$ a realization of $\Sigma|_L$, the transversal Levi-Civita connection $\nabla^T : \Gamma(\widehat{\mathbb{S}}) \rightarrow \Gamma(T^*L \otimes \widehat{\mathbb{S}})$ is given by

$$(3.4) \quad \nabla_X^T Y := \begin{cases} \text{pr}_{\widehat{\mathbb{S}}}(\widehat{\nabla}_X Y), & X \in \Gamma(\widehat{\mathbb{S}}), \\ \text{pr}_{\widehat{\mathbb{S}}}([X, Y]), & X \in \Gamma(\mathbb{V}), \end{cases}$$

for any $Y \in \Gamma(\widehat{\mathbb{S}})$, where $\widehat{\nabla}$ is the Levi-Civita connection of \widehat{g}^R (see Section 2.2). Since $\text{Ric}^g(V, \cdot) = 0$ and $g_{\widehat{\mathbb{S}} \times \widehat{\mathbb{S}}} = g^R|_{\widehat{\mathbb{S}} \times \widehat{\mathbb{S}}}$ we have that

$$(3.5) \quad \text{Ric}_{\widehat{\mathbb{S}} \times \widehat{\mathbb{S}}}^T = \text{Ric}^g|_{\widehat{\mathbb{S}} \times \widehat{\mathbb{S}}} \geq 0.$$

Since any class in $H_B^1(\mathcal{F})$ can be represented by a Δ_B -harmonic one-form $\alpha \in \Omega_B^1(L)$ [20, Theorem 7.51], we choose such a basic-harmonic α for each generator of $H_B^1(\mathcal{F})$. To apply a Bochner argument we need an appropriate Weizenböck formula which is given by [8, Proposition 6.7] and by integration reads

$$0 = \int_L \|\nabla^T \alpha\|^2 + \int_L \text{Ric}^T(\alpha^\sharp, \alpha^\sharp),$$

where α^\sharp is the dual of α w.r.t. \widehat{g}^R . Hence $\nabla_X^T \alpha = 0$ for all $X \in \Gamma(TL)$. This proves $\dim H_B^1(\mathcal{F}) \leq n = \dim \widehat{\mathbb{S}}$ and thus the second asserted inequality. For the first inequality we apply Lemma 3.1; this completes the proof. \square

We can now prove the key step for the proof of the main result. Namely, the – by the previous lemma – maximality of the dimension of the first Betti number of L and $\text{Ric}^g|_{TL \times TL} \geq 0$ already yield that L is the torus with the curvature of $\nabla^g|_L$ on L being light-like.

PROPOSITION 3.3. — *Let (\mathcal{M}^{n+2}, g) be an oriented $(n + 2)$ -dimensional Lorentzian manifold with parallel light-like vector field V such that $\text{Ric}^g|_{TL \times TL} \geq 0$ and the leaves L of the codimension one foliation induced by \mathbb{V}^\perp are compact. Then, $b_1(L) \leq n + 1$ and $b_1(L) = n + 1$ if and only if $\nabla^g|_L$ has light-like curvature and L is diffeomorphic to the torus.*

Proof. — The proof works as follows: Let $b_1(L) = n + 1$. We will define for every realization of the screen bundle another screen distribution along L which is horizontal and integrable. The associated Riemannian

metric h to this screen distribution then has non-negative Ricci curvature by Lemma 2.2 and hence (L, h) is the flat torus implying $\nabla^g|_L$ to have light-like curvature.

Step 1. — Fix a realization of the screen bundle and hence a Riemannian flow (L, \mathcal{F}, g^R) on L and denote by \widehat{g}^R the bundle-like metric s.t. the mean-curvature one form κ is basic-harmonic. W.l.o.g. we may additionally assume $\widehat{g}^R(V, V) = 1$. We define $\chi := \widehat{g}^R(V, \cdot)$.

Step 2. — Recall that $H_B^n(\mathcal{F}) \in \{0, \mathbb{R}\}$ (see Lemma 3.1) since (L, \mathcal{F}, g^R) is transversally orientable by Lemma 2.5. Assume that $H_B^n(\mathcal{F}) = 0$. Then we would have by the sequence (3.2) that $H_B^1(\mathcal{F}) \cong H_{\text{dR}}^1(L)$ which is impossible due to $b_1(L) = n + 1$ and Lemma 3.2. Hence $H_B^n(\mathcal{F}) = \mathbb{R}$ and [15, Theorem A] (see also [20, Theorem 6.17] for an English version) implies that $(L, \mathcal{F}, \widehat{g}^R)$ is an *isometric* Riemannian flow, i.e. V is a \widehat{g}^R -Killing field and therefore $\mathcal{L}_V\chi = 0$ and $\kappa = 0$.

Step 3. — By Lemma 3.2 and $b_1(L) = n + 1$ we infer $\dim H_B^1(\mathcal{F}) = n$. For $(L, \mathcal{F}, \widehat{g}^R)$ being an *isometric* Riemannian flow, the Gysin long exact sequence (3.2) is given as

$$(3.6) \quad 0 \longrightarrow H_B^1(\mathcal{F}) \longrightarrow H_{\text{dR}}^1(L) \xrightarrow{\Phi} H_B^0(\mathcal{F}) \xrightarrow{\delta} H_B^2(\mathcal{F}) \longrightarrow \dots$$

where $\Phi = (V \lrcorner \cdot)_*$ and $\delta = [d\chi \wedge \cdot]$, see [20, Theorem 6.13]. Note that since $H_\kappa^0(\mathcal{F}) \cong H_B^n(\mathcal{F})$ by [20, Theorem 7.54] and $\kappa = 0$ we have $H_B^0(\mathcal{F}) = H_\kappa^0(\mathcal{F}) \cong H_B^n(\mathcal{F}) = \mathbb{R}$. This together with (3.6) implies the short exact sequence

$$(3.7) \quad 0 \longrightarrow H_B^1(\mathcal{F}) \longrightarrow H_{\text{dR}}^1(L) \longrightarrow \text{im } \Phi \longrightarrow 0$$

with $\text{im } \Phi = \mathbb{R}$ since $b_1(L) = n + \dim \text{im } \Phi$. Hence, by the exactness of (3.6), $\ker \delta = \mathbb{R}$ and so $0 = \delta([1]) = [1 \cdot d\chi]$. Hence $[d\chi]$ vanishes in $H_B^2(\mathcal{F})$. In this case, $d\chi = d_B\alpha$ for some $\alpha \in \Omega_B^1(L)$. Then, $\omega := \chi - \alpha \in \Omega^1(L)$ is closed w.r.t. d and $\omega(V) = 1$. We define

$$(3.8) \quad \mathbb{S} := \ker \omega.$$

Obviously, \mathbb{S} is integrable and it is horizontal since $\mathcal{L}_V\omega = \mathcal{L}_V\chi - \mathcal{L}_V\alpha = 0$ as α is a basic 1-form and $\mathcal{L}_V\chi = 0$ by Step 2.

Step 4. — Define a Riemannian metric on L associated to g and \mathbb{S} by

$$h(X, Y) := \begin{cases} 1, & X = Y = V \\ g(X, Y), & X, Y \in \Gamma(\mathbb{S}) \\ 0, & X \in \Gamma(\mathbb{V}) \text{ and } Y \in \Gamma(\mathbb{S}) \\ & \text{or } Y \in \Gamma(\mathbb{V}) \text{ and } X \in \Gamma(\mathbb{S}) \end{cases}$$

and linear extension. Since \mathbb{S} is horizontal and integrable, Lemma 2.2 yields $\text{Ric}^h = \text{Ric}^g|_{TL \times TL} \geq 0$. Therefore, (L, h) turns into a compact, orientable Riemannian manifold with non-negative Ricci-curvature and is thus isometric to the flat torus [16, Ch. 7, Corollary 19]. In particular, equation (2.3) in Lemma 2.2 implies that $\nabla^g|_L$ has light-like curvature. \square

The last lemma of this section now relates the dimension of $H_{\text{dR}}^1(\mathcal{M})$ with the dimension of $H_{\text{dR}}^1(L)$ for a compact leaf of the codimension one foliation \mathbb{V}^\perp .

LEMMA 3.4. — *Let (\mathcal{M}, g) be an $(n + 2)$ -dimensional Lorentzian manifold with parallel light-like vector field V such that the leaves L of the codimension one foliation induced by \mathbb{V}^\perp are compact. Then,*

- (i) $b_1(\mathcal{M}) \leq b_1(L) + 1$.
- (ii) $\pi_1(\mathcal{M}) \cong \mathbb{Z} \rtimes_\varphi \pi_1(L)$ for some homomorphism $\varphi \in \text{Hom}(\mathbb{Z}, \text{Aut}(\pi_1(L)))$ and hence $H_1(\mathcal{M}, \mathbb{Z}) \cong \mathbb{Z} \oplus H_1(L, \mathbb{Z})/K$, for the subgroup K of $\pi_1(L)$ generated by the elements $\varphi(k)([\gamma]) \cdot [\gamma]^{-1}$, $k \in \mathbb{Z}$, $[\gamma] \in \pi_1(L)$.

Proof. — Since \mathbb{V}^\perp is defined as $\mathbb{V}^\perp = \ker \sigma$ for $\sigma := g(V, \cdot)$ and $d\sigma = 0$, all leaves have trivial leaf holonomy (see Section 2.2). The compactness assumption for the leaves then implies, that \mathcal{M} fibers over S^1 or \mathbb{R} with each leaf being given as a fiber of the fibration [19, Corollary 8.6]. Hence, if \mathcal{M} is not compact, \mathcal{M} fibers over \mathbb{R} and thus $\mathcal{M} \simeq \mathbb{R} \times L$ for a fixed leaf. Otherwise, if \mathcal{M} is compact, \mathcal{M} fibers over S^1 and the long exact sequence of homotopy groups for fibrations [9, Theorem 4.41] yields

$$0 \longrightarrow \pi_1(L) \longrightarrow \pi_1(\mathcal{M}) \longrightarrow \mathbb{Z} \longrightarrow 0$$

and hence $\pi_1(\mathcal{M}) \cong \mathbb{Z} \rtimes_\varphi \pi_1(L)$ for some $\varphi \in \text{Hom}(\mathbb{Z}, \text{Aut}(\pi_1(L)))$ since \mathbb{Z} is free. By the Hurewicz theorem we have $H_1(\mathcal{M}, \mathbb{Z}) \cong \pi_1(\mathcal{M})_{\text{ab}}$, where for any group G we denote with $G_{\text{ab}} := G/[G, G]$ its Abelianization. However, for any semi-direct product $G = A \rtimes_\psi B$ for some $\psi \in \text{Hom}(A, \text{Aut}(B))$ we have that

$$G_{\text{ab}} = A_{\text{ab}} \oplus B_{\text{ab}}/H$$

with H denoting the subgroup of B generated by elements $\psi(a)b \cdot b^{-1}$ with $a \in A$ and $b \in B$, cf. [7, Proposition 3.3]. Therefore, we see that

$$H_1(\mathcal{M}, \mathbb{Z}) \cong \pi_1(\mathcal{M})_{\text{ab}} \cong \mathbb{Z} \oplus H_1(L, \mathbb{Z})/K$$

for the subgroup K generated by the elements $\varphi(k)([\gamma]) \cdot [\gamma]^{-1}$, $k \in \mathbb{Z}$, $[\gamma] \in \pi_1(L)$. In particular, $b_1(\mathcal{M}) = 1 + \text{rank } H_1(L, \mathbb{Z}) - \text{rank } K \leq b_1(L) + 1$. \square

4. Proof of the Main Theorem

Let us first proof part (i) of the Main Theorem.

With $b_1(\mathcal{M}) = n + 2$ we infer $b_1(L) = n + 1$ by Lemma 3.4(i) taking into account that $b_1(L) \leq \dim L = n + 1$ by Lemma 3.2. Hence, all assumptions of Proposition 3.3 are satisfied. We obtain, that the connection $\nabla^g|_L$ on each leaf L induced by the Levi-Civita connection of g has light-like curvature. By [11, Proposition 6] this is equivalent for (\mathcal{M}, g) to have light-like hypersurface curvature.

Since $L = \mathbb{T}^{n+1}$ and hence \mathcal{M} fibers over S^1 with toric fibers, the long exact sequence of homotopy groups for fibrations implies that for $k > 1$ all homotopy groups $\pi_k(\mathcal{M})$ vanish. Therefore, \mathcal{M} is a $K(\pi_1(\mathcal{M}), 1)$ -space⁽⁸⁾, while $\pi_1(\mathcal{M}) \cong \mathbb{Z}_\varphi \rtimes \pi_1(L) \cong \mathbb{Z}_\varphi \rtimes \mathbb{Z}^{n+1}$ by Lemma 3.4(ii) and since $\pi_1(L) = \mathbb{Z}^{n+1}$. By assumption, $H_1(\mathcal{M}, \mathbb{Z}) = \mathbb{Z}^{n+2} \oplus \text{Tor}$ and hence

$$\mathbb{Z}^{n+2} \oplus \text{Tor} = H_1(\mathcal{M}, \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z}^{n+1}/K$$

by Lemma 3.4(ii). But this equation can only hold, if and only if K is trivial. Namely, comparing the ranks of the left and the right hand side, we observe

$$n + 2 = 1 + (n + 1) - \text{rank } K \iff \text{rank } K = 0.$$

As K is a subgroup of $\pi_1(L) \cong \mathbb{Z}^{n+1}$, K is trivial. Therefore, \mathcal{M}^{n+2} is a $K(\mathbb{Z}^{n+2}, 1)$ -space and hence homotopy-equivalent to the torus [9, Theorem 1B.8]. But in the case of the torus, for $\dim \mathcal{M} \leq 3$ this is even equivalent for \mathcal{M} itself⁽⁹⁾ or some finite cover (if $\dim \mathcal{M} > 4$) to be diffeomorphic to the standard torus [22, Page 236]. For $\dim \mathcal{M} = 4$ we can only conclude that \mathcal{M} is homeomorphic to \mathbb{T}^4 [5, Chapter 11.5].

It remains to prove part (ii). This is straightforward since $\mathcal{M} \simeq \mathbb{R} \times L$ for a fixed leaf L by [19, Corollary 8.6] (see the proof of Lemma 3.4 for details) and hence $n + 1 = b_1(\mathcal{M}) = b_1(L)$ so Proposition 3.3 applies. \square

⁽⁸⁾ With $K(G, 1)$ -spaces we denote the Eilenberg–MacLane spaces with first fundamental group equal to G and all other homotopy groups vanishing.

⁽⁹⁾ For the 3-dimensional case, Waldhausen [21] proved this for *Haken manifolds* and thus in particular for 3-dimensional closed manifolds fibering over the circle. Note that for 3-dimensional manifolds classifications up to diffeomorphism and homeomorphism coincide [14].

BIBLIOGRAPHY

- [1] H. BAUM, “Holonomy groups of Lorentzian manifolds: a status report”, in *Global differential geometry*, Springer Proc. Math., vol. 17, Springer, Heidelberg, 2012, p. 163-200.
- [2] R. G. BRASHER, “The homology sequence of the double covering: Betti numbers and duality”, *Proc. Amer. Math. Soc.* **23** (1969), p. 714-717.
- [3] L. CONLON, “Transversally parallelizable foliations of codimension two”, *Trans. Amer. Math. Soc.* **194** (1974), p. 79-102, erratum, *ibid.* 207 (1975), 406.
- [4] D. DOMÍNGUEZ, “Finiteness and tenseness theorems for Riemannian foliations”, *Amer. J. Math.* **120** (1998), no. 6, p. 1237-1276.
- [5] M. H. FREEDMAN & F. QUINN, *Topology of 4-manifolds*, Princeton Mathematical Series, vol. 39, Princeton University Press, Princeton, NJ, 1990, viii+259 pages.
- [6] A. S. GALAEV & T. LEISTNER, “Holonomy groups of Lorentzian manifolds: classification, examples, and applications”, in *Recent developments in pseudo-Riemannian geometry*, ESI Lect. Math. Phys., Eur. Math. Soc., Zürich, 2008, p. 53-96.
- [7] D. L. GONÇALVES & J. GUASCHI, “The lower central and derived series of the braid groups of the sphere”, *Trans. Amer. Math. Soc.* **361** (2009), no. 7, p. 3375-3399.
- [8] G. HABIB & K. RICHARDSON, “Modified differentials and basic cohomology for Riemannian foliations”, *J. Geom. Anal.* **23** (2013), no. 3, p. 1314-1342.
- [9] A. HATCHER, *Algebraic topology*, Cambridge University Press, Cambridge, 2002, xii+544 pages.
- [10] K. LÄRZ, “Global Aspects of Holonomy in Pseudo-Riemannian Geometry”, PhD Thesis, Humboldt-Universität zu Berlin, 2011.
- [11] T. LEISTNER, “Screen bundles of Lorentzian manifolds and some generalisations of pp-waves”, *J. Geom. Phys.* **56** (2006), no. 10, p. 2117-2134.
- [12] T. LEISTNER & D. SCHLIEBNER, “Completeness of compact Lorentzian manifolds with Abelian holonomy”, <http://arxiv.org/abs/1306.0120>, 2013.
- [13] A. MASON, “An application of stochastic flows to Riemannian foliations”, *Houston J. Math.* **26** (2000), no. 3, p. 481-515.
- [14] E. E. MOISE, “Affine structures in 3-manifolds. V. The triangulation theorem and Hauptvermutung”, *Ann. of Math. (2)* **56** (1952), p. 96-114.
- [15] P. MOLINO & V. SERGIESCU, “Deux remarques sur les flots riemanniens”, *Manuscripta Math.* **51** (1985), no. 1-3, p. 145-161.
- [16] P. PETERSEN, *Riemannian geometry*, second ed., Graduate Texts in Mathematics, vol. 171, Springer, New York, 2006, xvi+401 pages.
- [17] G. REEB, “Sur la courbure moyenne des variétés intégrales d’une équation de Pfaff $\omega = 0$ ”, *C. R. Acad. Sci. Paris* **231** (1950), p. 101-102.
- [18] J. I. ROYO PRIETO, “The Gysin sequence for Riemannian flows”, in *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, Contemp. Math., vol. 288, Amer. Math. Soc., Providence, RI, 2001, p. 415-419.
- [19] R. SHARPE, *Differential geometry: Cartan’s generalization of Klein’s Erlangen program*. Foreword by S. S. Chern., Berlin: Springer, 1997 (English), xix + 421 pages.
- [20] P. TONDEUR, *Geometry of foliations*, Monographs in Mathematics, vol. 90, Birkhäuser Verlag, Basel, 1997, viii+305 pages.
- [21] F. WALDHAUSEN, “On irreducible 3-manifolds which are sufficiently large”, *Ann. of Math. (2)* **87** (1968), p. 56-88.
- [22] C. T. C. WALL, *Surgery on compact manifolds*, second ed., Mathematical Surveys and Monographs, vol. 69, American Mathematical Society, Providence, RI, 1999, Edited and with a foreword by A. A. Ranicki, xvi+302 pages.

Manuscrit reçu le 24 juin 2014,
révisé le 30 septembre 2014,
accepté le 23 octobre 2014.

Daniel SCHLIEBNER
Humboldt-Universität zu Berlin
Institut für Mathematik,
Rudower Chaussee 25
D-12489 Berlin (Germany)
schliebn@mathematik.hu-berlin.de