

# ANNALES DE L'I. H. P., SECTION A

YVONNE CHOQUET BRUHAT

**Global existence theorems for hyperbolic harmonic maps**

*Annales de l'I. H. P., section A*, tome 46, n° 1 (1987), p. 97-111

[http://www.numdam.org/item?id=AIHPA\\_1987\\_\\_46\\_1\\_97\\_0](http://www.numdam.org/item?id=AIHPA_1987__46_1_97_0)

© Gauthier-Villars, 1987, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
http://www.numdam.org/*

---

# Global existence theorems for hyperbolic harmonic maps

by

**Yvonne CHOQUET BRUHAT**

Institut de Mécanique, Université Paris VI

---

## INTRODUCTION

Harmonic maps between properly riemannian manifolds have a long mathematical history, with global results of great interest in geometry and physics (cf. [1] [2] [16]). The harmonic maps with source a riemannian manifold of hyperbolic (lorentzian) signature have been studied more recently, however such maps appear in numerous problems of physics, from the harmonic gauge of General Relativity to the non-linear  $\sigma$  models and many others, as pointed out in [15] [17] and [14].

The natural problem for such «hyperbolic harmonic maps» is the Cauchy problem, that is the determination of the map from its value, and the value of its first derivative, on a space like submanifold of the source. A local existence theorem of a solution of the problem for harmonic maps from  $M^{n+1}$ , Minkowski space time of arbitrary dimension, into various compact riemannian manifold has been proved by Ginibre and Velo [10], using I. Segal's theory of non linear semi-group [12]. They also prove for such problems a global existence theorem when  $n = 1$ , by using energy estimates. A global existence from  $M^2$  into a complete riemannian has been proved, by a different method, for smooth data by Gu Chao Hao [11].

In this article we prove a local existence theorem for harmonic maps from a globally hyperbolic manifold  $(M, g)$  into a riemannian manifold  $(N, h)$  both arbitrary except for some regularity conditions. The proof uses the embedding of  $(N, h)$  in an euclidean space  $(\mathbb{R}^q, e)$ , like Ginibre and Velo [10]. Another proof, which used only the standard local existence and uniqueness results for hyperbolic equations had been indicated in [3], which treated harmonic gauges in General Relativity.

We prove in § 5 a global existence theorem in the case  $n = 1$ , using the second order equation satisfied by the differential of an harmonic mapping and, like Ginibre and Velo, the resulting *a priori* estimates.

In § 6 we prove a global existence theorem for  $(M, g) = M^{n+1}$ , with  $n$  odd, arbitrary if the Cauchy data are sufficiently near from those of a constant map. We use the method of conformal transformation as in [6] [7] and [8]. The theorem is valid for  $n = 1$ , because of conformal invariance, and for  $n = 3$  because the operator satisfies an analogue of the condition indicated as sufficient by Christodoulou in [8] (which treats scalar systems).

It results from counter examples constructed by Shatah [9] that this last theorem cannot be true for arbitrary, large, data.

## 1. DÉFINITIONS

Let  $(M, g)$  and  $(N, h)$  be two smooth riemannian manifolds of arbitrary signature and dimensions. Let

$$f : M \rightarrow N$$

be a smooth map. The *differential* of  $f$  at  $x \in M$  is a linear map

$$\nabla f(x) : T_x M \rightarrow T_{f(x)} N$$

it is therefore an element of  $T_x^* M \otimes T_{f(x)} N$ . The differential itself,  $\nabla f$  is a mapping  $x \rightarrow \nabla f(x)$ , that is a section of the vector bundle with base  $M$  and fiber at  $x$  the vector space  $T_x^* M \otimes T_{f(x)} N$ . This vector bundle—one forms on  $M$  with values at  $x$  in  $T_{f(x)} N$ —is denoted  $T^* M \otimes f^{-1} N$ . The vector bundle with base  $M$  and fiber  $T_{f(x)} N$  at  $x$  is denoted  $f^{-1} TN$ . If  $(x^\alpha)$  and  $(y^\alpha)$  are respectively local coordinates in  $M$  and  $N$ , and  $f$  is represented in these coordinates by

$$y^\alpha = f^\alpha(x^\beta)$$

the derivative  $f$  is represented by

$$(x^\alpha) \rightarrow \left( \frac{\partial f^\alpha}{\partial x^\beta}(x^\beta) \right).$$

The metrics  $g$  on  $M$  and  $h$  on  $N$  endow the fiber at  $x$  of the vector bundle  $E = T^* M \otimes f^{-1} TN$  with a scalar product  $G(x) = g^\#(x) \otimes h(f(x))$ , where  $g^\#$  is the contravariant tensor canonically associated with  $g$ . In coordinates, if  $u$  and  $v$  are two sections of  $E$  :

$$(1) \quad G(x)(u, v) = g^{\alpha\beta}(x^\lambda) h_{ab}(f^c(x^\lambda)) u_a^\alpha(x^\lambda) v_b^\beta(x^\lambda).$$

The vector bundle  $E \equiv T^* M \otimes f^{-1} TN$  is endowed with a linear connection  $\nabla$ , mapping sections of  $E$  into sections of  $T^* M \otimes E$ , by the usual rules: if  $s$  is a section of  $f^{-1} TN$  and  $t$  a section of  $T^* M$  we have:

$$\nabla_v(t \otimes s) = {}^g \nabla_v t \otimes s + t \otimes f^* {}^h \nabla_v s$$

with  ${}^g\nabla$  and  ${}^h\nabla$  the riemannian covariant derivatives in the metrics  $g$  and  $h$  respectively. In local coordinates if  $(x^\alpha) \mapsto (u^\alpha(x^\alpha))$  is a section of  $E$ , we have:

$$(2) \quad \nabla_\alpha u_\beta^\alpha(x^\lambda) = \partial_\alpha u_\beta^\alpha(x^\lambda) + \frac{\partial f^b}{\partial x^\alpha} \Gamma_b{}^\alpha_c(f^d(x^\lambda)) u_\beta^c(x^\lambda) - \Gamma_\alpha{}^\mu_\beta(x^\lambda) u_\mu^\alpha(x^\lambda)$$

where  $\Gamma_b{}^\alpha_c$  and  $\Gamma_\alpha{}^\lambda_\beta$  denote respectively the riemannian connexions of  $g$  and  $h$ . The mapping  $f$  is called *harmonic* if

$$(3) \quad \text{tr}_g \nabla^2 f = 0$$

that is, in local coordinates

$$g^{\alpha\beta} \nabla_\alpha \partial_\beta f^\alpha \equiv g^{\alpha\beta} (\partial_{\alpha\beta}^2 f^\alpha - \Gamma_\alpha{}^\lambda_\beta \partial_\lambda f^\alpha + \Gamma_b{}^\alpha_c \partial_\alpha f^b \partial_\beta f^c) = 0.$$

If  $f$  satisfies (3) it is a critical point of the functional

$$f \mapsto E(f) = \int_M G(\nabla f, \nabla f) d\mu(g) = \int_M g^{\alpha\beta}(x^\lambda) h_{ab}(f^c(x^\lambda)) \partial_\alpha f^a \partial_\beta f^b d\mu(g).$$

## 2. HYPERBOLIC HARMONIC MAPS. ENERGY INTEGRAL

When the metrics  $g$  and  $h$  are properly riemannian the integral (4) is called the energy of the mapping  $f$ . When  $g$  is of hyperbolic signature we will define another integral as the energy of  $f$ , like for usual wave equations.

We define the stress energy tensor of the map  $f$  as the covariant 2-tensor on  $M$  given by

$$T = (h \circ f)(\nabla f, \nabla f) - \frac{1}{2} g(g^\# \otimes h \circ f)(\nabla f \otimes \nabla f)$$

that is

$$(2.1) \quad T_{\alpha\beta} = (h_{ab} \circ f) \partial_\alpha f^a \partial_\beta f^b - \frac{1}{2} g_{\alpha\beta} g^{\lambda\mu} (h_{ab} \circ f) \partial_\lambda f^a \partial_\mu f^b$$

We have on  $M$

$$(2.2) \quad \nabla_\alpha T^\alpha_\beta \equiv (h_{ab} \circ f) \partial_\beta f^a \nabla^\alpha \partial_\alpha f^b$$

that is

$$\nabla \cdot T = (h \circ f)(\nabla f, \text{tr}_g \nabla^2 f).$$

Thus  $\nabla \cdot T = 0$  if  $f$  is a harmonic map.

We suppose that  $(M, g)$  is a hyperbolic manifold with  $M = S \times \mathbb{R}$ ,  $S_t \equiv S \times \{t\}$  space-like, we denote by  $n$  their unit time like normal. Let  $X$  be a time like vector field. We define the energy density of  $f$  relative to  $S_t$  and  $X$  by:

$$e(f) = T(X, n) = X^\alpha n^\beta T_{\alpha\beta}$$

we have

$$(2.3) \quad e(f) = \frac{1}{2} \gamma^{\alpha\beta} (h_{ab} \circ f) \partial_\alpha f^a \partial_\beta f^b = \frac{1}{2} \gamma^\# \otimes (h \circ f)(\nabla f, \nabla f)$$

where  $\gamma^*$  is the quadratic form

$$\gamma^{\alpha\beta} = n^\alpha X^\beta + n^\beta X^\alpha - g^{\alpha\beta} X^\lambda n_\lambda.$$

It is well known that this form is positive definite if  $g$  of hyperbolic signature  $(+ - - \dots)$  with  $X$  and  $n$  time like.

We deduce from

$$\nabla_\alpha T^{\alpha\beta} = 0$$

when  $f$  is a harmonic map that

$$(2.4) \quad \nabla_\alpha (X_\beta T^{\alpha\beta}) = \frac{1}{2} T^{\alpha\beta} (\nabla_\alpha X_\beta + \nabla_\beta X_\alpha).$$

By integration of 2.4 on  $S \times [0, t]$  we obtain the following:

**PROPOSITION.** — Let  $f$  be a smooth map such that  $\nabla f|_{S_t}$  has a compact support for  $0 \leq \tau \leq t$  then, if  $f$  is harmonic it satisfies the identity:

$$(2.5) \quad \int_{S_t} N e(f) d\mu_t \equiv \int_{S_0} N e(f) d\mu_0 + \frac{1}{2} \int_0^t \int_{S_\tau} N (T \cdot L X) d\mu_\tau$$

$d\mu_t$  denotes the volume element of the metric  $\bar{g}_t$  induced on  $S_t$  by  $g$ ,  $N$  is the lapse function,  $N = g(X, n)$ , that is  $N = (g^{00})^{-1/2}$  if  $X$  is the tangent vector to the curves  $\{x\} \times \mathbb{R}$  and the coordinates are adapted to the product  $S \times \mathbb{R}$ : the volume element of  $(M, g)$  is

$$d\mu(g) = N d\mu_{x^0} dx^0.$$

**DEFINITION 1.** — The manifold  $(M, g)$  is said *regularly hyperbolic* if

1)  $M$  and  $g$  are smooth and  $M = S \times \mathbb{R}$ , the metrics  $\bar{g}_t$  induced on  $S_t = S \times \{t\}$  are (properly) riemannian <sup>(1)</sup>, and uniformly equivalent to the metric  $\bar{g}_0$  which is complete.

2) The tangent vector  $X$  to the lines  $\{x\} \times \mathbb{R}$  is time like, and there exists strictly positive numbers  $a$  and  $b$  such that

$$\inf_M g(X, X) \geq a \geq 0 \quad \text{and} \quad \sup_M N \leq b$$

we then have also, since  $N = g(X, n) \geq (g(X, X))^{1/2}$

$$\inf_M N \geq a^{1/2} \quad \text{and} \quad \sup_M g(X, X) \leq b^2.$$

If  $(M, g)$  is regularly hyperbolic the metric  $\gamma$  on  $M = S \times \mathbb{R}$  defined by 2.3 is uniformly equivalent to the metric  $\Gamma = (dx^0)^2 - \bar{g}_0$ .

We shall then add to the definition of regular hyperbolicity the following.

3) The riemann curvature of  $g$ , together with as many of its covariant derivatives as is relevant, is bounded in  $\Gamma$ -norm.

<sup>(1)</sup> These metrics are negative definite:  $-\bar{g}_t$  is positive definite.

**DEFINITION 2.** — A (properly) riemannian manifold is said to be regular if it is smooth, has a non zero injectivity radius (thus is complete), and has a bounded riemannian curvature, as well as its covariant derivatives up to the relevant order.

Tensor products of the metrics  $\Gamma$  and  $h$  give scalar products and norms in the fiber at  $x \in M$  of vector bundles  $(\otimes T_x M)^p (\otimes f^{-1}TN)^q$  or their duals. We denote this norm by  $| \cdot |$ . We have, for instance,

$$|\nabla f|^2 = \Gamma^{\alpha\beta}(h_{ab} \circ f) \partial_\alpha f^\alpha \partial_\beta f^\beta.$$

If  $s$  and  $u$  are two sections of such vector bundles we have, at a point  $x \in M$

$$|s \otimes u| = |s| |u|, \quad |s \cdot u| \leq |s| |u|$$

if  $s \cdot u$  is some contracted tensor product

Therefore, in particular

$$|LX \cdot T| \leq |LX| |T|.$$

It results from the definition that  $|g|$  and  $|g^\#|$  are uniformly bounded if  $(M, g)$  is regularly hyperbolic (cf. [5]). Thus, due to the expression of  $T$ , there exists a constant  $C$  such that

$$(2.6) \quad |T| \leq Ce(f)$$

and also, if  $|X|$  is uniformly bounded on  $M$ , a constant still denoted  $C$  such that

$$|LX \cdot T| \leq Ce(f).$$

From the equality (2.5) results then the inequality ( $C_0$  and  $C$  positive constants)

$$(2.7) \quad y(t) \leq C_0 y(0) + C \int_0^t y(\tau) d\tau$$

with

$$y(t) = \int_{S_t} |\nabla f|^2 d\mu_0.$$

If  $y$  is a continuous function of  $t$  we deduce from (2.7), by the Gromwall lemma

$$(2.8) \quad y(t) \leq K(t)y(0)$$

with  $K(t)$  the continuous function of  $t$

$$K(t) = C_0 e^{Ct}.$$

### 3. SECOND ORDER EQUATION FOR $f$

**PROPOSITION.** — Every smooth harmonic map  $f: (M, g) \rightarrow (N, h)$  satisfies the equation

$$(\nabla \cdot \nabla)(\nabla f) - \text{Ricc}(g)\nabla f + \text{tr}_g(f^* \text{Riem}(h) \cdot \nabla f) = 0$$

that is, in local coordinates

$$(3.1) \quad \nabla^\lambda \nabla_\lambda \partial_\alpha f^\alpha - R_\alpha^\beta \partial_\beta f^\alpha + R_{cd}{}^a_b \partial_\alpha f^\alpha \partial_\beta f^\beta \partial_\mu f^\mu = 0.$$

The proof, independant of signature, is straightforward and given in [1]. We now consider the case  $g$  hyperbolic and  $h$  properly riemannian. We set ( $h_{ab}$  stands always for  $h_{ab} \circ f$ )

$$(3.2) \quad T_{\alpha\beta}^{(1)} = e^{\lambda\mu} h_{ab} \left\{ \nabla_\alpha \partial_\lambda f^\alpha \nabla_\beta \partial_\mu f^\beta - \frac{1}{2} g_{\alpha\beta} g^{\rho\sigma} \nabla_\rho \partial_\lambda f^\alpha \nabla_\sigma \partial_\mu f^\beta \right\}$$

We have identically, after use of the Ricci identity

$$(3.3) \quad \begin{aligned} \nabla_\alpha T_{(1)}^{\alpha\beta} &\equiv e^{\lambda\mu} h_{ab} \nabla^\alpha \nabla_\alpha \partial_\lambda f^\alpha \nabla_\beta \partial_\mu f^\beta \\ &+ e^{\lambda\mu} h_{ab} \nabla^\alpha \partial_\lambda f^\alpha (-R_{\alpha\beta}{}^\mu \partial_\mu f^\beta + \partial_\alpha f^\alpha \partial_\beta f^\beta \partial_\mu f^\mu R_{cd}{}^b{}_e) \\ &+ h_{ab} (\nabla^\alpha e^{\lambda\mu}) \left\{ \nabla_\alpha \partial_\lambda f^\alpha \nabla_\beta \partial_\mu f^\beta - \frac{1}{2} g_{\alpha\beta} g^{\rho\sigma} \nabla_\rho \partial_\lambda f^\alpha \nabla_\sigma \partial_\mu f^\beta \right\}. \end{aligned}$$

Using (3.1) we see that, for a harmonic map,  $\nabla_\alpha T_{(1)}^{\alpha\beta}$  is a polynomial  $Q(f, \nabla f, \nabla^2 f)$  of degree 2 in  $\nabla^2 f$ , with coefficients of degree 1 or 3 [respectively 0] in  $\nabla f$  for the terms of degree 1 [respectively 2] in  $\nabla^2 f$ . The mapping  $f$  itself appears through  $h \circ f$  and  $\text{Riem}(h) \circ f$ .

On the other hand we have:

$$e_{1,f} \equiv T_{\alpha\beta}^{(1)} X_\beta n^\beta = \frac{1}{2} \gamma^{\alpha\beta} \gamma^{\lambda\mu} h_{ab} \nabla_\alpha \partial_\lambda f^\alpha \nabla_\beta \partial_\mu f^\beta \geq 0.$$

Integrating the identity:

$$\nabla_\alpha (X_\beta T_{(1)}^{\alpha\beta}) \equiv X_\beta \nabla_\alpha T_{(1)}^{\alpha\beta} + \frac{1}{2} T_{(1)}^{\alpha\beta} (\nabla_\alpha X_\beta + \nabla_\beta X_\alpha)$$

with  $T_{\alpha\beta}^{(1)}$  and  $\nabla_\alpha T_{(1)}^{\alpha\beta}$  given by (3.2) and (3.3) gives for a harmonic map, with compact support in space, using (3.1), an equality:

$$(3.4) \quad \int_{S_t} N e_{1,f} d\mu(\bar{g}_t) = \int_{S_0} N e_{1,f} d\mu(\bar{g}_0) + \int_0^t \int_{S_\tau} N Q_1(f, \nabla f, \nabla^2 f) d\mu(\bar{g}_\tau) d\tau$$

with  $Q_1$  of the type

$$Q_1(f, \nabla f, \nabla^2 f) \equiv \Sigma k \nabla^2 f \cdot \{ \text{Riem}(g) \cdot \nabla f + \text{Riem}(h) \circ f \cdot (\otimes \nabla f)^3 + \nabla^2 f \}$$

with  $k$  polynomial in  $g^\#$ ,  $h$ ,  $X$  and  $\nabla X$ .

If  $(M, g)$  is regularly hyperbolic, and  $(N, h)$  regularly riemannian we deduce from (3.4) an inequality, as in § 2; with  $C_0, C_1, C_2, C_3$  positive constants:

$$(3.5) \quad y_1(t) \leq C_0 y_1(0) + C_1 \int_0^t y_1(\tau) d\tau + \int_0^t \int_{S_\tau} (C_2 |\nabla^2 f| |\nabla f| + C_3 |\nabla^2 f| |\nabla f|^3) d\mu_0 d\tau$$

where, by definition

$$y_1(t) = \int_{S_t} |\nabla^2 f|^2 d\mu_0$$

and

$$|\nabla^2 f|^2 = e^{\lambda\mu} e^{\alpha\beta} (h_{ab} \circ f) \nabla_\alpha \partial_\lambda f^a \nabla_\beta \partial_\mu f^b.$$

#### 4. LOCAL EXISTENCE

Let  $N$  be a submanifold of the riemannian manifold  $(Q, q)$  and  $h$  be the metric induced on  $N$  by  $q$ . We shall suppose that  $N$  is defined by  $p$  equations

$$N : \Phi^I(z) = 0, \quad z \in Q, \quad I = 1, \dots, p$$

where  $\phi = (\Phi^I) : Q \rightarrow \mathbb{R}^p$  is a smooth map of rank  $p$  at each point of  $N$ . The matrix  $m = (m^{IJ})$  given by

$$m = q(\nabla\phi, \nabla\phi), \quad \text{i. e. } m^{IJ} = (q^{AB} \partial_A \phi^I \partial_B \phi^J) \circ f$$

( $x^A$  coordinates in  $Q$ ) is then positive definite on  $M$  when  $f$  takes its values in  $N$ . We denote by  $m^{-1} = m_{IJ}$  the inverse matrix.

**LEMMA 1.** — A necessary and sufficient condition for the mapping  $f : M \rightarrow N \subset Q$  to be a harmonic map from  $(M, g)$  into  $(N, h)$  is that, as a mapping  $M \rightarrow Q$  it satisfies the equations which read in local coordinates  $x^\alpha$  in  $M$  and  $x^A$  in  $Q$ :

$$(4.1) \quad \hat{\nabla}^\alpha \nabla_x f^A + \lambda_I (q^{AB} \partial_B \Phi^I) \circ f = 0$$

where  $(\hat{\nabla}^\alpha \nabla_x f^A)$  is the tension field of the map  $f : (M, g) \rightarrow (Q, q)$  and

$$(4.2) \quad \lambda_I = m_{IJ} g^{\alpha\beta} \partial_\alpha f^A \partial_\beta f^B (\hat{\nabla}_B \partial_A \phi^J) \circ f$$

together with the conditions

$$\phi \circ f = 0.$$

*Proof* (cf. a particular case in Ginibre and Velo [10]).

A mapping  $f : M \rightarrow N$  defines a mapping  $F : M \rightarrow Q$  by

$$(4.3) \quad F = i \circ f$$

where  $i$  denotes the embedding (identity map)  $N \rightarrow Q$ .

The integral constructed with  $F : M \rightarrow Q$

$$E(F) = \int_M (g^\# \otimes q)(\nabla F, \nabla F) d\mu(g)$$

is equal to the integral (1.4) constructed with  $f$  since

$$\nabla F = \nabla i \cdot \nabla f, \quad \text{i. e. } \partial_\alpha F^A = \partial_\alpha i^A \partial_\alpha f^A$$

and  $h$  is the metric induced by  $i$  on  $N$ ; that is  $h_{ab} = q_{AB}\partial_a i^A \partial_b i^B$ , and

$$(4.4) \quad E(f) = E(F).$$

Thus a critical point of  $E(f)$  is a critical point of  $E(F)$  with the constraint  $\phi(F) = 0$ , that is a solution of equations of the form:

$$(4.5) \quad \hat{V}^\alpha \nabla_\alpha F^A + \lambda_I q^{AB} \partial_B \Phi^I = 0$$

where the  $\lambda_I$  (Lagrange multipliers) are determined by derivating twice the conditions

$$\Phi^I \circ F = 0$$

with  $\Phi^I \circ F$  considered as a mapping  $M \rightarrow Q \rightarrow \mathbb{R}^p$ , and contracting with  $g$ :

$$(4.6) \quad \nabla^\alpha \nabla_\alpha (\Phi^I \circ f) \equiv \partial_A \Phi^I \hat{V}^\alpha \partial_\alpha F^A + g^{\alpha\beta} \partial_\alpha F^A \partial_\beta F^B \nabla_B \partial_A \Phi^I = 0$$

comparing (4.5) and (4.6) gives

$$(4.7) \quad \lambda_I q^{AB} \partial_B \Phi^I = g^{\alpha\beta} \partial_\alpha F^A \partial_\beta F^B \nabla_B \partial_A \Phi^I$$

(4.7) is equivalent to (4.2) ( $F = i \circ f = f$ , since  $i$  is the identity mapping  $N \rightarrow N \subset Q$ ).

**LEMMA 2.** — The equations (4.1) satisfied by a mapping  $M \rightarrow Q$ , with  $\lambda_I$  given by (4.2), imply that the mapping  $\phi \circ f : M \rightarrow \mathbb{R}^p$  satisfies the homogeneous wave equation on  $M$

$$(4.8) \quad \nabla^\alpha \nabla_\alpha (\phi \circ f) = 0.$$

*Proof.* — (4.6) implied by (4.1) and (4.2).

**DEFINITION 3.** — A submanifold  $N$  of  $\mathbb{R}^q$  given by  $N = \{y \in \mathbb{R}^q, \phi(y)=0\}$  with  $\phi$  a smooth map  $\mathbb{R}^q \rightarrow \mathbb{R}^p$  is said to be regularly defined by  $\phi$  if there exists  $a > 0$  and  $\varepsilon > 0$  such that

$$\inf_{y \in N_\varepsilon} |\det m^{IJ}(y)| \geq a > 0, \quad N_\varepsilon = \{y \in \mathbb{R}^q, d(y, N) < \varepsilon\}$$

$d$  denotes the euclidean distance. The definition means that  $\phi$  is uniformly of rank  $p$  in some uniform neighbourhood of  $N$ .

**DEFINITION 4.** — We denote by  $H_s(S)$  the Sobolev space of  $\mathbb{R}^q$ -valued functions on the regularly riemannian manifold  $(S, \bar{g}_0)$ , closure of  $C^\infty$ ,  $\mathbb{R}^q$  valued functions with compact support on  $S$  in the norm:

$$\|\varphi\|_{H_s}^2 = \int_S \sum_{k=0}^s |D^k \varphi|^2 d\mu_0$$

where  $D$  is the covariant derivative in the metric  $\bar{g}_0$  for each scalar valued

map  $\varphi^A : S \rightarrow \mathbb{R}$ , and  $| \cdot |_e$  is the  $\bar{g}_0$  and  $e$  norm of the set  $D^k\varphi = (D^k\varphi^A)$ , for instance

$$| D^2\varphi |_e^2 = e_{AB} \bar{g}_0^{ij} \bar{g}_0^{hk} D_{ih}^2 \varphi^A D_{jk}^2 \varphi^B, \quad e_{AB} = \delta_{AB}.$$

**THEOREM (local existence).** — Let  $(M, g)$ ,  $M = S \times \mathbb{R}$ , be a regularly hyperbolic manifold of dimension  $n + 1$  (definition 1).

Let  $(N, h)$  be a regular riemannian manifold, regularly defined by a mapping  $\phi : \mathbb{R}^q \rightarrow \mathbb{R}^p$ . Let  $\varphi, \dot{\varphi}$  be mappings  $S \rightarrow \mathbb{R}^q$ ,  $\varphi \in H_s(S)$ ,  $\dot{\varphi} \in H_{s-1}(S)$ ,  $s > \frac{n}{2} + 1$  such that  $\Phi \circ \varphi = 0$ ,  $(\nabla \phi \circ \varphi) \cdot \dot{\varphi} = 0$ .

Then there exists  $l > 0$  and on  $S \times (-l, l)$  a harmonic map  $f : (M, g) \rightarrow (N, h)$ , with  $h$  the metric induced on  $N$  by the euclidean metric  $e$  of  $\mathbb{R}^q$ , such that

$$(4.9) \quad f|_{S_0} = \varphi, \quad \partial_0 f|_{S_0} = \dot{\varphi}.$$

*Proof.* — We apply lemma 1 with  $(Q, q) = (\mathbb{R}^q, e)$ . Equations (4.1) reads then:

$$(4.10) \quad \nabla^\alpha \nabla_\alpha f^A + (\nabla \phi^T \nabla \phi)^{-1}_{IJ}(f) g^{\alpha\beta} \partial_\alpha f^A \partial_\beta f^B (\partial_{AB}^2 \Phi^J)(f) = 0;$$

they are a system of  $q$ , numerical, quasi-linear, quasi-diagonal second order hyperbolic equations on  $M$ , with smooth coefficients if  $d(f, N) < \varepsilon$ . The local existence theorem, on  $S \times (-l, l)$  is a standard result, since  $d(\varphi, N) = 0$ . The solution  $f : S \times (-l, l) \rightarrow \mathbb{R}^q$  satisfies  $\phi \circ f = 0$  because  $\phi \circ f$  satisfies the homogeneous wave equation (4.8) with zero Cauchy data:

$$\phi \circ f|_S = \phi \circ \varphi|_{=0}, \quad \partial_0(\phi \circ f)|_S = (\nabla \phi \circ f) \cdot \partial_0 f|_S = (\nabla \phi \circ \varphi) \cdot \dot{\varphi} = 0.$$

**REMARK 1.** — The local existence theorem for a numerical hyperbolic system gives that the interval of existence depends continuously on the  $H_{s_0} \times H_{s_0-1}$  norm,  $s_0$  smallest integer such that  $s_0 > \frac{n}{2} + 1$ , of the Cauchy data, and tends to infinity when these norms tend to zero. If  $N$  is a submanifold of  $\mathbb{R}^q$ , we can always, by translation, take the origin of  $\mathbb{R}^q$  at some arbitrary given point  $y_0$  of  $N$ . The  $H_s(S_0)$  norm of a map  $\varphi : S_0 \rightarrow \mathbb{R}^q$  is by definition

$$\| \varphi \|_s = \left\{ \sum_{k=0}^s \int_S |\nabla^k \varphi|^2 d\mu_0 \right\}^{1/2}$$

with

$$|\varphi|^2 = \sum_{A=1}^q |\varphi^A|^2.$$

Small  $H_s$  norm for  $\varphi$  means then nearness of  $\varphi$  from the constant map  $M \rightarrow y_0$ .

**REMARK 2.** — Every riemannian manifold  $(N, h)$  can be isometrically embedded in a space  $(\mathbb{R}^q, e)$ -we have supposed moreover that  $N$  is given by equations  $\Phi^I = 0$ . We could have proceeded without this hypothesis, either inspired by techniques used by Eells and Sampson in the elliptic case, either by using atlases on  $M$  and  $N$ , together with local existence and uniqueness theorems (cf. an indication of such a proof in [3]).

## 5. GLOBAL EXISTENCE WHEN $n = 1$

The global existence of a solution of the Cauchy problem for harmonic maps from 2-dimensional Minkowski space time  $M^2$  into a complete riemannian manifold  $(N, h)$  has been proved by Gu Chao Hao [11] for smooth initial data. It has been proved by Ginibre and Velo [10] from the two dimensional Minkowski space into various compact riemannian manifolds for  $H_2 \times H_1$  Cauchy data. This result can be generalized:

**THEOREM.** — Let  $(M, g)$ ,  $M = S \times \mathbb{R}$ , be a regularly hyperbolic manifold of dimension 2, and  $(N, h)$  be a regular riemannian manifold, regularly defined as a submanifold of  $(\mathbb{R}^q, e)$  by mapping  $\phi : \mathbb{R}^q \rightarrow \mathbb{R}^p$ ,  $N = \{y \in \mathbb{R}^q, \phi(y) = 0\}$ .

Let

$$\varphi \in H_s(S), \quad \dot{\varphi} \in H_{s-1}(S), \quad s \geq 2$$

be given maps  $\varphi : S \rightarrow \mathbb{R}^q$ ,  $\dot{\varphi} : S \rightarrow \mathbb{R}^q$ , such that  $\phi \circ \varphi = 0$ ,  $(\nabla \phi \circ \varphi) \cdot \dot{\varphi} = 0$ . Then there exists on  $M$  a harmonic map  $f : (M, g) \rightarrow (N, h)$  taking on  $S_0 = S \times \{0\}$  these Cauchy data.

*Proof.* — In the case  $n = 1$  the local existence theorem is valid with  $s \geq 2$ . The solution  $f : S \times (-l, l) \rightarrow N \subset \mathbb{R}^q$  admits a restriction on each  $S_t = S \times \{t\}$ ,  $|t| < l$  which is a mapping  $f_t : S_t \rightarrow N \subset \mathbb{R}^q$  which belongs to  $H_2(S)$  (definition 4). The derivative  $\partial_0 f$  admits a restriction  $(\partial_0 f)_t : S_t \rightarrow f^{-1}TN$  by  $x \rightarrow T_{f_t(x)}N \subset \mathbb{R}^q$ ,  $(\partial_0 f)_t \in H_1(S)$ . The energy inequality (2.9) implies, when  $f = i \circ f$  is considered as a mapping into  $N$  :

$$(5.1) \quad y(t) \leqq K(t)y(0)$$

with

$$y(t) = \int_S |\nabla f|_h^2 d\mu_0$$

with, due to the definition of  $\Gamma$ :

$$(5.2) \quad |\nabla f|_h^2 = (h_{ab} f_a)(\hat{c}_0 f^a \hat{c}_0 f^b - \bar{g}_0^{ij} \hat{c}_i f^a \hat{c}_j f^b)$$

but we have, since  $f = i \circ f$  and  $h = i^* e$

$$(5.3) \quad |\nabla f|_h^2 = |\nabla f|_e^2 = e_{AB} (\hat{c}_0 f^A \hat{c}_0 f^B - \bar{g}_0^{ij} \partial_i f^A \partial_j f^B).$$

The inequality (2.9) implies therefore the non-blow up of the  $L^2(S, \bar{g}_0)$  norm of  $Df_t$  and of  $(\partial_0 f)_t$ , as mappings  $S \rightarrow \mathbb{R}^q$ , and thus also of  $f_t$ : the norms  $\|f\|_{H_1(S)}$  (and thus  $\|f_t\|_{C_0(S)}$ ) are bounded by continuous functions of  $t$  which extend to  $t = +\infty$ . To prove the non blow up of the second derivatives we look at the identity (3.4). Due to the regularity hypothesis we have, with  $C$  a constant

$$(5.4) \quad \int_{S_t} Q_1 d\mu_t \leq C \int_{S_t} |\nabla^2 f|_h (|\nabla f|_h + |\nabla f|_h^3 + |\nabla^2 f|_h) d\mu_0$$

using the fact that  $f = i \circ f$  we find inequalities of the form ( $C_1$  and  $C_2$  positive constants)

$$(5.5) \quad \begin{aligned} |\nabla^2 f|_h^2 &\leq |\hat{\nabla}^2 f|_e^2 + C_1 |\nabla f|^4 \\ |\hat{\nabla}^2 f|_e^2 &\leq |\nabla^2 f|_h^2 + C_2 |\nabla f|^4 \end{aligned}$$

(recall that  $\hat{\nabla}$  denotes the covariant derivative of  $f$  as a mapping  $(M, g) \rightarrow (\mathbb{R}^q, e)$  and  $\nabla$  as mapping  $M \rightarrow N$ ).

We deduce from (5.4), by the Cauchy-Schwartz inequality

$$(5.6) \quad \begin{aligned} \int_{S_t} Q_1 d\mu_t &\leq C \int_{S_t} |\nabla^2 f|_h^2 d\mu_0 \\ &+ \left\{ \int_{S_t} |\nabla^2 f|_h^2 d\mu_0 \right\}^{1/2} \left\{ \left( \int_{S_t} |\nabla f|^2 d\mu_0 \right)^{1/2} + \left( \int_{S_t} |\nabla f|^6 d\mu_0 \right)^{1/2} \right\}. \end{aligned}$$

Considering  $f$  as mapping  $M \rightarrow \mathbb{R}^q$ , and setting

$$z(t) = \int_{S_t} |\hat{\nabla}^2 f|_e^2 d\mu_0$$

we obtain, using (3.5) (5.5) and (5.6), with  $C$  some constant

$$z(t) \leq C_0 y_1(0) + C \int_0^t \left\{ z(\tau) + \int_{S_\tau} (|\nabla f|^2 + |\nabla f|^4 + |\nabla f|^6) d\mu_0 \right\} d\tau.$$

By the inequality (2.9) we know that

$$\int_{S_t} |\nabla f|^2 d\mu_0 = y(t) \leq K(\tau)y(0).$$

We bound the integrals  $\int_{S_t} |\nabla f|^4 d\mu_0$  and  $\int_{S_t} |\nabla f|^6 d\mu_0$  by using the following Sobolev inequality, valid in any dimension if  $a = \frac{n(p-1)}{p}$  and  $S$  admits a uniformly locally finite atlas:

$$\|u\|_{L^p(S)} \leq C \|Du\|_{L^1(S)}^2 \|u\|_{L^1(S)}^{1-a}$$

by taking

$$u = |\nabla f|^2.$$

If  $n = 1$ ,  $a = \frac{p-1}{p}$ ,  $p = 1, 2$  or  $3$  we have therefore:

$$\int_{S_t} |\nabla f|^{2p} d\mu_0 = (\|u\|_{L^p})^p \leq C \|Du\|_{L^1(S)}^{(p-1)/p} \|u\|_{L^1(S)}^{1/p}$$

we have

$$\|u\|_{L^1(S)} = y(t)$$

and

$$\|Du\|_{L^1(S)}^2 \leq Cy(t)z(t).$$

We then obtain an integral inequality for  $z(t)$ , with coefficients continuous functions of  $t$  extendable for all  $t$ , and at most of degree one in  $z(t)$ . The non-blow up of  $z(t)$  follows.

## 6. GLOBAL EXISTENCE WHEN $(M, g) = M^{n+1}$ , SMALL DATA

Met  $M^{n+1} = (\mathbb{R}^{n+1}, \eta)$  be  $n + 1$  dimensional Minkowski space time.

Let  $\Sigma^{n+1} = (S^n \times \mathbb{R}, g)$  be the Einstein cylinder with its canonical metric.  $M^{n+1}$  is known to be conformal to a subset  $V$  of  $\Sigma^{n+1}$ , that is:

$$(6.1) \quad g = \Omega^2 \eta \quad \text{on} \quad V = \mathbb{R}^{n+1}$$

the identification of  $V \subset \Sigma^{n+1}$  with  $\mathbb{R}^{n+1}$  being given in canonical polar coordinates respectively  $(t, r, \dots)$  on  $\mathbb{R}^{n+1}$  and  $(T, \alpha, \dots)$  on  $\Sigma^{n+1}$  by (cf. [7])

$$(6.2) \quad \begin{aligned} T &= \operatorname{Arctg}(t+r) + \operatorname{Arctg}(t-r) \\ \alpha &= \operatorname{Arctg}(t+r) - \operatorname{Arctg}(t-r) \\ V : \alpha - \Pi < T < \Pi - \alpha \end{aligned}$$

We have

$$(6.3) \quad g = dT^2 - d\alpha^2 - \sin^2 \alpha d\sigma^2 = \Omega^2 \eta = \Omega^2 (dt^2 - dr^2 - r^2 g_{S^{n-1}})$$

$$\Omega = \cos T + \cos$$

$g_{S^{n-1}}$  metric of the sphere  $S^{n-1}$ .

We remark that  $\Omega$  extends to an analytic function on  $\Sigma^{n+1}$ , which vanishes on  $\partial V$ ; the submanifold  $t = 0$  (i.e.  $\mathbb{R}^n \times \{0\}$ ) is mapped diffeomorphically onto  $T = 0$  (i.e.  $S^n \times \{0\}$ ), minus its north pole  $\alpha = \Pi$ . On  $S^n \times \{0\}$  we have ( $\alpha = \Pi$  is  $r = +\infty$ )

$$\Omega|_{T=0} = 1 + \cos \alpha = 2(1 + r^2)^{-1}.$$

To a mapping  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^q$  corresponds, by the previous diffeomorphism a mapping, still denoted  $\varphi$ , defined almost everywhere on  $S^n$ .

**THEOREM.** — Let  $(N, h)$  be a riemannian submanifold of  $(\mathbb{R}^q, e)$  regularly defined by a smooth map  $\phi : \mathbb{R}^q \rightarrow \mathbb{R}^p$

$$N = \{ y \in \mathbb{R}^q, \phi(y) = 0 \}.$$

Let  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^q$  and  $\dot{\varphi} : \mathbb{R}^n \rightarrow \mathbb{R}^q$  be given mappings such that

$$(6.4) \quad \phi \circ \varphi = 0, \quad (\nabla \phi \circ \varphi) \cdot \dot{\varphi} = 0$$

with

$$(1 + \cos \alpha)\varphi \in H_s(S^n), \quad (1 + \cos \alpha)\dot{\varphi} \in H_{s-1}(S^n), \quad s > \frac{n}{2} + 1$$

then there exists a harmonic map  $f : M^{n+1} \rightarrow (N, h)$  such that

$$f|_{\mathbb{R}^n} = \varphi, \quad \delta_0 f|_{\mathbb{R}^n} = \dot{\varphi}$$

if the mappings  $\varphi$  and  $\dot{\varphi}$  are sufficiently near in the relevant norms respectively from a constant map and zero.

*Proof.* — It is inspired from the proofs of [7] and [8].

We do not restrict the generality by supposing that  $N$  passes through the origin of  $\mathbb{R}^q$ , that is  $\phi(0) = 0$ .

To a mapping  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^q$  we associate  $\tilde{f} : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^q$  defined on  $V$  by:

$$(6.5) \quad \tilde{f} = \Omega^{(1-n)/2} f.$$

We deduce then from (6.1), since the scalar curvature of  $\Sigma^{n+1}$  is  $n(n-1)$  that

$$(6.6) \quad \square_g \tilde{f} - \frac{(n-1)^2}{4} \tilde{f} = \Omega^{-(3+n/2)} \square_\eta f$$

where  $\square_g$  and  $\square_\eta$  are the wave operators in the metrics  $g$  and  $\eta$  respectively. We see that  $f$  satisfies (4.1) (4.2), with  $(Q, q) = (\mathbb{R}^q, e)$  that is

$$(6.6a) \quad \square_\eta f^A + \lambda_I [(\partial_A \phi^I) \circ f] = 0,$$

$$(6.7b) \quad \phi(f) = 0$$

with

$$\lambda_I = m_{IJ}(f) \eta^{\alpha\beta} \partial_\alpha f^A \partial_\beta f^B [(\partial_{AB}^2 \phi^I) \circ f], \quad m^{IJ} = \sum_{A=1}^q (\partial_A \phi^I \partial_B \phi^J) \circ f$$

if and only if

$$(6.8) \quad \square_g \tilde{f} - \frac{(n-1)^2}{4} \tilde{f} + \Omega^{-(3+n/2)} \lambda_I [(\partial_A \phi^I) \circ \Omega^{(n-1)/2} \tilde{f}] = 0$$

with

$$\lambda_I = m_{IJ} \Omega^2 g^{\alpha\beta} \partial_\alpha (\Omega^{(n-1)/2} \tilde{f}^A) \partial_\beta (\Omega^{(n-1)/2} \tilde{f}^B) [(\partial_{AB}^2 \phi^I) \circ (\Omega^{(n-1)/2} \tilde{f})]$$

and

$$\phi(\Omega^{(n-1)/2} \tilde{f}) = 0.$$

We have

$$\partial_\alpha (\Omega^{(n-1)/2} \tilde{f}^A) = \Omega^{(n-1)/2} \partial_\alpha \tilde{f}^A + (n-1)/2 \Omega^{(n-3)/2} \partial_\alpha \Omega \tilde{f}^A$$

$\nabla \Omega$  extends to a bounded function on  $\Sigma^{n+1}$ , and so does  $\Omega^{-1} g(\nabla \Omega, \nabla \Omega)$  (cf. [8]) since

$$g(\nabla \Omega, \nabla \Omega) = g^{\alpha\beta} \partial_\alpha \Omega \partial_\beta \Omega = \cos^2 \alpha - \cos^2 T = \Omega (\cos \alpha - \cos T).$$

Therefore the equation (6.8) extends to a semi-linear, semi-diagonal second order system with smooth coefficients for a mapping  $\tilde{f}$  from an open set  $U$  of  $\Sigma^{n+1}$  into  $\mathbb{R}^p$  if on the one hand  $n$  is odd and if, on the other hand,  $\tilde{f}$  is such that

$$d((\Omega^{(n-1)/2} \tilde{f})(X), N) < \eta \quad \forall X \in U$$

this last property will be *a fortiori* satisfied since  $0 \in N$  if

$$\sup_{X \in U} |\Omega^{(n-1)/2}(X) \tilde{f}(X)|_e < \eta$$

thus if

$$\sup_{X \in U} |\tilde{f}(X)|_e < \eta.$$

The existence of an open set  $U = S^n \times (-l, l)$  where the equation (6.8) has a solution  $\tilde{f}$  taking the Cauchy data:

$$\begin{aligned} \tilde{f}|_{S^n \times \{0\}} &= (1 + \cos \alpha)^{(1-n)/2} \varphi \\ \partial_0 \tilde{f}|_{S^n \times \{0\}} &= (1 + \cos \alpha)^{-(1+n)/2} \dot{\varphi} \end{aligned}$$

is then a consequence of the local existence theorem, and Sobolev inequalities, if  $s > \frac{n}{2} + 1$ . The length depends continuously on the norms of the Cauchy data, and we have  $l > \Pi$  if these norms are small enough.

The mapping  $f = \Omega^{(n-1)/2} f$  is defined on  $M^{n+1}$ , satisfies (4.1), and also (4.2) (lemma 2).

**REMARK.** — The hypothesis (6.4) on  $\varphi$  implies that  $\varphi$  tends to the constant map  $\mathbb{R}^n \rightarrow 0 \in N$ , when  $r$  tends to infinity.

From the theorem follow decay estimates for  $f$  on  $M^{n+1}$  (i. e. rate of approximating the constant map  $M^{n+1} \rightarrow 0 \in N$ ) when  $t$  or  $r$  tend to infinity.

## REFERENCES

- [1] J. EELLS and H. SAMPSON, *Amer. J. of Maths*, t. **86**, 1964, p. 109-160.
- [2] A. LICHNEROWICZ, *Symposia Mathematica*, vol. III, 1970, p. 341-402.
- [3] Y. CHOQUET-BRUHAT et C. GILAIN, *C. R. Ac. Sc. Paris*, t. **279**, 1974, p. 827.
- [4] J. LERAY, *Hyperbolic differential equations*, I. A. S. Princeton, 1952.
- [5] Y. CHOQUET-BRUHAT, D. CHRISTODOULOU et M. FRANCAVIGLIA, *Ann. I. H. P.*, Vol. XXXI, n° 4, 1979, p. 399-414.
- [6] D. CHRISTODOULOU, *C. R. Acad. Sc. Paris*, t. **292**, 1981, p. 139-141.
- [7] Y. CHOQUET-BRUHAT et D. CHRISTODOULOU, *Ann. E. N. S.*, t. **14**, 1981, p. 481-500.
- [8] D. CHRISTODOULOU, *Comm. in pure and appl. Maths.*, à paraître.
- [9] J. SHATAH, *Amer. Math. Soc. meeting*, New Orléans, 1986.
- [10] J. GINIBRE et G. VELO, *Ann. of Phys.*, t. **142**, n° 2, 1982, p. 393-415.
- [11] GU CHAO HAO, *Comm. on pure and applied maths XXXIII*, 1980, p. 727-737.
- [12] I. SEGAL, *Ann. of Maths*, t. **78**, 1963, p. 339-378.
- [13] Y. CHOQUET-BRUHAT et F. CAGNAC, *J. Maths pures et appliquées*, t. **63**, 1984, p. 377-390.
- [14] A. P. WHITMAN, R. J. KNILL, W. R. STROEGER, *Some harmonic maps on pseudo riemannian manifolds*. Preprint.
- [15] C. W. MISNER, *Phys. Rev. D.*, 1978, p. 4510-4524.
- [16] J. EELLS and L. LEMAIRE, *Bull. London Math. Soc.*, t. **10**, 1978, p. 1-68.
- [17] Y. NUTKU, *Ann. Inst. Poincaré*, t. **A 21**, 1974, p. 175-183.

(Manuscrit reçu le 12 mai 1986)