

ANNALES DE L'I. H. P., SECTION B

VICTOR H. DE LA PEÑA

Erratum / Correction to : “A bound on the moment generating function of a sum of dependent variables with an application to simple random sampling without replacement”

Annales de l'I. H. P., section B, tome 31, n° 4 (1995), p. 703-704

http://www.numdam.org/item?id=AIHPB_1995__31_4_703_0

© Gauthier-Villars, 1995, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section B » (<http://www.elsevier.com/locate/anihpb>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ERRATUM

Correction to:

A bound on the moment generating function of a sum of dependent variables with an application to simple random sampling without replacement

by

Victor H. de la PEÑA

Department of Statistics, Columbia University,
NY, NY 10027, USA.

ABSTRACT. – In this note we provide a corrected statement for Theorem 2 of our article *Ann. Inst. Henri Poincaré*, Vol. 30, n° 2, 1994, pp. 197-211.

RÉSUMÉ. – Dans cette note nous donnons une version corrigée du Théorème 2 de notre article *Ann. Inst. Henri Poincaré*, Vol. 30, n° 2, 1994, p. 197-211.

1. In this note we correct an error found in [2]. We are thankful to Amir Dembo for pointing us to the problem. First of all, due to a transcription oversight, we gave incorrect bounds on $P(|Z| \geq t)$, where Z is a standard normal variable. We refer the reader to page 49 of [1] for the correct bounds. The statement of Theorem 2 of [2] (wich is not used anywhere else in that paper) should be replaced by the following similar statement.

THEOREM 2. – Let $\{x_i\}$ be an arbitrary sequence of random variables. Assume that for a σ -field \mathcal{G} , $\{y_i\}$ is a \mathcal{G} -conditionally independent sequence tangent to $\{x_i\}$. Let Z be a normal random variable with mean zero and variance 1. Then, the inequality

$$P\left(\sum_{i=1}^n y_i \geq x\right) \leq AP(Z \geq x)$$

for some universal constant $0 < A < \infty$, and all $x \geq 0$, implies that for all $x \geq 0$,

$$P\left(\sum_{i=1}^n x_i \geq x\right) \leq \sqrt{A} \exp -(x^2/4).$$

The proof of this is easy and follows in spirit the original proof given, after eliminating the use of the bounds on $P(|Z| \geq t)$ and using Corollary 1 instead of Corollary 2.

REFERENCES

- [1] Y. S. CHOW and H. TEICHER, *Probability Theory*. Springer-Verlag, New York, 1978.
- [2] V. H. de la PEÑA, A bound on the moment generating function of a sum of dependent variables with an application to simple random sampling without replacement. *Ann. Inst. Henri Poincaré*, Vol. **30**, n° 2, 1994, pp. 197-211.