

Limit laws for the energy of a charged polymer

Xia Chen¹

Department of Mathematics, University of Tennessee, Knoxville, TN 37996-1300, USA. E-mail: xchen@math.utk.edu

Received 14 July 2006; revised 12 January 2007; accepted 1 March 2007

Abstract. In this paper we obtain the central limit theorems, moderate deviations and the laws of the iterated logarithm for the energy

$$H_n = \sum_{1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j = S_k\}}$$

of the polymer $\{S_1, \dots, S_n\}$ equipped with random electrical charges $\{\omega_1, \dots, \omega_n\}$. Our approach is based on comparison of the moments between H_n and the self-intersection local time

$$Q_n = \sum_{1 \leq j < k \leq n} 1_{\{S_j = S_k\}}$$

run by the d -dimensional random walk $\{S_k\}$. As partially needed for our main objective and partially motivated by their independent interest, the central limit theorems and exponential integrability for Q_n are also investigated in the case $d \geq 3$.

Résumé. Cet article est consacré à l'étude du théorème central limite, des déviations modérées et des lois du logarithme itéré pour l'énergie

$$H_n = \sum_{1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j = S_k\}}$$

du polymère $\{S_1, \dots, S_n\}$ doté de charges électriques $\{\omega_1, \dots, \omega_n\}$. Notre approche se base sur la comparaison des moments de H_n et du temps local de recouvrements

$$Q_n = \sum_{1 \leq j < k \leq n} 1_{\{S_j = S_k\}}$$

de la marche aléatoire d -dimensionnelle $\{S_k\}$. L'étude du théorème central limite et de l'intégrabilité exponentielle de Q_n (dans le cas $d \geq 3$) est également menée, tant pour comme outil pour notre principal objectif que pour son intérêt intrinsèque.

MSC: 60F05; 60F10; 60F15

Keywords: Charged polymer; Self-intersection local time; Central limit theorem; Moderate deviation; Laws of the iterated logarithm

¹Research partially supported by NSF Grant DMS-0704024.

1. Introduction

In the physics literature, the geometric shape of certain polymers is often described as an interpolation line segment with the vertices given as the n -step lattice (simple) random walk

$$\{S_1, S_2, \dots, S_n\}.$$

By placing independent, identically distributed electric charges $\omega_k = \pm 1$ to each vertex of the polymer, Kantor and Kardar [16] consider a model of polymers with random electrical charges associated with the Hamiltonian

$$H_n = \sum_{1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j = S_k\}}. \tag{1.1}$$

In the physics literature, H_n is called the energy of the polymer. To understand the physics intuition of H_n , we assign an electrical charge ω_k to the random site S_k for all $k = 1, 2, \dots$. Assume that when two charges meet, the pair with opposite signs gives negative contribution while the pair with the same sign gives positive contribution. Thus, H_n represents the total electrical interaction charge of the polymer $\{S_1, S_2, \dots, S_n\}$.

We point out some other works by physicists in this direction. In [10], the charges are i.i.d. Gaussian variables. In [11], the charges take 0–1 values. We also refer the reader to [4,18] for the continuous versions of the polymer with random charges. Finally, we mention the survey paper by van der Hofstad and König [12] for a long list of mathematical models connected to polymers.

As for other connections, we cite the comment by Martínez and Petritis [18]: “It is argued that a protein molecule is very much like a random walk with random charges attached at the vertices of the walk; these charges are interacting through local interactions mimicking Lennard–Jones or hydrogen-bond potentials”.

We study the asymptotic behaviors of H_n given in (1.1). In the rest of the paper, $\{S_n\}_{n \geq 1}$ is a symmetric random walk on \mathbb{Z}^d with covariance matrix Γ (or variance σ^2 as $d = 1$). We assume that the smallest group that supports $\{S_n\}_{n \geq 1}$ is \mathbb{Z}^d . Throughout, $\{\omega_k\}_{k \geq 1}$ is an i.i.d. sequence of symmetric random variable with

$$\mathbb{E}\omega_1^2 = 1 \quad \text{and} \quad \mathbb{E}e^{\lambda_0 \omega_1^2} < \infty \quad \text{for some } \lambda_0 > 0. \tag{1.2}$$

Our first result is on the central limit theorems.

Theorem 1.1. *As $d = 1$,*

$$\frac{1}{n^{3/4}} H_n \xrightarrow{d} (2\sigma)^{-1/2} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^{1/2} U, \tag{1.3}$$

where U is a random variable with standard normal distribution, $L(t, x)$ is the local time of the 1-dimensional Brownian motion $W(t)$ such that U and $W(t)$ are independent.

As $d = 2$,

$$\frac{1}{\sqrt{n \log n}} H_n \xrightarrow{d} \frac{1}{\sqrt{2\pi} \sqrt[4]{\det \Gamma}} U. \tag{1.4}$$

As $d \geq 3$,

$$\frac{1}{\sqrt{n}} H_n \xrightarrow{d} \sqrt{\gamma} U, \tag{1.5}$$

where

$$\gamma = \sum_{k=1}^{\infty} \mathbb{P}\{S_k = 0\}. \tag{1.6}$$

Here is our explanation on the dimensional dependence appearing in Theorem 1.1. The higher the dimension is, the less likely the random walk is to have long-range interaction (self-intersection). In the multi-dimensional case ($d \geq 2$), therefore, H_n is a sum of random variables with weak dependence and yields a Gaussian limit when properly normalized. It should be pointed that the low level of long-range interaction is vital for the chaos

$$\sum_{1 \leq j < k \leq n} a_{j,k} \omega_j \omega_k$$

to have a Gaussian limit when properly normalized. A simple example is when $a_{j,k} \equiv 1$. In this case

$$\sum_{1 \leq j < k \leq n} \omega_j \omega_k = \frac{1}{2} \left\{ \left[\sum_{j=1}^n \omega_j \right]^2 - \sum_{j=1}^n \omega_j^2 \right\}.$$

By the classic law of large numbers and classic central limit theorem,

$$\frac{1}{n} \sum_{1 \leq j < k \leq n} \omega_j \omega_k \xrightarrow{d} \frac{1}{2} (U^2 - 1)$$

which sharply contrasts to the statements in Theorem 1.1.

Our next theorem describes the moderate deviation behaviors of H_n .

Theorem 1.2. As $d = 1$,

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{P} \{ \pm H_n \geq \lambda (nb_n)^{3/4} \} = -\frac{1}{2} \sigma^{2/3} (3\lambda)^{4/3}, \quad \lambda > 0, \tag{1.7}$$

for any positive sequence $\{b_n\}$ satisfying

$$b_n \rightarrow \infty \quad \text{and} \quad b_n = o(\sqrt[7]{n}), \quad n \rightarrow \infty. \tag{1.8}$$

As $d = 2$,

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{P} \{ \pm H_n \geq \lambda \sqrt{n(\log n)b_n} \} = -\pi \sqrt{\det(\Gamma)} \lambda^2, \quad \lambda > 0, \tag{1.9}$$

for any positive sequence $\{b_n\}$ satisfying

$$b_n \rightarrow \infty \quad \text{and} \quad b_n = o(\log n), \quad n \rightarrow \infty. \tag{1.10}$$

As $d \geq 3$,

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{P} \{ \pm H_n \geq \lambda \sqrt{nb_n} \} = -\frac{\lambda^2}{2\gamma}, \quad \lambda > 0, \tag{1.11}$$

for any positive sequence $\{b_n\}$ satisfying

$$b_n \rightarrow \infty \quad \text{and} \quad b_n = o\left(\left(\frac{n}{\log n}\right)^{1/4}\right), \quad n \rightarrow \infty. \tag{1.12}$$

Our moderate deviations applied to the law of the iterated logarithm:

Theorem 1.3. As $d = 1$,

$$\limsup_{n \rightarrow \infty} \frac{\pm H_n}{(n \log \log n)^{3/4}} = \frac{2^{3/4}}{3} \sigma^{-1/2} \quad a.s. \tag{1.13}$$

As $d = 2$,

$$\limsup_{n \rightarrow \infty} \frac{\pm H_n}{\sqrt{n \log n \log \log n}} = \frac{1}{\sqrt{\pi \det(\Gamma)^{1/4}}} \quad a.s. \tag{1.14}$$

As $d \geq 3$,

$$\limsup_{n \rightarrow \infty} \frac{\pm H_n}{\sqrt{n \log \log n}} = \sqrt{2\gamma} \quad a.s. \tag{1.15}$$

We compare the results and treatments between the present paper and some recent works on self-intersection local times such as [3,6]. On the one hand, we shall see that the asymptotic behaviors of H_n described in our main theorems are closely related to those of the self-intersection local time

$$Q_n = \sum_{1 \leq j < k \leq n} 1_{\{S_j = S_k\}}. \tag{1.16}$$

Indeed, our approach is based on the moment comparisons between H_n and Q_n (see Proposition 2.1). In particular, the difference in limiting distribution between the case $d = 1$ and the case $d \geq 2$ in Theorem 1.1 is caused by the fact that in the case $d = 1$, Q_n converges (in distribution) to the Brownian self-intersection local times when properly normalized (see [8]), while as $d \geq 2$, Q_n is asymptotically close to its expectation (see [3] for $d = 2$ and the Section 5 for $d \geq 3$).

On the other hand, the fact that Q_n is close to the quadratic form

$$\sum_{x \in \mathbb{Z}^d} l^2(n, x)$$

of the local time $l(n, x)$ plays a crucial role in the study of the self-intersection local time Q_n (see e.g., [3,8]). It allows, for example, some technologies developed along the line of probability in Banach space. Unfortunately, this idea does not work in our setting. Indeed, the fact (in view of Theorem 1.1) that the second term in the decomposition

$$\sum_{x \in \mathbb{Z}^d} \left[\sum_{j=1}^n \omega_j 1_{\{S_j=x\}} \right]^2 = 2H_n + \sum_{j=1}^n \omega_j^2 \tag{1.17}$$

is the dominating term shows that H_n is not even in the same asymptotic order as the quadratic form on the left-hand side.

The key estimations are carried out in Proposition 2.1. Our approach relies on the following crucial observation. By (1.17) we have

$$H_n = \frac{1}{2} \sum_{x \in \mathbb{Z}^d} \left\{ \left[\sum_{j=1}^n \omega_j 1_{\{S_j=x\}} \right]^2 - \sum_{j=1}^n \omega_j^2 1_{\{S_j=x\}} \right\}. \tag{1.18}$$

Conditioned on the random walk $\{S_k\}$, the random variables

$$\left[\sum_{j=1}^n \omega_j 1_{\{S_j=x\}} \right]^2 - \sum_{j=1}^n \omega_j^2 1_{\{S_j=x\}}, \quad x \in \mathbb{Z}^d,$$

form an independent family and, for each fix $x \in \mathbb{Z}^d$,

$$\left[\sum_{j=1}^n \omega_j 1_{\{S_j=x\}} \right]^2 - \sum_{j=1}^n \omega_j^2 1_{\{S_j=x\}} \stackrel{d}{=} \left[\sum_{j=1}^{l(n,x)} \omega_j \right]^2 - \sum_{j=1}^{l(n,x)} \omega_j.$$

By a classic estimate for independent sums, and by some combinatorial computation, a conditional moment estimate given in Proposition 2.1 links H_n with Q_n . Another fact repeatedly used in this work is that the self-intersection occurring at the frequently visited sites does not make a significant contribution to the quantities H_n and Q_n . Consequently, the pairs H_n and \tilde{H}_n (defined in (2.3)); Q_n and \tilde{Q}_n (defined in (2.4) below) are exchangeable in our setting.

Beyond mathematical technicality, the creation of the present paper is based on our belief that H_n resembles, in the limiting behaviors described in our main theorems, the random quantity

$$H'_n = \sum_{1 \leq j < k \leq n} 1_{\{S_j=S_k\}} U_{j,k},$$

where $U_{j,k}$ are i.i.d. standard normal random variables independent of $\{S_k\}$. Notice that H'_n is conditionally normal with conditional variance Q_n . Our observation explains why and how the limiting behaviors of H_n depend on its conditional variance Q_n . It should be pointed out, however, that the replacement of $\omega_j \omega_k$ by $U_{j,k}$ is highly non-trivial and should not be taken for granted, in view of the example given next to Theorem 1.1.

In Section 2, we establish a comparison (Proposition 2.1) between the moments of H_n and Q_n , and then apply it to prove Theorem 1.1. Our approach relies on combinatorial and conditioning methods. In Section 3, Proposition 2.1 is further applied to prove Theorem 1.2 through a Laplacian argument. In Section 4, the laws of the iterated logarithm given in Theorem 1.3 are proved as a consequence of our moderate deviations. The non-trivial part of this section is a maximal inequality (Lemma 4.1) of Lévy type. In Section 5, we investigate the weak laws and exponential integrabilities for the renormalized self-intersection local time $Q_n - \mathbb{E}Q_n$ in the high dimensions ($d \geq 3$). The central limit theorem given in Theorem 5.1 and the exponential integrability given in Theorem 5.2 provide sharp bounds on $Q_n - \mathbb{E}Q_n$, which constitute the replacement of Q_n by $\mathbb{E}Q_n$ carried out in our argument for Theorem 1.1 and for Theorem 1.2 (the estimate of $Q_n - \mathbb{E}Q_n$ needed in the case $d = 2$ was established in [3,20]). In addition, the results given in Section 5 are of independent interest as a part of the study of the self-intersection local times in high dimensions and are partially motivated by some recent works of Asselah and Castell [1] and Asselah [2].

2. Moment comparison and laws of weak convergence

We begin with the following classic lemma.

Lemma 2.1. *Assume (1.2). Then*

$$\mathbb{E} \left\{ \left(\sum_{j=1}^n \omega_j \right)^2 - \sum_{j=1}^n \omega_j^2 \right\}^2 = 2n(n-1). \tag{2.1}$$

More generally, there is a constant $C > 0$ such that for any integers $n \geq 1$ and $m \geq 2$,

$$\mathbb{E} \left| \left(\sum_{j=1}^n \omega_j \right)^2 - \sum_{j=1}^n \omega_j^2 \right|^m \leq m!(Cn(n-1))^{m/2}. \tag{2.2}$$

Proof. The first part follows from the following straightforward computation:

$$\mathbb{E} \left\{ \left(\sum_{j=1}^n \omega_j \right)^2 - \sum_{j=1}^n \omega_j^2 \right\}^2 = 4 \mathbb{E} \left\{ \sum_{1 \leq j < k \leq n} \omega_j \omega_k \right\}^2 = 4 \sum_{1 \leq j < k \leq n} \mathbb{E}(\omega_j^2 \omega_k^2) = 2n(n-1).$$

For the second part, we only need to show

$$\mathbb{E} \left| \left(\sum_{j=1}^n \omega_j \right)^2 - \sum_{j=1}^n \omega_j^2 \right|^m \leq m! C^{m/2} n^m.$$

By the inequality

$$\left(\mathbb{E} \left| \left(\sum_{j=1}^n \omega_j \right)^2 - \sum_{j=1}^n \omega_j^2 \right|^m \right)^{1/m} \leq \left(\mathbb{E} \left(\sum_{j=1}^n \omega_j \right)^{2m} \right)^{1/m} + \left(\mathbb{E} \left(\sum_{j=1}^n \omega_j^2 \right)^m \right)^{1/m}$$

all we need is that

$$\mathbb{E} \left(\sum_{j=1}^n \omega_j \right)^{2m} \leq C^{m/2} m! n^m$$

and that

$$\mathbb{E} \left| \sum_{j=1}^n \omega_j^2 \right|^m \leq C^{m/2} m! n^m.$$

Due to similarity we only prove the first inequality. Notice that by symmetry

$$\mathbb{E} \left(\sum_{j=1}^n \omega_j \right)^{2m} = \sum_{\substack{k_1 + \dots + k_n = m \\ k_1, \dots, k_n \geq 0}} \frac{(2m)!}{(2k_1)! \dots (2k_n)!} \mathbb{E} \omega^{2k_1} \dots \mathbb{E} \omega^{2k_n}.$$

By the integrability given in (1.2) there is a constant $c_1 > 0$ such that

$$\mathbb{E} \omega^{2k} \leq k! c_1^k, \quad k = 0, 1, 2, \dots$$

Notice also the very rough estimate

$$c_2^k (k!)^2 \leq (2k)! \leq c_3^k (k!)^2, \quad k = 0, 1, 2, \dots$$

So we have

$$\mathbb{E} \left(\sum_{j=1}^n \omega_j \right)^{2m} \leq C^{m/2} m! \sum_{\substack{k_1 + \dots + k_n = m \\ k_1, \dots, k_n \geq 0}} \frac{m!}{k_1! \dots k_n!} = C^{m/2} m! n^m. \quad \square$$

Let K_n be a positive sequence which may vary in different settings and will later be specified in each specific setting. Recall that Q_n is given in (1.16) and define the local time

$$l(n, x) = \sum_{k=1}^n 1_{\{S_k = x\}}, \quad x \in \mathbb{Z}^d, n = 1, 2, \dots$$

The asymptotic behaviors of the local times of the random walks have been studied extensively. We cite the book by Révész [19] for an overview.

The following two random quantities play important roles in this paper:

$$\tilde{H}_n = H_n 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}}, \tag{2.3}$$

$$\tilde{Q}_n = Q_n 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}}. \tag{2.4}$$

In addition, we introduce the deterministic quantity

$$A_m(n) = \frac{1}{2^m} \sum_{(y_1, \dots, y_m) \in B_m} \mathbb{E} \left(1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m l(n, y_k) (l(n, y_k) - 1) \right),$$

where $m, n = 1, 2, \dots$ and

$$B_m = \{(y_1, \dots, y_m) \in (\mathbb{Z}^d)^m; y_1, \dots, y_m \text{ are distinct}\}. \tag{2.5}$$

An easy observation gives that

$$\begin{aligned} A_m(n) &\leq \frac{1}{2^m} \sum_{y_1, \dots, y_m \in \mathbb{Z}^d} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m l(n, y_k) (l(n, y_k) - 1) \right) \\ &= \mathbb{E} \tilde{Q}_n^m. \end{aligned} \tag{2.6}$$

Some more substantial comparisons are given in the following.

Proposition 2.1. *There is a constant $C > 0$ independent of n, m and the choice of K_n , such that*

$$\mathbb{E} \tilde{H}_n^m \leq \frac{m!}{2^m} \sum_{l=1}^{\lfloor 2^{-1}m \rfloor} \frac{1}{l!} K_n^{m-2l} 2^l C^{(m-2l)/2} \binom{m-l-1}{m-2l} \mathbb{E} \tilde{Q}_n^l. \tag{2.7}$$

On the other hand, for any integers $m, n \geq 1$,

$$\mathbb{E} \tilde{H}_n^{2m} \geq \frac{(2m)!}{2^m m!} A_m(n), \tag{2.8}$$

$$\mathbb{E} \tilde{Q}_n^m \leq \sum_{l=1}^m \binom{m}{l} \left(\frac{lK_n^2}{2} \right)^{m-l} A_l(n). \tag{2.9}$$

Proof. Notice that

$$H_n = \frac{1}{2} \sum_{x \in \mathbb{Z}^d} \left\{ \left(\sum_{j=1}^n \omega_j \mathbf{1}_{\{S_j=x\}} \right)^2 - \sum_{j=1}^n \omega_j^2 \mathbf{1}_{\{S_j=x\}} \right\} = \frac{1}{2} \sum_{x \in \mathbb{Z}^d} \Lambda_n(x) \quad (\text{say}). \tag{2.10}$$

Hence,

$$\mathbb{E} \tilde{H}_n^m = 2^{-m} \sum_{x_1, \dots, x_m \in \mathbb{Z}^d} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right).$$

For each $1 \leq l \leq m$, let

$$A_l = \{(x_1, \dots, x_m) \in (\mathbb{Z}^d)^m; \#\{x_1, \dots, x_m\} = l\}.$$

Then,

$$\mathbb{E} \tilde{H}_n^m = 2^{-m} \sum_{l=1}^m \sum_{(x_1, \dots, x_m) \in A_l} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right). \tag{2.11}$$

Write

$$C_l = \{F \subset \mathbb{Z}^d; \#(F) = l\}$$

and for any $\{y_1, \dots, y_l\} \in C_l$, set

$$A_l(y_1, \dots, y_l) = \{(x_1, \dots, x_m) \in (\mathbb{Z}^d)^m; \{x_1, \dots, x_m\} = \{y_1, \dots, y_l\}\}.$$

Notice that

$$1_{A_l}(x_1, \dots, x_m) = \sum_{\{y_1, \dots, y_l\} \in \mathcal{C}_l} 1_{A_l(y_1, \dots, y_l)}(x_1, \dots, x_m).$$

Thus

$$\begin{aligned} & \sum_{(x_1, \dots, x_m) \in A_l} \mathbb{E} \left(1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right) \\ &= \sum_{\{y_1, \dots, y_l\} \in \mathcal{C}_l} \sum_{(x_1, \dots, x_m) \in A_l(y_1, \dots, y_l)} \mathbb{E} \left(1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right). \end{aligned}$$

For any $(x_1, \dots, x_m) \in A_l(y_1, \dots, y_l)$, let i_k be the number of x_1, \dots, x_m which are equal to y_k , where $k = 1, \dots, l$. Then

$$\mathbb{E} \left(1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right) = \mathbb{E} \left\{ 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \Lambda_n(y_k)^{i_k} \right\}.$$

Consequently,

$$\begin{aligned} & \sum_{(x_1, \dots, x_m) \in A_l(y_1, \dots, y_l)} \mathbb{E} \left(1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right) \\ &= \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{m!}{(i_1)! \dots (i_l)!} \mathbb{E} \left\{ 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \Lambda_n(y_k)^{i_k} \right\}. \end{aligned}$$

Summarizing the above discussion,

$$\begin{aligned} & \sum_{(x_1, \dots, x_m) \in A_l} \mathbb{E} \left(1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right) \\ &= \sum_{\{y_1, \dots, y_l\} \in \mathcal{C}_l} \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{m!}{(i_1)! \dots (i_l)!} \mathbb{E} \left\{ 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \Lambda_n(y_k)^{i_k} \right\}. \end{aligned}$$

Notice that the quantity

$$f(y_1, \dots, y_l) \equiv \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{m!}{(i_1)! \dots (i_l)!} \mathbb{E} \left\{ 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \Lambda_n(y_k)^{i_k} \right\}$$

is invariant under the permutations over $\{y_1, \dots, y_l\}$. So we have

$$\begin{aligned} & \sum_{(x_1, \dots, x_m) \in A_l} \mathbb{E} \left(1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \Lambda_n(x_k) \right) \\ &= \frac{1}{l!} \sum_{(y_1, \dots, y_l) \in B_l} \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{m!}{(i_1)! \dots (i_l)!} \mathbb{E} \left\{ 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \Lambda_n(y_k)^{i_k} \right\}, \end{aligned}$$

where B_l is defined by (2.5). By (2.11)

$$\begin{aligned} \mathbb{E}\tilde{H}_n^m &= 2^{-m} \sum_{l=1}^m \frac{1}{l!} \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 1}} \frac{m!}{(i_1)! \cdots (i_l)!} \\ &\times \sum_{(y_1, \dots, y_l) \in B_l} \mathbb{E} \left\{ 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \Lambda_n(y_k)^{i_k} \right\}. \end{aligned} \tag{2.12}$$

We adopt the notation “ \mathbb{E}^ω ” for the expectation with respect to $\{\omega_k\}_{k \geq 1}$ and for each $y \in \mathbb{Z}^d$, write $D(y) = \{1 \leq k \leq n; S_k = y\}$. Then

$$\Lambda(y) = \left(\sum_{j \in D(y)} \omega_j \right)^2 - \sum_{j \in D(y)} \omega_j^2$$

and for distinct y_1, \dots, y_l , the sets $D(y_1), \dots, D(y_l)$ are disjoint. Hence, by independence,

$$\mathbb{E}^\omega \prod_{k=1}^l \Lambda_n(y_k)^{i_k} = \prod_{k=1}^l \mathbb{E}^\omega \Lambda_n(y_k)^{i_k}.$$

In particular, the above quantity is zero if any of i_1, \dots, i_l is 1. Consequently, the terms in (2.12) with $l > m/2$ are equal to zero,

$$\mathbb{E}\tilde{H}_n = 0 \tag{2.13}$$

and for any integer $m \geq 2$,

$$\begin{aligned} \mathbb{E}\tilde{H}_n^m &= 2^{-m} \sum_{l=1}^{\lfloor 2^{-1}m \rfloor} \frac{1}{l!} \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 2}} \frac{m!}{(i_1)! \cdots (i_l)!} \\ &\times \sum_{(y_1, \dots, y_l) \in B_l} \mathbb{E} \left\{ 1_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \mathbb{E}^\omega \Lambda_n(y_k)^{i_k} \right\}. \end{aligned} \tag{2.14}$$

Notice that

$$\mathbb{E}^\omega \Lambda_n(y_k)^{i_k} = \mathbb{E}^\omega \left\{ \left(\sum_{j=1}^{l(n, x)} \omega_j \right)^2 - \sum_{j=1}^{l(n, x)} \omega_j^2 \right\}^{i_k}. \tag{2.15}$$

By Lemma 2.1 we have

$$\begin{aligned} \prod_{k=1}^l \mathbb{E}^\omega \Lambda_n(y_k)^{i_k} &\leq \prod_{k=1}^l i_k! C_{i_k}^{i_k/2} \{l(n, y_k)(l(n, y_k) - 1)\}^{i_k/2} \\ &= (i_1! \cdots i_l!) (C_{i_1}^{i_1/2} \cdots C_{i_l}^{i_l/2}) \prod_{k=1}^l \{l(n, y_k)(l(n, y_k) - 1)\}^{i_k/2}, \end{aligned}$$

where $C_i = 1$ as $i = 2$ and C_i is the constant C given in (2.2) as $i \geq 3$. We may assume that $C \geq 1$ in the rest of the proof.

Hence,

$$\begin{aligned}
 \mathbb{E} \tilde{H}_n^m &\leq \frac{m!}{2^m} \sum_{l=1}^{\lfloor 2^{-1}m \rfloor} \frac{1}{l!} \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 2}} C_{i_1}^{i_1/2} \dots C_{i_l}^{i_l/2} \\
 &\quad \times \sum_{(y_1, \dots, y_l) \in B_l} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \{l(n, y_k)(l(n, y_k) - 1)\}^{i_k/2} \right) \\
 &\leq \frac{m!}{2^m} \sum_{l=1}^{\lfloor 2^{-1}m \rfloor} \frac{1}{l!} K_n^{m-2l} \left\{ \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 2}} C_{i_1}^{i_1/2} \dots C_{i_l}^{i_l/2} \right\} A_l(n) \\
 &\leq \frac{m!}{2^m} \sum_{l=1}^{\lfloor 2^{-1}m \rfloor} \frac{1}{l!} K_n^{m-2l} 2^l \left\{ \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 2}} C_{i_1}^{i_1/2} \dots C_{i_l}^{i_l/2} \right\} \mathbb{E} \tilde{Q}_n^l,
 \end{aligned} \tag{2.16}$$

where the last step follows from (2.6).

For each (i_1, \dots, i_l) with $i_1 + \dots + i_l = m$, write

$$k = k(i_1, \dots, i_l) = \#\{1 \leq j \leq l; i_j = 2\}.$$

We have

$$m = i_1 + \dots + i_l \geq 2k + 3(l - k)$$

which leads to $l - k \leq m - 2l$. Thus,

$$\begin{aligned}
 \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 2}} (C_{i_1}^{i_1/2} \dots C_{i_l}^{i_l/2}) &= \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 2}} C^{(l-k)/2} \leq C^{(m-2l)/2} \sum_{\substack{i_1+\dots+i_l=m \\ i_1, \dots, i_l \geq 2}} 1 \\
 &= C^{(m-2l)/2} \binom{m-l-1}{m-2l}.
 \end{aligned}$$

Hence, (2.7) follows from (2.16).

To prove (2.8), we come to (2.14) and we notice that the symmetry of $\{\omega_k\}$ implies that for any integer $l \geq 1$,

$$\mathbb{E}^\omega A_n(y_k)^{i_k} = 2^{i_k} \mathbb{E} \left(\sum_{1 \leq j_1 < j_2 \leq l(n, x)} \omega_{j_1} \omega_{j_2} \right)^{i_k} \geq 0. \tag{2.17}$$

Replacing m by $2m$ in (2.4) and only keeping the term with $l = m$ on the right-hand side, we obtain

$$\begin{aligned}
 \mathbb{E} \tilde{H}_n^{2m} &\geq \frac{(2m)!}{2^{2m} m!} 2^{-m} \sum_{(y_1, \dots, y_m) \in B_m} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m \mathbb{E}^\omega A_n(y_k)^2 \right) \\
 &= \frac{(2m)!}{2^m m!} A_m(n),
 \end{aligned}$$

where the second step follows from (2.1) in Lemma 2.1 and (2.15).

To prove (2.9), we adopt the argument used for (2.12).

$$\begin{aligned}
 \mathbb{E} \tilde{Q}_n^m &= 2^{-m} \sum_{x_1, \dots, x_m \in \mathbb{Z}^d} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^m l(n, x_k) (l(n, x_k) - 1) \right) \\
 &= 2^{-m} \sum_{l=1}^m \frac{1}{l!} \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{m!}{i_1! \dots i_l!} \\
 &\quad \times \sum_{(y_1, \dots, y_l) \in B_l} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l \{l(n, y_k) (l(n, y_k) - 1)\}^{i_k} \right) \\
 &\leq \frac{m!}{2^m} \sum_{l=1}^m \frac{1}{l!} \left\{ \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{1}{i_1! \dots i_l!} \right\} K_n^{2(m-l)} \\
 &\quad \times \sum_{(y_1, \dots, y_l) \in B_l} \mathbb{E} \left(\mathbf{1}_{\{\sup_{x \in \mathbb{Z}^d} l(n, x) \leq K_n\}} \prod_{k=1}^l l(n, y_k) (l(n, y_k) - 1) \right) \\
 &= m! \sum_{l=1}^m \frac{1}{l!} K_n^{2(m-l)} 2^{-(m-l)} A_l(n) \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{1}{i_1! \dots i_l!}.
 \end{aligned}$$

Finally, (2.9) follows from the following estimate:

$$\begin{aligned}
 \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{1}{i_1! \dots i_l!} &\leq \sum_{\substack{i_1 + \dots + i_l = m \\ i_1, \dots, i_l \geq 1}} \frac{1}{(i_1 - 1)! \dots (i_l - 1)!} \\
 &= \sum_{\substack{i_1 + \dots + i_l = m-1 \\ i_1, \dots, i_l \geq 0}} \frac{1}{i_1! \dots i_l!} = \frac{l^{m-l}}{(m-l)!}.
 \end{aligned}$$

□

Proof of Theorem 1.1. We start with the case $d = 1$. Notice that

$$Q_n = \frac{1}{2} \left(\sum_{x \in \mathbb{Z}} l^2(n, x) - n \right).$$

By Theorem 1.2 of [6],

$$n^{-3/2} Q_n \xrightarrow{d} \frac{1}{2\sigma} \int_{-\infty}^{\infty} L^2(1, x) dx. \tag{2.18}$$

Fix $0 < \delta < 1/2$ and let $K_n = n^{(1+\delta)/2}$. By the classic fact (see, for example, [19]) that

$$n^{-1/2} \sup_{x \in \mathbb{Z}} l(n, x) \xrightarrow{d} \sigma^{-1} \sup_{x \in \mathbb{R}} L(1, x) \tag{2.19}$$

we have that

$$n^{-3/2} \tilde{Q}_n \xrightarrow{d} \frac{1}{2\sigma} \int_{-\infty}^{\infty} L^2(1, x) dx, \tag{2.20}$$

which gives

$$\lim_{n \rightarrow \infty} n^{-3m/2} \mathbb{E} \tilde{Q}_n^m = \frac{1}{(2\sigma)^m} \mathbb{E} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^m, \quad m = 1, 2, \dots \tag{2.21}$$

Replacing m by $2m + 1$ in (2.7) we have

$$\begin{aligned} \mathbb{E} \tilde{H}_n^{2m+1} &\leq \frac{(2m+1)!}{2^{2m+1}} \sum_{l=1}^m \frac{1}{l!} n^{(1+\delta)(m-l)+(1+\delta)/2} 2^l C^{(2m-2l+1)/2} \binom{2m-l}{2m-2l+1} \mathbb{E} \tilde{Q}_n^l \\ &= O \left(\sum_{l=1}^m n^{(1+\delta)(m-l)+(1+\delta)/2} n^{3l/2} \right) = o \left(n^{3(2m+1)/4} \right), \quad n \rightarrow \infty, \end{aligned} \tag{2.22}$$

for all $m = 0, 1, 2, \dots$

Replacing m by $2m$ in (2.7) and by (2.21) we have

$$\begin{aligned} \mathbb{E} \tilde{H}_n^{2m} &\leq \frac{(2m)!}{2^{2m}} \sum_{l=1}^m \frac{1}{l!} n^{(1+\delta)(m-l)} 2^l C^{m-l} \binom{2m-l-1}{2m-2l} \mathbb{E} \tilde{Q}_n^l \\ &\sim \frac{(2m)!}{2^{2m}} \sum_{l=1}^m \frac{1}{l!} n^{(1+\delta)(m-l)} (2\sigma)^{-l} n^{3l/2} 2^l \\ &\quad \times \mathbb{E} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^l C^{m-l} \binom{2m-l-1}{2m-2l}, \quad n \rightarrow \infty. \end{aligned}$$

Clearly, the right-hand side is dominated by the term with $l = m$. Consequently,

$$\limsup_{n \rightarrow \infty} n^{-3m/2} \mathbb{E} \tilde{H}_n^{2m} \leq \frac{1}{(2\sigma)^m} \frac{(2m)!}{2^m m!} \mathbb{E} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^m \tag{2.23}$$

for all $m = 1, 2, \dots$. In particular, combining this with (2.8) we have

$$A_m(n) = O(n^{3m/2}), \quad n \rightarrow \infty, m = 1, 2, \dots \tag{2.24}$$

On the other hand, by (2.24) and (2.21), the right-hand side of (2.9) is dominated by the term with $l = m$. Hence,

$$\liminf_{n \rightarrow \infty} n^{-3m/2} A_m(n) \geq \frac{1}{(2\sigma)^m} \mathbb{E} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^m, \quad m = 1, 2, \dots \tag{2.25}$$

From (2.8),

$$\liminf_{n \rightarrow \infty} n^{-3m/2} \mathbb{E} \tilde{H}_n^{2m} \geq \frac{1}{(2\sigma)^m} \frac{(2m)!}{2^m m!} \mathbb{E} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^m. \tag{2.26}$$

In summary of (2.22), (2.23) and (2.26), and noticing that

$$\mathbb{E} U^{2m} = \frac{(2m)!}{2^m m!} \quad \text{and} \quad \mathbb{E} U^{2m+1} = 0$$

we have that for every $m = 0, 1, 2, \dots$,

$$\lim_{n \rightarrow \infty} n^{-3m/4} \mathbb{E} \tilde{H}_n^m = \frac{1}{(2\sigma)^{m/2}} (\mathbb{E} U^m) \mathbb{E} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^{m/2}. \tag{2.27}$$

Notice the fact that for any $\theta \in \mathbb{R}$,

$$\mathbb{E} \exp \left\{ \theta \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^{1/2} U \right\} = \mathbb{E} \exp \left\{ \frac{\theta^2}{2} \int_{-\infty}^{\infty} L^2(1, x) dx \right\} < \infty,$$

where the last step follows from Theorem 1.1 (with $m = 1$ and $p = 2$) in [8]. Therefore, (2.27) implies that

$$\frac{1}{n^{3/4}} \tilde{H}_n \xrightarrow{d} 2^{-1} \sigma^{-1/2} \left(\int_{-\infty}^{\infty} L^2(1, x) dx \right)^{1/2} U.$$

By (2.11) and by our choice of K_n we have

$$\mathbb{P} \left\{ \sup_{x \in \mathbb{Z}} l(n, x) > K_n \right\} \rightarrow 0 \tag{2.28}$$

as $n \rightarrow \infty$. Thus, we have proved Theorem 1.1 in the case $d = 1$.

The proof in the multi-dimensional cases is essentially the same. Instead of (2.12), we have that

$$\frac{Q_n}{n \log n} \xrightarrow{P} (2\pi)^{-1} (\det \Gamma)^{-1/2} \quad \text{as } d = 2, \tag{2.29}$$

$$\frac{Q_n}{n} \xrightarrow{P} \gamma \quad \text{as } d \geq 3. \tag{2.30}$$

Indeed, (2.29) and (2.30) follow from the weak convergence of the sequences $(Q_n - \mathbb{E}Q_n)/n$ when $d = 2$ (see [20]), $(Q_n - \mathbb{E}Q_n)/\sqrt{n \log n}$ when $d = 3$ (see Theorem 5.1) and $(Q_n - \mathbb{E}Q_n)/\sqrt{n}$ when $d \geq 4$ (see Theorem 5.1); and from the well-known fact that

$$\mathbb{E}Q_n \sim \begin{cases} (2\pi)^{-1} (\det \Gamma)^{-1/2} n \log n, & d = 2, \\ \gamma n, & d \geq 3. \end{cases} \tag{2.31}$$

In addition, it is well known [19] that

$$\sup_{x \in \mathbb{Z}^2} \frac{l(n, x)}{(\log n)^2} \quad \text{and} \quad \sup_{x \in \mathbb{Z}^d} \frac{l(n, x)}{\log n}$$

are almost surely bounded in the case $d = 2$ and the case $d \geq 3$, respectively. Thus, if we define $K_n = M(\log n)^2$ as $d = 2$, and $K_n = M \log n$ as $d \geq 3$. Then (2.28) holds as the constant $M > 0$ is sufficiently large.

Therefore, a modification of the proof for (2.27) gives that

$$\lim_{n \rightarrow \infty} (n \log n)^{-m/2} \mathbb{E} \tilde{H}_n^m = 2^{-m/2} (2\pi)^{-m/2} (\det \Gamma)^{-m/4} \mathbb{E} U^m \quad \text{as } d = 2, \tag{2.32}$$

$$\lim_{n \rightarrow \infty} n^{-m/2} \mathbb{E} \tilde{H}_n^m = 2^{-m/2} \gamma^{m/2} \mathbb{E} U^m \quad \text{as } d \geq 3. \tag{2.33}$$

So the multi-dimensional part of Theorem 1.1 follows from (2.32) and (2.33). □

3. Moderate deviations

Recall that $K_n = M(\log n)^2$ as $d = 2$, where $M > 0$ is a large but fixed constant. Take $K_n = (n/\log n)^{1/4}$ as $d \geq 3$. In the case $d = 1$, (1.8) implies that there is a positive sequence M_n such that

$$M_n \rightarrow \infty \quad \text{and} \quad M_n^2 \left(\frac{b_n^7}{n} \right)^{1/4} \rightarrow 0, \quad n \rightarrow \infty. \tag{3.1}$$

So in this section we take $K_n = M_n \sqrt{nb_n}$ as $d = 1$.

An important fact is that under our choice,

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{P} \left\{ \sup_{x \in \mathbb{Z}^d} l(n, x) > K_n \right\} = -\infty \tag{3.2}$$

in all dimensions. We refer to [5] for (3.2) under $d = 1$, and [19] for (3.2) under $d \geq 2$.

Another important fact is that

$$\mathbb{E} \tilde{H}_n^{2m+1} \geq 0, \quad m = 0, 1, \dots, \tag{3.3}$$

which follows from (2.13), (2.14) and (2.17).

We claim that Theorem 1.2 holds if we can prove that for any $\theta > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \pm \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n \right\} = \frac{\theta^4}{96\sigma^2} \quad \text{as } d = 1, \tag{3.4}$$

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \pm \theta \sqrt{\frac{b_n}{n \log n}} \tilde{H}_n \right\} = \frac{\theta^2}{4\pi \sqrt{\det \Gamma}} \quad \text{as } d = 2, \tag{3.5}$$

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \pm \theta \sqrt{\frac{b_n}{n}} \tilde{H}_n \right\} = \frac{\gamma \theta^2}{2} \quad \text{as } d \geq 3. \tag{3.6}$$

Indeed, according to the Gärtner–Ellis theorem (Theorem 2.3.6, p. 44 in [9]), (3.4)–(3.6) imply that \tilde{H}_n satisfies the moderate deviations given in Theorem 1.2. By Theorem 4.2.13, p. 130 in [9], the moderate deviations pass from \tilde{H}_n to H_n through the exponential equivalence given by

$$\limsup_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{P} \{ \tilde{H}_n \neq H_n \} = \lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{P} \left\{ \sup_{x \in \mathbb{Z}^d} l(n, x) > K_n \right\} = -\infty,$$

where the second step follows from (3.2).

In the rest of this section, we prove (3.4), (3.5) and (3.6) in three separate parts.

Case $d = 1$.

By (2.7) in Proposition 2.1,

$$\begin{aligned} & \sum_{m=2}^{\infty} \frac{\theta^m}{m!} \left(\frac{b_n}{n^3} \right)^{m/4} \mathbb{E} \tilde{H}_n^m \\ & \leq \sum_{m=2}^{\infty} \frac{\theta^m}{m!} \left(\frac{b_n}{n^3} \right)^{m/4} \frac{m!}{2^m} \sum_{l=1}^{\lfloor 2^{-1}m \rfloor} \frac{1}{l!} K_n^{m-2l} 2^l C^{(m-2l)/2} \binom{m-l-1}{m-2l} \mathbb{E} \tilde{Q}_n^l \\ & = \sum_{l=1}^{\infty} \frac{\theta^{2l}}{2^l l!} \left(\frac{b_n}{n^3} \right)^{l/2} \mathbb{E} \tilde{Q}_n^l \sum_{m=2l}^{\infty} \left(\frac{\theta}{2} \right)^{m-2l} K_n^{m-2l} \left(\frac{b_n}{n} \right)^{(m-2l)/4} C^{(m-2l)/2} \binom{m-l-1}{m-2l}. \end{aligned}$$

Notice that

$$\begin{aligned} & \sum_{m=2l}^{\infty} \left(\frac{\theta}{2} \right)^{m-2l} K_n^{m-2l} \left(\frac{b_n}{n^3} \right)^{(m-2l)/4} C^{(m-2l)/2} \binom{m-l-1}{m-2l} \\ & = \sum_{m=0}^{\infty} \left(\frac{\theta}{2} \right)^m K_n^m \left(\frac{b_n}{n^3} \right)^{m/4} C^{m/2} \binom{m+l-1}{m} \\ & = \left(1 - \frac{\sqrt{C} \theta K_n b_n^{1/4}}{2n^{3/4}} \right)^{-(l-1)}, \end{aligned}$$

where the last step follows from the Taylor expansion:

$$(1-x)^{-(l-1)} = \sum_{m=0}^{\infty} \binom{m+l-1}{m} x^m, \quad |x| < 1. \quad (3.7)$$

Combining the above estimate with (2.13) gives

$$\mathbb{E} \exp \left\{ \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n \right\} \leq \mathbb{E} \exp \left\{ \frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} \left(1 - \frac{\sqrt{C} \theta K_n b_n^{1/4}}{2n^{3/4}} \right)^{-1} \tilde{Q}_n \right\}.$$

In view of (2.13), by the Taylor expansion one can easily see that

$$\mathbb{E} \exp \left\{ -\theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n \right\} \leq \mathbb{E} \exp \left\{ \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n \right\}.$$

So we have

$$\mathbb{E} \exp \left\{ \pm \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n \right\} \leq \mathbb{E} \exp \left\{ \frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} \left(1 - \frac{\sqrt{C} \theta K_n b_n^{1/4}}{2n^{3/4}} \right)^{-1} \tilde{Q}_n \right\}. \quad (3.8)$$

Notice that

$$Q_n \leq \frac{1}{2} \sum_{x \in \mathbb{Z}} l^2(n, x) \leq \frac{1}{2} n \sup_{x \in \mathbb{Z}} l(n, x). \quad (3.9)$$

For any $\lambda > 0$,

$$\mathbb{E} \exp \left\{ \lambda \frac{b_n^{1/2}}{n^{3/2}} Q_n \right\} \leq \mathbb{E} \exp \left\{ \frac{\lambda}{2} \sqrt{\frac{b_n}{n}} \sup_{x \in \mathbb{Z}} l(n, x) \right\}.$$

By the fact (see Lemmas 11 and 12 in [15]) that

$$\limsup_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \frac{\lambda}{2} \sqrt{\frac{b_n}{n}} \sup_{x \in \mathbb{Z}} l(n, x) \right\} < \infty,$$

we have that for any $\lambda > 0$

$$\limsup_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \lambda \frac{b_n^{1/2}}{n^{3/2}} Q_n \right\} < \infty. \quad (3.10)$$

Recall (Theorem 1.3 in [8], with $m = 1$ and $p = 2$) that for any $\lambda > 0$

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{P} \{ Q_n \geq \lambda n^{3/2} b_n^{1/2} \} = -6\sigma^2 \lambda^2. \quad (3.11)$$

According to Varadhan's integral lemma (Theorem 4.3.1, p. 137 in [9]), (3.10) and (3.11) imply that for any $\lambda > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \lambda \frac{b_n^{1/2}}{n^{3/2}} Q_n \right\} = \sup_{y > 0} \{ y \lambda - 6\sigma^2 \lambda^2 \} = \frac{\lambda^2}{24\sigma^2}. \quad (3.12)$$

This, together with (3.8), gives the desired upper bound for (3.4):

$$\limsup_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \pm \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n \right\} \leq \frac{\theta^4}{96\sigma^2}. \quad (3.13)$$

On the other hand, by (2.8)

$$\sum_{m=1}^{\infty} \frac{\theta^{2m}}{(2m)!} \left(\frac{b_n}{n^3}\right)^{m/2} \mathbb{E} \tilde{H}_n^{2m} \geq \sum_{m=1}^{\infty} \frac{1}{m!} \frac{\theta^{2m}}{2^m} \left(\frac{b_n}{n^3}\right)^{m/2} A_m(n). \tag{3.14}$$

Write

$$\bar{\theta} = \theta \exp \left\{ -\frac{\theta^2 K_n^2}{8} \sqrt{\frac{b_n}{n^3}} \right\}.$$

By (2.9),

$$\begin{aligned} & \sum_{m=1}^{\infty} \frac{1}{m!} \frac{\bar{\theta}^{2m}}{2^m} \left(\frac{b_n}{n^3}\right)^{m/2} \mathbb{E} \tilde{Q}_n^m \\ & \leq \sum_{m=1}^{\infty} \frac{\bar{\theta}^{2m}}{2^m} \left(\frac{b_n}{n^3}\right)^{m/2} \sum_{l=1}^m \binom{m}{l} \left(\frac{l K_n^2}{2}\right)^{m-l} A_l(n) \\ & = \sum_{l=1}^{\infty} \frac{1}{l!} \left(\frac{\bar{\theta}^2}{2}\right)^l \left(\frac{b_n}{n^3}\right)^{l/2} A_l(n) \sum_{m=l}^{\infty} \frac{1}{(m-l)!} \left(\frac{\bar{\theta}^2}{2}\right)^{m-l} \left(\frac{b_n}{n^3}\right)^{(m-l)/2} \left(\frac{l K_n^2}{2}\right)^{m-l} \\ & = \sum_{l=1}^{\infty} \frac{1}{l!} \left(\frac{\bar{\theta}^2}{2}\right)^l \left(\frac{b_n}{n^3}\right)^{l/2} A_l(n) \exp \left\{ l \frac{\bar{\theta}^2 K_n^2}{4} \sqrt{\frac{b_n}{n^3}} \right\} \\ & \leq \sum_{l=1}^{\infty} \frac{1}{l!} \left(\frac{1}{2}\right)^l \left(\bar{\theta} \exp \left\{ \frac{\theta^2 K_n^2}{8} \sqrt{\frac{b_n}{n^3}} \right\}\right)^{2l} \left(\frac{b_n}{n^3}\right)^{l/2} A_l(n) \\ & = \sum_{l=1}^{\infty} \frac{1}{l!} \frac{\theta^{2l}}{2^l} \left(\frac{b_n}{n^3}\right)^{l/2} A_l(n), \end{aligned} \tag{3.15}$$

where the second inequality follows from the fact that $\bar{\theta} \leq \theta$.

Combining (3.14) and (3.15),

$$\begin{aligned} \sum_{m=0}^{\infty} \frac{\theta^{2m}}{(2m)!} \left(\frac{b_n}{n^3}\right)^{m/2} \mathbb{E} \tilde{H}_n^{2m} & \geq \mathbb{E} \exp \left\{ \frac{\bar{\theta}^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n \right\} \\ & = (1 + o(1)) \mathbb{E} \exp \left\{ \frac{\theta^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n \right\}, \end{aligned} \tag{3.16}$$

where the second step follows from the estimate (notice that (3.9) implies that $\tilde{Q}_n \leq n K_n / 2$)

$$\begin{aligned} \frac{\bar{\theta}^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n & = \frac{\theta^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n - \frac{\theta^2}{2} \left[1 - \exp \left\{ \frac{\theta^2 K_n^2}{4} \sqrt{\frac{b_n}{n^3}} \right\} \right] \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n \\ & \geq \frac{\theta^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n - \frac{\theta^2}{2} \frac{\theta^2 K_n^2}{4} \sqrt{\frac{b_n}{n^3}} \frac{b_n^{1/2}}{n^{3/2}} \frac{n K_n}{2} = \frac{\theta^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n - o(1), \quad n \rightarrow \infty. \end{aligned} \tag{3.17}$$

In view of (3.3), by (3.16) we conclude that

$$\mathbb{E} \exp \left\{ \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n \right\} \geq (1 + o(1)) \mathbb{E} \exp \left\{ \frac{\theta^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \tilde{Q}_n \right\}. \tag{3.18}$$

Getting the lower bound for the negative coefficients $-\theta$ is harder. To do this we need to control the terms with odd powers. Replacing m by $2m + 1$ in (2.7) we obtain

$$\begin{aligned}
& \sum_{m=1}^{\infty} \frac{\theta^{2m+1}}{(2m+1)!} \left(\frac{b_n}{n^3}\right)^{(2m+1)/4} \mathbb{E} \tilde{H}_n^{2m+1} \\
& \leq \sum_{m=1}^{\infty} \frac{\theta^{2m+1}}{2^{2m+1}} \left(\frac{b_n}{n^3}\right)^{(2m+1)/4} \sum_{l=1}^m \frac{1}{l!} K_n^{2(m-l)+1} 2^l C^{(2m-2l+1)/2} \binom{2m-l}{2m-2l+1} \mathbb{E} \tilde{Q}_n^l \\
& = \frac{\theta \sqrt{C}}{2} K_n \left(\frac{b_n}{n^3}\right)^{1/4} \sum_{l=1}^{\infty} \frac{1}{l!} \frac{\theta^{2l}}{2^l} \left(\frac{b_n}{n^3}\right)^{l/2} \mathbb{E} \tilde{Q}_n^l \\
& \quad \times \sum_{m=l}^{\infty} \left(\frac{\theta}{2}\right)^{2(m-l)} C^{m-l} K_n^{2(m-l)} \left(\frac{b_n}{n^3}\right)^{(m-l)/2} \binom{2m-l}{2m-2l+1}. \tag{3.19}
\end{aligned}$$

Noticing that

$$\binom{2m-l}{2m-2l+1} = \frac{2m-l}{2m-2l+1} \binom{2m-l-1}{2m-2l} \leq l \binom{2m-l-1}{2m-2l}$$

we have

$$\begin{aligned}
& \sum_{m=l}^{\infty} \left(\frac{\theta}{2}\right)^{2(m-l)} C^{m-l} K_n^{2(m-l)} \left(\frac{b_n}{n^3}\right)^{(m-l)/2} \binom{2m-l-1}{2m-2l} \\
& = l \sum_{m=0}^{\infty} \left(\frac{\theta}{2}\right)^{2m} K_n^{2m} \left(\frac{b_n}{n^3}\right)^{m/2} C^m \binom{2m+l-1}{2m} \\
& \leq l \sum_{m=0}^{\infty} \left(\frac{\theta}{2}\right)^m K_n^m \left(\frac{b_n}{n^3}\right)^{m/4} C^m \binom{m+l-1}{m} \\
& = l \left(1 - \frac{\sqrt{C}\theta K_n b_n^{1/4}}{2n^{3/4}}\right)^{-(l-1)},
\end{aligned}$$

where the last step follows from (3.7).

By (3.17), therefore,

$$\begin{aligned}
& \sum_{m=1}^{\infty} \frac{\theta^{2m+1}}{(2m+1)!} \left(\frac{b_n}{n^3}\right)^{(2m+1)/4} \mathbb{E} \tilde{H}_n^{2m+1} \\
& \leq \sqrt{C} \frac{\theta^3}{4} K_n \left(\frac{b_n}{n^3}\right)^{3/4} \sum_{l=1}^{\infty} \frac{1}{(l-1)!} \left(\frac{\theta^2}{2}\right)^{l-1} \left(\frac{b_n}{n^3}\right)^{(l-1)/2} \\
& \quad \times \left(1 - \frac{\sqrt{C}\theta K_n b_n^{1/4}}{2n^{3/4}}\right)^{-(l-1)} \mathbb{E} \tilde{Q}_n^l \\
& \leq \sqrt{C} \frac{\theta^3}{4} K_n \left(\frac{b_n}{n^3}\right)^{3/4} \mathbb{E} \left(\tilde{Q}_n \exp \left\{ \frac{\theta^2}{2} \frac{b_n^{1/2}}{n^{3/2}} \left(1 - \frac{\sqrt{C}\theta K_n b_n^{1/4}}{2n^{3/4}}\right)^{-1} \tilde{Q}_n \right\} \right).
\end{aligned}$$

By the estimate $\tilde{Q}_n \leq \frac{1}{2} n K_n$ and by the assumption (1.8) we have that

$$K_n \left(\frac{b_n}{n^3}\right)^{3/4} Q_n \leq \frac{M^2 b_n^{7/4}}{2 n^{1/4}} \rightarrow 0, \quad n \rightarrow \infty.$$

In view of (2.13), we have proved that

$$\sum_{m=0}^{\infty} \frac{\theta^{2m+1}}{(2m+1)!} \left(\frac{b_n}{n^3}\right)^{(2m+1)/4} \mathbb{E} \tilde{H}_n^{2m+1} = o\left(\mathbb{E} \exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} \tilde{Q}_n\right\}\right), \quad n \rightarrow \infty. \tag{3.20}$$

This, together with (3.16), yields

$$\mathbb{E} \exp\left\{-\theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n\right\} \geq (1 + o(1)) \mathbb{E} \exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} \tilde{Q}_n\right\}, \quad n \rightarrow \infty.$$

Combining this with (3.18),

$$\mathbb{E} \exp\left\{\pm \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n\right\} \geq (1 + o(1)) \mathbb{E} \exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} \tilde{Q}_n\right\}, \quad n \rightarrow \infty. \tag{3.21}$$

To estimate the right-hand side of (3.21), notice that

$$\begin{aligned} \mathbb{E} \exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} \tilde{Q}_n\right\} &\geq \mathbb{E}\left(\exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} Q_n\right\} 1_{\{\sup_{x \in \mathbb{Z}} l(n,x) \leq K_n\}}\right) \\ &= \mathbb{E} \exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} Q_n\right\} - \mathbb{E}\left(\exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} Q_n\right\} 1_{\{\sup_{x \in \mathbb{Z}} l(n,x) > K_n\}}\right). \end{aligned}$$

Consequently, by (3.12)

$$\begin{aligned} &\max\left\{\liminf_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp\left\{\frac{\theta^2 b_n^{1/2}}{4 n^{3/2}} \tilde{Q}_n\right\}, \limsup_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E}\left(\exp\left\{\frac{\theta^2 b_n^{1/2}}{4 n^{3/2}} Q_n\right\} 1_{\{\sup_{x \in \mathbb{Z}} l(n,x) > K_n\}}\right)\right\} \\ &\geq \frac{\theta^4}{96\sigma^2}. \end{aligned} \tag{3.22}$$

By the Cauchy–Schwarz inequality,

$$\begin{aligned} &\mathbb{E}\left(\exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} Q_n\right\} 1_{\{\sup_{x \in \mathbb{Z}} l(n,x) > K_n\}}\right) \\ &\leq \left(\mathbb{E} \exp\left\{\theta^2 \frac{b_n^{1/2}}{n^{3/2}} Q_n\right\}\right)^{1/2} \left(\mathbb{P}\left\{\sup_{x \in \mathbb{Z}} l(n,x) > K_n\right\}\right)^{1/2}. \end{aligned}$$

Hence, (3.2) and (3.12) imply that

$$\limsup_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E}\left(\exp\left\{\frac{\theta^2 b_n^{1/2}}{2 n^{3/2}} Q_n\right\} 1_{\{\sup_{x \in \mathbb{Z}} l(n,x) > K_n\}}\right) = -\infty.$$

In view of (3.22),

$$\liminf_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp\left\{\frac{\theta^2 b_n^{1/2}}{4 n^{3/2}} \tilde{Q}_n\right\} \geq \frac{\theta^4}{96\sigma^2}. \tag{3.23}$$

Combining (3.21) and (3.23) gives the desired lower bound

$$\liminf_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp\left\{\pm \theta \frac{b_n^{1/4}}{n^{3/4}} \tilde{H}_n\right\} \geq \frac{\theta^4}{96\sigma^2}. \tag{3.24}$$

Therefore, (3.4) follows from (3.13) and (3.24).

Case $d = 2$.

Similar to (3.8) and (3.21), respectively,

$$\mathbb{E} \exp \left\{ \pm \theta \sqrt{\frac{b_n}{n \log n}} \tilde{H}_n \right\} \leq \mathbb{E} \exp \left\{ \frac{\theta^2}{2} \left(1 - \frac{\sqrt{C} \theta K_n}{2} \sqrt{\frac{b_n}{n \log n}} \right)^{-1} \frac{b_n}{n \log n} \tilde{Q}_n \right\} \tag{3.25}$$

and

$$\mathbb{E} \exp \left\{ \pm \theta \sqrt{\frac{b_n}{n \log n}} \tilde{H}_n \right\} \geq (1 + o(1)) \mathbb{E} \exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n \log n} \tilde{Q}_n \right\}, \quad n \rightarrow \infty. \tag{3.26}$$

Applying Jensen’s inequality on the right-hand side of (3.26),

$$\mathbb{E} \exp \left\{ \theta \sqrt{\frac{b_n}{n \log n}} \tilde{H}_n \right\} \geq (1 + o(1)) \exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n \log n} \mathbb{E} \tilde{Q}_n \right\}, \quad \theta \in \mathbb{R}. \tag{3.27}$$

By the fact (implied by (3.2)) that $\mathbb{P}\{\sup_{x \in \mathbb{Z}^2} l(n, x) > K_n\} \rightarrow 0, n \rightarrow \infty$, we have that

$$\mathbb{E} \tilde{Q}_n \sim \mathbb{E} Q_n \sim (2\pi)^{-1} \det \Gamma^{-1/2} n \log n, \quad n \rightarrow \infty,$$

where the second step follows from (2.31).

Consequently,

$$\liminf_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \pm \theta \sqrt{\frac{b_n}{n \log n}} \tilde{H}_n \right\} \geq \frac{\theta^2}{4\pi \sqrt{\det \Gamma}}. \tag{3.28}$$

On the other hand, recall the fact (Lemma 2.3 in [3]) that

$$\mathbb{E} \exp \left\{ \frac{\lambda}{n} |Q_n - \mathbb{E} Q_n| \right\} < \infty$$

for some $\lambda > 0$. By the assumption (1.10) we have

$$\begin{aligned} \mathbb{E} \exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n \log n} Q_n \right\} &\leq \exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n \log n} \mathbb{E} Q_n \right\} \mathbb{E} \exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n \log n} |Q_n - \mathbb{E} Q_n| \right\} \\ &= O \left(\exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n \log n} \mathbb{E} Q_n \right\} \right), \quad n \rightarrow \infty. \end{aligned} \tag{3.29}$$

Combining this with (3.25) gives

$$\limsup_{n \rightarrow \infty} \frac{1}{b_n} \log \mathbb{E} \exp \left\{ \pm \theta \sqrt{\frac{b_n}{n \log n}} \tilde{H}_n \right\} \leq \frac{\theta^2}{4\pi \sqrt{\det \Gamma}}. \tag{3.30}$$

Thus, (3.5) follows from (3.28) and (3.30).

Case $d \geq 3$.

The treatment in the case $d \geq 3$ is almost same as the one given in the case $d = 2$, except that here we use

$$\mathbb{E} \exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n} \tilde{Q}_n \right\} = O \left(\exp \left\{ \frac{\theta^2}{2} \frac{b_n}{n} \mathbb{E} Q_n \right\} \right), \quad n \rightarrow \infty, \tag{3.31}$$

instead of (3.29).

We end this section with the proof of (3.31). Notice that

$$\begin{aligned} \mathbb{E} \exp \left\{ \frac{\theta^2 b_n}{2n} \tilde{Q}_n \right\} &\leq 1 + \mathbb{E} \left(\exp \left\{ \frac{\theta^2 b_n}{2n} Q_n \right\} 1_{\{\sup_{x \in \mathbb{Z}^d} l(n,x) \leq K_n\}} \right) \\ &\leq 1 + \mathbb{E} \exp \left\{ \frac{\theta^2 b_n}{2n} \mathbb{E} \tilde{Q}_n \right\} \mathbb{E} \left(\exp \left\{ \frac{\theta^2 b_n}{2n} |Q_n - \mathbb{E} Q_n| \right\} 1_{\{\sup_{x \in \mathbb{Z}^d} l(n,x) \leq K_n\}} \right). \end{aligned}$$

Therefore, we need only to prove that

$$\sup_n \mathbb{E} \left(\exp \left\{ \frac{\theta^2 b_n}{2n} |Q_n - \mathbb{E} Q_n| \right\} 1_{\{\sup_{x \in \mathbb{Z}^d} l(n,x) \leq K_n\}} \right) < \infty. \tag{3.32}$$

By the fact that $Q_n \leq 2^{-1} n K_n$ on the event $\{\sup_{x \in \mathbb{Z}^d} l(n,x) \leq K_n\}$, we have $|Q_n - \mathbb{E} Q_n| \leq 2^{-1} n K_n$. Consequently,

$$\begin{aligned} &\mathbb{E} \left(\exp \left\{ \frac{\theta^2 b_n}{2n} |Q_n - \mathbb{E} Q_n| \right\} 1_{\{\sup_{x \in \mathbb{Z}^d} l(n,x) \leq K_n\}} \right) \\ &\leq \mathbb{E} \exp \left\{ \frac{\theta^2 b_n}{2n} (2^{-1} n K_n)^{1/3} |Q_n - \mathbb{E} Q_n|^{2/3} \right\}. \end{aligned}$$

Finally, (3.32) follows from Theorem 5.2 and our assumptions on $\{b_n\}$ given in Theorem 1.2.

4. Laws of the iterated logarithm

The following Lévy type inequality is needed in our proof of the upper bounds in Theorem 1.3.

Lemma 4.1. *For any $s, t > 0$ and integer $n \geq 2$,*

$$\min_{1 \leq k \leq n} \mathbb{P}\{|H_k| \leq s\} \mathbb{P}\left\{\max_{1 \leq l \leq n} |H_l| \geq s + t\right\} \leq 2\mathbb{P}\{|H_n| \geq t\}. \tag{4.1}$$

Proof. Write

$$\tau = \inf\{l \geq 1; |H_l| \geq s + t\}$$

and notice that for each $1 \leq l \leq n$,

$$\sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j = S_k\}} \stackrel{d}{=} H_{n-l},$$

where we follow the convention that both sides are zero if $l = n - 1$ or $n = l$. Thus

$$\begin{aligned} &\min_{1 \leq k \leq n} \mathbb{P}\{|H_k| \leq s\} \mathbb{P}\left\{\max_{1 \leq l \leq n} |H_l| \geq s + t\right\} \\ &= \sum_{l=1}^n \min_{1 \leq k \leq n} \mathbb{P}\{|H_k| \leq s\} \mathbb{P}\{\tau = l\} \\ &\leq \sum_{l=1}^n \mathbb{P}\left\{\left| \sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j = S_k\}} \right| \leq s\right\} \mathbb{P}\{\tau = l\} \\ &= \sum_{l=1}^n \mathbb{P}\left\{\tau = l, \left| \sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j = S_k\}} \right| \leq s\right\}, \end{aligned} \tag{4.2}$$

where the last step follows from independence between $\{\tau = l\}$ and

$$\sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j=S_k\}}.$$

For each $1 \leq l \leq n$, write

$$H_n^{(l)} = \sum_{1 \leq j < k \leq l} \omega_j \omega_k 1_{\{S_j=S_k\}} + \sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j=S_k\}} - \sum_{j=1}^l \sum_{k=l+1}^n \omega_j \omega_k 1_{\{S_j=S_k\}}.$$

Notice that

$$\begin{aligned} & \left\{ \tau = l, \left| \sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j=S_k\}} \right| \leq s \right\} \\ & \subset \left\{ \tau = l, \left| \sum_{1 \leq j < k \leq l} \omega_j \omega_k 1_{\{S_j=S_k\}} + \sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j=S_k\}} \right| \geq t \right\} \\ & \subset \mathbb{P}\{\tau = l, |H_n| + |H_n^{(l)}| \geq 2t\}. \end{aligned}$$

Hence,

$$\begin{aligned} & \mathbb{P}\left\{ \tau = l, \left| \sum_{l+1 \leq j < k \leq n} \omega_j \omega_k 1_{\{S_j=S_k\}} \right| \leq s \right\} \\ & \leq \mathbb{P}\{\tau = l, |H_n| \geq t\} + \mathbb{P}\{\tau = l, |H_n^{(l)}| \geq t\}. \end{aligned} \tag{4.3}$$

We now claim that

$$\mathbb{P}\{\tau = l, |H_n^{(l)}| \geq t\} = \mathbb{P}\{\tau = l, |H_n| \geq t\}. \tag{4.4}$$

Indeed, (4.4) follows from the fact that the random vectors

$$(\omega_1, \dots, \omega_n) \quad \text{and} \quad (\omega_1, \dots, \omega_l, -\omega_{l+1}, \dots, -\omega_n)$$

have the same distribution, and that replacing the first vector by the second does not change the event $\{\tau = l\}$ but changes H_n into $H_n^{(l)}$.

Finally, by (4.2), (4.3) and (4.4),

$$\begin{aligned} & \min_{1 \leq k \leq n} \mathbb{P}\{|H_k| \leq s\} \mathbb{P}\left\{ \max_{1 \leq l \leq n} |H_l| \geq s + t \right\} \\ & \leq 2 \sum_{l=1}^n \mathbb{P}\{\tau = l, |H_n| \geq t\} \leq 2\mathbb{P}\{|H_n| \geq t\}. \end{aligned} \quad \square$$

Proof of Theorem 1.3. Due to similarity we only consider the case $d = 1$. To prove the upper bound in (1.13), it suffices to show

$$\limsup_{n \rightarrow \infty} \frac{|H_n|}{(n \log \log n)^{3/4}} \leq \frac{2^{3/4}}{3} \sigma^{-1/2} \quad \text{a.s.} \tag{4.5}$$

Let $\theta > 0$ and

$$\lambda_1 > \frac{2^{3/4}}{3} \sigma^{-1/2}$$

be fixed but arbitrary. Write $n_k = \lceil \theta^k \rceil$ for $k = 1, 2, \dots$. Take $\varepsilon > 0$ small enough so

$$\lambda_1 - \varepsilon > \frac{2^{3/4}}{3} \sigma^{-1/2}.$$

By Theorem 1.1,

$$\min_{1 \leq m \leq n} \mathbb{P}\{|H_m| \leq \varepsilon (n_k \log \log n_k)^{3/4}\} \geq \frac{1}{2}$$

as k is sufficiently large. By Lemma 4.1, therefore,

$$\mathbb{P}\left\{\max_{1 \leq l \leq n_k} |H_l| \geq \lambda_1 (n_k \log \log n_k)^{3/4}\right\} \leq 4\mathbb{P}\{|H_{n_k}| \geq (\lambda_1 - \varepsilon)(n_k \log \log n_k)^{3/4}\}.$$

By (1.7) in Theorem 1.2 (with $b_n = \log \log n$),

$$\sum_k \mathbb{P}\left\{\max_{1 \leq l \leq n_k} |H_l| \geq \lambda_1 (n_k \log \log n_k)^{3/4}\right\} < \infty.$$

By the Borel–Cantelli lemma,

$$\limsup_{k \rightarrow \infty} \frac{1}{(n_k \log \log n_k)^{3/4}} \max_{1 \leq l \leq n_k} |H_l| \leq \lambda_1 \quad \text{a.s.}$$

For any large integer n , if $n_k \leq n \leq n_{k+1}$, then

$$\frac{|H_n|}{(n \log \log n)^{3/4}} \leq (\theta^{3/4} + o(1)) \frac{1}{(n_{k+1} \log \log n_{k+1})^{3/4}} \max_{1 \leq l \leq n_{k+1}} |H_l|.$$

So we have

$$\limsup_{n \rightarrow \infty} \frac{|H_n|}{(n \log \log n)^{3/4}} \leq \theta^{3/4} \lambda_1 \quad \text{a.s.}$$

Letting $\theta \rightarrow 1^+$ and $\lambda_1 \rightarrow 2^{3/4} 3^{-1} \sigma^{-1/2}$ gives (4.5).

We only prove the lower bound for H_n :

$$\limsup_{n \rightarrow \infty} \frac{H_n}{(n \log \log n)^{3/4}} \geq \frac{2^{3/4}}{3} \sigma^{-1/2} \quad \text{a.s.} \tag{4.6}$$

as the proof of the lower bound for $-H_n$ is analogous.

Let n_k be defined as above (but with large constant $\theta > 0$) and let the constant λ_2 satisfying

$$\lambda_2 < \frac{2^{3/4}}{3} \sigma^{-1/2}.$$

Let $\varepsilon > 0$ be small enough so

$$\lambda_2 + \varepsilon < \frac{2^{3/4}}{3} \sigma^{-1/2}.$$

Notice that

$$\sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j \mathbf{1}_{\{S_i=S_j\}} \stackrel{d}{=} H_{n_{k+1}-n_k}.$$

As $\theta > 0$ and k are sufficiently large,

$$\begin{aligned} & \mathbb{P} \left\{ \sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j 1_{\{S_i=S_j\}} \geq \lambda_2 (n_{k+1} \log \log n_{k+1})^{3/4} \right\} \\ & \geq \mathbb{P} \{ H_{n_{k+1}-n_k} \geq (\lambda_2 + \varepsilon) ((n_{k+1} - n_k) \log \log (n_{k+1} - n_k))^{3/4} \}. \end{aligned}$$

By (1.7) in Theorem 1.2 again,

$$\sum_k \mathbb{P} \left\{ \sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j 1_{\{S_i=S_j\}} \geq \lambda_2 (n_{k+1} \log \log n_{k+1})^{3/4} \right\} = \infty.$$

Notice that

$$\sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j 1_{\{S_i=S_j\}}, \quad k = 1, 2, \dots,$$

is an independent sequence. By the Borel–Cantelli lemma,

$$\limsup_{k \rightarrow \infty} \frac{1}{(n_{k+1} \log \log n_{k+1})^{3/4}} \sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j 1_{\{S_i=S_j\}} \geq \lambda_2 \quad \text{a.s.}$$

In addition, (4.5) implies that

$$\limsup_{k \rightarrow \infty} \frac{1}{(n_{k+1} \log \log n_{k+1})^{3/4}} |H_{n_k}| \leq \theta^{-3/4} \frac{2^{3/4}}{3} \sigma^{-1/2} \quad \text{a.s.}$$

Consequently,

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \frac{1}{(n_{k+1} \log \log n_{k+1})^{3/4}} \left\{ H_{n_k} + \sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j 1_{\{S_i=S_j\}} \right\} \\ & \geq \lambda_2 - \theta^{-3/4} \frac{2^{3/4}}{3} \quad \text{a.s.} \end{aligned} \tag{4.7}$$

Recall the notation

$$H_{n_{k+1}}^{(n_k)} = H_{n_k} + \sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j 1_{\{S_i=S_j\}} - \sum_{i=1}^{n_k} \sum_{j=n_k+1}^{n_{k+1}} 1_{\{S_i=S_j\}}.$$

We have

$$H_{n_k} + \sum_{n_k+1 \leq i < j \leq n_{k+1}} \omega_i \omega_j 1_{\{S_i=S_j\}} = \frac{H_{n_{k+1}} + H_{n_{k+1}}^{(n_k)}}{2}.$$

Therefore, by (4.7)

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \frac{H_{n_{k+1}}}{(n_{k+1} \log \log n_{k+1})^{3/4}} + \limsup_{k \rightarrow \infty} \frac{H_{n_{k+1}}^{(n_k)}}{(n_{k+1} \log \log n_{k+1})^{3/4}} \\ & \geq 2 \left(\lambda_2 - \theta^{-3/4} \frac{2^{3/4}}{3} \sigma^{-1/2} \right) \quad \text{a.s.} \end{aligned} \tag{4.8}$$

On the other hand, notice that for each k ,

$$H_{n_{k+1}}^{(n_k)} \stackrel{d}{=} H_{n_{k+1}}.$$

By Theorem 1.2 (with $b_n = \log \log n$)

$$\begin{aligned} & \sum_k \mathbb{P}\{H_{n_{k+1}}^{(n_k)} \geq \lambda(n_{k+1} \log \log n_{k+1})^{3/4}\} \\ &= \sum_k \mathbb{P}\{H_{n_{k+1}} \geq \lambda(n_{k+1} \log \log n_{k+1})^{3/4}\} < \infty \quad \forall \lambda > \frac{2^{3/4}}{3} \sigma^{-1/2}. \end{aligned}$$

By the Borel–Cantelli lemma,

$$\limsup_{k \rightarrow \infty} \frac{H_{n_{k+1}}^{(n_k)}}{(n_{k+1} \log \log n_{k+1})^{3/4}} \leq \frac{2^{3/4}}{3} \sigma^{-1/2} \quad \text{a.s.}$$

Combining this with (4.8) yields

$$\limsup_{k \rightarrow \infty} \frac{H_{n_{k+1}}}{(n_{k+1} \log \log n_{k+1})^{3/4}} \geq 2 \left(\lambda_2 - \theta^{-3/4} \frac{2^{3/4}}{3} \sigma^{-1/2} \right) - \frac{2^{3/4}}{3} \sigma^{-1/2} \quad \text{a.s.}$$

Consequently,

$$\limsup_{n \rightarrow \infty} \frac{H_n}{(n \log \log n)^{3/4}} \geq 2 \left(\lambda_2 - \theta^{-3/4} \frac{2^{3/4}}{3} \sigma^{-1/2} \right) - \frac{2^{3/4}}{3} \sigma^{-1/2} \quad \text{a.s.}$$

Letting $\theta \rightarrow \infty$ and $\lambda_2 \rightarrow 3^{-1} 2^{3/4} \sigma^{-1/2}$ on the right-hand side gives (4.6). □

5. Self-intersection in high dimension

From Theorem 1.1, we have seen that the multi-dimensional case ($d \geq 2$) is different from the case $d = 1$. Here is the reason: contrary to (2.18), a concentration phenomenon appearing as

$$Q_n / \mathbb{E}Q_n \xrightarrow{p} 1$$

takes over when $d \geq 2$. The concentration also plays a role in the moderate deviations (Theorem 1.2) as $d \geq 2$. In our treatment given in Sections 2 and 3, Q_n is replaced by $\mathbb{E}Q_n$ when $d \geq 2$. To justify such action, we need to show that Q_n and $\mathbb{E}Q_n$ are asymptotically close enough. More precisely, our concern in this section is the central limit theorem and the exponential integrability for the renormalized self-intersection local time $Q_n - \mathbb{E}Q_n$. The case $d = 2$ has been investigated. In [20], it was proved that

$$\frac{1}{n} (Q_n - \mathbb{E}Q_n) \xrightarrow{d} (\det \Gamma)^{-1/2} \gamma_1, \tag{5.1}$$

where γ_t is the renormalized self-intersection local times

$$\gamma_t = \iint_{0 \leq r < s \leq t} \delta_0(W(r) - W(s)) \, dr \, ds - \mathbb{E} \iint_{0 \leq r < s \leq t} \delta_0(W(r) - W(s)) \, dr \, ds, \quad t \geq 0,$$

run by a planar Brownian motion $W(t)$. In [3], it was proved that

$$\mathbb{E} \exp \left\{ \frac{\lambda}{n} |Q_n - \mathbb{E}Q_n| \right\} < \infty$$

for some $\lambda > 0$. In the following discussion, we focus our attention on the case $d \geq 3$. Apart from its role in the charged polymers, the study of self-intersection local time is an important subject for its own sake. Our involvement on the integrability problems is also motivated by the recent interest [1,2] in the large deviations for Q_n in the case $d \geq 3$.

In high-dimensional cases defined by $d \geq 3$, a related object is the range $\#\{S[1, n]\}$ given by

$$\#\{S[1, n]\} = \#\{S_1, \dots, S_n\}.$$

It has been known [13,14,17] that

$$\frac{1}{\sqrt{n \log n}} (\#\{S[1, n]\} - \mathbb{E}\#\{S[1, n]\}) \xrightarrow{d} c_1 U \tag{5.2}$$

as $d = 3$; and

$$\frac{1}{\sqrt{n}} (\#\{S[1, n]\} - \mathbb{E}\#\{S[1, n]\}) \xrightarrow{d} c_2 U \tag{5.3}$$

as $d = 4$, where $U \sim N(0, 1)$. It is now widely believed that as $d \geq 3$, Q_n and $\#\{S[1, n]\}$ have very similar behaviors. In particular, we have

Theorem 5.1. As $d = 3$,

$$\frac{1}{\sqrt{n \log n}} (Q_n - \mathbb{E}Q_n) \xrightarrow{d} \lambda_1 U; \tag{5.4}$$

as $d \geq 4$,

$$\frac{1}{\sqrt{n}} (Q_n - \mathbb{E}Q_n) \xrightarrow{d} \lambda_2 U, \tag{5.5}$$

where

$$\begin{aligned} \lambda_1 &= \frac{1}{\sqrt{2\pi^2 \det(\Gamma)}}, \\ \lambda_2 &= \sqrt{3G^2(0) + G(0) + 2 \sum_{x \in \mathbb{Z}^d} G^3(x)}, \\ G(x) &= \sum_{k=1}^{\infty} \mathbb{P}\{S_k = x\}, \quad x \in \mathbb{Z}^d. \end{aligned}$$

Proof. The proof is inspired by some ideas used in [13,17] in the setting of the ranges. Due to similarity we only consider the case $d = 3$. Let $\{\gamma_n\}$ be a positive sequence such that

$$\gamma_n \rightarrow \infty \quad \text{and} \quad \gamma_n = o(\sqrt{\log n}), \quad n \rightarrow \infty.$$

Let $0 = n_0 < n_1 < \dots < n_{\gamma_n} = n$ be an integer partition of $[0, n]$ such that for each $1 \leq i \leq \gamma_n$, $n - i - n_{i-1} = \lfloor n\gamma_n^{-1} \rfloor$ or $\lfloor n\gamma_n^{-1} \rfloor + 1$. Then

$$Q_n = \sum_{i=1}^{\gamma_n} \sum_{n_{i-1} < j < k \leq n_i} 1_{\{S_j=S_k\}} + \sum_{i=1}^{\gamma_n-1} \sum_{j=n_{i-1}+1}^{n_i} \sum_{k=n_i+1}^n 1_{\{S_j=S_k\}}. \tag{5.6}$$

For each $1 \leq i \leq \gamma_n - 1$,

$$\sum_{j=n_{i-1}+1}^{n_i} \sum_{k=n_i+1}^n 1_{\{S_j=S_k\}} \stackrel{d}{=} \sum_{j=1}^{n_i-n_{i-1}} \sum_{k=1}^{n-n_i} 1_{\{S_j=S'_k\}} \leq \sum_{j=1}^n \sum_{k=1}^n 1_{\{S_j=S'_k\}},$$

where $\{S'_k\}$ is an independent copy of $\{S_k\}$. Thus,

$$\mathbb{E} \left(\sum_{i=1}^{\gamma_n-1} \sum_{j=n_{i-1}+1}^{n_i} \sum_{k=n_i+1}^n 1_{\{S_j=S_k\}} \right)^2 \leq \gamma_n^2 \mathbb{E} \left(\sum_{j=1}^n \sum_{k=1}^n 1_{\{S_j=S'_k\}} \right)^2 = o(n \log n). \quad (5.7)$$

In addition, notice that the random variables

$$\sum_{n_{i-1} < j < k \leq n_i} 1_{\{S_j=S_k\}}, \quad i = 1, 2, \dots, \gamma_n,$$

are independent with

$$\sum_{n_{i-1} < j < k \leq n_i} 1_{\{S_j=S_k\}} \stackrel{d}{=} Q_{n_i-n_{i-1}}, \quad i = 1, 2, \dots, \gamma_n.$$

By Lemma 5.1 and by (5.7),

$$\text{Var} \left(\sum_{i=1}^{\gamma_n} \sum_{n_{i-1} < j < k \leq n_i} 1_{\{S_j=S_k\}} \right) \sim \lambda_1^2 n \log n.$$

By Theorem 5.2, we can check the Lederberg condition. Hence,

$$\sum_{i=1}^{\gamma_n} \sum_{n_{i-1} < j < k \leq n_i} 1_{\{S_j=S_k\}} / n \log n \xrightarrow{d} \lambda_1 U. \quad (5.8)$$

Finally, (5.4) follows from (5.6), (5.7) and (5.8). \square

Lemma 5.1. Let λ_1 and λ_2 be given in Theorem 5.1, As $d = 3$,

$$\text{Var}(Q_n) \sim \lambda_1^2 n \log n, \quad n \rightarrow \infty. \quad (5.9)$$

As $d \geq 4$,

$$\text{Var}(Q_n) \sim \lambda_2^2 n, \quad n \rightarrow \infty. \quad (5.10)$$

Proof. Notice that

$$\begin{aligned} Q_n &= \sum_{j=1}^{n-1} \sum_{k=j+1}^{\infty} 1_{\{S_k=S_j\}} - \sum_{j=1}^{n-1} \sum_{k=n+1}^{\infty} 1_{\{S_k=S_j\}} \\ &= \sum_{j=1}^{n-1} Z_j - \sum_{j=1}^{n-1} W_j^n \quad (\text{say}). \end{aligned} \quad (5.11)$$

Write $p_k(x) = \mathbb{P}\{S_k = x\}$ and recall that $\{S'_k\}$ is an independent copy of $\{S_k\}$. For any $i \leq j \leq k \leq n$,

$$\begin{aligned}
 \mathbb{E}W_i^n W_j^n &= \sum_{k,l=n+1}^{\infty} \mathbb{P}\{S_l - S_j = 0, S_k - S_j = S_i - S_j\} \\
 &= \sum_{k,l=n-j+1}^{\infty} \mathbb{P}\{S_l = 0, S_k = S'_{j-i}\} \\
 &= \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{k,l=n-j+1}^{\infty} \mathbb{P}\{S_l = 0, S_k = x\} \\
 &\leq \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{n-j+1 \leq k \leq l < \infty} \mathbb{P}\{S_l = 0, S_k = x\} \\
 &\quad + \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{n-j+1 \leq l \leq k < \infty} \mathbb{P}\{S_l = 0, S_k = x\}.
 \end{aligned} \tag{5.12}$$

For the first term on the right-hand side,

$$\begin{aligned}
 &\sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{n-j+1 \leq k \leq l < \infty} \mathbb{P}\{S_l = 0, S_k = x\} \\
 &= \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{n-j+1 \leq k \leq l < \infty} \mathbb{P}\{S_{l-k} = x\} \mathbb{P}\{S_k = x\} \\
 &\leq C \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{n-j+1 \leq k \leq l < \infty} p_{l-k}(x) \frac{1}{k^{d/2}} \\
 &= C \sum_{n-j+1 \leq k \leq l < \infty} p_{l-k+j-i}(0) \frac{1}{k^{d/2}},
 \end{aligned}$$

where the second step follows from the classic fact that $\sup_{x \in \mathbb{Z}^d} p_k(x) = O(k^{-d/2})$.

As for the second term, a similar estimate yields that

$$\begin{aligned}
 &\sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{n-j+1 \leq l \leq k < \infty} \mathbb{P}\{S_l = 0, S_k = -x\} \\
 &\leq C \sum_{n-j+1 \leq l \leq k < \infty} p_{k-l+j-i}(0) \frac{1}{l^{d/2}}.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 \mathbb{E}W_i^n W_j^n &\leq 2C \sum_{n-j+1 \leq k \leq l < \infty} p_{l-k+j-i}(0) \frac{1}{l^{d/2}} \\
 &= 2 \left(\sum_{k=n-j+1}^{\infty} p_k(0) \right) \left(\sum_{l=j-i}^{\infty} \frac{1}{l^{d/2}} \right) = O((n-j)^{1-d/2} (j-i)^{1-d/2}).
 \end{aligned}$$

Therefore, as $n \rightarrow \infty$,

$$\sum_{j=1}^n \sum_{i=1}^j \mathbb{E}W_i^n W_j^n = \begin{cases} O(n), & d = 3, \\ O((\log n)^2), & d = 4, \\ O(1), & d \geq 5. \end{cases} \tag{5.13}$$

For $1 \leq i \leq j \leq n$,

$$\begin{aligned}
 \text{Cov}(Z_i, Z_j) &= \text{Cov}\left(\sum_{k=j+1}^{\infty} 1_{\{S_k=S_i\}}, \sum_{k=j+1}^{\infty} 1_{\{S_k=S_j\}}\right) \\
 &= \text{Cov}\left(\sum_{k=1}^{\infty} 1_{\{S_k=S'_{j-i}\}}, \sum_{k=1}^{\infty} 1_{\{S_k=0\}}\right) \\
 &= \sum_{k,l=1}^{\infty} \{\mathbb{P}\{S_k = S'_{j-i} S_l = 0\} - p_{k+j-i}(0)p_l(0)\} \\
 &= \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{k,l=1}^{\infty} \mathbb{P}\{S_k = -x S_l = 0\} - G(0) \sum_{k=j-i+1}^{\infty} p_k(0).
 \end{aligned} \tag{5.14}$$

Write

$$\begin{aligned}
 &\sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{k,l=1}^{\infty} \mathbb{P}\{S_k = x S_l = 0\} \\
 &= \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{1 \leq k \leq l < \infty} \mathbb{P}\{S_k = x S_l = 0\} \\
 &\quad + \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{1 \leq l < k < \infty} \mathbb{P}\{S_k = x S_l = 0\}.
 \end{aligned}$$

We have

$$\begin{aligned}
 &\sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{1 \leq k \leq l < \infty} \mathbb{P}\{S_k = x S_l = 0\} \\
 &= \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{1 \leq k \leq l < \infty} p_k(x) p_{l-k}(x) \\
 &= \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) G(x) \sum_{l=0}^{\infty} p_l(x) \\
 &= p_{j-i}(0) G(0) (1 + G(0)) + \sum_{x \neq 0} p_{j-i}(x) G^2(x)
 \end{aligned}$$

and

$$\begin{aligned}
 &\sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{1 \leq l < k < \infty} \mathbb{P}\{S_k = x S_l = 0\} \\
 &= \sum_{x \in \mathbb{Z}^d} p_{j-i}(x) \sum_{1 \leq l < k < \infty} p_{k-l}(x) p_l(0) \\
 &= \sum_{1 \leq l < k < \infty} p_{k-l+j-i}(0) p_l(0) = G(0) \sum_{k=j-i+1}^{\infty} p_k(0).
 \end{aligned}$$

In summary of the argument since (5.14),

$$\text{Cov}(Z_i, Z_j) = p_{j-i}(0) G(0) (1 + G(0)) + \sum_{x \neq 0} p_{j-i}(x) G^2(x).$$

Consequently,

$$\begin{aligned} \text{Var}\left(\sum_{i=1}^n Z_i\right) &= \sum_{i=1}^n \text{Var}(Z_i) + 2 \sum_{1 \leq i < j \leq n} \text{Cov}(Z_i, Z_j) \\ &= nG(0)(1 + G(0)) + 2G(0)(1 + G(0)) \sum_{j=1}^{n-1} G_{n-j}(0) + 2 \sum_{x \neq 0} G^2(x) \sum_{i=1}^{n-1} G_{n-i}(x) \\ &= nG(0)(1 + G(0)) + 2G(0) \sum_{j=1}^{n-1} G_j(0) + 2 \sum_{x \in \mathbb{Z}^d} G^2(x) \sum_{j=1}^{n-1} G_j(x), \end{aligned} \tag{5.15}$$

where

$$G_j(x) = \sum_{i=1}^j p_i(x).$$

When $d \geq 4$,

$$\sum_{x \in \mathbb{Z}^d} G^3(x) < \infty.$$

By (5.15)

$$\text{Var}\left(\sum_{i=1}^n Z_i\right) \sim n \left\{ G(0)(1 + G(0)) + 2G^2(0) + 2 \sum_{x \in \mathbb{Z}^d} G^3(x) \right\}, \quad n \rightarrow \infty.$$

By (5.11) and (5.13), this implies (5.10).

We now consider the case $d = 3$. We use the fact that (p. 308, [21])

$$G(x) \sim (2\pi)^{-1} \det(\Gamma)^{-1/2} \langle x, \Gamma^{-1}x \rangle^{-1/2}, \quad |x| \rightarrow \infty. \tag{5.16}$$

By (5.16),

$$\sum_{\langle x, \Gamma^{-1}x \rangle > j} G(x)^2 G_j(x) = O\left(\frac{1}{j} \sum_{x \in \mathbb{Z}^d} G_j(x)\right) = O(1), \quad j \rightarrow \infty. \tag{5.17}$$

In addition,

$$\begin{aligned} &\sum_{\langle x, \Gamma^{-1}x \rangle \leq j} G(x)^2 (G(x) - G_j(x)) \\ &\leq (G(0) - G_j(0)) \sum_{\langle x, \Gamma^{-1}x \rangle \leq j} G^2(x) \\ &\leq C(G(0) - G_j(0)) \sum_{\langle x, \Gamma^{-1}x \rangle \leq j} \frac{1}{1 + |x|^2} = O(1), \quad j \rightarrow \infty, \end{aligned} \tag{5.18}$$

and

$$\begin{aligned} \sum_{\langle x, \Gamma^{-1}x \rangle \leq j} G^3(x) &\sim \frac{1}{(2\pi)^3 \det(\Gamma)^3} \int_{\{1 \leq \langle x, \Gamma^{-1}x \rangle \leq j\}} \langle x, \Gamma^{-1}x \rangle^{-3/2} dx \\ &= \frac{1}{(2\pi)^3 \det(\Gamma)} \int_{\{1 \leq |y| \leq \sqrt{j}\}} |y|^{-3} dy = \frac{1}{(2\pi)^2 \det(\Gamma)} \log j, \quad j \rightarrow \infty. \end{aligned} \tag{5.19}$$

Combining (5.17), (5.18) and (5.19) gives

$$\sum_{x \in \mathbb{Z}^d} G^2(x) G_j(x) \sim \frac{1}{(2\pi)^2 \det(\Gamma)} \log j, \quad j \rightarrow \infty.$$

By (5.14),

$$\text{Var} \left(\sum_{i=1}^n Z_i \right) \sim \frac{2}{(2\pi)^2 \det(\Gamma)} n \log n, \quad n \rightarrow \infty,$$

which, together with (5.11) and (5.13), implies (5.9). □

We now investigate the integrability of Q_n . Write

$$J_n = \sum_{j=1}^n \sum_{k=1}^n 1_{\{S_j = S'_k\}},$$

where $\{S'_n\}$ is an independent copy of $\{S_n\}$. J_n is known as the intersection local time between two independent trajectories.

Lemma 5.2. *As $d \geq 3$, there is a constant $C_d > 0$ such that*

$$\mathbb{E} J_n^m \leq C_d^m (m!)^{3/2} n^{m/2}, \quad m, n = 1, 2, \dots \tag{5.20}$$

Proof. Recall the fact (p. 3282, [6])

$$\mathbb{E} J_n^m \leq (m!) (1 + \mathbb{E} J_n)^m, \quad m, n = 1, 2, \dots, \tag{5.21}$$

and the fact that

$$\mathbb{E} J_n = \begin{cases} O(\sqrt{n}), & d = 3, \\ O(\log n), & d = 4, \\ O(1), & d \geq 5. \end{cases} \tag{5.22}$$

A trivial and rough summarization of (5.22) gives that $\mathbb{E} J_n = O(\sqrt{n})$ as $d \geq 3$. By (5.21), we obtain a weaker version of (5.20):

$$\mathbb{E} J_n^m \leq C^m (m!)^2 n^{m/2}, \quad m, n = 1, 2, \dots \tag{5.23}$$

To strengthen (5.23) into (5.20), recall (Theorem 5.1, of [6]) that for any integers $m, n_1, \dots, n_a \geq 1$,

$$(\mathbb{E} J_n^m)^{1/2} \leq \sum_{\substack{k_1 + \dots + k_a = m \\ k_1, \dots, k_a \geq 0}} \frac{m!}{k_1! \dots k_a!} (\mathbb{E} J_{n_1}^{k_1})^{1/2} \dots (\mathbb{E} J_{n_a}^{k_a})^{1/2}. \tag{5.24}$$

We first consider the case $n \geq m$. Write $l(m, n) = 1 + \lfloor \frac{n}{m} \rfloor$. By (5.23) there is a $C > 0$ independent of m and n such that

$$\mathbb{E} J_{l(m,n)}^m \leq C^m (m!)^2 \left(\frac{n}{m} \right)^{m/2}.$$

Taking $a = m$ in (5.24) gives

$$\begin{aligned} (\mathbb{E}J_n^m)^{1/2} &\leq \sum_{\substack{k_1+\dots+k_m=m \\ k_1,\dots,k_m \geq 0}} \frac{m!}{k_1! \dots k_m!} (\mathbb{E}J_{l(m,n)}^{k_1})^{1/2} \dots (\mathbb{E}J_{l(m,n)}^{k_m})^{1/2} \\ &\leq m! C^{m/2} \left(\frac{n}{m}\right)^{m/4} \sum_{\substack{k_1+\dots+k_m=m \\ k_1,\dots,k_m \geq 0}} 1 = m! C^{m/2} m^{-m/4} n^{m/4} \binom{2m-1}{m}. \end{aligned}$$

Since $m^m \geq m!$ and

$$\binom{2m-1}{m} \leq 4^m$$

we have established (5.20) with $C_d = 8C$ in the case $n \geq m$.

As for the case $n < m$, the trivial fact $J_n \leq n^2$ leads to the following trivial bound,

$$\mathbb{E}J_n^m \leq n^{2m} \leq m^{3m/2} n^{m/2} \leq C^m (m!)^{3/2} n^{m/2},$$

where the last step follows from the Stirling formula. □

As for the exponential integrability of the renormalized self-intersection local time $Q_n - \mathbb{E}Q_n$, we have the following theorem.

Theorem 5.2. *As $d = 3$,*

$$\sup_n \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n \log n}} |Q_n - \mathbb{E}Q_n|^{2/3} \right\} < \infty \quad \text{for every } \theta > 0. \tag{5.25}$$

As $d \geq 4$,

$$\sup_n \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n}} |Q_n - \mathbb{E}Q_n|^{2/3} \right\} < \infty \quad \text{for some } \theta > 0. \tag{5.26}$$

Proof. The proof given here is radically different from the approach used in the case $d = 2$ (Lemma 2.3 in [3]) where the treatment is Le Gall–Varadhan’s triangular approximation. Due to similarity, we only consider $d = 3$. We first prove that for any integer $m \geq 1$,

$$\mathbb{E}|Q_n - \mathbb{E}Q_n|^m = O((n \log n)^{m/2}), \quad n \rightarrow \infty. \tag{5.27}$$

We carry out induction on m . By Lemma 5.2, (5.27) holds as $m = 1, 2$. We let $m \geq 3$, assume that it is true for all $1 \leq j \leq m - 1$ and prove it is true for m .

Given n , write $n_1 = \lfloor n/2 \rfloor$, $n_2 = n - n_1$,

$$Q'_{n_2} = \sum_{n_1+1 \leq j < k \leq n} 1_{\{S_j=S_k\}} \quad \text{and} \quad \bar{J}_n = \sum_{j=1}^{n_1} \sum_{k=n_1+1}^n 1_{\{S_j=S_k\}}.$$

By Lemma 5.2 there is $C_0 > 0$ such that

$$\mathbb{E}|\bar{J}_n - \mathbb{E}\bar{J}_n|^m \leq C_0^m (m!)^{3/2} n^{m/2} m, \quad n = 1, 2, \dots \tag{5.28}$$

By independence between Q_{n_1} and Q'_{n_2} ,

$$\begin{aligned}
 & (\mathbb{E}|Q_n - \mathbb{E}Q_n|^m)^{1/m} \\
 & \leq \left\{ \mathbb{E}(|Q_{n_1} - \mathbb{E}Q_{n_1}| + |Q'_{n_2} - \mathbb{E}Q'_{n_2}|)^m \right\}^{1/m} + \left\{ \mathbb{E}|\bar{J}_n - \mathbb{E}\bar{J}_n|^m \right\}^{1/m} \\
 & \leq \left\{ \sum_{j=0}^m \binom{m}{j} \mathbb{E}|Q_{n_1} - \mathbb{E}Q_{n_1}|^j \mathbb{E}|Q'_{n_2} - \mathbb{E}Q'_{n_2}|^{m-j} \right\}^{1/m} \\
 & \quad + C_0 \sqrt{n} (m!)^{3/(2m)}.
 \end{aligned} \tag{5.29}$$

Combining the induction assumption and (5.29),

$$\begin{aligned}
 & (\mathbb{E}|Q_n - \mathbb{E}Q_n|^m)^{1/m} \\
 & \leq \left\{ O((n \log n)^{m/2}) + \mathbb{E}|Q_{n_1} - \mathbb{E}Q_{n_1}|^m + \mathbb{E}|Q'_{n_2} - \mathbb{E}Q'_{n_2}|^m \right\}^{1/m} + O(\sqrt{n}).
 \end{aligned}$$

Write

$$\alpha_k = \sup \left\{ (2^k \log 2^k)^{-1/2} (\mathbb{E}|Q_n - \mathbb{E}Q_n|^m)^{1/m}; 2^k \leq n \leq 2^{k+1} \right\}.$$

Then,

$$\alpha_{k+1} \leq \left\{ O(1) + 2^{-(m-2)/2} \alpha_k^m \right\}^{1/m} + o(1) \leq 2^{-(m-2)/(2m)} \alpha_k + O(1), \quad k \rightarrow \infty.$$

By the fact that $m \geq 3$ one can see that the sequence α_k is bounded. We have proved (5.27).

We now claim that there is $C > 0$ such that

$$\mathbb{E}|Q_n - \mathbb{E}Q_n|^m \leq C^m (m!)^{3/2} (n \log n)^{m/2}, \quad m, n = 1, 2, \dots \tag{5.30}$$

Indeed, take m_0 sufficiently large so that

$$(1 - 2^{-(m-2)/(2m)})^{-1} \frac{(m-1)^{1/m}}{\sqrt{2}} \leq \frac{1}{2}, \quad 1 - 2^{-(m-2)/(2m)} \geq \frac{1}{4}$$

for all $m \geq m_0$. By (5.27), there is a constant $C > 0$ such that for all $j = 1, \dots, m_0$,

$$\mathbb{E}|Q_n - \mathbb{E}Q_n|^j \leq C^j (j!)^{3/2} (n \log n)^{j/2}, \quad n = 1, 2, \dots$$

We may assume that $C \geq 8C_0$. (Recall that C_0 is given in (5.28).) By induction (on m), all we have to prove is that for any $m \geq m_0$, if

$$\mathbb{E}|Q_n - \mathbb{E}Q_n|^j \leq C^j (j!)^{3/2} (n \log n)^{j/2}, \quad n = 1, 2, \dots, \tag{5.31}$$

for every $j = 1, \dots, m-1$ then

$$\mathbb{E}|Q_n - \mathbb{E}Q_n|^m \leq C^m (m!)^{3/2} (n \log n)^{m/2}, \quad n = 1, 2, \dots \tag{5.32}$$

From (5.29) and (5.31) we have

$$\begin{aligned}
 (\mathbb{E}|Q_n - \mathbb{E}Q_n|^m)^{1/m} & \leq \left\{ 2^{-m/2} C^m (n \log n)^{m/2} \sum_{j=1}^{m-1} \binom{m}{j} (j!)^{3/2} ((m-j)!)^{3/2} \right. \\
 & \quad \left. + \mathbb{E}|Q_{n_1} - \mathbb{E}Q_{n_1}|^m + \mathbb{E}|Q'_{n_2} - \mathbb{E}Q'_{n_2}|^m \right\}^{1/m} + C_0 \sqrt{n} (m!)^{3/(2m)}.
 \end{aligned}$$

Notice that

$$\sum_{j=1}^{m-1} \binom{m}{j} (j!)^{3/2} ((m-j)!)^{3/2} = m! \sum_{j=1}^{m-1} (j!)^{1/2} ((m-j)!)^{1/2} \leq (m!)^{3/2} (m-1)$$

and that by (5.27)

$$\beta_m \equiv \sup_{n \geq 1} \{ (n \log n)^{-m/2} \mathbb{E} |Q_n - \mathbb{E} Q_n|^m \} < \infty.$$

We have

$$\begin{aligned} \beta_m^{1/m} &\leq \left\{ \frac{m-1}{2^{m/2}} (m!)^{3/2} C^m + 2^{-(m-2)/2} \beta_m \right\}^{1/m} + C_0 (m!)^{3/(2m)} \\ &\leq C \frac{(m-1)^{1/m}}{\sqrt{2}} (m!)^{3/(2m)} + 2^{-(m-2)/(2m)} \beta_m^{1/m} + C_0 (m!)^{3/(2m)}. \end{aligned}$$

Hence,

$$\begin{aligned} \beta_m^{1/m} &\leq (1 - 2^{-(m-2)/(2m)})^{-1} \left\{ \frac{(m-1)^{1/m}}{\sqrt{2}} C + C_0 \right\} (m!)^{3/(2m)} \\ &\leq \left(\frac{1}{2} C + 4C_0 \right) (m!)^{3/(2m)} \leq C (m!)^{3/(2m)}. \end{aligned}$$

Hence, (5.27) holds. By (5.27) and the Taylor expansion there is $\theta_0 > 0$ such that

$$\sup_n \mathbb{E} \exp \left\{ \frac{\theta_0}{\sqrt[3]{n \log n}} |Q_n - \mathbb{E} Q_n|^{2/3} \right\} < \infty. \quad (5.33)$$

It remains to extend (5.33) to any $\theta > 0$. Indeed, for any $\theta > \theta_0$, one can find an integer l such that for any n there is an integer partition $0 = n_0 < n_1, \dots, < n_l = n$ such that $n_i - n_{i-1} < n(\theta_0/\theta)^3$ ($i = 1, \dots, l$). Write

$$Q_n = \sum_{i=1}^l \sum_{n_{i-1} < j, k \leq n_i} 1_{\{S_j = S_k\}} + \sum_{i=2}^l \sum_{j=1}^{n_{i-1}} \sum_{k=n_{i-1}+1}^{n_i} 1_{\{S_j = S_k\}}. \quad (5.34)$$

Notice that for each $2 \leq i \leq l$,

$$\sum_{j=1}^{n_{i-1}} \sum_{k=n_{i-1}+1}^{n_i} 1_{\{S_j = S_k\}} \stackrel{d}{=} \sum_{j=1}^{n_{i-1}} \sum_{k=1}^{n_i - n_{i-1}} 1_{\{S_j = S'_k\}} \leq J_n.$$

By Lemma 5.2 and the Taylor expansion there is $\lambda > 0$ such that

$$\sup_{n \geq 1} \mathbb{E} \exp \left\{ \frac{\lambda}{\sqrt[3]{n}} \left(\sum_{j=1}^{n_{i-1}} \sum_{k=n_{i-1}+1}^{n_i} 1_{\{S_j = S_k\}} \right)^{2/3} \right\} < \infty.$$

By the triangular inequality and the Hölder inequality, therefore,

$$\sup_{n \geq 1} \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n \log n}} \left(\sum_{i=2}^l \sum_{j=1}^{n_{i-1}} \sum_{k=n_{i-1}+1}^{n_i} 1_{\{S_j = S_k\}} \right)^{2/3} \right\} < \infty.$$

Consequently, by the Jensen inequality

$$\sup_{n \geq 1} \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n \log n}} \left(\mathbb{E} \sum_{i=2}^l \sum_{j=1}^{n_{i-1}} \sum_{k=n_{i-1}+1}^{n_i} 1_{\{S_j=S_k\}} \right)^{2/3} \right\} < \infty.$$

By independence and by the triangular inequality

$$\begin{aligned} & \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n \log n}} \left(\sum_{i=1}^l \left| \sum_{n_{i-1} < j, k \leq n_i} 1_{\{S_j=S_k\}} - \mathbb{E} \sum_{n_{i-1} < j, k \leq n_i} 1_{\{S_j=S_k\}} \right| \right)^{2/3} \right\} \\ & \leq \prod_{i=1}^l \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n \log n}} |Q_{n_i-n_{i-1}} - \mathbb{E} Q_{n_i-n_{i-1}}|^{2/3} \right\} \\ & \leq \prod_{i=1}^l \mathbb{E} \exp \left\{ \frac{\theta_0}{\sqrt[3]{(n_i - n_{i-1}) \log(n_i - n_{i-1})}} |Q_{n_i-n_{i-1}} - \mathbb{E} Q_{n_i-n_{i-1}}|^{2/3} \right\}. \end{aligned}$$

Hence, the desired (5.25) follows from (5.33) and (5.34). □

By slightly modifying our argument, we can prove that Theorem 5.2 holds also for the range. We include this in our paper for future reference.

Theorem 5.3. *As $d = 3$,*

$$\sup_n \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n \log n}} |\#\{S[1, n]\} - \mathbb{E}\#\{S[1, n]\}|^{2/3} \right\} < \infty \quad \text{for every } \theta > 0. \tag{5.35}$$

As $d \geq 4$,

$$\sup_n \mathbb{E} \exp \left\{ \frac{\theta}{\sqrt[3]{n}} |\#\{S[1, n]\} - \mathbb{E}\#\{S[1, n]\}|^{2/3} \right\} < \infty \quad \text{for some } \theta > 0. \tag{5.36}$$

Proof. The argument used here is essentially the same as the one for Q_n . To carry it through, (5.21) needs to be replaced by Lemma 3.1 in [17], and (5.24) needs to be replaced by Lemma 6 in [7]. The rest of the proof follows an obvious modification of the argument for Theorem 5.2. □

Acknowledgments

The author thanks anonymous referee(s) for constructive comments and suggestions.

References

- [1] A. Asselah and F. Castell. Self-intersection local times for random walk, and random walk in random scenery in dimension $d \geq 5$. Preprint, 2005. Available at arXiv:math.PR/0509721.
- [2] A. Asselah. Large deviation estimates for self-intersection local times for simple random walk in \mathbb{Z}^3 . *Probab. Theory Related Fields*. To appear.
- [3] R. F. Bass, X. Chen and J. Rosen. Moderate deviations and laws of the iterated logarithm for the renormalized self-intersection local times of planar random walks. *Electron. J. Probab.* **11** (2006) 993–1030. MR2261059
- [4] E. Buffet and J. V. Pulé. A model of continuous polymers with random charges. *J. Math. Phys.* **38** (1997) 5143–5152. MR1471918
- [5] X. Chen. On the law of the iterated logarithm for local times of recurrent random walks. In *High Dimensional Probability II (Seattle, WA, 1999)* 249–259, 2000. MR1857326
- [6] X. Chen. Exponential asymptotics and law of the iterated logarithm for intersection local times of random walks. *Ann. Probab.* **32** (2004) 3248–3300. MR2094445

- [7] X. Chen. Moderate deviations and law of the iterated logarithm for intersections of the range of random walks. *Ann. Probab.* **33** (2005) 1014–1059. MR2135311
- [8] X. Chen and W. Li. Large and moderate deviations for intersection local times. *Probab. Theory Related Fields* **128** (2004) 213–254. MR2031226
- [9] A. Dembo and O. Zeitouni. *Large Deviations Techniques and Applications*, 2nd edition. Springer, New York, 1998. MR1619036
- [10] B. Derrida, R. B. Griffiths and R. G. Higgs. A model of directed walks with random self interactions. *Europhys. Lett.* **18** (1992) 361–366.
- [11] B. Derrida and P. G. Higgs. Low-temperature properties of directed walks with random self-interactions. *J. Phys. A* **27** (1994) 5485–5493. MR1295374
- [12] R. van der Hofstad and W. König. A survey of one-dimensional random polymers. *J. Statist. Phys.* **103** (2001) 915–944. MR1851362
- [13] N. C. Jain and W. E. Pruitt. The range of transient random walk. *J. Anal. Math.* **24** (1971) 369–393. MR0283890
- [14] N. C. Jain and W. E. Pruitt. Further limit theorem for the range of random walk. *J. Anal. Math.* **27** (1974) 94–117. MR0478361
- [15] N. C. Jain and W. E. Pruitt. Asymptotic behavior of the local time of a recurrent random walk. *Ann. Probab.* **11** (1984) 64–85. MR0723730
- [16] Y. Kantor and M. Kardar. Polymers with self-interactions. *Europhys. Lett.* **14** (1991) 421–426.
- [17] J.-F. Le Gall and J. Rosen. The range of stable random walks. *Ann. Probab.* **19** (1991) 650–705. MR1106281
- [18] S. Martínez and D. Petritis. Thermodynamics of a Brownian bridge polymer model in a random environment. *J. Phys. A* **29** (1996) 1267–1279. MR1385633
- [19] P. Révész. *Random Walks in Random and Non-Random Environments*. World Scientific, London, 1990. MR1082348
- [20] J. Rosen. Random walks and intersection local time. *Ann. Probab.* **18** (1990) 959–977. MR1062054
- [21] F. Spitzer. *Principles of Random Walk*. Van Nostrand, Princeton, New Jersey, 1964. MR0171290